

Original Research Article

Biosynthesis of Lovastatin, an anti-cholesterol drug by *Aspergillus wentii* NCIM 661 from Palm kernel cake via Solid-state fermentation

ABSTRACT

Over the past few years, the utilization of various agricultural residual wastes for the production of bioactive metabolites of industrial significance has been increased under solid-state fermentation in converting waste to wealth. In this context, present investigation presents the biosynthesis of an anti-cholesterol drug, lovastatin from palm kernel cake (PKC), a by-product obtained during the palm oil processing as a potential substrate, using *Aspergillus wentii* NCIM 661 under solid state fermentation (SSF). All the crucial process parameters such as initial moisture content, pH, incubation temperature, fermentation time and the effect of additional nutritional sources were optimized using single-parameter optimization to enhance the lovastatin production. A yield of 2.71 mg of lovastatin per gram dry substrate was obtained with palm kernel cake under the optimized fermentation parameters respectively. This study successfully and productively utilized the agro-waste and fungal strain for the biosynthesis of lovastatin at their best and demonstrated the feasibility of solid-state fermentation for the commercial production of metabolites of therapeutic significance. Findings from this study are very much promising for the economic utilization and value addition of these important agro residues, which are abundantly available in many developing countries like India.

Keywords: *Aspergillus wentii*; Mevinolin; Lovastatin; Optimization; Palm kernel cake; Solid-state fermentation.

1. INTRODUCTION

Lovastatin ($C_{24}H_{36}O_5$; also known Monacolin K or Mevinolin) belongs to the class of natural statins, which is most widely and effectively used to control hypercholesterolemia (accumulation of cholesterol in blood plasma). It competitively inhibits 3-hydroxy-3-methylglutaryl coenzyme-A (HMG-CoA) reductase, the rate-limiting enzyme of cholesterol biosynthesis [1-3]. Of all the statins available, lovastatin was the first natural statin approved by United States Food and Drug Administration (USFDA) in the year 1987 [4-5]. Moreover, lovastatin has also been reported to possess other therapeutic applications such as an anti-tumor agent against various forms of cancer, anti-inflammatory activity and also plays a vital role in the prevention of neurological disorders; bone disorders etc [6-8].

Lovastatin is mostly biosynthesized from various fungal genus and species as a secondary metabolite. Several fungal genera such as *Aspergillus*, *Monascus*, *Phoma*, *Penicillium* *Trichoderma*, *Pleurotus*, *Hypomyces* are reported as potential lovastatin producers [9, 10]. Earlier, commercial production of lovastatin by fungi was achieved by employing submerged fermentation (SmF) using *Aspergillus terreus* [11-14]. Over the past few years, solid-state fermentation (SSF) has emerged as an alternative to submerged fermentation, because of several advantages it offers such as easy control of process contamination, requires fewer processing and down-streaming stages, utilizes lesser power and generates lesser effluent [15,16]. Another important feature of SSF is its ability to use inexpensive substrates in the form of agro-waste residues for the production of valuable metabolites of industrial importance [17-19]. Very limited documented data is available on lovastatin production under solid-state fermentation using various microbial genera [20-32].

Palm kernel cake (PKC) is the by-product of oil palm industry; generated after the processing of oil from kernel. It is nutritionally rich containing (% w/w) dry matter 90; crude protein 16.1; ether extract 0.8; crude fibre 15.2; Ash 4; N-free extract 63; calcium 0.29; phosphorous 0.71 and metabolized energy 6.2 [33]. In the current work, the potentiality of palm kernel cake (PKC) was evaluated to see

whether the residual waste could be used as a promising substrate in SSF using the fungal strain *A. wentii* NCIM 661 for the biosynthesis of therapeutic drug-lovastatin. To the best of our knowledge, this is the first paper reporting palm kernel cake for lovastatin production using *A. wentii* NCIM 661.

2. MATERIALS AND METHODS

2.1 Chemicals

Used chemicals are of analytical grade and were purchased from Loba Chemie Pvt. Ltd, Mumbai, India. Culture media was purchased from Hi-media Laboratories, Mumbai, India.

2.2 Microorganism and Inoculum preparation

Fungal strain, *Aspergillus (A.) wentii* NCIM 661, received from National Collection of Industrial Microorganisms (NCIM), Pune was used in the present study. The culture was maintained on potato dextrose agar (PDA) slants at 28°C, stored at 4°C and sub-cultured monthly. To a well-sporulated slant of *A. wentii*, 10 ml of sterilized Tween-80 solution (0.1% v/v) was added. The spore surface was dislodged with an inoculation needle and agitated thoroughly using cyclomixer (REMI: CM-101 PLUS, Mumbai, India) to suspend the spores uniformly. This was used as inoculum throughout the study.

2.3 Substrate

Palm kernel cake (PKC) was collected from nearby palm oil processing mill in Guntur, Andhra Pradesh, India. Before use, the substrate was sun-dried to remove any extra moisture content and sieved to particle size of 0.5 mm. The substrate was used in SSF without any pre-treatment.

2.4 Solid-state fermentation

PKC (5g) was taken in to 250 ml Erlenmeyer flasks. The moisture content of the media was maintained at 60% using the moistening medium. The contents in the flasks were autoclaved at 121°C (15 lb) for 20 min, cooled to room temperature and inoculated with 1 ml of the fungal spore suspension. The contents were uniformly mixed thoroughly and incubated at 28°C in an incubator for desired period of time (i.e. one week).

2.5 Lovastatin extraction and assay

After completion of fermentation time, the flasks were dried at 40°C for 24 h and crushed into powder form. About 2g of the powdered material was taken and extracted with 100 ml of methanol: water (1:1, v/v) mixture (pH 7.7) in 250 ml Erlenmeyer flask and keeping the flasks at 30°C in rotary shaker at 180-200 rpm for 2 h. After 2 h, the mixture was centrifuged at 10,000 rpm for 10 min and the supernatant was filtered through 0.45 µm membrane filter. The obtained filtrate was collected in vials and preserved at 4°C for further analysis. Lovastatin in the clear extract was estimated by high performance liquid chromatography (HPLC) using a C₁₈ column (250 mm x 4.6 mm x 5 mm internal diameter). A mixture of 0.02 M phosphate buffer (pH 7.7) and acetonitrile in the ratio of 65:35 (v/v) was used as mobile phase. The mobile phase flow rate was maintained at 1.0 ml/min and lovastatin was detected at 238 nm with an injection volume of 20 µL [20]. The production of lovastatin is expressed in mg/g dry weight substrate (gds). The yield of lovastatin was calculated [34]. The obtained lovastatin yield was expressed as milligram per gram of the dry substrate (mg/gds).

2.6 Optimization of fermentation conditions

All the essential physicochemical and nutritional variables that influence the lovastatin yield were optimized using single-parameter optimization over a wider range. The parameters such as moisture content (40-80%), pH (4-11, adjusted with 1N HCl/NaOH), incubation temperature (22-40°C), fermentation time (24-168 h). In addition, the impact of various carbon sources (glucose, lactose, maltose, fructose, sucrose, soluble starch, xylose, and cellulose) and nitrogen sources (ammonium sulphate, ammonium nitrate, yeast extract, malt extract, urea and peptone) were also assessed. All

the SSF experiments along with analytical assays were run in three sets and the results reported were mean value of the three sets and standard deviation was $\pm < 5\%$.

3. RESULTS AND DISCUSSION

Since one of the primary motivations for SSF bioprocesses is its economical advantage in utilizing cost-effective agro-wastes for the production of valuable metabolites. So, based on the chemical and nutritional factors, cost and availability, palm kernel cake (PKC) was chosen as a promising substrate in this SSF to carry out optimization experiments for lovastatin enhancement.

3.1 Optimization of fermentation time

The maximum lovastatin yield of 1.19 mg/g of dry substrate was achieved after 72 h of fermentation time (Fig.1). The lovastatin yield increased up to 72 h which explained that lovastatin is a kind of fungal secondary metabolite and its accumulation in mycelia seems growth relatedness.




Fig. 1. Effect of fermentation time on lovastatin yield. The maximum yield obtained after 72 h of fermentation time.

After 96 h, there is a drastic decrease in lovastatin yield by the fungal strain. The reason for this decrement might be due to the onset of microbial death phase or the micro organism has attained a stage, from which it could not balance its steady growth with the available nutritional sources [35].

3.2 Optimization of moisture content

Optimal yield of lovastatin (1.62 mg/g dry substrate) was achieved at 60% moisture content (Fig. 2.). Moisture content is one of the critical factors that determine the success of SSF. Other than the optimized value, metabolite yield was lower on either side. With an increase in moisture level, there is a decrease in porosity, change in the substrate particle structure, lowers oxygen transfer, enhancement of bacterial growth and formation of aerial mycelia [36]. On the other hand, lower moisture content reduces the solubility of substrate nutrients, lowers degree of swelling and higher water tension [37]. The same 60% moisture content was also observed with *Aspergillus fischeri* under solid-state fermentation using coconut oil cake [25].




Fig. 2. Effect of moisture content on lovastatin yield. At 60%, the lovastatin yield was maximum

3.3 Optimization of initial pH

The profound effect of initial pH on the lovastatin production was as shown in Fig. 3. Maximum lovastatin yield (2.02 mg/g dry substrate) was recorded at pH 7.0.




Fig. 3. Effect of initial pH on lovastatin production. At pH 7.0, the yield of lovastatin was optimal

pH beyond 7.0 resulted in the reduction of lovastatin production gradually due to the inactivation of the fungal strain, because pH strongly influences the transport of various components across the cell

membrane and in turn support the cell growth and product formation. Basically agro-residual wastes utilized in SSF possess excellent buffering capacity. Most of the fungal species are active in the pH range of 3.5-7 and also lower pH avoids the contamination by other microbes [35,38].

3.4 Optimization of incubation temperature

Results indicated that maximum lovastatin production (2.47 mg/g dry substrate) was obtained when SSF was carried out at 30°C. However, lovastatin yield reduced after optimal incubation temperature of 30°C. Generally, most of the fungal strains grow well in a temperature range between 25-32°C and any variation in their growth temperature results in poor metabolites production of metabolites [39]. Moreover, higher temperatures could lead to poor growth of microorganisms during fermentation due to the thermal denaturation of microbial bio-actives for metabolic pathways [40]. These results are coinciding with those previously reported for lovastatin production by *Monascus ruber* [23].




Fig. 4. Effect of incubation temperature on lovastatin production. The lovastatin yield was high at 30°C

3.5 Effect of nutritional sources

The nutritional (both carbon and nitrogen) sources were supplemented to the SSF medium in the range of 0.25-2.0% (w/w). None of the nutritional sources, other than glucose as carbon source at 0.5% (w/w), had shown profound impact on the microbial growth and lovastatin yield (data not presented). The optimal yield of lovastatin reported was 2.71 mg/g dry substrate. The reason might be due to the reason that the utilized substrate, palm kernel cake is already a rich source of energy and protein which is self-sufficient in nourishing the fungal strain without any external nutrient requirement.

4. CONCLUSION

The finding from this study, clearly demonstrated the lovastatin production process based on palm kernel cake as a potential substrate in SSF is economically feasible and attractive as it is a cheap and readily available agro-residual byproduct in India. This result is of significant interest due to the productive utilization of low cost and abundant availability of residues for the production of value-added compounds of industrial importance. .

182 REFERENCES

- 183 1. Alberts AW, Chen J, Kuron G, Hunt V, Huff J, Hoffman C et al. Mevinolin: A highly potent
184 competitive inhibitor of hydroxy methylglutaryl-coenzyme A reductase and a cholesterol-lowering
185 agent. Proc Natl Acad Sci USA. 1980; 77(7): 3957-3961.
186
- 187 2. Alberts AW. Lovastatin and Simvastatin-inhibitors of HMG CoA reductase and cholesterol
188 biosynthesis. Cardiology 1990; 77: 14- 21.
189
- 190 3. Schimmel TG, Borneman WS, Conder MJ. Purification and characterization of a lovastatin
191 Esterase from *Clonostachys compactiuscula*. Appl Environ Microb.1997; 63(4): 1307-1311.
192
- 193 4. Manzoni M, Rollini M. Biosynthesis and biotechnological production of statins by filamentous
194 fungi and application of these cholesterol lowering drugs. Appl Microbiol Biotechnol 2002; 58(5):
195 555-564.
196
- 197 5. Radha KV, Lakshmanan D. A review: lovastatin production and applications. Asian J Pharma
198 Clinical Res. 2013; 6(3): 21-26.
199
- 200 6. Seenivasan A, Subhagar S, Aravindan R, Viruthagiri T. Microbial production and biomedical
201 applications of lovastatin. Ind J Pharm Sciences. 2009; 701-709.
202
- 203 7. Barrios GJ, Miranda RU. Biotechnological production and applications of statins. Appl Microbiol
204 Biotechnol. 2010; 85(4): 869-883.
205
- 206 8. Praveen VK, Savitha J. Solid State Fermentation: An Effective Method for Lovastatin Production
207 by Fungi-A Mini Review. The Open Tropical Medicine Journal. 2012; 5: 1-5
208
- 209 9. Chakravarti R, Sahai V. Compactin—a review. Appl Microbiol Biotechnol 2004; 64(5): 618-624.
210
- 211 10. Upendra RS, Pratima K, Amiri ZR, Shwetha L, Ausim M. Screening and molecular
212 characterization of natural fungal isolates producing lovastatin. J Microb Biochem Technol. 2013;
213 5(2): 25-30.
214
- 215 11. Kumar MS, Jana SK, Senthil V, Shashanka S, Kumar SV, Sadhukhan AK. Repeated fed batch
216 process for improving lovastatin production. Process Biochem 2000; 36(4):363-368.
217
- 218 12. Jia Z, Zhang X, Zhao Y, Cao X. Enhancement of lovastatin production by supplementing
219 polyketide antibiotics to the submerged culture of *Aspergillus terreus*. Appl Biochem Biotechnol
220 2010; 160(7): 2014-2025.
221
- 222 13. Subhan M, Faryal R, Macreadie I. Exploitation of *Aspergillus terreus* for the Production of Natural
223 Statins. J Fungi (Basel) 2016; 2(2): 1-13.
224
- 225 14. Azeem M, Saleem Y, Hussain Z, Javed MM. Optimization of Culture Conditions for the
226 Production of Lovastatin by *Aspergillus terreus* in Submerged Fermentation. Pharm Chem J
227 2018; 52: 284-289.
228
- 229 15. Bhargav S, Panda BP, Ali M, Javed S. Solid-state fermentation: an overview. Chem Biochem
230 Eng Q 2008; 22(1): 49-70.
231
- 232 16. Abdul Manan M, Webb C. Modern microbial solid state fermentation technology for future
233 biorefineries for the production of added-value products. Biofuel Research Journal 2017; 4(4):
234 730-740.
235
- 236 17. Lizardi-Jiménez MA, Hernández-Martínez R. Solid state fermentation (SSF): diversity of
237 applications to valorize waste and biomass. 3 Biotech. 2017 May 1;7(1):44.
238
- 239 18. Abu Yazid N, Barrena R, Komilis D, Sánchez A. Solid-state fermentation as a novel paradigm for
240 organic waste valorization: a review. Sustainability 2017; 9(2): 224.

- 241
242 19. Sadh PK, Duhan S, Duhan JS. Agro-industrial wastes and their utilization using solid state
243 fermentation: A review. *Bioresour Bioprocess* 2018; 5(1): 1-15.
244
- 245 20. Valera HR, Gomes J, Lakshmi S, Gurujara R, Suyanarayan S, Kumar D. Lovastatin production
246 by solid-state fermentation using *Aspergillus flavipes*. *Enzyme Microbiol Technol.* 2005, 37 (5):
247 521-526.
248
- 249 21. Xu BJ, Wang QJ, Jia XQ, Sung CK. Enhanced Lovastatin Production by Solid State
250 Fermentation of *Monascus ruber*. *Biotechnol Bioproc E.* 2005, 10 (1):78-84.
251
- 252 22. Wei P, Xu Z, Cen PJ. Lovastatin production by *Aspergillus terreus* in solid-state fermentation.
253 2007. *J Zhejiang Univ.* 8(9). 1521-1526.
254
- 255 23. Panda BP, Javed S, Ali M. Optimization of fermentation Parameters for higher lovastatin
256 production in Red mold rice through co-culture of *Monascus purpureus* and *Monascus ruber*.
257 *Food Bioprocess Technol.* 2008, 53:342-346.
258
- 259 24. Pansuriya RC, Singhal RS: Response surface methodology for optimization of production of
260 Lovastatin by solid-state fermentation. *Braz J Microbiol.* 2010, 41:164-172.
261
- 262 25. Madhu Latha P, Chanakya P, Srikanth M. Lovastatin production by *Aspergillus fischeri* under
263 solid-state fermentation from coconut oil cake. *Nepal J Biotechnol.* 2012; 2(1): 26-36.
264
- 265 26. Faseleh Jahromi M, Liang JB, Ho YW, Mohamad R, Goh YM, Shokryazdan P. Lovastatin
266 production by *Aspergillus terreus* using agro-biomass as substrate in solid state fermentation. *J*
267 *Biomed Biotech.* 2012: 196264.
268
- 269 27. Kavitha V, Janani B, Jayaraman A. Optimization of process parameters for lovastatin production
270 from red gram husk by solid-state fermentation. *Int J of Sci Tech.* 2014; 3(9): 1413-1418.
271
- 272 28. Kamath PV, Dwarakanath BS, Chaudhary A, Janakiraman S. Optimization of culture conditions
273 for maximal lovastatin production by *Aspergillus terreus* (KM017963) under solid state
274 fermentation. *HAYATI J Biosciences.* 2015; 22(4): 174-180.
275
- 276 29. Bhargavi S, Praveen V, Mariam S, Sreepriya M, Savitha J. Purification of Lovastatin from
277 *Aspergillus terreus* (KM017963) and Evaluation of its Anticancer and Antioxidant Properties.
278 Purification of Lovastatin from *Aspergillus terreus* (KM017963) and Evaluation of its Anticancer
279 and Antioxidant Properties. 2016; 17(8): 3797-3803.
280
- 281 30. Javed S, Meraj M, Mahmood S, Hameed A, Naz F, Hassan S, Irfan R. Biosynthesis of lovastatin
282 using agro-industrial wastes as carrier substrates. *Trop J Pharm Res* 2017; 16(2): 263-269.
283
- 284 31. Munir N, Asghar M, Murtaza MA, Akhter N, Rasool G, Shah SM, Tahir IM, Khan FS, Riaz M,
285 Sultana S, Rashid A. Enhanced production of Lovastatin by filamentous fungi through solid state
286 fermentation. *Pak J Pharm Sci.* 2018; 31(4): 1583-1589.
287
- 288 32. Balraj J, Jairaman K, Kalieswaran V, Jayaraman A. Bioprospecting lovastatin production from a
289 novel producer *Cunninghamella blakesleeana*. *3 Biotech* 2018; 8(8): 359.
290
- 291 33. Sabu A, Pandey A, Daud MJ, Szakacs G. Tamarind seed powder and palm kernel cake: two
292 novel agro residues for the production of tannase under solid state fermentation by *Aspergillus*
293 *niger* ATCC 16620. *Bioresource Technol.* 2005; 96(11): 1223-1228.
294
- 295 34. Muthumary J, Sashirekha S. Detection of taxol, an anticancer drug, from selected *coelomycetous*
296 fungi. *Indian J Sci Technol.* 2007; 1:1-10.
297
- 298 35. Nampoothiri KM, Baiju TV, Sandhya C, Sabu A, Szakacs G, Pandey A. Process optimization for
299 antifungal chitinase production by *Trichoderma harzianum*. *Process Biochem.* 2004; 39(11):
300 1583-1590.

- 301
302 36. Singhania RR, Patel AK, Soccol CR, Pandey A. Recent advances in solid-state fermentation.
303 Biochem Eng J. 2009; 44(1):13-18.
304
305 37. Rashid JI, Samat N, Yusoff WM. Optimization of temperature, moisture content and inoculum
306 size in solid state fermentation to enhance mannanase production by *Aspergillus terreus* SUK-1
307 using RSM. Pak J Biol Sci. 2011; 14(9): 533-539.
308
309 38. Raimbault M. General and microbiological aspects of solid substrate fermentation. Electron J
310 Biotechnol 1998. 1(3): 174-188.
311
312 39. Bharti AK, Kumar A, Kumar A, Dutt D. Exploitation of *Parthenium hysterophorous* biomass as
313 low-cost substrate for cellulase and xylanase production under solid-state fermentation using
314 *Talaromyces stipitatus* MTCC 12687. J Radiat Res App Sci. 2018; 11: 271-280.
315 40. Gautam A, Kumar A, Dutt D. Production of cellulase-free xylanase by *Aspergillus flavus* ARC-12
316 using pearl millet stover as the substrate under solid-state fermentation. J Adv Enzy Res. 2015;
317 1: 1-9.