

Diffuse Barium Associated Peritonitis as A Complication of Barium Enema Examination

Budhi Ida Bagus¹, Raharjo Suryo Wahyu², Metria Ida Bagus³

¹ Department of Surgery, Sebelas Maret University, Indonesia

² Surgery Department, Moewardi General Hospital, Indonesia

³ Medical Faculty, Sebelas Maret University, Indonesia

Abstract

Introduction : Barium peritonitis is a rare but life threatening complication of contrast examination of the gastrointestinal tract. The chemical peritonitis due to barium contamination is characteristically severe and difficult to treat. Such a complication has a high morbidity and mortality rate.

Case Report: A 64 years old, Indonesian male patient was referred immediately to our surgery department after extravasation was found during barium enema procedure. The patient underwent surgery after the diagnosis of acute abdomen was made. Exploratory laparotomy revealed perforation of the rectum with a diameter of 3 cm, located 12 cm above the anal verge and barium covering whole peritoneal cavity. There were no post operative complications found on this patients and the patients tolerate well with oral intake early on the post operative day. The post operative period was uneventful and the patient was discharged from hospital on the 7th post operative day. He was in good general health without any ileostomy related problems at 30 days after surgery and the closure of the ileostomy would be pending for the next 3 months.

Conclusion : Rectal perforation during barium enema examination with subsequent barium leakage into the peritoneal cavity, is a serious complication and an emergency condition. Barium-associated peritonitis associated with morbidity and with potential mortality. Early recognition and proper management are crucial for successful treatment.

Keywords: peritonitis, barium leak, rectal perforation, emergency

Introduction

Barium peritonitis is a rare but life threatening complication of contrast examination of the gastrointestinal tract. The incidence of peritonitis following barium enema is in the order of 2-8 per 10.000 investigations. Barium, is a silver-white compound that outlines the colon and rectal wall on X-ray and is used for the detection of filling defects and other abnormalities. While it was a relatively less invasive procedure, complications can occur during and after Barium Enema examination.¹

Generalised peritonitis in such circumstances was rare, since most perforations were limited to the retroperitoneum. The chemical peritonitis due to barium contamination is characteristically severe and difficult to treat. However, the most important and life-threatening complication of barium examination is rectal perforation, which is caused by air used for insufflation of the rectum and the colon during the procedure to enhance imaging. Such a complication has a high morbidity and mortality rate.²

Case Report

A 64 years old, Indonesian male patient was referred immediately to our surgery emergency department after leakage was found during barium enema procedure (**Figure 1**). The patient presented with complain of nausea, vomiting and abdominal pain. His vital signs included temperature of 37.4°C, blood pressure of 110/ 70 mmHg, pulse rate of 82 beats/min, respiratory rate was 24 breaths/min. On physical examination, the abdomen was distended and tenderness was noted to direct and muscular guarding in all quadrants. Routine hematology and biochemistry tests were normal except for raised of total leucocyte count (11.600/mm³). On biochemical examination showed decreased of the albumin

(2.0 g/dl). A chest X-ray did not reveal any pneumoperitoneum. The patient underwent surgery after the diagnosis of acute abdomen was made.

Exploratory laparotomy revealed perforation of the rectum with a diameter of 3 cm, located 12 cm above the anal verge and barium covering whole peritoneal cavity (Figure 2 and 3). After an effective abdominal irrigation with normal saline, ileostomy was performed. Oral intake was given during first 24 hours after removing the nasogastric tube on the first postoperative day. On the second postoperative day, the patient is in stable condition and is allowed to move to the ward and well tolerated oral intake. On 7th post operative day, the patients could discharged from hospital with no post operative complication and well torelated oral intake. On 30 days follow up as out patient, the patient was tolerate well with solid food and no stoma related complication and no surgical site infection.



Figure 1. Barium Enema examination results in the patient showed contrast in the rectum to caecum, no visible mass or contrast leakage



Figure 2. Barium leakage around the intestine and peritoneal cavity




Figure 3. Intraoperative photograph showing the site of rectal perforation

Discussion

Tadros and Watters suggested four mechanisms of injury : trauma from the enema, overinflation of the balloon, recent colonoscopic instrumentation especially associated with biopsy and the presence of rectal mucosal disease such as cancer, stricture, diverticulosis or inflammatory bowel disease.¹ Rarely the colon may burst due to excessive transmural pressure alone.² Different types of perforation have been described in the literature. One study classifies the perforation as either intramural (incomplete) or extramural (complete).³ Peterson et al. divided perforations into five categories: 1) perforations of the anal canal below the levator ani muscle, 2) incomplete perforations such as perforation of the rectal mucosa, 3) perforations into the retroperitoneum, 4) transmural perforations into the adjacent viscera and 5) perforations into the free intraperitoneal cavity. The clinical signs, radiological findings, treatment strategies and prognosis may vary in each category.⁴

Surgery is not always required. In intramural or small retroperitoneal perforations, good results are reported with conservative treatment consisting of bowel rest, combined with total parenteral nutrition, intravenous fluid treatment, and broad-spectrum antibiotics.^{5,6} Surgical debridement is only required in case of large amounts of leakage and intramural abscesses, or in patients not responding to conservative treatment.^{7,8}

In retroperitoneal perforations with considerable leakage of barium a subgroup with a much poorer prognosis the extravasation should be drained and a diverting colostomy is advised, however, treatment by rectosigmoid resection, primary anastomosis, and proximal colostomy also has been reported.^{9,10,11}

Our patient was suspected of having chronic dhiarrea before, in the history we get nausea, vomiting, and flatulence before entering the hospital. Overinflation of the balloon is not recorded during barium-enema by the doctor and trauma from the enema in the mucosa is not suspected because the perforation site was 12 cm above the anal verge. However, in our patients, a plain abdominal X-ray revealed an excessive amount of barium that dilates the rectum and sigmoid accompanied by

extravasation. Possible causes of excessive amount of intraluminal barium and rectal perforation are excessive pressure during the procedure and thinning of the intestinal wall due to inflammation of the previous intestine which makes the intestinal wall more fragile to become perforated.

The incidence of colorectal perforation during barium-enema radiography can be reduced by 1) performing proctoscopy before barium enema, 2) avoiding the use of anal bubbles in patients with known rectal lesions, using safe-balloon tip designs insert it after careful rectal examination 3) avoid barium studies in patients with active colitis, 4) delay examination for at least six days in case of biopsy in or polypectomy, 5) avoidance of pressure formation greater than that made by barium suspension column one meters, and 6) using lower barium concentrations whenever possible.^{11,12}

Peritonitis caused by the leakage of the barium could be worsen because when this barium leaked into the peritoneal cavity, barium was the good media for bacterial translocation, those made the peritonitis became severe.¹³ In this case, there was no delayed time of referral to our hospital, thus emergency operation could controlled the barium leakage into peritoneal cavity.

Conclusion

Rectal perforation during barium enema examination with subsequent barium leakage into the peritoneal cavity, is a serious complication and an emergency condition. Barium-associated peritonitis associated with morbidity and with potential mortality. Early recognition and proper management are crucial for successful treatment.

Disclaimer regarding consent and ethical approval:

As per university standard guideline participant consent and ethical approval has been collected and preserved by the authors.

Funding

No financial support was provided for this study

Conflict of interest

The authors have no conflicts of interest to report

REFERENCES

1. Tadros S, Watters JM: Retroperitoneal perforation of the rectum during barium enema examination. *Can J Surg* 1988, 31:49-50
2. Williams S, Harned R: Recognition and prevention of barium enema perforations. *Curr Probl Diagn Radiol* 1991, 20:121-151.
3. Vora P, Chapman A. Complications from radiographer-performed double contrast barium enemas. *Clin Radiol* 2004; 59: 364-8.
4. Peterson N, Rohrmann CA Jr, Lennard ES. Diagnosis and treatment of retroperitoneal perforation complicating the double-contrast barium-enema examination. *Radiology* 1982; 144: 249-52.
5. Fry RD, Shemesh EI, Kodner IJ, Fleshman JW, Timmcke AE. Perforation of the rectum and sigmoid colon during barium-enema examination. Management and prevention. *Dis Colon Rectum* 1989;32:759-64.
6. Saigusa S, Ohi M, Oki S, Ichikawa T, Kobayasi M, Inoue Y. Delayed awareness of the history of barium examination: perforated barium appendicitis. *Case Rep Gastrointest Med*. 2017; 2017: 631-635.
7. Pandit N, Singh H, Jaiswal LS. Barium peritonitis: a disastrous complication of unnecessary diagnostic study. *Trop Doct*. 2018 Apr;48(2): 171-173.
8. Matiasovic M, Nutt A, Parsons K, Doran I. Barium peritonitis with abdominal abscessation. *J Small Anim Pract*. 2018 Jun 19. doi: 10.1111/jsap.12851.
9. Rosenklint A, Buemann B, Hansen P, Baden H. Extraperitoneal perforation of the rectum during barium enema. *Scand J Gastroenterol* 1975;10:87 – 90.
10. Levy MD, Hanna EA. Extraperitoneal perirectal extravasation of barium during a barium enema examination: natural course and treatment. *Am Surg* 1980;46: 382-5.

11. Williams S, Harned R: Recognition and prevention of barium enema perforations. *Curr Probl Diagn Radiol* 1991, 20:121-151
12. Kojima H, Hojo S, Manabe T, Demura S, Sekine S, Shibuya K. The efficacy of steroids for postoperative persistent inflammatory reaction in a patient with bariumperitonitis: A case report. *Int J Surg Case Rep.* 2017;36:38-41.
13. Rehman ZU. Cocoon abdomen - A rare cause of intestinal obstruction. *Int J Surg Case Rep.* 2015;11:95-97.