

Evaluation of the antioxidant markers in *Clarias gariepinus* heart exposed to mercury and ammonium chloride

Abstract

Reactive Oxygen Species (ROS) parameters and antioxidant enzyme activities in heart of *Clarias gariepinus* weighing (15 ± 1.7 g) in toxic exposure to mercury concentrations, 0.1 mg/L group B, group C 10 mg/L ammonium chloride and group D 0.1 mg/L mercury and 10 mg/L ammonium chloride for 7 consecutive days at 26 ± 1 °C temperature. Group A was exposed to water alone and serve as control. At the end of the experimental period, Reduced glutathione (GSH), NO levels and the activities of Peroxidase, Superoxide dismutase (SOD) and Alanine Amino Transferase (ALT) in heart were assayed. The result showed a significant increase ($P < 0.05$) in the activities of ALT, Peroxidase, and SOD. However, a significant decrease in the levels of total protein content, GSH and NO were also observed. The results showed that the combined effect of mercury and ammonium had an adverse effect greater than that obtained individually.

Introduction

Fish constitutes an important aspect of human food due to the high level of quality protein and essential amino acids for the proper growth and functioning of body muscles and tissues. Fish are commonly situated at the top of the food chain and therefore, they can accumulate large amount of toxicants (Alaa *et al.*, 2010). Fish are considered as one of the most susceptible aquatic organisms to toxic substances present in water (Enkeleda *et al.*, 2013). Fish are also usually considered an organism of choice for assessing the effects of environmental pollution on the aquatic ecosystem. In nature, aquatic animals are constantly exposed to toxic substances (Ijeoma *et al.*, 2015). Rapid industrial development, as well as the use of metals in production processes has led to the increased discharges of toxic substances into the environment (Borham and Rahimeh, 2011).

Mercury occurs naturally as a mineral and is widely distributed throughout the environment as a result of natural and human activities. Inorganic mercury is the most common form of metal released by industries in the environment. Mercury (Hg) is a famous contaminant to aquatic life (fish and marine organisms), Which accumulates in many fish species and causes toxicity to various organs (Bano and Hasan 1990; Jalila *et al.*, 2012). On the other hand Ammonium Chloride (NH_4Cl) is an inorganic, mildly acidic, white crystalline salt compound that is highly soluble in water. The toxicity of ammonia (NH_3) and its compounds such as Ammonium chloride (NH_4Cl) have been ascribed to the fact that it can readily diffuse through the gill membranes due to its high solubility (Hegazi and Hasanein 2010). Free radicals and ROS are produced by a variety of oxidative enzymes. Metabolic processes and metal-catalyzed oxidation (Halliwell and Gutteridge, 1990). It is therefore of great significance to evaluate pollution effects on fish for both environmental protection and socio-economic reasons (Farombi *et al.*, 2007; Okomoda *et al.*, 2010). That is why this study investigated the toxicities of mercury and ammonium chloride on some antioxidant markers.

43

44 **Materials and methods**

45 **Chemicals**

46 Trishydroxyaminomethane, Hydrochloric acid, pyrogallol, hydrogen peroxide, sulphosalicylic
47 acid, DTNB, Bovine Serum Albumin.

48 **Fish husbandry**

49 *Clarias gariepinus* were obtained from the Niger Delta University fish farm. The fish were
50 transported in a plastic container. Fish of nearly equal size weighing (15 ± 1.7 g) were distributed
51 in 40 L plastic aquaria and acclimated to the laboratory conditions for one week. Four fish were
52 placed in every aquarium. Group A (Control) : Fish received palm fruits only as diet. Group B :
53 Fish received 10 mg/l of Ammonium chloride with palm fruits. Group C : Fish received 0.1 mg/l
54 of Mercury (II) nitrate with palm fruits. Group D : Fish received 10 mg/l of Ammonium chloride
55 and 0.1 mg/l Mercury II nitrate combined with palm fruits. The experiment lasted for 7 days. The
56 fish were then sacrificed by a sharp blow on the head. The fish were immediately dissected and
57 the heart was quickly removed and washed in ice-cold 1.15% KCl solution blotted and weighed.
58 They were then homogenized in 9 volumes of homogenizing buffer (50 mM Tris - HCl mixed
59 with 1.15% KCl and pH adjusted to 7.4), using Teflon Homogenizer. The resulting homogenate
60 was centrifuged at 10,000g for 20 min in a Beckman centrifuge at -40°C . The supernatant was
61 decanted and stored at -20°C until biochemical analysis.

62 **Protein content**

63 Protein precipitated from 0.1 ml of supernatant with an equal volume of 10% ice-cold trichloroacetic acid
64 was solubilized in a known volume of 0.1 N NaOH solution. The quantity of protein in the sample was
65 determined following the method of Lowry et al. (1951) using a standard curve prepared with bovine
66 serum albumin.

67 **Peroxidase activity**

68 The peroxidase activity was measured following the method of Wadhwa et al (1988). In the
69 experimental set, the reaction mixture included 2.0 ml of 0.1 M potassium phosphate buffer (pH
70 7.0), 1.0 ml pyrogallol (0.005 M), 1.0 ml of a 0.05 M hydrogen peroxide " H_2O_2 " solution and 1.0
71 ml heart supernatant (crude enzyme preparation) in a total volume of 5 ml. The reaction was
72 started by adding 1 ml of supernatant to each of experimental. After 5 min at 25°C , the reaction
73 was stopped by adding 0.5 ml H_2SO_4 (5% v/v). The optical density was measured at 430 nm in a
74 JASCO 7800 UV/visible spectrophotometer (Japan). After suitable corrections with the control,
75 the enzyme activity was expressed as optical density (OD) of purpurogallin formed/mg protein/5
76 min.

77

78 **Superoxide dismutase (SOD) activity**

79 The SOD activity was measured by the method of Marklund and Marklund (1974). The reaction
80 mixture consisted of 2.875 ml Tris-HCl buffer (50 mM, pH 8.5), pyrogallol (24 mM in 10 mM
81 HCl) and 100 μL of heart homogenate in a total volume of 3 ml. The enzyme activity was
82 measured at 420 nm and was expressed as units/mg protein. One unit of enzyme is defined as the
83 enzyme activity that inhibits auto-oxidation of pyrogallol by 50%.

84 **AST Activity**

85 Briefly, 0.1ml of heart was mixed with phosphate buffer (100mmol/L, pH 7.4), L-aspartate
86 (100mmol/L), and α -oxoglutarate (2mmol/L) and the mixture incubated for exactly 30min at
87 37°C . Then 0.5ml of 2, 4-dinitrphenylhydrazine (2mmol/L) was added to the reaction mixture
88 and allowed to stand for exactly 20min at 25°C . Then 0.5 ml of NaOH (0.4mol/L) was added
89 and the absorbance at 546nm read against the reagent blank (Reitman and Frankel 1957).

90
91

92 **Glutathione content**

93 The GSH content in liver homogenate was determined by the method of Jollow et al. (1974) in
94 which 1.0 ml of PMS fraction (10%) was mixed with 1.0 ml of sulphosalicylic acid (4%). The
95 samples were incubated at 4 °C for at least 1 h and then subjected to centrifugation at 1200g for
96 15 min at 4 °C. The assay mixture contained 0.4 ml filtered aliquot, 2.2 ml phosphate buffer (0.1
97 M, pH 7.4) and 0.4 ml DTNB (10 mM) in a total volume of 3.0 ml. The yellow colour developed
98 was read immediately at 412 nm on spectrophotometer. The GSH content was calculated as nmol
99 of DTNB conjugate formed/g tissue using molar extinction coefficient of $13.6 \times 10^3 \text{ M}^{-1} \text{ cm}$

100 **Nitrite estimation in plasma**

101 Nitrite assay was done using Griess reagent by the method of Green *et al* (1982) with some
102 modifications. In brief, 100 µL of Griess reagent (1:1 solution of 1% sulfanilamide in 5%
103 phosphoric acid and 0.1% naphthylethylene diamine dihydrochloride in water) was added to 100
104 µL of heart sample incubate for 10 min at room temperature protected from light.
105 Purple/magenta colour began to form immediately. Absorbance was measured at 546 nm, nitrite
106 concentration was calculated using a standard curve for sodium nitrite, and nitrite levels were
107 expressed as nmol/mg protein.

108 **Histopathology**

109 The heart tissues were excised, flushed with saline, cut open longitudinally along the main axis,
110 and then again washed with saline. These heart sections were fixed in 10% buffered formalin for
111 at least 24 h and after fixation, the specimens were dehydrated in ascending grades of ethanol,
112 cleared in benzene, and embedded in paraffin wax. Blocks were made and 5µm thick sections
113 were cut from the heart. The paraffin embedded liver tissue sections were deparaffinized using
114 xylene and ethanol. The slides were washed with phosphate buffered saline (PBS) and
115 permeabilized with permeabilization solution (0.1M citrate, 0.1% TritonX-100). These sections
116 stained with haematoxylin and eosin and were observed under a light microscope at 40x
117 magnifications to investigate the histoarchitecture of heart.

118 **Statistical analyses**

119 Results are presented as means±standard deviation. The statistical evaluation of all data was
120 done using one-way analysis of variance (ANOVA) followed by Dunnett's test. P values < 0.05
121 were regarded as statistically significant.

122 **Results**


123 **Table 1**

124 Glutathione and nitric oxide levels, Peroxidase, superoxide dismutase and ALT activities in heart
125 of *Clarias gariepinus* exposed to ammonium chloride and mercury nitrate.

Groups	Protein (mg/ml)	ALT (U/L)	GSH (nmol DTNB/mg protein)	Peroxidase O.D purpurogallin/mg protein	NO (nmol/mg protein)	SOD (U/mg protein)
A	1.03±0.08 ^a	66.92±7.84 ^a	4.08±0.107 ^a	6.13±0.16 ^a	22.19±3.19 ^a	13.45±0.10 ^a
B	0.76±0.16 ^b	89.46±8.61 ^b	1.18±0.28 ^b	6.41±0.70 ^a	44.48±14.71 ^b	7.88±0.19 ^b
C	0.66±0.00 ^b	92.2±5.83 ^b	1.17±0.03 ^b	6.57±0.096 ^a	44.98±0.62 ^b	9.56±0.16 ^b
D	0.49±0.01 ^b	81.20±5.33 ^b	0.63±0.06 ^b	8.52±0.08 ^b	58.16±1.78 ^a	5.65±0.03 ^b

128

129 Values are expressed as mean \pm standard deviation of triplet determination. Mean with same
130 superscript letters on a column are not significantly different ($P < 0.05$). The result showed a
131 significant change ($P < 0.05$) in the activities of ALT, Peroxidase, and SOD. However, a
132 significant decrease in the levels of total protein content, GSH and NO were also observed
133



134
135
136 **Figure 1:** Photomicrograph of Heart: Group A (Control) shows normal cardiac muscle fibres
137 (MF). Similarly, Groups B and C also displayed normal cardiac tissue architecture. On the
138 contrary group D show focal area of necrosis (NMF). H&E x100

139
140 **Discussion**

141 Due to the industrial development, many dangerous chemicals have been released directly or
142 indirectly which resulted in the polluted water bodies. The African cat fish (*Clarias gariepinus*)
143 shows high sensitivity towards dissolving toxicants, hence this fish has been utilized as a
144 biomarker to indicate the existence of toxicant exposure. The present study revealed a significant
145 variation in the activities of Alanine aminotransferase (ALT), reduced glutathione (GSH), Nitric
146 oxide (NO), peroxidase and Superoxide dismutase (SOD). The changes in these biochemical
147 parameters indicate that they can be used as indicators of related stress in fish on exposure to
148 toxic chemicals such as ammonium chloride and mercury (II) nitrate.

149
150 Peroxidases are haem-containing enzymes that use hydrogen peroxide as the electron acceptor to
151 catalyze a number of oxidative reactions in this experiment the activity of peroxidase decreases

152 although catalase is another enzyme that can carry out the same function the results are in line
153 with the work reported by Wadhwa et al (1988).

154
155 GSH is a tripeptide, a cellular antioxidant. It protects the heart cells from oxidative stress and
156 plays an important role in the scavenging reactions by acting both as a nucleophilic of various
157 undesired compounds and their toxic metabolites and as a specific substrate for the enzymes GPx
158 and GST. The present study reveals that there was a significantly different ($P < 0.05$) from group
159 A the levels of GSH were high as compared to the other groups but the decrease in the levels of
160 GSH is more pronounced in group D may be the combined effects of mercury and ammonium.
161 These results are in line with the research results of Hegazi et al (2010).

162
163 Elevated activities of heart ALT are a common sign of heart disease. The levels of heart enzymes
164 are used as diagnostic indicators of heart injury. ALT is one of the most sensitive tests employed
165 in the diagnosis of heart diseases. The activity of ALT increased significantly from group A –D
166 this shows that there is an injury and since the activity of ALT is more in group D the injury is
167 much in D.

168
169 Superoxide anion formed from aerobic respiration is scavenged by Super Oxide Dismutase
170 (SOD) leading to the formation of hydrogen peroxide (H_2O_2) and in the presence of reduced
171 transition metals (such as iron) reactive hydroxyl radical ($OH\bullet$) is produced through the Fenton
172 reaction. The decreased activity of SOD in groups B-D indicates that there are more ROS in the
173 form of super oxide that has not been scavenged by this enzyme due to the toxicity of mercury or
174 ammonium or both. These findings are also in tune with the recent works of Hegazi et al (2010).

175
176 Nitric Oxide (NO) is another important mediator in the pathogenesis of inflammatory diseases.
177 The levels of this molecule are higher in groups B-D these results show that many inflammatory
178 enzymes are produced but that less of these molecules are produced in group A. This shows that
179 the toxicity of mercury, ammonium or the combination of both is lined with inflammation. These
180 are inconsonant with the work reported by Eboh et al (2016) even though nitric oxide was
181 assayed on the skin of rats.

182
183

184 **Conclusion**

185 In conclusion, at the end of the experimental period, Reduced glutathione (GSH), NO levels and
186 the activities of Peroxidase, Superoxide dismutase (SOD) and Alanine Amino Transferase (ALT)
187 in heart were assayed. The result showed a significant increase ($P < 0.05$) in the activities of
188 ALT, Peroxidase, and SOD. However, a significant decrease in the levels of total protein
189 content, GSH and NO were also observed. The results showed that the combined effect of
190 mercury and ammonium had an adverse effect greater than that obtained individually.

191

192 **References**

193 Eboh A.S, Frank-Oputu A, Tonkiri A, Wodu E, Ere D, Arhoghro E.M. (2016) Inhibitory Effect of
194 a Biflavonoid Antioxidant (Kolaviron) on Benzoyl Peroxide Induced Free Radical Generation in Rat
195 Skin. *The Pharmaceutical and Chemical Journal*, 3(4), 144-149.
196
197 Hegazi, M.M. Attia, Z.I. Ashour O.A. (2010). Oxidative stress and antioxidant enzymes in liver
198 and white muscle of Nile tilapia juveniles in chronic ammonia exposure *Aquatic Toxicology* 99,
199 118–125
200
201 Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. (1982).
202 Analysis of nitrate, nitrite, and [¹⁵N] nitrate in biological fluids. *Anal. Biochem.* 126, 131–138.
203
204 Jollow, D.J. Mitchell, J.R. Zampaglione, N. Gillette, J.R. (1974). Bromobenzene induced liver
205 necrosis: protective role of glutathione and evidence for 3,4- bromobenzene oxide as the
206 hepatotoxic metabolite, *Pharmacology* 11, 151–169.
207
208 Marklund, S. Marklund, G. (1974) Involvement of the superoxide anion radical in the
209 autoxidation of pyrogallol and a convenient assay for superoxide dismutase, *Eur. J. Biochem.* 47,
210 469–474.
211 Wadhwa, R., Kaur, M., & Sharma, S. P. (1988). An antioxidant induced alterations in peroxidase
212 activity in ageing *Zaprionus paravittiger*. *Mech Ageing Dev* 45, 277–283.
213
214 Halliwell, B., Gutteridge, J.M.C. (1990). Role of free radicals and catalytic metal ions in human
215 disease: an overview. *Methods Enzymol.* 186, 1–85.
216
217 Hegazi, M.M., Hasanein, S.S. (2010). Effects of chronic exposure to ammonia concentrations on
218 brain monoamines and ATPases of Nile tilapia (*Oreochromis niloticus*). *Comp. Biochem.*
219 *Physiol. C* 151, 420–425.
220
221 Bano Y, Hasan M. (1990). Histopathological lesions in the body organs of catfish,
222 *Heteropneustes fossilis* following mercury intoxication. *J Environ Sci Health* 25:67–85.
223
224 Okomoda, J., Ayuba, V. O. & Omeji, S. (2010). Hematological changes of *Clarias gariepinus*
225 fingerlings exposed to acute toxicity of formalin. *Production, Agriculture and Technology* 6(1),
226 92-101
227
228 Alaa, G. M. Osman, E. & Kloas, W. (2010). Water Quality and Heavy Metal Monitoring in
229 Water, Sediments, and Tissues of the African Catfish *Clarias gariepinus*, from the River Nile,
230 Egypt. *Journal of Environmental Protection*, 1(2), 389-400.
231
232 Enkeleda, L. D, Jetmira, A., Muhamet, Z., Imer, H., Dariana, B. & Fatgzim, L. C. (2013). Heavy
233 metals in fish for public consumption and consumer protection, *Natural Montenegrina*
234 *Podgorica*, 9(3), 843-851.
235
236 Farombi, E. O., Adelowo, O. A. & Ajimoko, Y. R. (2007). Biomarkers of oxidative stress and
237 heavy metal levels as indicators of environmental pollution in African Cat fish (*Clarias*)

238 *gariiepinus*) from Nigerian Ogunriver. *International Journal of Environmental Research and*
239 *Public Health*, 4(2), 158-165.

240

241 Ijeoma, K. H., Onyoche, N. E., Uju, O. V. & Chukwuene, I. F. (2015). Determination of Heavy
242 Metals in Fish (*Clarias gariiepinus*) Organs from Asaba Major Markets, Delta State, Nigeria,
243 *American Chemical Science Journal* 5(2), 135-147.

244

245 Borham, M. & Rahimeh, B. (2011). Influence of water hardness and pH on Acute Toxicity of Hg
246 on fresh water fish, *Capoeta fusca*. *World Journal of Fish and Marine Science*, 3(2), 132-136.

247

248 Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951). Protein measurement with
249 Folin phenol reagent. *J Biol Chem* 193, 265–275.

250

251

252 Reitman, S. and Frankel, S. (1957). Colorimetric method for the determination of serum glutamic
253 oxaloacetic and glutamic pyruvic transaminase. *American Journal Clinical Pathology*.28: 56 –
254 61.

255

UNDER PEER REVIEW