

# **Management of Paediatric Pain: How knowledgeable are Healthcare Providers at a Tertiary Centre, Southern Nigeria**

## **ABSTRACT**

**Background:** Many disease processes and interventional procedures in paediatrics are associated with pain. In spite of its frequency, it is undertreated globally, exposing children to the risks of long-term physical and psychological sequelae, which can have adverse effects on future treatments.

**Objectives:** To ascertain the knowledge and attitude of healthcare providers towards management of paediatric pain at the University of Port Harcourt Teaching Hospital.

**Methods:** In this cross-sectional survey, a semi-structured and self-administered questionnaire was distributed amongst a convenient sample of physicians and nurses attached to clinical departments/units where children are cared for, and those in educational units, between June and August 2017. Their knowledge and attitude toward pain management were analysed using Epi Info v7 software (CDC, USA).

**Results:** One hundred and ninety seven subjects participated in the study, 95(48.2%) physicians and 102(51.8%) nurses, all of various cadres. Seventy-four percent of them had more than 5 years working experience. One third of participants(30.4%) could name up to 3 non-pharmacological methods of pain management, of which the commonest were cold compress (48(24.4%) with nurses>physicians), followed by hot compress(38(19%)) and distraction(24(12%)). The commonest procedures for which non-pharmacological interventions of pain management were usually applied included immunisation of infants (20%) and venepuncture (17%) while lumbar puncture was the least (1.5%). Half of respondents(52%) had never prescribed/administered morphine to children.

**Conclusion:** Healthcare providers had poor knowledge of paediatric pain management. There is an urgent need to build their capacity to enable optimal relief of pain among paediatric patients.

**Keywords:** *Management, Paediatric pain, Knowledge, Healthcare providers, Southern Nigeria*

## **1. INTRODUCTION**

The International Association for the Study of Pain (IASP) defines pain as an unpleasant sensory and emotional experience associated with actual or potential tissue damage.[1] It is sometimes described as “what the patient says hurts”. [2] Uncontrolled pain is multi-faceted and impacts on the physical, psychological, social and spiritual well-being of the individual.[2,3,4] Therefore, if not addressed it could negatively affect the patient’s quality of life. But in the paediatric patient, many injuries, disease processes and most interventional procedures are associated with pain, which becomes of great concern.[5,6] **There are**

different types of pain, including acute and chronic pain. While the acute pain last a short time and has a specific cause, the chronic one persists beyond the initial insult that triggered it and the expected time for healing, which is assumed to be 3 months or longer, with no ongoing tissue damage identified.[7,21] The overall prevalence rates for different childhood pains range from 4-88%.[9] However, in spite of its frequency, pain is inadequately assessed and managed, exposing children to the risks of long-term physical and psychological sequelae, including among others, anticipatory anxiety during future procedures, a lowering of the pain threshold and sensitization to future pain.[2,5]

In sub-Saharan Africa, children are faced with a higher disease burden and poorer access to quality healthcare.[10] Regardless of disease processes, sufferings and pains they experience, management of the pain is sub-optimal and hindered by several factors, including certain cultures and assumptions claiming among others, that pain perception in children is unreal and transient.[10-12] On the other hand, availability and accessibility to analgesic medicines are also important factors why paediatric patients who often present to the hospital with pain are mostly under treated.[10,12] Furthermore, pain management takes low priority in most low- and middle-income countries due to limited and poor allocation of resources.[7]

Various protocols for pain management that employ both pharmacologic and non-pharmacologic strategies have been developed, and are being used as standard practice, especially in developed countries. Whereas, they are less understood or implemented among African health care workers.[10,13] Among others, little or no pre-service training and self-development; attitude, culture and believes of health practitioners have been identified as major factors why medical staff often lack sufficient knowledge for management of paediatric pain.[7,10]

Healthcare providers who care for children are mainly responsible for relieving pain and suffering when required. However, clinicians' approach to pain management in children has not been studied in our institution. Thus, this survey was conducted to ascertain the knowledge and attitude of healthcare providers towards management of paediatric pain in our institution, a tertiary centre in southern Nigeria.

## 2. MATERIAL AND METHODS

The study was conducted at the University of Port Harcourt Teaching Hospital (UPTH), an 800-bed tertiary care facility located in the southern part of Nigeria, and a major referral centre for patients from within Rivers State, with its under-15 population of 2,437,138 and neighbouring states.[14]

In this cross-sectional survey, a semi-structured and self-administered questionnaire, was used as instrument for data collection. The questionnaires were administered to a convenient sample of physicians and nurses attached to clinical departments/units where children are cared for, and those in educational units, through their unit heads during various departmental activities and retrieved that same or following day.

Departments and units involved included: Departments of Paediatrics (85 beds, with an average of 2,000 hospital admissions yearly), Surgery (Paediatric Surgery- 18 beds with an average of 10 surgeries weekly; Burns and plastics; Orthopedics), Ear-Nose & Throat, Ophthalmology, Paediatric Dentistry and Nursing services (Ward Nurses, Nurse Tutors and Nurses in the Post Basic Nursing Education Unit). The services provided by these health care professionals include treatments of common and specialised conditions in children as well as supportive care.

75 Approval for the study was obtained from the Ethics Committee of the hospital and from the  
76 respondents.

77 Descriptive statistics was used to present demographic data. Chi-square was used to  
78 compare the responses of the doctors and nurses to each question at a 95% confidence  
79 interval and a *P*-value less than .05 was considered significant. All tests were done with the  
80 Epi Info v7 software (CDC, USA).

81

### 82 3. RESULTS

83

84 Out of 220 questionnaires distributed, 197(90% response rate) were returned from  
85 95(48.5%) physicians and 102(51.8%) nurses. One third of the respondents, 66(33.5%) had  
86 5 to 10 years working experience post graduation while 49(25%) had more than 15 years.  
87 (Table 1)

88 Table 1. Characteristics of the study population

	Frequency	Percentage
<b>Role</b>		
Doctor	95	48.2
Nurse	102	51.8
<b>Total</b>	<b>197</b>	<b>100</b>
<b>Years of work experience</b>		
5 years	52	26.4
5 - 10 years	66	33.5
10 - 15 years	30	15.2
>15 years	49	24.9
<b>Total</b>	<b>197</b>	<b>100.0</b>

89

90 The commonest procedures for which non-pharmacological interventions of pain  
91 management were usually employed included immunisation of infants (20%) and  
92 venepuncture (17%). Nurses significantly applied them while changing dressings (*P*= .001)  
93 compared to doctors (Table 2). Whereas lumbar puncture was the least procedure for which  
94 non-pharmacological methods were considered, and by nurses only. On the other hand,  
95 pharmacological methods were mainly used for bone marrow aspiration (40%), even though  
96 significantly higher proportions of nurses had applied them for immunisation of infants (*P*=  
97 .03) and venepuncture (*P*= .007). Four percent of the respondents would make use of  
98 pharmacological measures of pain control while changing dressings.

99

100 Table 2. Medical procedures and indicated methods of pain management

	Doctors (%)	Nurses (%)	Total	<i>P</i> -value
<b>Procedures for which non-pharmacological methods of pain management are indicated</b>				
Immunization of infants	14 (14.7)	26 (25.4)	<b>40 (20.3)</b>	.06
Venepuncture	19 (20)	15 (14.7)	<b>34 (17.3)</b>	.32
Dressing changes	7 (7.4)	24 (23.5)	<b>31 (15.7)</b>	.001*
Insertion of venous catheter	3 (3.2)	9 (8.8)	<b>12 (6.1)</b>	.09
Lumbar puncture	0 (0.0)	3 (2.9)	<b>3 (1.5)</b>	.09
<b>Procedures for which pharmacological methods of pain management are indicated</b>				
Bone marrow aspiration	31 (32.6)	48 (47.1)	<b>79 (40.1)</b>	.03*
Lumbar puncture	7 (7.4)	9 (8.8)	<b>16 (8.1)</b>	.70

Immunization of infants	3 (3.2)	12 (11.7)	<b>15 (7.6)</b>	.02*
Insertion of venous catheter	6 (6.3)	6 (5.8)	<b>12 (6.1)</b>	.89
Venepuncture	1 (1.1)	10 (9.8)	<b>11 (5.6)</b>	.007*
Dressing changes	2 (2.1)	6 (5.8)	<b>8 (4.1)</b>	.17

One third of participants ( $n=60$ , 30.4%) could name up to 3 non-pharmacological methods of pain management, of which the commonest they had used in the 3 months preceding the survey were cold compress ( $n=48$ , 24.4%), hot compress ( $n=38$ , 19%) and distraction ( $n=24$ , 12%). A significantly higher proportion of nurses identified cold compress as a measure for managing pain compared to physicians ( $P < .001$ ) (Figure 1).






Fig. 1. Non-pharmacological methods of pain management used during the 3 months preceding the survey

Ibuprofen was the commonest drug known to be useful for treating chronic pain in children, significantly higher with nurses ( $P < .001$ ). On the other hand, gabapentin and amitriptyline were not known as useful medication for childhood chronic pain (Figure 2).



117

118 Figure 2. Useful drugs for the treatment of chronic pain in children  
 119 The large majority of respondents ( $n=103$ , 52%) had never prescribed/administered  
 120 morphine to children, whereas about 10% had done that sometimes (Figure 3).  
 121



122

123 Fig. 3. Attitude of respondents towards prescription/ administration of morphine to children  
 124  
 125 Majority of respondents are unaware whether a standard protocol for management of pain in  
 126 children exist in the hospital. For a third of them, there is none (Figure 4).  
 127  
 128




Fig. 4. Existence of a standard protocol for the management of paediatric pain in the hospital

#### 4. DISCUSSION

In the course of receiving health care, children are exposed to multiple invasive medical procedures as part of their preventive, curative or palliative treatment. The pain associated with these procedures has been reported as one of the most significant and distressing cause of pain for hospitalised children.[15] In this study, majority of respondents failed to associate painful procedures with interventions for pain control, implying that children are frequently subjected to unnecessary pain. Similarly Offiong found that less than 10% of Nigerian paediatric health care workers prescribed analgesics for procedural pain. [16] This is worrisome as these children are exposed to the risks of adverse sequelae of untreated procedural pain in childhood that can have a lifelong impact, including needle phobia among others, in which the fear may become generalized to individuals, objects, and situations related to needles, such as doctors, nurses, syringes and examination rooms.[17-19]

Non-pharmacological measures have been found highly effective, have excellent safety profiles and are easy to learn. They are recommended for use whenever possible in conjunction with pharmacological options to help lower levels of anxiety, pain and distress.[12,13,20] In this study, few respondents resolved to non-pharmacological methods of pain control while caring for their patients, especially nurses, even for procedures performed by physicians. This may be due to fact that nurses spend more time with patients and by virtue of their training in providing comfort. Our finding differs from the report of De Freitas *et al* which showed that all healthcare professionals who participated in their survey and were working in some pediatric units in southern Brazil adopted non-pharmacologic measures of pain management.[21] Respondents who had some kind of training in pain treatment, during or after their education were found to be better informed about pain management in children.[21] The disparity with our finding could be due to the fact that pre-service training, especially for doctors, lays greater emphasis on pharmacology than other means of pain alleviation while in-service training opportunities on the subject are not readily available in our setting.[7] Whereas use of hot water bottle was the commonest non-



pharmacological measure stated in the afore mentioned survey, use of cold compress was the most common option applied by respondents in this study, while prayers and fluid therapy were the only strategies administered to children with sickle cell anaemia and acute pain in a tertiary hospital in south-western Nigeria.[21,22] Surprisingly, prayers were not mentioned as an option in this study, despite the fact that we live in a very religious setting. It is possible that respondents were not aware that this simple strategy could be useful in pain management.

Chronic pain is associated with poor quality of life, serves no biological purpose, and is more difficult to treat.[4,7,12] A multidisciplinary approach has been recommended for its care, including Non-steroidal anti-inflammatory drugs (NSAIDs) for mild pain while morphine is still advocated for moderate to severe pain, to which adjuvant medications can be added for optimal pain control. [2,3,6,7,13] In this study, healthcare professionals mainly preferred ibuprofen, for treating chronic pain in children while less than 10% believed that morphine could be useful. This result is in consonance with previous reports in Nigeria.[7,22] While exploring how clinicians treated chronic pain in 3 tertiary centres in middle and northern Nigeria, Sanya *et al* found that ibuprofen was the most used NSAID(73%), especially for patients with sickle cell disease which was the most common cause of chronic pain, which could be explained by the fact that Nigeria has the highest burden of sickle cell disease in Africa.[7,23] Whereas less than 10% of clinicians were familiar with the use of opioids, only 2% also prescribe antiepileptics as adjuvant analgesics to treat chronic pain.

Adjuvant analgesics are used in combination with the primary analgesics to improve outcome and maintain the balance between relief and side effect, and have been found useful for treating chronic pain even in children.[2-4,6,13,20] Respondents in this study were not familiar with some of the commonly used ones, including anticonvulsants such as gabapentin and tricyclic antidepressants such as amitriptyline. This gap in knowledge of healthcare professionals regarding pain in clinical practice, which could limit the ability to achieve adequate pain relief in children has been previously documented and attributed among others to a lack of knowledge about various classes and dosages of analgesics found to be safe in the paediatric population and inadequate pain education in medical/nursing schools and during specialty trainings.[7,10,22]

Most physicians in this study had never prescribed morphine, which is considered the drug of choice for moderate to severe pain in children.[4,6,22] However, a significantly higher proportion of nurses had administered it, possibly because in the study setting, nurses are not licensed to prescribe morphine, but are usually the ones who administer the morphine prescribed by the physician. Moreover, some of them had either rotated through the children emergency wards or children wards where oncology patients are being managed and for whom morphine had been requested by the paediatric oncologist. The finding in this study is in consonance with previous reports showing that opioids were rarely prescribed by Nigerian healthcare providers.[7,16,22,24] Oshikoya *et al* in south western part of the country found that morphine and its derivatives, except hydromorphone, were rarely used to treat acute pain in children with sickle cell anaemia during emergency admission, despite the recommendation in our National guideline to do so.[20,23] This is worrisome as poorly treated acute pain could progress to chronic pain, which patients tend to eventually live with. [25] In the same vein, even though post operative pain was not singled out in the present survey, studies exploring its practice in Nigeria showed that the use of morphine was practically nonexistent as commonly used postoperative analgesia included paracetamol, NSAIDs, tramadol and pentazocine. [24,26]

Problems of unavailability, high cost, fear of serious adverse effects, in particular respiratory depression, masking symptoms, addiction and analgesic abuse by parents were pointed out in the afore mentioned studies as reasons why morphine was not used, but this was not explored in the present survey.[16,22,24] This may be subject to future studies, as well as chronification of pain among our patients. On the other hand, our finding is at disparity with

the study by Freitas *et al*, in which 40% of health professionals in 3 paediatric units in Brazil had administered opioids.[7,21] This is not surprising as about half of the respondents in the later study had some kind of training in pain management during or after their education.[7,21]

The majority of respondents were unaware of the existence of a protocol for management of pain in children in the study setting, which is similar to the report of Nasir *et al*. in Nigeria.[24] It is more likely because treatment of pain has not been given priority in the clinical management of patients in our institution, which could reflect deficiencies at both individual and institutional levels. Individual deficiencies, stemming from little pre-/in-service training or none, or lack of organised pain management workshops, while a lack of policy and will from the hospital management to insure implementation of guidelines for the treatment of pain in the paediatric population are also possible reasons.

## 5. CONCLUSION

Healthcare workers in our Institution had poor knowledge of non-pharmacological and pharmacological interventions, including adjuvant medications recommended for analgesia in children. Development and distribution of standard treatment protocols for paediatric pain, as well as setting of minimum standards and organizational support are advocated to enable professionals offer optimum pain management to their patients

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## ETHICAL APPROVAL

Approval for the study was obtained from the Ethics Committee of the University of Port Harcourt Teaching Hospital and from the respondents.

Disclaimer: - This manuscript was presented in a Conference.

Conference name: 3rd International Conference of Children's Palliative in Durban, South Africa, in May 2018

Available link: - <http://www.icpcnconference.org/wp-content/uploads/2018/06/130.-Management-of-Paediatric-Pain-How-Knowledgeable-are-Healthcare-Providers-at-a-Tertiary-centre-Southern-Nigeria-Dr-Gracia-Eke.pdf>

## REFERENCES

1. Mersky H. A list with definition and notes on usage: Recommendations by the IASP Subcommittee on Taxonomy. Pain 1979; 6: 249-52



- 261 2. Downing J, Atieno M, Debere S, Mwangi-Powell F, Ddungu H, Kiyange F, et al.  
262 Beating Pain: A Pocket Guide for Pain Management in Africa. Kiwanuka R, Leng M,  
263 Meiring M, Ulaya P, Williams S, Ddungu H, et al., (eds). Kampala, Uganda: African  
264 Palliative Care Association (APCA); 2010: 1-105 p.
- 265 3. Amery J. A Really Practical Handbook of Children's Palliative Care for doctors and  
266 nurses anywhere in the World: Part 6; 44-122; Lulu Publishing services; 2016
- 267 4. Meiring M. Management of common symptoms and problems in paediatric palliative  
268 care. Clinical guidelines. USAID, HPCA 2012: 13-27
- 269 5. Taddio A, Chambers CT, Halperin SA, Ipp M, Lockett D, Rieder MJ, et al.  
270 Inadequate pain management during routine childhood immunizations: The nerve of  
271 it. Clin Ther. 2009; 31: S152-67
- 272 6. World Health Organization. WHO guidelines on the pharmacological treatment of  
273 persisting pain in children with medical illnesses. WHO Library Cataloguing-in-  
274 Publication Data. WHO geneva. 2012;1-172.
- 275 7. Sanya EO, Kolo PM, Makusidi MA. A survey on doctors' knowledge and attitude of  
276 treating chronic pain in three tertiary hospitals in Nigeria. Niger Med J 2014; 55(2):  
277 106-10
- 278 8. International Association for the Study of Pain, Subcommittee on Taxonomy.  
279 Classification of chronic pain. Descriptions of chronic pain syndromes and definitions  
280 of pain terms. Pain Suppl. 1986; 3: S1-226. PMID: 3461421
- 281 9. Zeltzer LK, Krane EJ, Palermo TM. Pediatric pain management. In: Kliegman RM,  
282 Stanton BF, St Geme JW, Schor NF. Nelson Textbook of Pediatrics. 20<sup>th</sup> edition,  
283 Philadelphia. Elsevier, Inc 2016; pp 430-47
- 284 10. Albertyn R, Rode H, Millar AJW, Thomas J. Challenges associated with paediatric  
285 pain management in Sub Saharan Africa. Int J Surg. 2009;7:91-3
- 286 11. Elusiyan JB, Senbanjo IO. Management of pain in children: a review of the literature.  
287 Niger J Med. 2005; 14(4): 363-67
- 288 12. Adeboye M, Fakayode E, Adeniran M. Paediatrics pain management. Symp Niger J  
289 Paed. 2013; 40(1): 97-104
- 290 13. Kahsay H. Assessment and treatment of pain in pediatric patients. Curr Pediatr Res.  
291 2017; 21(1): 148-57
- 292 14. Legal notice: Federal Republic of Nigeria Official Gazette No 2, Abuja 15<sup>th</sup> May 2009  
293 Vol. 96. Page B 39- 40. Legal Notice on Publication of 2006 Census Final Results
- 294 15. Birnie KA, Chambers CT, Fernandez CV, Forgeron PA, Latimer MA, McGrath PJ, et  
295 al., Hospitalized children continue to report undertreated and preventable pain. Pain  
296 Res Manag 2014; 19(4): 198-204
- 297 16. Offiong UM. Pain management in paediatric practice: The view of paediatric health  
298 care provider on the use of analgesics. WJPMR 2017; 3(11): 44-7
- 299 17. Taddio A, McMurtry CM, Shah V, Riddell RP, Chambers CT, Noel M, et al. Reducing  
300 pain during vaccine injections: clinical practice guideline. CMAJ 2015; 187(13): 975-  
301 82
- 302 18. Weisman SJ, Bernstein B, Schechter NL. Consequences of Inadequate analgesia  
303 during painful procedures in children. Arch Pediatr Adolesc Med. 1998;152:147-49
- 304 19. Hamilton JG. Needle phobia: A neglected diagnosis. J Fam Pract 1995; 41:169-175
- 305 20. Amery J, Meiring M, Albertyn R, Jassal S. Pain. In: Children's Palliative Care in  
306 Africa. Amery J (ed). Oxford University Press 2009: 97-124
- 307 21. De Freitas GRM, De Castro CG, Castro SMJ, Heineck I. Degree of knowledge of  
308 Health Care professionals about pain management and use of opioids in Pediatrics.  
309 Pain Medicine 2014; 15: 807-19
- 310 22. Oshikoya K, Oreagba I. Acute pain management in children with sickle cell anaemia  
311 during emergency admission to a teaching hospital in Lagos, Nigeria. SAJCH 2015;  
312 9(4): 119-23

- 313 23. The Federal Ministry of Health, Nigeria. National Guideline for the Control and  
314 Management of Sickle Cell Disease. 2014
- 315 24. Nasir AA, Ameh EA, Abdur-Rahman LO, Kolawole IK, Oyedepo OO, Adeniran JO.  
316 Postoperative pain management in children: A survey of practices of pediatric  
317 surgeons in Nigeria. J Clin Sci 2017;14:138-43.
- 318 25. Pergolizzi JV, Raffa RB, Taylor R. Treating acute pain in the light of the  
319 chronification of pain. Pain Management Nursing 2014; 15(1): 380-90
- 320 26. Osifo OD, Aghahowa ES. Safety profile and efficacy of commonly used analgesics  
321 in surgical neonates in Benin City, Nigeria. Am J Perinatol 2008;25:617-22  
322  
323  
324

UNDER PEER REVIEW