

**Prevalence and Intensity of Gastrointestinal Parasitic Infection of Goats in
Belo Sub Division, Boyo Division, North West Region of Cameroon**

ABSTRACT

Background: Goats provide milk, meat, fiber and are also companion pets. They are easy to rear though production is affected by environmental conditions and parasitism. The aim of this study was to investigate the prevalence and intensity of gastrointestinal (GI) parasitic infections in goats.

Materials and Methods: A study was conducted in Belo Sub Division from July 2016 to October 2016. A total of 499 fecal samples were randomly collected directly from the recta of 499 goats in six villages and analyzed for the detection of any parasitic ova or oocysts using standard saturated sodium chloride flotation technique, while fecal egg/oocyst count were estimated using the modified McMaster technique.

Results: The study found that all 499 goats with a mean EPG value of $494,3 \pm 374,8$) were found to harbor at least two gastrointestinal parasites. The prevalence and intensity of various parasites encountered respectively were: *Eimeria spp* (86%), ($455,2 \pm 400,8$), *Haemonchus spp* (74,5%), ($1282,9 \pm 1244,4$), *Toxocara spp* (72,5%) ($953,3 \pm 814,3$), *Charbertia spp* (55,9%), ($448,2 \pm 416,0$), *Fasciola spp* (45,4%), ($475,0 \pm 338,1$), *Moniezia spp* (42,2%), ($828,6 \pm 793,9$), *Oesophagostomum spp* (33,1%), ($638,3 \pm 463,5$), *Strongyloides spp* (32,5%), ($200,0 \pm 00$), *Trichostrongylus spp* (28,3%) ($200,0 \pm 00$), *Trichuris spp* (23,7%) ($200,0 \pm 00$), *Teladorsagia spp* (14,6%), ($200,0 \pm 00$) and *Nematodorus spp* (8,1%), ($50,0 \pm 0,0$). There was no significant difference in prevalence (100%) in the different age groups, type of husbandry management system and locality ($P > 0.05$) except for gender where there was significant difference.

Conclusion: Gastrointestinal parasitic infections in goats from Belo Sub Division are common, with a very high prevalence. This high prevalence of gastrointestinal parasitism among the goats possibly reflected grazing, low immunity due to malnutrition and lack of anthelmintic treatment programs.

Key Words: Prevalence, Intensity, Gastrointestinal Parasites, Goats, Belo Sub Division.

36 INTRODUCTION

37 Livestock increases economic status of the rural population and plays a crucial role in the
38 economic well-being of populations Worldwide. Goats are the oldest domesticated animals by
39 man [1]. Evolutionary biology indicates that goats were domesticated about 10,000 years ago
40 at the dawn of the Neolithic age [2]. The West African dwarf goats are popular as hobby goats
41 due to their easy maintenance resilience and small stature. In rearing them, they do not require
42 as much space as the larger dairy goat counterparts. Their gentle and friendly natures make
43 them good companion pets [3]. Goats are important to man in different spheres and aspects of
44 life. They provide milk which is more easily digestible than cow milk [4]. Their milk is also
45 used in industries in the production of cheese .The rearing of goats provides
46 employment and income to rural populations. In order to rear goats, a minimum investment
47 of money is required, even without specific arrangement for housing and homemade supplied
48 feed. Grazing is mostly done on road-side grass lands and fields [5].

49 According to Gadahi et al [6], improper care, unhygienic environment, extreme
50 climate and close contact with infected animals, goats get infected with a variety of parasites
51 Parasitism in goat is a substantial problem plaguing farmers across the nation and it has a
52 highly detrimental effect on the goat industry [7]. Production potential of livestock
53 development programs is plagued in tropical and subtropical areas by prevalence of
54 helminthiasis which causes high mortality and great economic losses [8].

55 Goat production and rearing in Belo is challenged with gastrointestinal parasitism
56 being one of the main obstacles. The prevalence of gastrointestinal parasites is related to agro-
57 climatic conditions like quantity and quality of pasture, temperature, humidity and grazing
58 behavior of the host [9]. Infection with gastrointestinal parasites of goats depends on the
59 quantity and species of goats present, general health, age, nutritional and immunological
60 status of the animal. These infections occur mostly as mixed infections of different GIT
61 parasites. Emaciation, persistent diarrhea and weight loss are usually the main symptoms
62 [10].Villous atrophy causes impaired digestion and malabsorption of nutrients, leading to
63 decrease in live-weight gain, fiber and milk production as well as reproductive performance
64 of goats and therefore has a serious impact on animal health and productivity. Hence, GIT
65 parasitism of goats represents the greatest economic constraint and the most important
66 limiting factor of small ruminant production [11, 12,13] .

67 **Monitoring of research on parasitic invasions, particularly have wide biological**
68 **importance as well as practical. As a result of these studies it is possible to establish the**
69 **population of infected animals, and in several cases, to determine the composition of species**
70 **of parasites. Monitoring studies are also useful to determine the prevalence of parasites in**

71 ruminants. The invasion of parasites in adult animals runs mainly subclinical form, and are
72 not noticeable to owners of animals and very often also for veterinary services. Adult animals,
73 however, are a source of infection for young animals especially in small ruminants, sheep and
74 goats. Amongst the gastrointestinal parasitic diseases of greatest importance in goats are:
75 Nematodes (roundworms), Cestodes (tapeworms), Trematodes (liverflukes) and Coccidia [14,
76 15]. Therefore this study was designed to determine the prevalence and intensity of
77 gastrointestinal parasitic infestation of goats in Belo Sub division.

78 MATERIALS AND METHOD

79 Study area Description

80 This study was carried out in Belo Sub Division, Boyo Division, North West Region,
81 Cameroon from July 2016 to October 2016. Belo Sub Division is located about 50 km from
82 Bamenda. It is found between latitude $6^{\circ}4'$ and $6^{\circ}20'$ North, between longitude $10^{\circ}11'$ and
83 $10^{\circ}30'$ East. The entire Sub Division covers a surface area of about 46.068 square kilometers
84 and situated within part of the most mountainous sections of the Western highlands of
85 Cameroon.

86 Characteristics of sampled animals

87 The goats are grazed in open spaces, along the road, yard, and garbage sites and
88 around houses in the municipalities. The age of the goats considered for the study ranged
89 between 0–5 years, characterized as young goats (Less than 6 months old), adult goats (6 to
90 24 months old inclusive), and old goats (more than 24 months, but Less than 5 years old).
91 Goats of both sexes were involved in the study.

92 2.3. Parasitological Techniques

93 2.3.1. Collection of samples

94 Coprologic analysis was done to have a quantitative and qualitative appreciation of
95 the prevalence of infection of the parasites. For the qualitative analysis, faeces were analysed
96 by the double-centrifugal flotation technique using saturated sodium chloride solution. For
97 quantitative analysis or determination of the number of eggs per gram of faeces, the Mc
98 Master technique [16].

99 Classification of GI parasitic infections by virtue of mean EPG.

100 The animals were categorized as lightly, moderately and severely (heavily) infected
101 according to their egg per gram of feces (EPG) counts. Egg counts from 50-799, 800-1200
102 and over 1200 eggs per gram of feces were considered as light, moderate and heavy infection,
103 respectively [17].

104 **Statistical Analysis**

105 The collected data was stored in Excel 2007, later transferred to Statistical Packages
 106 for Social Science (SPSS version 19.0) for statistical analysis. The prevalence of
 107 gastrointestinal parasites was compared using Chi square test. Mann-Whitney test was used
 108 to evaluate parasite intensity between sex and Kruskal-Wallis test was used to compare
 109 parasite intensity between age and locality. A critical probability of (P < 0.05) was adopted
 110 throughout as a cut-off point for statistical significance between groups compared. All
 111 statistical tables retrieved from analysis with SPSS.

112 **RESULTS.**

113 **Overall Prevalence and Intensity of gastrointestinal parasites.**

114 The analysis of fecal samples (Table1) revealed that all 499 samples examined, were
 115 positive with mixed gastrointestinal parasite infections. There was an overall prevalence of
 116 100 percent and a mean EPG value of (494,3 ± 374,8).

117 **Table1: Prevalence and Intensity of gastrointestinal parasites**

Parasites	Number examined	Number of infested animals	Prevalence % of infestation	Intensity (mEPG/OPD ±SD)*	
Nematodes	<i>Nematodirus spp</i>	40	8.1	50.0 ± 0.0	
	<i>Haemonchus spp</i>	499	74.5	1282.9 ± 1244.4	
	<i>Oesophagostomum spp</i>	163	33.1	638.3 ± 463.5	
	<i>Chabertia spp</i>	279	55.9	448 ± 416.0	
	<i>Trichuris spp</i>	117	23.7	200.0 ± 00	
	<i>Strongyloides spp</i>	162	32.5	200.0 ± 00	
	<i>Teladorsagia spp</i>	73	14.6	200.0 ± 00	
	<i>Toxocara spp</i>	362	72.5	953.3 ± 814.3	
	<i>Trichostrongylus spp</i>	141	28.3	200.0 ± 00	
Trematodes	<i>Fasciola spp</i>	499	224	45.4	475.0 ± 338.1
Cestodes	<i>Moniezia spp</i>	499	208	42.2	828.6 ± 793.9
Protozoa	<i>Eimeria spp</i>	499	429	86	455.2 ± 400.8

118

119 **Influence of gender on Prevalence and Intensity infections**

120 Globally, out of the 499 goats examined, 236 were males, while 263 were females
 121 Both sexes each had 100% prevalence of GI parasitic infections with statistical significance
 122 difference (P<0.05). Multiple infections were more prevalent in female goats than male goats.
 123 (Table 2). Female goats had the highest mean EPG value of 526,5 ± 388,3 compared to 462,0
 124 ± 283,6 in male goats with no significant difference (P>0.05) (Table 3)..

125 **Table 2: Prevalence of infections by gender**

Parasite	Gender		Total N(%)	P-value
	Males	Females		

	N0. examined	N0. infected	Prevalence (%)	N0. examined	N0. infected	Prevalence (%)		
<i>Nematodirus spp</i>		12	2.4		28	7.7	40 (8.1)	0.028
<i>Haemonchus spp</i>		206	41.3		166	33.3	372 (74.6)	0.001
<i>Oesophagostomum spp</i>		31	6.3		132	26.8	163 (33.1)	0.000
<i>Chabertia spp</i>		86	17.2		193	38.7	279 (55.9)	0.000
<i>Trichuris spp</i>		38	7.7		79	16.0	117 (23.7)	0.000
<i>Eimeria spp</i>		194	38.9		235	47.1	429 (86.0)	0.022
<i>Fasciola spp</i>	236	60	12.2	263	164	33.3	224 (45.4)	0.000
<i>Moniezia spp</i>		68	13.8		140	28.4	208 (42.2)	0.000
<i>Strongyloides spp</i>		49	9.8		113	22.6	162 (32.5)	0.000
<i>Teladorsagia spp</i>		19	3.8		54	10.8	73 (14.6)	0.000
<i>Toxocara spp</i>		170	34.1		192	38.5	362(72.5)	0.80
<i>Trichostrongylus</i>		43	8.6		98	19.6	141 (28.3)	0.000

126

127 **Table 3: Gender related intensity (mEPG/OPG)**

128

Parasite	Gender				Total (mEPG/OPG±SD)	P- value
	Males		Females			
	N0 examined	Intensity (mEPG/OPG±SD)*	N0 examined	Intensity (mEPG/OPG±SD)		
<i>Nematodirus spp</i>		50.0 ± 0.0		50.0 ± 0.0	50.0 ± 0.0	1
<i>Haemonchus spp</i>		798.4 ± 680.1		1767.3 ± 1808.4	1282.9 ± 1244.4	0.00
<i>Oesophagostomum spp</i>		600.0 ± 0.0		676.6 ± 463.5	638.3 ± 463.5	0.47
<i>Chabertia spp</i>		400.0 ± 0.0		496.3 ± 416.0	448 ± 416.0	0.00
<i>Trichuris spp</i>		200.0 ± 0.0		200.0 ± 00	200.0 ± 0.0	1
<i>Eimeria spp</i>	236	463.9 ± 374.8	263	446.4 ± 426.7	455.2 ± 400.8	0.08
<i>Fasciola spp</i>		460.0 ± 393.7		489.9 ± 282.4	475.0 ± 338.1	0.04
<i>Moniezia spp</i>		823.5 ± 810.0		833.6 ± 777.8	828.6 ± 793.9	0.57
<i>Strongyloides spp</i>		200.0 ± 0.0		200.0 ± 00	200.0 ± 0.0	1
<i>Teladorsagia spp</i>		200.0 ± 0.0		200.0 ± 00	200.0 ± 0.0	1
<i>Toxocara spp</i>		1148.2 ± 1144.4		758.3 ± 484.2	953.3 ± 814.3	0.02
<i>Trichostrongylus</i>		200.0 ± 0.0		200.0 ± 00	200.0 ± 0.0	1

129

131 **Influence of age on Prevalence and Intensity of infection**

132 Table 4 shows the prevalence of infection by age group of the goats examined. 70 were
 133 young goats, 303 were adult goats, while 126 were old goats. A prevalence of 100% was
 134 recorded in each of the 3 age groups with no significant difference (P>0.05). Multiple
 135 infections were more prevalent in adult goats than young and old goats Table 5 shows the
 136 intensity of GI parasites by age group of the study. The highest mean EPG was recorded by
 137 the young goats (558,1 ± 331,2), followed by the adult goats (529,3 ± 349,5) and old goats
 138 (463,0 ± 330,7) with no significant difference (P>0.05).

139 **Table 4: Age related Prevalence of infections.**

Parasite	Age						Total (N) Prevalence (%)	P- value
	Young		Adults		Old			
	N0 infected	Prevalence (%)	N0 infected	Prevalence (%)	N0. infected	Prevalence (%)		
	13	2.6	7	1.4	20	4.1	40 (8.1)	0.00

<i>Nematodirus spp</i>								
<i>Haemonchus spp</i>	40	8.0	240	48.1	92	18.4	372 (74.5)	0.001
<i>Oesophagostomum spp</i>	33	6.7	86	17.4	44	8.9	163 (33.1)	0.001
<i>Chabertia spp</i>	34	6.8	168	33.7	77	15.4	279 (55.9)	0.23
<i>Trichuris spp</i>	13	2.6	85	17.2	19	3.9	117 (23.7)	0.01
<i>Eimeria spp</i>	44	8.8	291	58.3	94	18.8	429 (86.0)	0.00
<i>Fasciola spp</i>	27	5.5	121	24.5	76	15.4	224 (45.4)	0.00
<i>Moniezia spp</i>	20	4.1	130	26.4	58	11.8	208 (42.2)	0.14
<i>Strongyloides spp</i>	21	4.2	98	19.6	43	8.6	162 (32.5)	0.84
<i>Teladorsagia spp</i>	0	0	47	9.4	26	5.2	73 (14.6)	0.00
<i>Toxocara spp</i>	58	11.6	210	4.1	94	18.8	362 (72.5)	0.06
<i>Trichostrongylus</i>	12	2.4	85	17.0	44	8.8	141 (28.3)	0.03

140

141

142 **Table 5: Age related intensity of infection**

143

Parasite	Age						P value
	Young		Adult		Old		
	N0 examined	Intensity (mEPG/OPG)	N0 examined	Intensity (mEPG/OPG)	N0 examined	Intensity (mEPG/OPG)	
<i>Nematodirus spp</i>		50.0 ± 0.0		50.0±0.0		50.0 ± 0.0	0.06
<i>Haemonchus spp</i>		1300.7 ± 1220.1		1473.9±1434.2		1072.9 ± 1079.6	0.001
<i>Oesophagostomum spp</i>		566.7 ± 196.6		1052.2±614.7		296.0 ± 102.0	0.001
<i>Chabertia spp</i>		450.0 ± 227.7		750.0±385.9		509.4 ± 437.8	0.23
<i>Trichuris spp</i>		200 ± 00		200.0±0.0		200.0 ± 0.0	0.07
<i>Eimeria spp</i>	70	404.5 ± 281.2	303	563.2±422.2	126	397.9 ± 499.0	0.00
<i>Fasciola spp</i>		850.2 ± 498.8		418.2±272.0		475.1 ± 243.5	0.00
<i>Moniezia spp</i>		840.0 ± 409.3		578.0±594.0		1067.8 ± 972.0	0.14
<i>Strongyloides spp</i>		200.0 ± 0.0		200.0±0.0		200.0 ± 0.0	0.84
<i>Teladorsagia spp</i>		200.0 ± 0.0		200.0±0.0		200.0 ± 0.0	0.70
<i>Toxocara spp</i>		1435.6 ± 1140.7		710.6±470.8		714.3 ± 531.4	0.06
<i>Trichostrongylus</i>		200.0 ± 0.0		200.0±0.0		200.0 ± 0.0	0.07

144

145

146 **Influence of husbandry systems on infection**

147 The prevalence of GI parasites by type of husbandry management system of the goats
 148 is shown in (Table 6). 210 goats were on free range, while 289 goats were tethered. Both
 149 types of husbandry management systems recorded each 100% prevalence of GI parasitic
 150 infections with no significant difference (P>0.05). Multiple infections were more prevalent in
 151 tethered goats than free range goats. The highest mean EPG value (513,5± 412,4) was
 152 recorded by tethered goats compared to (446,2 ± 333,1) on free range system with no
 153 significant difference (P>0.05).(Table 7)

154 **Table 6: Influence of husbandry systems on infection**

Parasite	Husbandry system				Total N(%)	P-value
	Tethered		Free range			
	N0. infected	Prevalence (%)	N0. infected	Prevalence (%)		
<i>Nematodirus spp</i>	20	4.1	20	4.1	40 (8.1)	0.323
<i>Haemonchus spp</i>	209	41.9	163	32.7	372 (8.1)	0.180

					(74.5)	
					163	
<i>Oesophagostomum spp</i>	104	21.1	59	12.0	(33.1)	0.101
					73	
<i>Charbertia spp</i>	32	6.4	41	8.2	(14.6)	0.008
					117	
<i>Trichuris spp</i>	78	15.8	39	7.9	(23.7)	0.043
					429	
<i>Eimeria spp</i>	239	47.9	190	38.1	(86.0)	0.014
					224	
<i>Fasciola spp</i>	133	27.0	91	18.5	(45.4)	0.756
					208	
<i>Moniezia spp</i>	109	22.1	99	20.1	(42.2)	0.170
					162	
<i>Strongyloides spp</i>	71	14.2	91	18.2	(32.5)	0.000
					73	
<i>Teladorsagia spp</i>	32	6.4	41	8.2	(14.6)	0.008
					362	
<i>Toxocara spp</i>	196	39.3	166	33.3	(72.5)	0.006
					141	
<i>Trichostrongylus</i>	95	19.0	46	9.2	(28.2)	0.007

155

156 **Table 7: Influence of husbandry system on intensity of infections**

Parasite	Husbandry system				Total Intensity (mEPG/OPG) ±SD	P- value
	Tethered		Free range			
	N0 examined	Intensity (mEPG/OPG)±SD	N0 examined	Intensity (mEPG/OPG)±SD		
<i>Nematodirus spp</i>		50.00±0.0		50.0±0.00	50.0 ± 0.0	1
<i>Haemonchus spp</i>		1283.9±1253.3		1331.9±1237.5	1282.9 ± 1244.4	0.774
<i>Oesophagostomum spp</i>		642.5±485.1		674.1±441.9	638.3 ± 463.5	0.512
<i>Charbertia spp</i>		505.7±445.5		390.7±386.5	448 ± 416.0	0.001
<i>Trichuris spp</i>		200.0±0.0		200.0±0.0	200.0 ± 0.0	1
<i>Eimeria spp</i>	236	591.6±525.1	263	318.8±278.7	455.2 ± 400.8	0.022
<i>Fasciola spp</i>		515.1±345.2		435.1±331.5	475.0 ± 338.1	0.001
<i>Moniezia spp</i>		614.7±848.2		1042.5±739.6	828.6 ± 793.9	0.000
<i>Strongyloides spp</i>		200.0±0.0		200.0±0.0	200.0 ± 0.0	1
<i>Teladorsagia spp</i>		200.0±0.0		200.0±0.0	200.0 ± 0.0	1
<i>Toxocara spp</i>		1158.5±1046.5		748.1±582.1	953.3 ± 814.3	1
<i>Trichostrongylus</i>		200.0±0.0		200.0±0.0	200.0 ± 0.0	0.908

157

158 **Influence of locality on Prevalence and Intensity of GI parasite infections**


159 The spectrum of gastrointestinal parasites presented in figure 1 shows the prevalence
 160 of GI parasites by locality of sampled goats. 62 goats were examined from Anjin, 219 from
 161 Belo, 60 from Baingo, 47 from Kitchu, 90 from Mbessa, and 21 from Njinikejem. All 6
 162 Villages recorded 100% prevalence each with no significant difference (P>0.05) of GI-
 163 parasitic infections. Multiple infections were also more prevalent in Belo goats than goats in
 164 the other village. The highest mean EPG value was recorded in Belo with no significance
 165 difference (p>0.05).

166

167

168

169



170
171
172
173
174
175
176
177

Figure 1: Prevalence and intensity of GI parasitic infection in some Localities

UNDER PEER REVIEW

178 DISCUSSION

179 Prevalence and Intensity of GI parasite infections

180 Goats harbor a variety of gastrointestinal (GI) parasites. Data from this study indicated
181 that gastrointestinal parasitic infections in goats from Belo Sub Division were common, with
182 an overall prevalence of 100%. All the 499 goats examined were infected with at least two
183 gastrointestinal parasites amongst which were *Haemonchus spp*, *Nematodirus spp*,
184 *Oesophagostomum spp*, *Chabertia spp*, *Strongyloides spp*, *Teladorsagia spp*, *Toxocara spp*,
185 *Trichostrongylus spp*, *Trichuris spp*, *Moniezia spp*, *Fasciola spp*. and coccidian (*Eimeria spp*)
186 giving a total of twelve parasites (9 Nematodes, 1 cestode, 1 trematode and 1 protozoan).
187 Sathaporn et al. [18], Nuraddis et al. [19] and Choubisa et al. [20] also reported slightly
188 similar types of GI parasites. The gastrointestinal parasitic infection rate of 100% recorded in
189 goats during this study agrees with the 100% prevalence reported by Dogo et al. [20] in Vom
190 and 90.4 reported by Ntonifor et al. [21] in Jakiri. This is higher than the (87.2%) prevalence
191 reported by Nuraddis et al. [19] and (72%) reported by Paul et al. [22] in Maiduguri. This is
192 quite high and shows that the agro-ecological and geo-climatic conditions of the study area
193 favor the growth and multiplication of these parasites. Climatic conditions, particularly
194 rainfall, are frequently associated with differences in the prevalence of GI parasitic infections,
195 because free-living infective stages (eggs, larvae, cysts, and oocysts) survive longer in moist
196 conditions [19]. Belo Sub Division experiences about eight months of rainy season from mid
197 March to mid November and about four months of dry season from mid November to mid
198 March. Since the study was conducted from July to October towards the end of the rainy
199 season, higher parasitic infections might be related to the availability of browse and a longer
200 browsing time in the warm-rainy season by the host, sufficient moisture and optimum
201 temperature. These create favorable conditions allowing for the larval development, oocyst
202 sporulation and survival of the infective larvae stage [23]. The high prevalence in this study
203 could also be attributed to illiteracy on the side of the goat keepers and their ignorance or
204 avoidance tendency of preventive measures [24]. For example, effective pasture management,
205 applied knowledge about host-parasite interactions and interrelations building the base for
206 low pasture infection rates for grazing animals, stocking rate reduction and regular intensive
207 monitoring of animal condition that can help optimize animal health status and anthelmintic
208 treatments [25]. The overall higher prevalence of GI parasitic infections in this study area
209 could also be attributed to lower immunity of hosts as a result of malnutrition [23, 24].
210 Among other factors that may have further contributed to these discrepancies observed

211 are host breeds and different husbandry practices. The physiological status of the animals
212 like parturition, lactation stage and pasture contamination can also influence the prevalence
213 of GI parasites in different areas [23].

214 Most important to the findings of Nuraddis et al.[19] compared to the present study,
215 *Monezia spp.* and *Eimeria spp.* were the only cestode and protozoa types found
216 respectively, a finding similar to Kanyari et al.²⁷ Encountered in this study were
217 *Nematodorus spp* and *Toxocara spp*, that Nuraddis et al.[19] did not encounter in Jimma,
218 Ethiopia. This difference may be due to variation in climate, parasite evolution or mixed
219 rearing that affect parasitic infection. The most prevalent and commonly observed parasite
220 was *Eimeria spp*, with a significant infection rate of (86%), which is higher compared to the
221 low prevalence (48%) reported by Kanyari et al. [26] in Kenya and (20.6%) reported by
222 Nuraddis et al.[19] in Jimma, Ethiopia. Similarly, low prevalence of (18.6) was reported by
223 Dogo et al. [21] , and Gebeyehu et al.[24] for *Eimeria spp* in Daegu, Korea. This high
224 prevalence of *Eimeria spp* in Belo Sub Division may be associated to the fact that *Eimeria*
225 oocysts are much resistant to disinfectants, and can remain in the environment (particularly
226 moist, shady areas) for long periods of time and maintain their infectivity. Stress factors such
227 as tethering, post weaning, dietary changes and other problems can precipitate an outbreak of
228 coccidiosis. In this study, the severity of GI parasitic infection depended on the number of
229 eggs per gram of feces. The intensity of infection measured by fecal egg or oocyst count
230 varied from light to heavy infection. In a high percentage of animals, light parasitic
231 infections were found, while heavy infections were less common. Among these
232 gastrointestinal parasites observed, *Haemonchus spp* had the highest overall mean EPG value
233 of 1445,2± 1594,4 which is higher than that reported by Ntonifor et al [22].

234 Female goats had higher multiple infections and mean EPG value than male goats
235 from our study and this agrees with the findings of Paul et al.[23] In a study by Sathaporn et
236 al.[20], male goats actually had a higher prevalence than female goats which disagrees with
237 our findings. This could be because most of the goats that are tethered in Belo Sub Division
238 are females.

239 In age related infections, multiple infections and mean EPG value was higher in
240 adults goats than the old and the young goats similar to the report of Gebeyehu et al.[24]
241 However, this result did not agree with the reports of Kanyari et al.[27], Gwaze et al. [28] and
242 Sathaporn et al.[20] who showed that young goats had higher prevalence of GI parasites than

243 adult goats. This middle age group had a significant higher prevalence of *Eimeria spp*
244 infections (58,3%) and higher oocyst numbers compared to other age groups in the present
245 study. This did not also agree with Sathaporn et al. [20] in Satun, Thailand who reported that
246 young goats had a higher prevalence (94,9%) of coccidial infections and higher oocyst
247 numbers in young goats (< 1 year) than older goats (> 2 years). This higher prevalence of GI
248 parasites and of coccidial infections in this age group might be due to the fact that a higher
249 incidence occurs during post weaning stress (since, coccidia is most frequently observed in
250 kids 2 to 4 weeks post weaning), tethering stress and stress related to dietary changes [29, 30]
251 in addition to the fact that immunity is low. The low prevalence of coccidial infections in the
252 young goats is probably due to the absence of this stress factors and in old goats probably
253 because of acquired immunity. Although natural immunity develops with repeated exposure³¹
254 younger goats remain highly susceptible. The Institute for International Cooperation in Animal
255 Biologics [32] reported that most ruminants stop shedding *Toxocara spp* eggs by the time
256 they are 2 to 4 months old and that *T.vitulorum* infections can be controlled by
257 eliminating patent infections, which occur only in 3 to 10 week old ruminants. Young goats
258 (<6 months old) had the highest mean EPG value of (630,8± 268,3) than other age groups,
259 with the highest parasitic intensity (2137,9 ± 4493,0) shown by *Toxocara spp* in this age
260 group. This high *Toxocara spp* intensity might probably be due to Transcolostral transmission
261 in the life cycle and sanitation standards related to *Toxocara spp*. [32] This finding even
262 though was not consistent with the reports of Nuraddis et al. [19], was not surprising because
263 naive young and old carriers frequently graze the same areas, coupled with the fact that young
264 goats have low immunity. The intensity of infection is also reportedly related to hygiene level
265 [33].

266 Goats examined in this study were either on free range or tethered systems all under
267 extensive management (grazing). Tethered goats actually had a higher multiple infections and
268 mean EPG than the free range goats. The highest infection rate of (47, 9%) was recorded by
269 *Eimeria spp.* in tethered goats. This high infection rate and intensity in tethered goats could be
270 explained by the fact that tethering is a stress factor [29,30]. Again most people in Belo Sub
271 Division tether goats in the same area throughout the tethering period with little rotation.
272 Consequently, the grazing environment becomes contaminated with various GI parasites eggs
273 and oocysts which infect the goats [25].

274 All Villages recorded 100% prevalence each of GI parasitic infection with no
275 statistical significance. These results differed from those of Sathaporn et al.[20] who reported

276 in Satun, that the prevalence of GI parasites of goats in seven Districts statistically varied
277 from 60% to 86.4% ($P < 0.05$). Belo had a higher multiple infections and mean EPG of
278 1233,6± 1145,3 compared to other five villages. Geographical consistence of prevalence in
279 Belo Sub Division might be due to the climatic conditions that are consistent in this area.
280 *Eimeria spp* recorded the highest prevalence of 38,2% and *Haemonchus spp* had the highest
281 mean EPG of 4467.3±4396.2 in Belo. Only Belo town can be classified as being a semi urban
282 town. The rest of the villages are rural. These geographical differences in the prevalence of
283 coccidial infections and other infections and high mean EPG value in Belo might be due to
284 the high population density and unhygienic conditions of the area compared to other Villages,
285 which leads to the high infection rates. Inadequate nutrition, however, which is common in
286 this area, may exacerbate the course of GI parasitic infections. The animals are generally
287 malnourished and suffer from other diseases, and are thus not resistant to nematode infection
288 [34].

289

290 CONCLUSION

291 Goats in Belo Sub Division are infested by gastrointestinal parasites. The adult goats
292 recorded higher multiple gastrointestinal parasites and mean EPG value than the young goats
293 and the old goats. Female goats recorded higher multiple gastrointestinal parasites and mean
294 EPG value than male goats. Tethered goats recorded higher multiple gastrointestinal parasites
295 and mean EPG value than free range. Belo recorded higher multiple gastrointestinal parasites
296 and mean EPG value than Njinikejem, Anjin, Kitchu, Baingo, Mbessa. Prevailing agro-
297 ecological and geo-climatic conditions, illiteracy on the side of goat keepers, avoidance
298 tendency of preventive measures and lack of anthelmintic treatments provide an ideal
299 condition for the transmission of the GI parasitic infections.

300 ETHICAL APPROVAL

301 All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85- 23,
302 revised 1985) were followed, as well as specific national laws where applicable. All experiments have
303 been examined and approved by the appropriate ethics committee.

304

305 Competing interests

306 We declare that we have no conflict of interest.

307 **Acknowledgements**

308 The authors wish to express sincere gratitude to the Laboratory of Applied Biology and
309 Ecology (LABEA).

310

311 **REFERENCES**

- 312 1. Hirsto, K. K. The History of the domestication of goat. 2008, 84.
- 313 2. Zeder M.A. & Hesse B. The initial domestication of goats (*Capra hircus*) in the Zagros
314 Mountains 10,000 years ago. *Science*.2000, 28 : 2254 – 2257 .
- 315 3. Mcloed L.A. Goats as pets. 1998, 25.
- 316 4. Edwards B.A.& Charlotte J.C. Fresh goat milk for infants: Myths and Realities. *London*
317 *Daily Telegraph*.2005, 5-7.
- 318 5. Devendra, C. Studies in the nutrition of the indigenous goat at Malaya and
319 requirement of liveweight gain. *Malay. Agri. J.*, 1992, 46: 98-118.
- 320 6. Gadahi, J. A., Arshed M. J., Ali Q., Javaid S. B. & Shah S. I. Prevalence of gastrointestinal
321 parasites of sheep and goat in and around Rawalpindi and Islamabad, Pakistan. *Vet.*
322 *World*.2009, 2:51–53.
- 323 7. Jones, R. Sheep Parasites and Diseases .[http:// www. kt. iger. bbsrc. ac. uk/ FACT %20](http://www.kt.iger.bbsrc.ac.uk/FACT%20sheet%20PDF%20files/kt36.pdf)
324 [sheet % 20 PDF % 20 files / kt 36 . pd f .2001, p.2 .](http://www.kt.iger.bbsrc.ac.uk/FACT%20sheet%20PDF%20files/kt36.pdf)
- 325 8. Al-Quaisy, H.H.K., Al-Zubaidy A.J., Altaf K.I. & Makkawi T.A. The pathogenecity of
326 haemonchosis in sheep and goat in Iraq. Clinical, parasitological and haematological
327 findings. *Vet. Parasitol*.1987, 24:221-228.
- 328 9. Pal R.A. & Qayyum M. Prevalence of gastrointestinal nematodes of sheep and goats in
329 upper Punjab, Pakistan. *Pak. Vet. J.* 1993, 13 (3): 138-141.
- 330 10. Scheuerle M. Anthelmintic resistance of *Haemonchus contortus* and the Famacha -
331 Method as a tool to delay the development of Anthelmintic resistance.*Inaugural-*
332 *Dissertation*.1st edition Parasitology. 2009, 46: 33-53
- 333 11. Perry B.D., Randolph T.F. Improving the assessment of the economic impact of
334 parasitic diseases and of their control in Production animals. *Vet. Parasitol*. 1999, 84, 145-
335 168.

- 336 12. Harper C. K.& Penzhorn B. L. Occurrence and diversity of coccidia in indigenous, Saanen
337 and crossbred goats in South Africa. *Veterinary Parasitology*. 1999, 82: 1-9
- 338 13. Kagira J. M. & Kanyari P. W. N. The role of parasitic diseases in causing mortalities in
339 small ruminants in a highly productive area of Central Province, Kenya. *Journal of South*
340 *Africa Veterinary Association*, 2001, 72: 147-149
- 341 14. Nuraddis I., Mulugeta T., Mihreteab B.& Sisay A. Prevalence of Gastrointestinal Parasites
342 of Small Ruminants in and Around Jimma Town, Western Ethiopia. *Acta Parasitologica*
343 *Globalis*. 2014, 5 (1): 26-32
- 344 15. Bagley C.V. Internal Parasites. Utah State University Extension, Logan UT. 84322-5600.
345 Available from: <http://www.tvsp.org/pdf/sheep/internal-parasites.pdf>. 1997.
- 346 16. Thienpont, D., Rochette, F.R., and Vanperijis, O.F.J. Diagnosis of verminosis by
347 coprological examinations. *Beerse, Belgium, Janssen Research Foundation*, 1979, P. 48-67.
- 348 19. Nuraddis I., Mulugeta T., Mihreteab B.& Sisay A. Prevalence of Gastrointestinal Parasites
349 of Small Ruminants in and Around Jimma Town, Western Ethiopia. *Acta Parasitologica*
350 *Globalis*. 2014., 5 (1): 26-32
- 351 20 Choubisa, S.L, Jaroli, V.J. Gastrointestinal parasitic infections in diverse species of
352 domestic ruminants inhabiting tribal rural areas of southern Rajasthan, India. *Journal of*
353 *Parasitic Diseases*. 2013. 37(2): 271-275.
- 354 20. Sathaporn J. Sinsamut S. Nongnuch P. Wissanuwat C. Witaya K.& Roger W.
355 Gastrointestinal Helminthic and Protozoal Infections of Goats in Satun, Thailand. *Trop Med*
356 *Parasitol*. 2012, 35:48-54.
- 357 21. Dogo G. I., Karaye P., Patrobas M.G., Galadima M.& Gosomji I.J.,(). Prevalence of
358 Gastrointestinal Parasites and their impact in Domestic animals in Vom, Nigeria. *Saudi*
359 *Journal of Medical and Pharmaceutical Sciences*. 2017,3, 211-216
- 360 22. Ntonifor H. N., Shei S. J., Ndaleh N. W.& Mbunkur G.N., Epidemiological studies of
361 gastrointestinal parasitic infections in ruminants in Jakiri, Bui Division, North West Region
362 of Cameroon. *Journal of Veterinary Medicine and Animal*. 2013. 5 (12), 344-352,
- 363 23. Paul B.T., Biu A.A., Gadzam M.A., Ali M., Mana H.P., Jairus Y. Point prevalence
364 and intensity of gastrointestinal parasite ova/oocysts and its association with body

- 365 condition score(BCS) of Sheep and goats in Maiduguri, Nigeria. The journal of
366 Advances in Parasitology. 2016. 3-88.
- 367 24. Gebeyehu E. B., Seo M. G., Jung B. Y., Byun J. WOem., J. K., Kim H. Y. & Kwak D.
368 Prevalence of Gastrointestinal Parasites in Korean Native Goats.*The Journal of Animal &*
369 *Plant Sciences*. 2013. 23(4): 986-989.
- 370 25. William J.A.P. An Introduction to Animal Husbandry in the Tropics. Bailliere Tindall,
371 London. 2001. 1:92-97.
- 372 26. Gerold R. & Hannah S. Alternative management strategies to prevent and control
373 endoparasite diseases in sheep and goat farming systems - a review of the recent scientific
374 knowledge.*Landbauforschung Völkenrode*. 2007. 2 (57):193-206
- 375 27. Kanyari P.W.N., KagiraJ.M. & MhomaR.J. Prevalence and intensity of Endoparasites
376 in small ruminants kept by farmers in Kisumu Municipality, Department of Veterinary
377 Pathology, Microbiology and Parasitology, Faculty of Veterinary Medicine, University of
378 Nairobi, Kenya. 2009.
- 379 28. Gwaze F.R., Chimonyo & DzamaK. Prevalence and loads of gastrointestinal parasites
380 of goats in the communal areas of the Eastern Cape Province of South Africa. *Small*
381 *Rumin. Res*. 2009. 84:132- 134.
- 382 29. John M. Ucd Vet Views .*California Cattlemen's Magazine* School of Veterinary
383 Medicine, University of California-Davis fluke vaccines. *Parasitology Today*. 2007. .,14, 224-
384 228.
- 385 30. Leite-Browning M. Coccidiosis of Goats and Prevention. Alabama Cooperative Extension
386 System Alabama, A&M University. 2009.
- 387 31. Newton S.E.& Munn E.A. The development of vaccines against gastrointestinal nematode
388 parasites particularly *Haemonchus contortus*.*Parasitol Today*. 1999. 15: 162-22.
- 389 32. Institute For International Cooperation in Animal Biologics (IICAB) Toxocariasis.
390 Toxocarosis, Visceral Larva Migrans, Ocular Larva Migrans, Larval Granulomatosis,
391 Toxocaral Retinitis. Iowa State University. College Of Veterinary Medicine
392 www.cfsph.iastate.edu/IICAB.2005.
- 393 33. Foreyt W.J. Coccidiosis and cryptosporidiosis in sheep and goats. *Vet Clin North Am*
394 *Food Anim Pract*. 1990. 6:655-70.
- 395 34. Maichomo M.W., Kagira J.M & Walker T point prevalence of gastrointestinal parasites in

396 calves, sheep and goats in Magadi Division, South-Western Kenya. Onderstepoort. Journal of
397 Veterinary Research. 2004. 71:257–261
398

UNDER PEER REVIEW