

Influence of Variety and Sowing Date on Incidence of *Cercospora* Leaf Spots Disease of Groundnut (*Arachis hypogaea* L.) in Makurdi, Benue state of Nigeria.

Richard Isa Bwala¹, Chigoziri Ekhuemelo², and Dauda Zacharia¹

¹Department of Crop Protection, Faculty of Agriculture, P. M. B.1069 University of Maiduguri, Nigeria.

³Department of Crop and Environment Protection, Federal University of Agriculture, P. M. B. 2373 Makurdi, Benue State, Nigeria.

ABSTRACT

Leaf spots disease of groundnut caused by *Cercospora* pathogens is one of the major economic production constraints militating against groundnut production in Nigeria. Field experiments were conducted during the 2011 and 2012 rainy seasons at the Teaching and Research Farm of the Federal University of Agriculture, Makurdi Nigeria to assess the effect of sowing dates on the incidence of *Cercospora* leaf spot of groundnut. The 2 x 4 x 3 factorial (2 groundnut varieties/ 4 sowing dates/ 3 replications) experiment was laid out in a Randomized Complete Block Design (RCBD) and replicated three times. Results indicated no significant ($P > 0.05$) effect of *Cercospora* leaf spot incidence on the two varieties in 2011 but in 2012 Ex-Dakar recorded significantly ($P \leq 0.05$) higher leaf spot incidence at 54 DAS and 61 DAS compared to Borno-Red variety. Sowing groundnut seeds from 14th June to 29th June recorded significantly higher ($P \leq 0.05$) leaf spot disease incidence compared with sowing groundnut seeds in May. Ex-Dakar variety recorded higher leaf defoliation compared with Borno-Red variety in 2011 and 2012 seasons. Results indicated that Borno-Red had significantly ($P \leq 0.05$) higher 100 seed weight in 2011, while Ex-Dakar recorded higher 100 seed weight in 2012. The results has proved that early sowing of groundnut in May can be employed as alternative strategy for the management of *Cercospora* leaf spot disease of groundnut in Makurdi, Nigeria

Key word: Disease incidence, Groundnut, leaf defoliation, leaf spot, Sowing dates,

1. INTRODUCTION

Leaf spots disease is one of the major biotic production constraint of groundnut in Nigeria and other parts of the World particularly where the crop is not grown under any protection umbrella. *Cercospora arachidicola* Hori (Early leaf spot) and *Phaeoisariopsis personata* (Berk & Curt) (Late leaf spot) are the two main pathogenic fungal micro-organisms responsible for the disease. The destructive nature of the two diseases on groundnut crop has given it a significant recognition worldwide, including other parts of Africa [22]. The disease is prevalent in almost all groundnut growing areas of the world and become endemic frequently ([16], [10]). Farmers in the developing countries have reported huge yield losses as a result of the negative impact of the disease attack on their crops. Infected soil, debris and shells are the three potential sources of leaf spots **inocula**. The pathogens usually **remain** in the soil and can infect almost every parts of the crop causing lesions on leaves, petioles, pegs, main stems and lateral branches [21]. The primary initial inoculum is responsible for the onset of the disease epidemic, while the rain-splash and wind-blown moist air helps in dispersing the secondary and tertiary spores to the adjacent susceptible plants. The environmental conditions and genetic make-up of the groundnut varieties plays a major role in the level of disease incidence and carry-over of the disease from seasons to season in different agro-ecological locations. In West Africa, about 50 to 70% yield loss have been reported [24] and because of the destructive nature of the disease, international attention has been given to the disease causal pathogens [12].

The used of unilateral chemicals for the management of the disease have been practiced in the developing countries since 1970s. Incidentally, these chemicals were associated with unwanted and unintended human and environmental consequences such as pesticides persistence, resistance, residual, pest resurgence and environmental pollution apart from their exorbitant cost and not readily available sometimes (Richard **et al.** [18]).

The manipulation of sowing date is another very important disease management strategy that has to be put into consideration in any groundnut production; but because of variation in weather conditions in different locations it may not auger well to adopt a specific sowing date. [6], reported that sowing date influences vegetative and reproductive growth period and the availability of weather parameters. And when climatic conditions are not suitable for need of one of yield components, it would negatively affect the seed yield [4].

Therefore, the objective of this study was to assess the influence of sowing dates on incidence of groundnut leaf spot disease in Makurdi, Benue State of Nigeria.

2. MATERIALS AND METHODS

The study was conducted at the Teaching and Research Farm of the Federal University of Agriculture Benue State, Nigeria during the 2011 and 2012 cropping seasons. The location lies between the Latitude 7.41⁰N and Longitude 8.35⁰E; at an elevation of 95m above sea level located within the Southern Guinea Savanna of Nigeria [2]. The experimental field has been under intensive cultivation of groundnut crops for more than fifteen years which ensures availability of adequate natural source of disease inoculum. The site received total annual rainfall of 955.74mm and 1492.80mm in 2011 and 2012 respectively, and relative humidity of 69.45% and 72.83% in the first and second year respectively. It had maximum and minimum temperatures of 32.98⁰C and 21.71⁰C; 32.8⁰C and 20.69⁰C in 2011 and 2012 respectively [9].

The experimental site was cleared and ridged manually using cutlass and hoe. The total experimental area used measured 49m x 8m = 392m² (0.0392ha). The 2 x 4 x 3 factorial experiment (2 groundnut varieties / 4 sowing dates / 3 replications) was arranged in a Randomized Complete Block Design (RCBD). The two groundnut varieties (Borno Red and Ex-Dakar) and four different sowing dates (D1 = 15th May, D2 = 30th May, D3 = 14th June and D4 29th June) formed the 24 treatment combinations. The groundnut varieties were assigned in the main plots each measuring 15m x 2m (30m²) whereas the four different sowing dates were assigned in the sub-plots each measuring 3 x 2m (6m²) with four ridges and replicated 3 times leaving 1m walking alley between the main plots and sub plots. Groundnut seeds were sown at a spacing of 20cm within rows and 75cm between rows at the rate of 2 seeds per hole. The experimental plots were kept weed-free throughout the study period and no chemical was used.

2.1 Data collection

2.1.1 Leaf spot incidence (%)

Disease incidence was assessed at 47, 54 and 61 DAS. The disease incidence was taken by counting the number of plants infected in the net plot and divided by the total number of plants per net plot and multiply by 100 using the disease incidence formula according to Turaki *et al.* [23].

$$Z = \frac{K}{Y} \times 100$$

Where:

Z = Disease incidence (%)

K = Number of plant stands infected by the disease in the net – plot

Y = Number of plant stands infected and uninfected by the disease in the net – plot

2.1.2 Leaf defoliation (%)

The leaf defoliation was assessed at 70, 80, 90 and 100 DAS. Eight plants were selected at random, tagged in the net plot of each plot and scored for leaf defoliation using the leaf defoliation scale according to [20].

1 = No leaf fall

2 = less than 10% leaf fall

3 = 10 – 25% leaf fall

4 = 25 – 50% leaf fall

5 = More than 50% leaf fall

$$\text{Leaf defoliation} = \frac{\sum n \times 100}{N \times 5}$$

Where: $\sum n$ = summation of individual assessments

N = Total number of plants assessed

5 = Highest score of the defoliation scale.

2.1.3 One hundred seeds weight (g/plot): At 130 DAS of physiological maturity stage, all the groundnut stands in the net plot of each plot were carefully lifted up, the pods were picked, sun dry and shelled separately. One hundred seeds were randomly hand-picked from the net plot of each plot and weighed in gram using a sensitive electronic weighing scale model (Sartorius 6MBH Gottingen-Type Fabr-Nr.) in the Plant Pathology laboratory of the Department of Crop and Environmental Protection Department, University of Agriculture, Makurdi Benue State.

2.2 Data Analysis

Data collected were subjected to analysis of variance {ANOVA} using [19] version. Two-Way analysis of variance was used and means were separated using the Duncan's New Multiple Range {DNMRT} at 5% probability level [17].

3. RESULTS

Results on incidence of leaf spots of groundnut as influenced by **variety**, sowing date and their interaction at 47, 54 and 61 DAS in 2011 and 2012 cropping seasons are presented in Table 1. In 2011, the effect **variety** on incidence of leaf **spots** disease **was** not significantly different ($P > 0.05$) from 47 to 61 DAS. However, in 2012, the effect of **variety** on disease incidence **was** significantly higher ($P \leq 0.05$) at 54 and 61 DAS but was not significant at 47 DAS. In 2011 and 2012, sowing date significantly influenced disease incidence at 47, 54 and 61 DAS. The interaction between **variety** and sowing date on disease incidence were not significantly different at 47 DAS and 61 DAS in 2011 and **at 47 and 61 DAS** in 2012, but was significant at 54 DAS in 2011.

In 2011, results show that groundnut sown between Early and 29th June had higher leaf spot incidence while the lowest leaf spot incidence of 14.51% was recorded from those groundnut plants sown in 15th May followed by those sown on 30th May which had disease incidence of 22.18% at 47 DAS. Similarly at 54 DAS, higher disease incidence **values** of 44.45% and 50.29% were recorded from those groundnut plants sown in 14th June and 29th June respectively. Those groundnuts sown in 15th May and 30th May had lower leaf spot disease spot incidence of 38.76% and 36.79% respectively at 54 DAS. Groundnut sown in 30th May, Early and 29th June had significantly higher leaf spot disease incidence of 64.69%, 68.43% and 67.80% respectively at 61

DAS. In 2012, among the varieties, Ex-Dakar recorded significantly ($P<0.05$) higher leaf spot incidence of 59.41% and 69.51% at 54 and 61 DAS respectively compared to Borno Red. In 2012, results indicated that those groundnut plants sown in 29th June recorded significantly higher

Table 1: Effect of Varieties, Sowing Dates and their Interaction on Incidence of Leaf Spot of Groundnut at 47, 54 and 61 DAS in 2011 and 2012 Cropping Seasons

Variety/Sowing Dates/Interactions	2011 Cropping Season			2012 Cropping Season		
	47 DAS	54 DAS	61 DAS	47 DAS	54 DAS	61 DAS
Variety (V)						
Borno Red	25.16±2.30	41.56±2.73	61.14±1.91	20.57±2.45	44.04±2.50 ^b	59.96±3.11 ^b
Ex-Dakar	24.06±2.42	43.58±2.01	66.91±2.42	24.88±2.07	59.41±1.91 ^a	69.51±1.99 ^a
<i>P-value</i>	0.74NS	0.55NS	0.07NS	0.19NS	≤0.05	≤0.05
<i>CV</i>	33.20	19.50	11.90	34.60	14.90	14.00
Sowing Dates (S)						
15th May	14.51±1.48 ^c	38.76±3.34 ^b	55.18±1.98 ^b	15.06±1.26 ^c	43.98±4.39 ^b	51.77±4.27 ^b
30th May	22.18±1.79 ^b	36.79±1.40 ^b	64.69±3.57 ^a	18.45±2.01 ^c	46.37±4.19 ^b	67.77±2.15 ^a
14th June	29.16±1.78 ^a	44.45±3.38 ^{ab}	67.80±2.54 ^a	25.44±2.75 ^b	57.92±3.14 ^a	69.02±2.17 ^a
29th June	32.58±1.53 ^a	50.29±2.16 ^a	68.43±1.70 ^a	31.94±1.42 ^a	58.63±3.04 ^a	70.39±2.27 ^a
<i>P-value</i>	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05
<i>CV</i>	16.40	15.50	9.80	21.00	17.70	10.80
Interactions (VX S)						
Borno Red						
15th May	17.63	32.20 ^b	54.00	13.00	34.52	44.23
30th May	20.71	36.10 ^b	60.60	14.37	37.75	63.67
14th June	27.56	49.70 ^{ab}	65.00	24.31	51.30	65.50
29th June	34.74	48.20 ^{ab}	65.00	30.59	52.59	66.45
Ex-Dakar						
15th May	11.40	39.20 ^b	56.40	17.11	53.45	59.31
30th May	23.65	37.50 ^b	68.80	22.52	54.99	71.86
14th June	30.76	45.30 ^{ab}	70.60	26.58	64.53	72.54
29th June	30.42	52.40 ^a	71.90	30.59	64.68	74.33
<i>P-value</i>	0.08NS	≤0.05	0.84NS	0.65NS	0.34NS	0.46NS
<i>CV</i>	14.80	12.00	9.20	19.40	7.10	7.40

Mean values within each column followed by the same letter (s) are not significantly different ($P \leq 0.05$) from each other according to Duncan's New Multiple Rang Test (DNMRT). CV = coefficient of variation, Ns = Not significant

disease incidence of 31.94%, followed by those sown in Early June which had 25.44% disease incidence, while those groundnut plants sown in 15th May and 30th May recorded lower disease incidence of 15.06% and 18.45% respectively at 47 DAS. Results at 54 DAS revealed that higher disease incidence of 57.92% and 58.63% were recorded on those groundnut plants sown in 29th June and 14th June respectively, while those sown on 15th May and 30th May recorded lower leaf spot incidence of 43.98% and 46.37% which was not significantly ($P > 0.05$) different. Similarly at 61 DAS higher disease incidence of 70.39%, 69.02% and 67.77% were recorded on those groundnut plants sown in 29th June, 14th June and 30th May respectively, while those sown in 15th May had the lowest leaf spot incidence of 51.77%. The results of interaction between varieties and sowing date on disease incidence at 54 DAS revealed that disease incidence of 52.40% was recorded from Ex-Dakar sown in 29th June which was not significantly different from disease incidence of same variety sown in 14th June, Borno Red sown in 14th June and 29th June respectively. The interaction between varieties and sowing dates resulted in lower disease incidence in both Borno Red and Ex-Dakar varieties sown in 15th May and 30th May at 54 DAS. The interaction of variety and sowing dates was not significant at 47 DAS and 61 DAS in 2011 and throughout the period of 2012 season.

Results on leaf defoliation of groundnut as influenced by variety, sowing date and their interaction at 70, 80, 90 and 100 DAS in 2011 cropping season are presented in Table 2.

The effect of variety on leaf defoliation was significant ($P \leq 0.05$) at 70, 80, 90 and 100 DAS. The effect of sowing date on leaf defoliation was significant from 80 to 100 DAS. The effect of interaction between varieties and sowing date on leaf defoliation was not significant ($P \leq 0.05$) at 70 DAS but were significant at 80, 90 and 100 DAS. Ex-Dakar recorded significantly higher leaf defoliation 19.10%, 39.94%, 64.07% and 80.38% compared with Borno Red had lower leaf defoliation of 14.94%, 33.14%, 54.34% and 75.56% at 70, 80, 90 and 100 DAS respectively. Result revealed that at 80 DAS, higher leaf defoliation of 39.45%, 37.87% and 34.89% were recorded on those groundnut plants sown on 29th June, 14th June and 15th May accordingly compared to those groundnut plants sown in 30th May. At 90 DAS, the leaf defoliation trend was similar to those of 80 DAS. Those groundnut plants sown in 14th June and 29th June recorded higher leaf defoliation of 80.79% and 82.82% respectively compared to those groundnut plants sown in mid and 30th May at 100 DAS. Sowing groundnut in June resulted in significantly higher leaf defoliation compared with sowing groundnut in May.

Table 2: Effect of Varieties, Sowing Dates and their Interaction on Leaf Defoliation of Groundnut at 70, 80, 90 and 100 DAS at 2011 Cropping Season

Variety/Sowing Dates/Interactions	70 DAS	80 DAS	90 DAS	100 DAS
Variety(V)				
Borno Red	14.94±0.42 ^b	33.14±0.82 ^b	54.34±1.20 ^b	75.56±1.36 ^b
Ex-Dakar	19.10±0.16 ^a	39.94±1.02 ^a	64.00±1.52 ^a	80.38±1.68 ^a
<i>P-value</i>	≤0.05	≤0.05	≤0.05	0.03
<i>CV</i>	9.90	8.80	8.00	6.80
Sowing Dates (S)				
15th May	17.26±0.94	34.89±1.87 ^{ab}	56.37±2.60 ^{ab}	74.95±2.02 ^b
30th May	15.57±1.06	33.96±1.70 ^b	55.13±2.13 ^b	73.33±1.97 ^b
14th June	17.33±1.23	37.87±1.77 ^{ab}	61.98±2.64 ^{ab}	80.79±1.69 ^a
29th June	18.06±1.20	39.45±1.82 ^a	63.19±2.70 ^a	82.82±1.39 ^a
<i>P-value</i>	0.46NS	0.04	0.04	≤0.05
<i>CV</i>	16.00	12.00	10.50	5.60
Interactions (V X S)				
Borno Red				
15th May	15.72	31.25	51.40	72.74
30th May	13.66	30.98	51.47	71.44
14th June	14.70	34.26	56.27	78.13
29th June	15.71	36.10	58.20	79.93
Ex-Dakar				
15th May	18.80	38.53	61.35	77.16
30th May	17.47	36.94	58.80	75.21
14th June	19.96	41.47	67.69	83.44
29th June	20.41	42.80	68.17	85.72
<i>P-value</i>	0.63NS	0.96NS	0.78NS	0.96NS
<i>CV</i>	9.10	7.10	5.90	5.00

Mean values within each column followed by the same letter (s) are not significantly different ($P \leq 0.05$) from each other according to Duncan's New Multiple Rang Test (DNMRT). CV = coefficient of variation, Ns = Not significant

Results on effect of **variety**, sowing date and their interaction on leaf defoliation at 70, 80, 90 and 100 DAS in 2012 cropping season are presented in Table 3. The effect of **variety** on leaf defoliation was **significant** ($P \leq 0.05$) at 80 and 90 DAS but was not at 70 and 100 DAS. The effect of sowing dates on leaf defoliation was **significant** at 70, 90 and 100 DAS but was not at 80 DAS. The interactive effect of sowing dates and varieties on leaf defoliation was not significant ($P > 0.05$) from 70 to 100 DAS

Results indicated that Ex-Dakar recorded significantly higher leaf defoliation of 39.61% and 66.13% compared to Borno Red which had the lower leaf defoliation of 32.44% and 56.88% at 80 DAS and 90 DAS respectively. Those groundnut plants sown in 29th June, 14th June and 15th May recorded higher leaf defoliation of 19.07%, 18.31% and 17.54% respectively while those groundnut plants sown in 30th May recorded the lowest leaf defoliation of 16.35% at 70 DAS. Results showed that at 90 DAS those groundnut plants sown on 15th May, 14th June and 29th June recorded higher leaf defoliation of 59.75%, 64.72% and 64.74% respectively compared with leaf defoliation of those groundnut plants sown on 30th May. Similarly, at 100 DAS, higher leaf defoliation of 86.37% and 83.58% which did not differ significantly from each other was recorded from those groundnut plants sown in 29th June and 14th June respectively. Result indicated that those groundnut plants sown in 15th May and 30th May recorded lower leaf defoliation of 77.41% and 78.72% respectively.

Table 3: Effect of Varieties, Sowing Dates and their Interaction on Leaf Defoliation of Groundnut at 70, 80, 90 and 100 DAS at 2012 Cropping Season

Variety/Sowing Dates/Interactions	70 DAS	80 DAS	90 DAS	100 DAS
Varieties (V)				
Borno Red	17.52±0.54	32.44±1.22 ^b	56.88±0.90 ^b	80.98±1.46
Ex-Dakar	18.11±0.64	39.61±0.55 ^a	66.13±1.25 ^a	82.06±1.19
<i>P-value</i>	0.49NS	≤0.01	≤0.01	0.57NS
<i>CV</i>	11.60	9.10	6.20	5.70
Sowing Dates (S)				
15th May	17.54±0.94 ^{ab}	33.74±2.01	59.75±1.91 ^{ab}	77.41±1.31 ^b
30th May	16.35±0.58 ^b	34.17±2.43	56.82±1.82 ^b	78.72±1.53 ^b
14th June	18.31±0.61 ^{ab}	36.94±1.70	64.72±2.60 ^a	83.58±0.92 ^a
29th June	19.07±0.89 ^a	39.26±1.16	64.74±2.18 ^a	86.37±0.57 ^a
<i>P-value</i>	0.04	0.16NS	0.04	≤0.01
<i>CV</i>	10.70	12.80	8.60	3.40
Interactions (V X S)				
Borno Red				
15th May	17.51	29.62	55.58	74.80
30th May	16.41	29.19	52.99	78.30
14th June	18.37	33.78	58.96	84.93
29th June	17.81	37.19	60.01	85.89
Ex-Dakar				
15th May	17.59	37.87	63.92	80.01
30th May	16.29	39.15	60.65	79.13
14th June	18.25	40.10	70.48	82.23
29th June	20.33	41.33	69.48	86.86
<i>P-value</i>	0.60NS	0.21NS	0.10NS	0.10NS
<i>CV</i>	11.10	6.60	2.20	3.10

Mean values within each column followed by the same letter (s) are not significantly different ($P \leq 0.05$) from each other according to Duncan's New Multiple Rang Test (DNMRT). CV = coefficient of variation, Ns = Not significant

Results on effect of **variety**, sowing date and their interaction on 100 seed weight (g) in 2011 and 2012 cropping seasons are presented in Table 4.

The effect of **variety** on 100 seed weight (g) was **significant** in 2011 and 2012. The effect of sowing dates, interaction between the varieties and sowing dates on 100 seeds weight **was** not **significant** ($P>0.05$) in 2011 and 2012. Results of 2011 shows that Borno Red recorded significantly the highest 100 seed weight of 40.95g compared with Ex-Dakar which had 39.59g, but on contrary, Ex-Dakar recorded significantly the highest 100 seed weight of 39.63g compared to Borno Red which recorded 35.64g in 2012.

Table 4: Effect of Varieties, Sowing Dates and their Interaction on 100 Seed Weight (g) in 2011 and 2012 Cropping Seasons

Variety/Sowing Dates/Interactions	100 Seed wt(g)	
Variety (V)		
Borno Red	40.95±0.26 ^a	35.64±1.23 ^b
Ex-Dakar	39.59±0.45 ^b	39.63±0.46 ^a
<i>P-value</i>	≤0.01	≤0.01
<i>CV</i>	3.20	8.60
Sowing Dates (S)		
15th May	39.84±0.72	39.37±1.03
30th May	40.60±0.64	37.40±1.70
14th June	40.31±0.63	37.07±1.64
29th June	40.33±0.45	36.70±1.83
<i>P-value</i>	0.85NS	0.64NS
<i>CV</i>	3.80	10.30
Interactions (V X S)		
Borno Red		
15th May	40.67s	37.97
30th May	41.55	34.77
14th June	41.29	35.10
29th June	40.31	34.71
Ex-Dakar		
15th May	39.02	40.77
30th May	39.66	40.03
14th June	39.33	39.05
29th June	40.35	38.68
<i>P-value</i>	0.57NS	0.94NS
<i>CV</i>	3.40	

Mean values within each column followed by the same letter (s) are not significantly different ($P \leq 0.05$) from each other according to Duncan's New Multiple Rang Test (DNMRT). CV = coefficient of variation, Ns = Not significant

4. DISCUSSION

Sowing date has a significant effect on plant growth and disease development in any agricultural production system. The performance of crop in relation to date of sowing would enable the researcher to validate recommendation of sowing date at that agro-ecological location (Azamali *et al.*[3])

The finding from this study indicates that early sowing significantly lowered leaf spot disease incidence compared with the late sown plants in the two cropping seasons. Ex-Dakar was more susceptible and exhibited significantly higher disease incidence compared to Borno Red at 54 and 61 DAS in 2012. 15th May sown plants had lower leaf spot incidence compared to 30th May sown crops which were moderate in their disease incidence, while the 29th June and 14th June sown crops recorded significant higher leaf spot incidence from 47 to 61 DAS in 2011 and 2012. Relatively groundnut sown in 15th May and 30th May had significantly lower leaf spot incidence in 2011 and 2012 compared with those sown in 14th June and 29th June of both years. Likewise, higher disease incidence was recorded on groundnut plants sown on 14th June and 29th June from 54 to 61 DAS in 2012 than in 2011 which could be attributed to higher annual rainfall (1,492.8mm) and favorable average relative humidity (72.85%) resulting to higher disease incidence in 2012 (Table 1). This result agrees with the finding of [7] which reported that leaf spots is generally more severe on late sown groundnut plants than the early sown plants due to warm temperature later in the season that are more favorable for the growth and spread of the leaf spot pathogens. Similarly, significantly higher percentage of leaf defoliation was observed on Ex-Dakar compared to local Borno-Red in the two cropping seasons. This may be due to the higher leaf spot disease incidence recorded on Ex-Dakar which could have resulted in subsequent higher leaf defoliation. This result agrees with the report of [5] in which *Cercospora* leaf spots varied among the three groundnut varieties they studied. Also [5] reported a higher leaf spot disease incidence in Ex-Dakar compared with RMP 12 and Damboa local varieties. The higher leaf defoliation recorded on plants sown in June could have been attributed to late sowing and intensive buildup of disease inoculum which led to higher disease incidence and consequently higher leaf defoliation. This observation is in agreement with the report of [7] which stated that groundnut plant sown late began leaf shading early because of early severe disease infection and attack on the tender leaves. Similarly, Adipala *et al.* [1] reported that late

sown groundnut showed high disease incidence and pest infestation resulting in lower yield whereas Waliyar *et al.*[25] reported that high leaf losses of up to 25 - 43% could result in the disruption of the photosynthetic process, lesser pods and lower fruit quality.

The variation in seed weight recorded among the groundnut varieties in the two cropping seasons implies that seed weight is a genotypic trait and could be equally influenced by environmental factors which is in conformity with the report of ([14, 15, 13]). This result agreed with report of Gorbet *et al.* [11] that number of spots per leaf and leaf defoliation both have a negative correlation with yield whereas [8] revealed in their finding that *Cercospora* leaf spot disease reduced yield by every 1% increase in disease severity. The management of groundnut leaf spot through the manipulating of sowing date was very effective in reduction of disease incidence and leaf defoliation.

5. CONCLUSION

This study has shown that early sowing of groundnut in May could be another alternative fruitful option for the management of *Cercospora* leaf spot disease of groundnut in Makurdi considering its cost effectiveness on the part of the farmers and environmental friendliness in terms of biodiversity.

REFERENCES

1. Adipala EC, Mukankusi S, Kyamanywa G, Epier HW, Willson H. Integrated Management of Groundnut Insect Pests and Diseases. *Seventh Annual Report of Integrated Pest Management Collaborative Res.* 2000; Support Program, Uganda (Abstract).
2. Agboola SA. An Agricultural Atlas of Nigeria, Oxford University Press. 1979; PP. 33-49.
3. Azamali SNRC, Nageswara Rao J, Craigon DR Wadis, Williams JH. Methods of calculating the population/yield relations of groundnut (*Arachis hypogaea* L.) in semi-arid climates. *Journal of Agricultural Science Cambridge.* 1993; 120: pp.23-222.
4. Baldwin J. Seeding rate, row patterns and planting dates. Peanut Home Page. 2005; Available at <http://www.caes.vga.edu/commodities/fieldcrop/peanut/index.html> (verified on 26 August 2008)

5. Bdliya BS, Mohammad AS. Effect of Inter-cropping millet with groundnut on the control of *Cercospora* leaf spot of groundnut in the Sudan Savanna of North Eastern Nigeria. *J. Sust. Agric.* 2006; 29(2): 19-14.
6. Canavar O, Kaynak MA. Growing degree day and sunshine radiation effects on peanut pod yield and growth. *African J. of Biotech.* 2010; 9(15):293-301.
7. Cooperative Extension. Peanut Production Update. College of Agriculture and Environmental Sciences and Family and Sciences. The University of Georgia. 2013; Pp. 70 – 73.
8. Das S, Roy TK. Assessment of losses in groundnut due to early and late leaf spots. *IAN.* 1995; 15: 34 – 36.
9. Department of Meteorological Services Makurdi, Benue State of Nigeria; Weather data of 2011 and 2012 cropping seasons.
10. Ghewande MP, Misra DP. Groundnut rust: A challenge to meet. *Seeds and Farms*, 1983; 9(6):12-15
11. Gorbet DW, Shokes FM, Jackson LF. Control of peanut leaf spot with combination of resistance and fungicide treatment. *Peanut Sci.* 1982; 9: 87-90.
12. ICRISAT. Information Bulletin no. 21. International Crop Research Institute for the Semi-Arid Tropics, Patancheru, Andhra Pradesh, India. 1985; pp 1-8.
13. Khatun A, Kabir G, Bhuiyan AH. Effect of harvesting stages on the seed quality of lentil (*Lens culinaris* L.) during storage. *Bangladesh J. Agric. Res.* 2009; 34:565-576.
14. Kola S, Gupta K. The timing of physiological maturity of seeds sunflower: Evaluation through multiple tests. *Seed Sci Tech.* 1982; 10: 457-467.
15. Mahesha CR, Channayeeraswami CR, Kurdikeri MB, Shekhargroude M, Merwade MN. Seed maturation studies in sunflower genotypes. *Seed Res.* 2001; 29: 95-97.
16. Meena B. Effect of *Pseudomonas fluorescens* Pflta based formulation under different storage periods against late leaf spot of groundnut. *International J. Plant Protec.* 2011; 4(1): 137-139.
17. Obi IU. Statistical methods of detecting differences between treatment means and research methodology issues in laboratory and field experiments. 2nd edition AP Express Publishers, Nsuka. 2002; 117pp

18. Richard BI, Degri MM, Dauda Z. Use of Organic Pesticides as Component of Integrated Pest Management (IPM) in the Nigeria Sudan Savanna. In. Kwari JD, Dugie IY, Gwary MM, Alhassan AB, Raji AO, Sotannde OA, Mallafiya DM, Sastawa BM. (eds): *Organic Agriculture: A Panacea for Sustainable Environmental and Food Security. Proceedings of the 6th National Conference on Organic Agriculture held at Faculty of Agriculture, University of Maiduguri, 21st - 24th November, 2010*; 195-205p
19. SAS. *Statistical Analysis Software Package Version 9.1*, SAS Institute Inc., Cary, NC, USA, 2009.
20. Schiller JM, Sampoapol R. Foliar Disease and Their Control in Rain fed Groundnut Production in Thailand. *Tropical Pest Management*. 1981; 27(1):13-23.
21. Shokes FM, Culbreath AK. Early and late leaf spots. Pages 17 – 20 in; compendium of Peanut Disease, Second Edition. N. Kokalis-Burelle, N, DM Porter, R. Rodriguez-Kabana, DH. Smith, P. Subrahmanyam, eds. *American Phytopathology Society*; St. Paul. USA. 1997; Pp. 17-20.
22. Subrahmanyam P, Smith DH. Influence of Temperature, Leaf Wetness period, Leaf Maturity, and Host Genotype on Web Blotch of Peanut *Oleagineux*. 1989; *Vol. 44*, pp. 27-31.
23. Turaki ZGS, Richard BI, Degri MM. Assessment of the effect of fungicide and seed rate on the incidence of leaf blast (*magnaphorth grisea*), on the growth of foxtail millet (*setaria italic* (L.) P. BEAUV) in North-eastern Nigeria. *Asian Journal of Agriculture and Rural Development*. 2014; 4 (2): pp 142-148.
24. Waliyar F. "Evaluation of Yield Losses Due to Groundnut Leaf Diseases in West Africa, Summary Proceedings of the second *ICRISAT Regional Groundnut Meeting for West Africa*. Niame.1990; 11 – 14 September 1990.
25. Waliyar F, Adomou M, Traore A. Rational use of fungicide application to maximize peanut yield under foliar disease pressure in West Africa. *Plant Disease*. 2000; 84: 120-1211.