

Original Research Article

Climatic Effects on Quality Parameters and Their Relationships of Bread Wheat Genotypes (*Triticum aestivum* L.) Grown Under Semi-Arid Region

Ayşe Gulgun OKTEM^{1*}, Abdullah OKTEM¹

*Corresponding author : gulgunoktem@harran.edu.tr

¹University of Harran, Faculty of Agriculture, Department of Field Crops, Sanliurfa, Turkey

ABSTRACT

Aims: This study was conducted to determine climatic effects to quality parameters of bread wheat genotypes grown in the semi-arid region.

Study design: The experiment was carried out in a randomized complete block design with three replications. Thirty-three bread wheat genotypes (*Triticum aestivum* L.) were grown in a field trial and kernel samples analyzed for thousand kernel weight, test weight, protein content, wet gluten, dry gluten and SDS-sedimentation value.

Place and Duration of Study: The research was carried out during the 2008 and 2009 growing seasons at Sanliurfa, Turkey.

Methodology: For analyses 20 main spikes that contained fully developed kernels were chosen randomly from each plot and taken to the laboratory for analyses. The nitrogen content of kernels was determined using the Kjeldahl method and the result was multiplied by the factor 5.7 to calculate the protein content of kernels. Test weight and Sodium Dodecyl Sulphate (SDS)-sedimentation values were determined using standard procedures. Wet and dry gluten values were determined using a glutomatic system after separating gluten from the soluble starch and protein fractions.

Results: Genotypes were significant ($P \leq 0.01$) for all tested characteristics. Thousand kernel weight ranged from 25.8 to 42.3 g, test weight from 73.7 to 81.7 kg hl⁻¹, protein content from 9.7 to 14.8%, wet gluten from 28.5 to 42.2%, dry gluten from 9.4 to 14.1% and SDS-sedimentation value from 19 to 39 ml. Bezostaya-I, Kutluk-94, Altay-85, Harmankaya-99, Lirasa, Kirgiz-95 and Dagdas-94 genotypes had the best quality among tested genotypes in the semi-arid climatic conditions. Protein content was positively correlated with wet gluten, dry gluten and SDS-sedimentation value, respectively. Relationships were significant between wet gluten and both dry gluten and SDS-sedimentation value. A positive significant correlation was found between thousand kernel weight and SDS-sedimentation value and between dry gluten and SDS-sedimentation value.

Conclusion: Protein content, wet and dry gluten and SDS-sedimentation value were affected by climatic conditions of semi-arid region. Protein content, wet and dry gluten value were high but SDS-sedimentation value were low due to high temperature and low precipitation in semi-arid region.

Keywords: Climatic effect, bread wheat, protein, wet and dry gluten, SDS, correlation

*E-mail address: gulgunoktem@harran.edu.tr.

16 1. INTRODUCTION

17 Wheat (*Triticum aestivum* L.) which is considered to be a good source of energy and
18 nutrients is used primarily for human consumption. Bread and bakery products have an
19 important role in human nutrition. Wheat is one of the most important crops in the world. It is
20 grown both in arid and semi-arid regions of the world as a rain-fed conditions. Turkey is one
21 of the largest producers of wheat in the world with about 7.7 million ha sown area and 21.5
22 million tons of annual production. Sowing area of wheat in the southeast region of Turkey is
23 about 759 717 ha and production is about 2 456 204 tons [1].

24 Wheat products are considered to be a good source of energy and nutrients for the human
25 body. The major use of wheat is bulgur, pasta and noodles, and various types of breads.
26 Bread and bakery products have an important role in human nutrition. Bakery products,
27 supplemented with various nutrients, have been gaining popularity worldwide.

28 The wheat processing industry requires grain lots which are consistent for moisture, test
29 weight and protein content. Wheat quality is a concept in continuous evolution in response to
30 market demands and consumer preferences for specific attributes of different end-products
31 [2]. The technological quality of wheat for milling and baking use varies widely. Growing
32 conditions, climate and variety characteristics are the most important factors affecting quality
33 and affects changes in protein and starch quality. About 13.5% protein content in Canada
34 and 11-13% in USA are acceptable standards for wheat, respectively [3, 4]. A thousand
35 kernel weight of 35-40 g is required in USA [3]. Gangadharappa et al. [5] stated that the
36 required quality parameters of wheat are a test weight of 79.6 kg hl⁻¹, gluten values in the
37 range of 7.93-9.60%, SDS-sedimentation value of 46 ml and protein concentration of 9.5%.
38 About 74 kg hlt⁻¹ test weight is required in Australia [4]. Pasha et al. [6] reported 19.67-36 mL
39 SDS-sedimentation volume value, 13.82-43.13% wet gluten content and 4.46 -14.55% dry
40 gluten values.

41 Wheat production under abiotic stress conditions has become important in recent years.
42 Grain composition and the quality of the wheat kernel are affected by both variety and
43 environment [7, 8, 9, 10]. The environment (climate, soil, agronomic practices, etc.) exerts a
44 strong influence on the expression of the technological quality of different cultivars [8, 11].

45 Rharrabti et al. [12] reported that thousand kernel weight and test weight are greatly affected
46 by climatic parameters. Grain protein content, perhaps the most important quality feature for
47 wheat, is known to be influenced by climatic factors such as rainfall and temperature, cultivar
48 and available moisture during grain filling [12, 13, 14, 15]. The protein content in wheat
49 kernels is influenced by climatic conditions [14, 16, 17, 18, 19, 20, 21]. After anthesis, heat
50 or drought may increase grain protein content [22, 23]. Faergestad et al. [24] emphasis
51 climatic conditions affect gluten composition of wheat kernel.

52 The availability of soil water is a major factor limiting wheat production in most regions of the
53 world. Not only is the amount of precipitation usually small, but there is often the problem of
54 poor and unpredictable distribution. Especially under semiarid and arid environments water
55 deficits often limit grain yields and quality. Effect of high temperatures and deficit water on
56 grain protein composition during grain filling period was reported by Oktem [25].

57 Genotype is also one of the most important quality factors. Wheat quality has implications for
58 human health and nutrition. The present investigation was undertaken with thirty-three wheat
59 genotypes to determine some quality parameters of bread wheat genotypes grown in semi-
60 arid region.

The objectives of this study were: (i) to determine some quality parameters of bread wheat genotypes grown in semi-arid region; (ii) to investigate the influence of climatic parameters on the expression of different grain quality characteristics; (iii) to study the relationships between quality traits; (iv) to evaluate the effect of environmental conditions on these relationships.

2. MATERIAL AND METHODS

This study was conducted during 2008 and 2009 in the Harran Plain, Sanliurfa, South-east Anatolia region of Turkey (altitude: 465 m; 37°08' N and 38°46' E). Climate varies from arid to semi-arid depending on the year. Total precipitation was 314 and 448 mm for 2008 and 2009 growing seasons, respectively. Monthly average temperature and total precipitation values were given in the Table 1. The soil texture of the experimental field was clay. Field capacity, permanent wilting point and bulk density of the soil was 33.8% (dry basis), 22.6% and 1.41 Mg m⁻³, respectively.

Table 1. Average temperature and total precipitation values of experiment years.

Months	Average Temperature (°C)		Minimum Temperature (°C)		Maximum Temperature (°C)		Total precipitation (mm)	
	2008-09	2009-10	2008-09	2009-10	2008-09	2009-10	2008-09	2009-10
October	20.5	21.9	9.6	12.5	35.3	34.8	22.5	76.6
November	14.1	12.2	5.8	4.7	28.5	24.0	35.3	35.5
December	7.0	10.0	3.0	2.0	22.1	18.7	37.7	121.2
January	5.7	8.3	-4.7	-3.2	15.7	18.8	29.8	95.7
February	8.0	9.1	0.1	-1.9	17.3	19.7	54.5	23.5
March	10.0	13.8	1.5	1.1	23.0	25.2	55.3	42.7
April	15.8	17.4	5.9	6.6	27.5	29.2	48.8	26.2
May	22.7	24.0	10.0	11.0	37.0	36.8	4.7	7.1
June	29.6	29.4	17.8	17.5	40.0	42.2	9.2	0.5
July	32.0	33.9	20.3	20.0	41.5	45.2	3.2	-
August	30.6	33.6	20.9	23.0	41.2	43.6	-	-
September	25.0	28.5	11.3	18.5	39.4	40.0	6.9	0.2

Thirty-three bread wheat genotypes (*Triticum aestivum* L.) were used in the study. The experiment was carried out in a randomized complete block design with three replications. Plot sizes were 6 m by 1.2 m (7.2 m²) and each plot consisted of six rows with a row spacing of 20 cm. The seeds were sown at 30-40 mm depth with a density of 500 plants m². At sowing, 60 kg ha⁻¹ of pure P and N was applied to each plot; this was followed by 60 kg ha⁻¹ of N when the plants reached 25-30 cm in height. As a first fertilizer Compose (20, 20, 0 NPK) and secondary Ammonium Nitrate (26% N) fertilizers were used at experiment.

For analysis of the kernel, 20 spikes that contained fully developed kernels were chosen randomly from each plot and taken to the laboratory for analysis. The nitrogen content of kernels was determined using the Kjeldahl method [26] and the result was multiplied by the factor 5.7 [27] to calculate the protein content of kernels, this was expressed on dry weight basis. Test weight of wheat samples were determined using standard procedures [28]. Sodium Dodecyl Sulphate (SDS)-sedimentation value [29] was determined for the wheat samples. Wet and dry gluten values were determined using a glutomatic system after separating gluten from the soluble starch and protein fractions [30].

*E-mail address: gulgunoktem@harran.edu.tr.

An analysis of variance (ANOVA) [31] was performed on the two years combined for the physico-chemical characteristics to evaluate statistical differences between genotypes. Differences among means were assessed by the Duncan's multiple range test ($P \leq 0.05$) [32]. A correlation analysis was performed to determine relationship among tested quality characteristics [33].

3. RESULTS AND DISCUSSION

Genotypes were statistically significant ($P \leq 0.01$) for thousand kernel weight, test weight, protein content, wet gluten, dry gluten and SDS-sedimentation traits.


3.1. Thousand Kernel Weight and Test Weight

The Pehlivan genotype gave the highest thousand kernel weight value whereas the lowest value was obtained from Sultan-95 genotype (Fig. 1.). One thousand kernel weight ranged from 25.8 to 42.3 g, and thousand kernel weight of some genotypes such as Momtchill, Bezostaya-I, Kutluk-94, Yüregir-89, Kinaci-97 and Marmara-86 were higher than the other genotypes (Table 2). Maddonni et al. [34] stated that genotypic difference might affect kernel biomass accumulation. Quality of wheat grains is likely to be reduced with the effect of climate in semi-arid cropping regions [35, 36]. Increasing environmental stress on wheat production associated with climate affects quality of wheat [37]. Increased temperature and water deficiency [25] limit the duration of the grain filling period and starch biosynthesis of grains [38, 39].

Genotypes were different from each other for test weight. Test weight values of bread wheat genotypes were between 73.7 (Kutluk-94) and 81.7 (Pehlivan) kg hl⁻¹. Average test weight of 79.6 kg hl⁻¹ was reported by Gangadharappa et al. [5]. Test weight was the highest at Bezostaya-1, Gonen-98, Ikizce, Lirasa, Cham-4, Orso and Marmara-86 genotypes. A higher test weight infers larger, higher quality grain, whereas low test weights are associated with either small or pinched kernels, or climatic damaged grain. Test weight is influenced by both genotype and environment [37].

Rharrabti et al. [12] reported that thousand kernel weight and test weight are greatly affected by climatic parameters, particularly high temperature during the final phase of grain filling. Water deficiency during grain growth, results lower test weights due to reduced accumulation rate.

*E-mail address: gulgunoktem@harran.edu.tr.



125

126 **Fig. 1. Thousand weight and test weight values of some bread wheat genotypes**
 127 **grown in South-eastern Anatolia region of Turkey. Vertical bars indicate standard**
 128 **errors of the mean.**

129 3.2. Protein Content

130 Protein content was the lowest for the Orso genotype (9.7%) while the highest value was
 131 found for Bezostaya-I (14.8%). The quality of wheat grain is dependent on the characteristics
 132 of starch and protein present. Variety composition is also one of the most important factors
 133 and it causes protein quality changes [7, 14]. Different levels of wheat kernel protein content
 134 values were reported 9.7-14.3% [40], 7.1-11.6% [41], 9.5% [5] and 14.9-21.54% [42]. Protein
 135 content of Kutluk-94, Dagdas-94, Altay-85, Harmankaya-99, Kirgiz-95, Lirasa and Aytin-98
 136 genotypes were higher than others (Fig. 2.), thus genotype had an effect on grain protein
 137 concentration. Genetic background is the most important factor for wheat protein quality and
 138 grain protein concentration [38]. The protein concentration is determined by the genetic
 139 background but is also influenced to a large extent by environmental factors such as rainfall
 140 and temperature [16, 18, 19].

141 The environment (climate, soil, agronomic practices, etc.) exerts a strong influence on the
 142 expression of the quality of different cultivars [8, 11]. Wheat kernel quality depends on
 143 precipitation amount in the rain fed conditions. Under rain-fed conditions the developing
 144 grains are frequently exposed to mild to severe stress at different stages of grain
 145 development. High temperatures and deficit water during grain filling period had a greater
 146 positive effect on grain protein composition [25]. Protein content increase under high
 147 temperature conditions [38]. The research area for this study, South-eastern Anatolia, is
 148 semi-arid region and characterized by warm winters, hot and dry summers with an
 149 inadequate and irregular rainfall distribution pattern.

150 An effective drought and hot climate in grain filling period, results high protein content in
 151 wheat grains under rainfed conditions (Table 1). Influences of the environment on protein
 152 content have been shown by other authors [14, 17, 20, 21]. Climatic factors such as

*E-mail address: gulgunoktem@harran.edu.tr.

temperature and amount of precipitation during the wheat growing terms have an important role for quality of kernel. Prior to anthesis, yield and grain protein content are influenced by effects of genetics, environment and other aspects of crop management [44]. But after anthesis, kernel growth is directly impacted by air temperature and water [25].

Protein ratio was high at the most of wheat genotypes in this study. It is seen climatic data from Table 1. that air temperature was high and precipitation was very low in the May month. Generally May month covers both milky and starch filling stages at wheat plant in the Harran Plain which is located in the southeast Anatolia region. In the semi-arid regions such as research area, air temperature increases suddenly and precipitation is very low in May month (Table 1) at the early starch filling period of kernel. High temperature and low water affects wheat plants negatively in this term. The duration of starch accumulation period ends in a short time due to high temperature and low water. Maturation begins at the most of the plants. Thus, plants mature more quickly at high temperature. Generally, the protein amount is stable in the milky stage, but the protein ratio can change according to the amount of starch filling in the kernel. If there is a decrease in the amount of starch in the kernel, the protein content percentage increases [25]. Frequently there is a negative relationship between grain yield and protein content [45]. Post-anthesis heat or drought may increase grain protein content but reduce yield because of their effects on starch production [22, 23, 45].




Fig. 2. Protein content and SDS-sedimentation values of some bread wheat genotypes grown in South-eastern Anatolia region of Turkey. Vertical bars indicate standard errors of the mean.

In the present study, the protein content was high due to low starch content in the kernel. Rao et al. [14] emphasis heat stress during the grain-filling stage influenced the protein content of wheat kernel.

Generally, a rise in temperature resulted in higher protein contents. Climatic factors significantly influence protein levels in wheat [8, 11]. Daniel and Triboi [46] stated that protein percent in wheat increased with the increase of air temperature. Topal et al. [47] reported that the protein content of the kernel increased with water stress. Mallikarjunaswamy et al. [48] reported that decreased irrigation water negatively affects the quality of kernel.

Table 2. Average thousand kernel weight, test weight, protein content, wet gluten, dry gluten and SDS-sedimentation values of some bread wheat genotypes grown in South-eastern Anatolia region of Turkey.

Genotypes	Thousand kernel weight** (g)	Test weight** (kg hl ⁻¹)	Protein content** (%)	Wet Gluten** (%)	Dry Gluten** (%)	SDS** (ml)
1. Kasifbey-95	30.5 m-p	78.5 efg	10.3 mn	30.1 n†	10.0 mn	21.0 r
2. Basribey-95	29.1 qrs	78.9 def	11.1 klm	32.4 m	10.8 j-m	23.5 p
3. Seyhan-95	29.4 q-s	79.7 cde	11.3 jkl	34.1 kl	11.0 i-m	26.0 n
4. Marmara-86	35.8 ef	80.2 bcd	13.2 b-g	38.9 cde	12.9 b-e	36.0 cde
5. Kirkpinar-79	34.6 gh	77.7 fgh	12.2 hij	35.3 ij	11.7 f-j	27.5 m
6. Ozdemirbey	28.6 rs	76.6 h	12.3 ghi	36.1 hi	11.9 e-i	33.0 g
7. Lirasa	31.2 lmn	80.5 abc	13.6 bcd	40.0 bc	13.2 abc	37.0 c
8. Cham-4	29.6 q-r	80.5 abc	12.6 d-i	39.6 cd	12.1 d-h	36.0 cde
9. Yuregir-89	36.8 de	79.7 cde	10.9 klm	30.8 n	10.4 klm	22.5 q
10. Atilla-12	30.4 m-p	80.1 bcd	12.5 e-i	35.2 ijk	11.9 e-i	27.5 m
11. Mohac-23	32.8 jk	77.7 fgh	11.8 ijk	33.4 lm	11.2 h-l	25.5 no
12. Momtchill	39.9 b	80.1 bcd	13.4 b-f	38.3 ef	12.7 cde	35.5 de
13. Pehlivan	42.3 a	81.7 a	13.5 b-e	38.2 ef	12.9 b-e	35.0 e
14. Kate A-1	30.2 n-q	79.7 cde	10.8 lm	30.6 n	10.3 lmn	22.0 q
15. Orso	29.7 o-r	81.3 ab	9.7 n	28.5 o	9.4 n	19.0 s
16. Fatima-2	30.2 n-q	79.7 cde	13.5 b-e	38.6 def	12.9 b-e	31.5 ij
17. Golia	29.1 qrs	78.9 def	12.1 hij	34.1 kl	11.3 g-l	28.0 m
18. Prostor	34.2 ghi	78.9 def	11.3 jkl	32.5 m	10.9 l-m	24.5 o
19. Kutluk-94	37.7 cd	73.7 i	14.2 ab	41.0 b	13.8 ab	38.0 b
20. Kirgiz-95	34.1 ghi	76.5 h	13.6 bcd	39.8 c	13.2 abc	36.5 cd
21. Aytin-98	33.4 hij	77.7 fgh	13.5 b-e	38.4 ef	12.9 b-e	34.0 f
22. Harmankaya-99	34.0 g-j	78.9 def	13.8 bc	39.1 cde	13.1 bcd	37.0 c
23. Sultan-95	25.8 t	74.1 i	13.1 c-h	36.9 gh	12.3 c-g	31.0 jk
24. Gonen-98	33.1 ij	80.7 abc	12.7 d-i	36.0 hi	11.6 f-j	30.5 k
25. Izmir-85	28.3 s	80.1 bcd	12.4 f-i	34.5 jkl	11.6 f-j	25.5 no
26. Dogankent-I	31.8 kl	78.9 def	13.3 b-f	37.5 fg	12.4 c-f	29.5 l
27. Kinaci-97	36.3 ef	77.7 fgh	10.9 klm	30.6 n	10.1 mn	22.0 q
28. Dagdas-94	33.6 hij	78.1 fg	14.0 abc	38.2 ef	13.0 bcd	29.5 l
29. Seri-82	31.5 im	78.9 def	12.2 g-j	33.8 l	11.4 f-k	25.0 no
30. Altay-85	31.1 lmn	77.3 gh	13.9 abc	39.9 bc	13.2 abc	32.5 gh
31. Ikizce	30.7 l-o	80.5 abc	13.4 b-f	38.8 cde	12.9 b-e	32.0 hi
32. Bezostaya-I	38.7 c	80.9 abc	14.8 a	42.2 a	14.1 a	39.0 a
33. Suzen-97	35.2 fg	80.1 bcd	10.9 klm	35.8 hi	10.5 klm	27.5 m
Average	37.7	78.9	12.5	36.0	11.9	29.7

† : There are no statistical differences among the genotypes in the same column having the same letter at 0.05 level according to Duncan test.

** : Denotes significant difference among genotypes $P \leq 0.01$.

3.3. Wet and Dry Gluten

Gluten is the major component of flour protein that determines processing quality. Wet gluten reflects the gluten quality and quantity. The highest wet gluten content was found in Bezostaya-I (42.2%), while Orso genotype gave the lowest value (28.5%) (Table 2). Kutluk-94, Lirasa, Altay-85, Kirgiz-95, Cham-4, Harmankaya-99, Marmara-86 and Ikizce genotypes recorded higher wet gluten value than the other genotypes. Pasha et al. [6] reported 13.82-43.13% wet gluten content values.

Dry gluten values varied from 9.4% (Orso) to 14.1% (Bezostaya-I). It was shown that the content of dry gluten of Kutluk-94, Altay-85, Lirasa, Kirgiz-95, Harmankaya-99 and Dagdas-94 genotypes were slightly higher than the other genotypes (Fig. 3.). The present findings are in collaboration with the previous studies conducted by Curic et al. [49] who reported the range of dry gluten from 8.44 to 11.77% in flours of different wheat varieties, and Lin et al. [50] found the range of dry gluten from 7.0 to 16.7%.

Gangadharappa et al. [5] stated that dry gluten values were in the range of 7.93-9.60%. Indrani et al. [51] reported 10.3% dry gluten value, and Pasha et al. [6] reported 4.46 - 14.55% dry gluten values. Gluten amount in wheat grain may affected by genetic and climatic factors [6, 52, 53].




Fig. 3. Dry gluten and wet gluten values of some bread wheat genotypes grown in South-eastern Anatolia region of Turkey. Vertical bars indicate standard errors of the mean.

The contents of protein and dry gluten reflect the quality of wheat varieties [52]. Gluten amount in the kernel is firmly related with protein amount in the kernel. Protein ratio in the kernel effects positively wet and dry gluten amount. Gluten amount in the kernel effects dough and bread quality. In the present study, gluten values were found high

correspondingly to protein content. Hence, effect of high temperature and low water in grain filling stages in semi-arid region result high protein content and gluten values. Faergestad et al. [24] emphasis climatic conditions affect kernel quality, protein and gluten composition of wheat kernel.

3.4. SDS-Sedimentation Value

SDS-sedimentation values of bread wheat genotypes ranged between 19.0 ml (Orso) and 39.0 ml (Bezostaya-I) (Table 2). The highest SDS-sedimentation values were found in Kutluk-94, Lirasa, Kirgiz-95, Harmankaya-99, Cham-4, Marmara-86 and Momtchill genotypes (Fig. 2.). Sedimentation value reflects the quality of protein [28]. Pasha et al. [6] reported 19.67-36 mL SDS-sedimentation volume values. Gangadharappa et al. [5] measured a 46 ml SDS-sedimentation value in wheat. Tonk et al. [42] reported higher SDS-sedimentation values of 46-95 ml in wheat.

The quality of the wheat kernel is affected by both variety and environment [10]. Balkan and Genctan [53] stated that SDS values should be between 30 and 43 ml. SDS values were lower than expected in the study. All of varieties gave lower SDS value than 40 ml. Most of SDS values were below 30 ml. SDS values can be reduced in dry and hot environments [20, 54]. SDS values increases with increasing temperature during grain filling up to about 30°C and then decreases as temperatures rise above 30 °C [20, 55]. Temperature during grain-filling period was higher than 30 °C in the present study (Table 1). Thus, it appears that increasing protein content due to high temperature and low water input during the grain filling period could lead to a decrease in SDS value under the conditions of our study. Water input during grain filling also had a negative influence on SDS volume [12].

3.5. Correlation Coefficients

Relationships between quality traits have been investigated in **most** studies on bread wheat [7, 24, 56]. Correlation coefficients for some quality parameters are given in Table 3. According to correlation analysis; a positive significant correlation was found between thousand kernel weight and SDS-sedimentation value (0.347*). Protein content was positive correlated with wet gluten (0.941**), dry gluten (0.986**) and SDS-sedimentation value (0.888**) at the $P \leq 0.01$ level, respectively. Some researchers reported a correlation between protein and wet gluten [9, 21, 40, 57, 58]. A positive correlation between protein and dry gluten value is emphasized by Anjum and Walker [59]. An inverse relationship between protein content and SDS volume was reported by Rharrabti et al. [12].

Positive correlations between wet gluten and both dry gluten (0.960**) and SDS-sedimentation value (0.956**) were great and significant at level of 1%. The **significant correlation in positive direction** between SDS-sedimentation value wet gluten content was reported by Pasha et al. [6] and Ozturk and Aydin [9]. There was a positive significant correlation between dry gluten and SDS-sedimentation value (0.920**) at the 1% level. Pasha et al. [6] emphasized a positive significant correlation of SDS-sedimentation value with dry gluten values.

*E-mail address: gulgunoktem@harran.edu.tr.

Table 3. Correlation coefficients among 1000 kernel weight, test weight, protein content, wet gluten, dry gluten and SDS-sedimentation values.

Traits	1000 kernel weight	Test weight	Protein content	Wet gluten	Dry gluten	SDS-sedimentation value
1000 kernel weight	1	0.158	0.289	0.278	0.298	0.347*
Test weight	-	1	-0.173	-0.119	-0.178	-0.110
Protein content	-	-	1	0.941**	0.986**	0.888**
Wet gluten	-	-	-	1	0.960**	0.956**
Dry gluten	-	-	-	-	1	0.920**

*: $P \leq 0.05$, **: $P \leq 0.01$.

4. CONCLUSION

The data obtained from our study indicate that quality parameter values of bread wheat genotypes were different from each other. Bezostaya-I, Kutluk-94, Lirasa, Altay-85, Kirgiz-95, Cham-4, Harmankaya-99, Marmara-86, Ikizce, Pehlivan, Momtchill, Fatima-2, Dagdas-94 and Aytin-98 genotypes were the best in quality among the tested other genotypes in semi-arid region. Differences in quality of bread wheat could be associated with differences in adaptation ability of genotypes, genotypic structure and reacted differently to soil and climate conditions. Climatic conditions during grain filling appear to be crucially important in determining grain quality in semi-arid environments. Protein content, wet and dry gluten and SDS-sedimentation values were affected by climatic factors in the semi-arid region. Protein content, wet and dry gluten values were high but SDS sedimentation values were low due to high temperature and low precipitation in semi-arid region. Although the hot and dry conditions of semi-arid region cause a large fluctuation in yield, they often provide the opportunity for a good expression of quality parameters such as high protein and gluten values.

A positive significant correlation was found between thousand kernel weight and SDS-sedimentation value. Protein content was positively correlated with wet gluten, dry gluten and SDS-sedimentation, respectively. Relationship between wet gluten and both dry gluten and SDS-sedimentation value were great and significant. There was a positive significant correlation between dry gluten and SDS-sedimentation value.

REFERENCES

1. Anonymous. Statistical values in cereals. The Turkish Statistic Institute, 2017, Ankara.
2. Troccoli A, Borrelli GM, Vita P, Fares C, Fonzo N. Durum wheat quality: a multidisciplinary concept (review). Journal of Cereal Science 2000;32:99-113.
3. Abaye AO, Brann DE, Alley MM, Griffey CA. Winter durum wheat: do we have all answers? College of Agriculture and life sciences, Virginia Cooperative extension. Crop and Soil Environmental Sciences Pub. Num. 424–802. Virginia Polytechnic Institute and Virginia State University, Virginia, USA. 1997.
4. Lee M, Lerohl M, Unterschultz J. Buyer preference for durum wheat: a stated preference approach. International Food and Agribusiness Management Review 20003;353-366.

*E-mail address: gulgunoktem@harran.edu.tr.

5. Gangadharappa GH, Ramakrishna R, Prabhasankar P. Chemical and scanning electron microscopic studies of wheat whole-meal and its streams from roller flour mill. *Journal of Food Engineering* 2008;85:366-371.
6. Pasha I, Anjum FM, Butt MS, Sultan JI. Gluten quality prediction and correlation studies in spring wheat. *Journal of Food Quality* 2007;30:438-449.
7. Peterson CJ, Graybosch PS, Baenziger PS, Grombacher AW. Genotype and environment effects on quality characteristics of hard red winter wheat. *Crop Sci.* 1992;32:98-103.
8. Blumenthal CS, Bekes F, Gras PW, Barlow EWR, Wrigley CW. Identification of wheat genotypes tolerant to the effects of heat stress on grain quality. *Cereal Chem.* 1995;72(6):539-544.
9. Ozturk A, Aydin F. Effect of water stress at various growth stages on some quality characteristics of winter wheat. *J. Agronomy & Crop Science* 2004;190:93-99.
10. Svihus B, Gullord M. Effect of chemical content and physical characteristics on nutritional value of wheat, barley and oats for poultry. *Animal Feed Science and Technology* 2002;102:71-92.
11. Randall PJ, Moss HJ. Some effects of temperature regime during grain filling on wheat quality. *Aust. J. Agric. Res.* 1990;41:603-617.
12. Rharrabti Y, Villegas D, Royo C, Martos-Nunez V, Moral GLF. Durum wheat quality in Mediterranean environments II. Influence of climatic variables and relationships between quality parameters. *Field Crops Research* 2003;80:133-140.
13. Campbell CA, Davidson H.R, Winkelman GE. Effect of nitrogen, temperature, growth stage and duration of moisture stress on yield components and protein content of Manitou spring wheat. *Can. J. Plant Sci.* 1981;61:549-563.
14. Rao ACS, Smith JL, Jandhyala VK, Papendick RI, Parr JF. Cultivar and climatic effects on the protein content of soft white winter wheat. *Agronomy Journal* 1992;85(5):1023-1028.
15. Uhlen KA, Hafskjold R, Kalhovd AH, Sahlstrom S, Longva A, Magnus EM. Effects of cultivars and temperature during grain filling on wheat protein content, composition, and dough mixing properties. *Cereal Chem.* 1998;75:460-465.
16. Crosbie GB, Fisher H. Variation in wheat protein content - the effect of environment. *Journal of Agriculture of Western Australia* 1987;28:124-127.
17. Naes T. The design of calibration in near infra-red reflectance analysis by clustering. *J Chemometrics* 1987;1:121-134.
18. Stapper M, Fischer RA. Genotype, sowing date and plant spacing influence on high-yielding irrigated wheat in southern New South Wales. II. Growth, yield and nitrogen use. *Aust J Agric Res.* 1990;41:1021-1041.
19. McDonald GK. Effects of nitrogenous fertilizer on the growth, grain yield and grain protein concentration of wheat. *Aust J Agric Res.* 1992;43:949-967.
20. Graybosch RA, Peterson CJ, Baenziger PS, Shelton DR. Environmental modification of hard red winter wheat flour protein composition. *J Cereal Sci.* 1995;22:45-51.
21. Johansson E, Nilsson H, Mazhar H, Skerriitt J, MacRitchie F, Svensson G. Seasonal effects on storage proteins and gluten strength in four Swedish wheat cultivars. *Sci. Food Agric.* 2002;82:1305-1311.
22. Bhullar SS, Jenner CF. Differential responses to high temperatures of starch and nitrogen accumulation in the grain of four cultivars of wheat. *Aust. J. Plant Physiol.* 1985;12:363-375.
23. Cassman KG, Bryant DC, Fulton AE, Jackson LF. Nitrogen supply effects on partitioning of dry matter and nitrogen to grain of irrigated wheat. *Crop Sci.* 1992;32:1251-1258.
24. Faergestad EM, Flaete NES, Magnus EM, Hollung K, Martens H, Uhlen AK. Relationships between storage protein composition, protein content, growing season and flour quality of bread wheat. *J Sci. Food Agric.* 2004;84:877-886.

25. Oktem A. Effect of water shortage on yield, and protein and mineral compositions of drip-irrigated sweet corn in sustainable agricultural systems. *Agricultural Water Management* 2008;95(9):1003-1010.
26. Bremner JM. Determination of nitrogen in soil by the Kjeldahl method. *J. Agric. Sci.* 1960;55:11–33.
27. Anonymous. Determination of crude protein in cereals and cereal products for food and for feed. Standard methods of the international association for cereal science and technology (ICC), ICC Standard No:105/2, 2002a, Vienna.
28. Pomeranz Y. Wheat II, In: *Wheat chemistry and technology* St. Paul, MN, USA: American of Association of Cereal Chemists 1998;1:17-21.
29. Anonymous. Determination of sedimentation value. SDS test of durum wheat. ICC Standard No:151, 2002b, Vienna.
30. Anonymous. Determination of wet and dry gluten in durum wheat. ICC Standard No:137, 2002c, Vienna.
31. Snedecor GW, Cochran WG. *Statistical Methods*, Eighth Edition. Iowa State University, Ames, Iowa. 1989.
32. Tallarida RJ, Murray RB. Duncan Multiple Range Test. In: *Manual of Pharmacologic Calculations*. Springer, New York, NY. 1987.
33. Kendall MG, Gibbons JD. *Rank Correlation Methods* (5th ed.). London: Edward Arnold. 1990.
34. Maddonni GA, Otegui ME, Bonhomme R. Grain yield components in maize: II. Postsilking crop growth and kernel weight. *Field Crops Res.* 1998;56(3):257- 264.
35. Asseng S, Foster I, Turner NC. The impact of temperature variability on wheat yields. *Global Change Biol.* 2011;17:997–1012.
36. Panozzo J, Walker CK, Partington DL, Neumann NC, Tausz M, Seneweera S, Fitzgerald GL. Elevated carbon dioxide changes grain protein concentration and composition and compromises baking quality. A FACE study. *J. Cereal Sci.* 2014;60 (3):461–470.
37. Nuttall JG, Leary GJO, Panozzo JF, Walker CK, Barlow KM, Fitzgerald GJ. Models of grain quality in wheat. *Field Crops Research* 2017;202:136-145.
38. Farooq M, Bramley H, Palta JA, Siddique KHM. Heat stress in wheat during reproductive and grain-filling phases. *Crit. Rev. Plant Sci.* 2011;30:1–17.
39. Ferreira MSL, Martre P, Mangavel C, Girousse C, Rosa NN, Samson M, Morel M. Physicochemical control of durum wheat grain filling and glutenin polymer assembly under different temeprture regimes. *J. Cereal Sci.* 2012;56:58–66.
40. Pena RJ, Zarco-Hernandez J, Mujeeb-Kazi A. Glutenin subunit compositions and bread making quality characteristics hexaploid wheat derived from *Triticum turgidum* x *Triticum tauschii* (coss.) schmal crosses. *J. Cereal Sci.* 1995;21:15-23.
41. Kindred DR, Gooding MJ, Ellis RH. Nitrogen fertiliser and seed rate effects on Hagberg falling number of wheat hybrids and their parents are associated with alpha-amylase activity, grain cavity size and dormancy. *Journal of the Science of Food and Agriculture* 2005;85(5):727-742.
42. Tonk FA, Ilker E, Tosun M. A study to incorporate high protein content from tetraploid wheat (*T. turgidum dicoccoides*) to hexaploid wheat (*T. aestivum vulgare*). *Turkish Journal of Field Crops* 2010;15(1):69-72.
43. DuPont FM, Altenbach S.B. Molecular and biochemical impacts of environmental factors on wheat grain development and protein synthesis. *Journal of Cereal Science* 2003;38:133-146.
44. Bahrman N, Gouis J, Negroni L, Amilhat L, Leroy P, Laine AL, Jaminon O. Differential protein expression assessed by two dimensional gel electrophoresis for two wheat varieties grown at four nitrogen levels. *Proteomics* 2004;4:709–719.
45. Fowler DB. Crop nitrogen demand and grain protein concentration of spring and winter wheat. *Agron. J.* 2003;95:260–265.

46. Daniel C, Triboi E. Effects of temperature and nitrogen nutrition on the grain composition of winter wheat: Effects on Gliadin Content and Composition. *Journal of Cereal Science* 2000;32:45–56.
47. Topal A, Yalvac K, Akgun N. Efficiency of top dresses nitrogen sources and application times in fallow-wheat cropping system. *Commun. In. Soil Sci. and Plant Anal.* 2003;34(9–10):1211–1224.
48. Mallikarjunaswamy SN, Ramachandrappa BK, Nanjappa HV. Effect of scheduling irrigation at different phenophases of sweet corn (*Zea mays saccharata*) and delineation of critical stages based on stress day index. *J. Agron. Crop Sci.* 1999;182:161–166.
49. Curic D. Gluten as a standard of wheat flour quality. *Food Technol. Biotechnol.* 2001;39(4):353–361.
50. Lin P, Chiang SH, Chang CY. Comparison of rheological properties of dough prepared with different wheat flours. *J. Food Drug Anal.* 2003;11(3):220–225.
51. Indrani D, Prabhasankar P, Rajiv J, Venkateswara RG. Influence of whey protein concentrate on the rheological characteristics of dough, microstructure and quality of unleavened flat bread (Parotta). *Food Research International* 2007;40:1254-1260.
52. Zhang L, Zhang Y, Song Q, Zhao H, Yu H, Zhang C, Xin W, Mao Z. Study on the quality of NILS of wheat cv. Long fumai 3 possessing HMWGS Null and 1 Subunits. *Agricultural Sciences in China* 2008;7(2):140-147.
53. Balkan A, Genctan T. Determination of yield and quality components of some bread wheat cultivars, which are added to mixed wheat for increasing flour quality in Tekirdag Conditions. *Turkey VI. Field Crops Congress* 1: 149-154, 5-9 September 2005, Antalya, Turkey.
54. Blumenthal CS, Bekes F, Batey IL, Wrigley CW, Moss HJ, Mares DJ, Barlow EWR. Interpretation of grain quality results from wheat variety trials with reference to high temperature stress. *Aust. J. Agric. Res.* 1991;42:325–334.
55. Martens H, Martens M. Modified jack-knife estimation of parameter uncertainty in bilinear modelling by partial least squares regression (PLSR). *Food Quality and Preference* 2000;11:5–16.
56. Matsuo RR, Dexter JE. Relationship between some durum wheat physical characteristics and semolina milling properties. *Can. J. Plant. Sci.* 1980;60:49–53.
57. Payne LM, Holt AF, Carrillo JM. Relationships between seed quality characteristics and hmw glutenin subunit composition determined using wheat grown in Spain. *Journal of Cereal Science* 1988;7:229-235.
58. Basset LM, Allan RE, Rubenthaler GL. Genotype x environment interactions on soft white winter wheat quality. *Agron J.* 1989;81:955-960.
59. Anjum FM, Walker CE. Electrophoretic identification of hard white spring wheats grown at different location in Pakistan in different years. *J. Sci. Food Agric.* 2000;80:1155–1161.