

LONG-TERM CONSUMPTION OF KOLA-NUT (*COLA NITIDA*) DIET DOES NOT INCREASE ANXIETY RELATED BEHAVIOUR IN MICE.

ABSTRACT

Following long-term consumption of kola nut (*Cola nitida*) diet, anxiety related behaviour were studied in 16 Swiss white mice (18-28g body weight). The open field (OF) test, elevated plus maze (EPM) and the light/dark transition box (LD) tests were used. Swiss white mice were fed either control diet (rodent chow; n=8) or kola nut diet (50% w/w kola-nut diet; n=8) for 28 days. All animals were allowed free access to clean drinking water. Results showed that the frequency of rearing in the kola nut diet group was lower ($p < 0.05$) compared to control. The non-exploratory behaviours like grooming and genital licking were also lower in the test group compared to control ($p < 0.001$, 0.05 respectively). In the EPM test, the duration in the open arm in the kola diet group was higher compared to control ($p < 0.01$). The duration of grooming in the test group was however higher in the closed arm compared with control ($p < 0.01$). The frequency of downward dips only correlated positively with the duration in the open arm in the control [$r(16) = 0.855$; $p < 0.01$]. The kola fed animals spent more time in the light region of the LD test ($p < 0.01$) rearing and walling ($p < 0.05$), and spent less time in the dark region when compared with their control. In conclusion, long-term consumption of kola nut diet decreased anxiety-related behaviour in the mice.

Key words: Kola nut, anxiety, Swiss white mice

INTRODUCTION

Kola is a tropical tree crop (family - *sterculiaceae*) which has socio-economic importance in Nigeria. Kola nut is used traditionally for ceremonies related to marriages, child naming, funerals and in other festivals and also chewed as a masticant¹. It is commercially grown in the West where it is known as "Obi" in Yoruba, consumed by the Northerners where it is known as "Goro" in Hausa, and revered in the East where it is called "Oji" in Igbo. In Cross River and Akwa Ibom it is called "Ibong" in Efik language. It is of great importance in the traditional institution. In fact, the Igbos in Nigeria liken kola nut as a fruit that brings with it good fortunes and life.

Whichever way the nuts are consumed (chewed raw or used in powdered form), the nuts produce a mild stimulates effect on the central nervous system and thus producing a tentative feeling of increased energy and reduction of hunger and fatigue². Fresh and cured kola nut chewed in small doses increase mental activity, reduce the need to sleep and also dispel hunger and thirst¹. Therefore, kola nut chewing has become very popular among students, drivers and many other consumers who need to remain active for unusually long periods.

In some developed countries, however, kola nut extracts are used industrially for the manufacturing of many cola-type soft drinks flavours³, as a source of caffeine for manufacture of many pharmaceutical products and essential oils⁴, and as a main ingredient in production of heat-tolerant chocolate bars⁵. In addition, caffeine is known to be a fat burner and therefore beneficial in assisting weight loss⁶. As a result of the commercial importance of kola nuts, a lot of research work has been done on *Cola nitida*, the kola of commerce, in Nigeria⁷. Presently, the bulk of kola nuts being produced in Nigeria are either consumed fresh locally or exported as sun-dried to some drier areas of Africa, where they are used as masticant or as sources of colourant for cloth dyeing but with little or no industrial use locally⁸.

There are over forty kola species but the most common species with major economic importance in Nigeria are *Cola nitida* and *Cola accuminata*⁹. Phytochemistry of *Cola nitida* show that it has 9.73-9.81% water, 2.72-2.21% ash, 3.02-2.20% fat, 19.14-15.24% protein, 7.30-4.18% crude fibre and 58.09 to 66.45% carbohydrate. *C. nitida* also has alkaloids (2.22%), tannin (6.46%) and saponin (8.06%), phenol (0.27%), flavonoid (0.12-0.14%)¹⁰. Caffeine is one of the most important alkaloids of *Cola nitida* and it forms 2.4% of its content¹¹.

Caffeine, a major alkaloid of *Cola nitida*, has been reported in some studies in mice to increase anxiety¹² while in other instances was reported to decreased anxiety¹³. In a human study on the effect of caffeine consumption among secondary school children in South West England, caffeine caused increased anxiety related behaviour¹⁴.

Since kola nut contains caffeine, and there are still controversies as to whether chronic use of caffeine reduces or increases anxiety, this study saw the need to investigate the effect long-term consumption of kola nut on anxiety-related behaviour. Given the chronic consumption of kola nuts among locals in Nigeria, we studied the effect of long-term consumption of kola nut on locomotor activity and anxiety-related behaviour in CD1 mice using the open field test, elevated plus maze test and light dark transition test.

MATERIALS AND METHODS

Male Swiss white mice purchased from the animal house of the Department of Physiology, University of Calabar. Mice were kept in standard well ventilated animal facility in the Department of Physiology at temperature of 26±2°C, 12/12 light/dark cycle. Mice were grouped into two: Control (n=8) given normal rodent chow (Vital feed Nigeria), and test (n=8) given 50% w/w kola-nut diet. Kola-nut diet was prepared by slicing, drying and grinding fresh kola-nut (*Cola nitida*) bought from Bogobiri (Hausa Market) in Calabar, Nigeria. Equal portions of the grinded kola-nut powder (10g) and rodent chow (10g) were used to constitute 50% w/w kola-nut diet¹⁵. All animals had access to clean drinking water and food *ad libitum*. This feeding was done for 28 days. All animals were weighed before and after the feeding period.

Internationally acceptable ethics for laboratory animal used were strictly adhered to. Ethical approval was duly obtained from the Faculty of Basic Medical Sciences Animal Ethics Committee with Protocol number 014PHS015007

The **open field apparatus** test used by Bisong *et al* (2006, 2018) was employed in this study. The test apparatus measured 72 x 72 x 32cm (l x b x h) with a floor divided into sixteen 18 x 18cm squares, and centre square of 36 x 36cm. Mice were allowed 5 minutes to explore the apparatus while behaviour are scored. Thirty minutes later, mice were tested for another 5 minutes¹⁶.

99

100 The **elevated plus maze** as described by Lister¹⁷ and used by Bisong *et al*^{18,19} was
101 employed in this study. The apparatus has 2 open arms (each measuring
102 30x5x15cm) and two closed arms (each measuring 30x5x15cm) extending from a
103 central, open square (5x5cm). The maze was elevated on a pedestal to a height of
104 45cm above the floor. Each mouse was placed in the centre square of the elevated
105 plus maze facing an open arm and its behaviour scored.

106

107 The **light/dark transition box** test as described by Hascoët and Bourini²⁰ and
108 used by Bisong *et al*¹⁸ was used. It is box measuring 46x30x27cm high (l x b x h
109 cm), divided into two compartments; a small 18x30cm area and a large 27x30cm
110 area with a 7x7cm door on floor of the partition linking the two chambers. The small
111 compartment was painted black to mimic darkness, whereas the large compartment
112 was painted white. Mice were given 5 minutes to explore the apparatus during which
113 behaviours were scored.

114

115 **Statistical Analysis**

116 Data obtained from the study were analysed using the Student T- test.
117 Associations between data were tested using the Pearson's correlation. Data were
118 presented as means \pm standard error of mean. Probability level of $P < 0.05$ was
119 accepted as significant.

120

121

122 **RESULTS**

123

124 ***Exploratory behaviour in the open field test***


125 The Exploratory behaviour, Rearing and walling, are forms of vertical
126 locomotor activity. The frequency of Rearing in the test group within the first five
127 minutes was $3.6 \pm 1.1/5\text{min}$, significantly lower when compared with control ($5.9 \pm$
128 $1.5/5\text{min}$; $P < 0.05$). At the end of 30 minutes, the frequency of rearing in the kola
129 group ($43.4 \pm 11.1/30\text{min}$) was also lower than that in control ($76.4 \pm 11.2/30\text{min}$; $P <$
130 0.05). Although the frequency of walling in the kola group seemed lower both at 5
131 minutes ($16.4 \pm 3.6/5\text{min}$) and 30 minutes ($58.3 \pm 14.0/30\text{min}$) when compared with
132 control (19.6 ± 2.3 at 5 minutes and 80.4 ± 10.7 at 30minutes; Figure 1) **it did not**
133 **differ significantly.**

134

135

136

137
138



157 * - Significant at P < 0.05 compared to control

158

159 Fig. 1: Exploratory behaviour; rearing (a) and walling (b) of mice fed kola diet in the
160 open field test.

161


162

163 **Grooming in the open fields test**


164

165 The frequency and duration of grooming in the open field apparatus were
166 significantly lower in the test group than in the control at the end of the 30 minutes
167 session. This was $16.4 \pm 2.8/30\text{min}$ in the kola group, which was less than that in the
168 control group of animals, which was 30.1 ± 2.9 ; ($P < 0.01$). The duration of the
169 grooming at the end of 30minutes ($192.5 \pm 25.4 \text{ sec}$) was also significantly lower in
170 the kola fed group compared to control ($352 \pm 54 \text{ sec}$; $P < 0.05$). This is shown in
171 Figure 2.

171



172



(b)

NS – Not significant compared to control
 * - Significant at $P < 0.05$ compared to control.

Fig. 2: Frequency (a) and Duration (b) of grooming in mice fed kola diet in five minutes and thirty minutes in the open field test.

Comparison of activities in the Elevated plus maze between kola fed and control mice.

The frequency of entry into the open arm was not significantly different between control and kola fed mice. The duration in the open arm was however, significantly higher in the test group compared to control ($p < 0.01$). The frequency and duration of downward dipping of mice did not differ significantly between test and control groups of mice. The frequency of entry into the closed arm of the elevated plus maze in the kola fed group was significantly lower compared to control ($P < 0.05$). The duration in the closed arm was also lower in the test group compared to control ($P < 0.05$). Although the frequency of grooming in the closed arm was not significantly different between test and control groups, the duration differed ($P < 0.01$). The frequency of rearing and the number of faecal boli at the end of the 5minutes session did not differ from control values. Table 1 below shows a summary of these comparisons.

198
199
200
201
202

Table 1: Summary of comparison of activities in the Elevated plus maze between kola fed and control mice.

Parameters	Control group	Kola fed group	Level of significance
Frequency of open arm entry (/5minutes)	2.8 ± 0.3	2.6 ± 0.6	NS
Duration in open arm (seconds)	45.3 ± 10.4	136 ± 18.7	0.01
Frequency of downward head dipping (/5minutes)	11.4 ± 2.9	12 ± 2.2	NS
Duration in closed arm (seconds)	218.1 ± 15.2	158.9 ± 25.5	0.05
Frequency of grooming in the closed arm (/5minutes)	3.3 ± 0.8	4 ± 0.5	NS
Duration of grooming in the closed arm (Seconds)	18.3 ± 5.7	44.9 ± 9.3	0.01
Frequency of genital licking in the closed arm (/5 minutes)	1.3 ± 0.4	2.3 ± 0.5	0.05
Frequency of rearing in the closed arm	19 ± 2.0	20 ± 2.1	NS
Number of faecal boli	0.4 ± 0.3	0.3 ± 0.2	NS

203 NS – Not significantly different compared to control.

204
205
206
207
208
209
210
211
212
213
214
215

Correlation between duration in the open arm and frequency of downward head dipping

There was a positive correlation between the duration in the open arm of the elevated plus maze and the frequency of downward head dipping in the control group [$r(16) = 0.855$; $p < 0.01$], Figure 3.




Fig. 3: Correlation between frequency of downward head dipping and duration in the open arm in the control group

The Effect of long-term feeding with kola diet on Activities in the light region of the light/ dark transition box.

Activities in the light region were generally higher for the kola fed group compared to control. Although the frequency of entry into the light region did not differ, the time spent (duration) in the light region was significantly higher in the test group (111.5 ± 9.43 seconds) compared to control (77.63 ± 12.7 seconds; $p < 0.05$). The frequency of line crossing did not differ between the test and control groups. The frequency of rearing and walling were higher in the test groups compared to control ($p < 0.01$; $p < 0.05$ respectively), Table 2.

Table 2: Activities in the light region of the light/ dark transition box.

Activity in the light region.	Control group	Kola group	fed	Level of significance
Frequency of entry (/5mins)	6.75 ± 0.90	6.0 ± 0.95		NS
Duration in light region (Sec)	77.63 ± 12.7	111.5 ± 9.43		0.05
Line crossing (/5mins)	59.0 ± 10.29	51.3 ± 6.43		NS
Frequency of rearing (/5mins)	3.13 ± 0.82	9.25 ± 1.65		0.01
Frequency of walling (/5mins)	7.75 ± 1.75	14.88 ± 2.76		0.05

NS – not significant compared to control

The Effect of chronic feeding with kola diet on Activities in the dark region of the light/ dark transition box.

Although the test group showed less activity in the dark region of the light/dark transition box, the exploratory activities were significantly higher compared to control. The frequency of entry and the time spent (duration) in the dark region did not differ in the test group when compared to control. Frequency of rearing in the test group was higher than the value for the control group ($p < 0.05$). Walling also followed a similar trend with the test group of animals walling more than control ($p < 0.05$). The frequency of grooming did not differ significantly but the duration of grooming was lower in the test group compared to control ($p < 0.05$). The frequency of genital licking in the test group of mice was also significantly lower compared to that in the control group ($p < 0.05$); Table 3.

Table 3: Activities in the dark region of the light/ dark transition box.

Activity in the dark region.	Control group	Kola fed group	Level of significance
Frequency of entry (/5mins)	7.75 \pm 0.90	5.13 \pm 0.83	NS
Duration in dark region (Sec)	222.5 \pm 12.37	197 \pm 13.97	NS
Frequency of rearing (/5mins)	8.5 \pm 1.56	14.0 \pm 3.1	0.05
Frequency of walling (/5mins)	12.88 \pm 2.53	20.4 \pm 2.74	0.05
Frequency of grooming (/5mins)	6.38 \pm 6.0	6.0 \pm 1.12	NS
Duration of grooming (sec)	63.88 \pm 13.73	32.6 \pm 6.43	0.05
Frequency of genital licking (/5mins)	4.5 \pm 1.09	2.0 \pm 0.61	0.05

NS – not significant compared to control

DISCUSSION AND CONCLUSION

The open field apparatus has been used to assess the emotionality of animals in a novel environment, as well as locomotion and exploration²¹. Although the frequency of line crossing did not differ significantly between the test and control groups, other forms of locomotor activity differed. This implies that there was no significant change in horizontal locomotor patterns.

The frequency of rearing was significantly lower in the kola fed group (test) when compared to control in the open field. This trend was also similar in the frequency of walling. The vertical locomotor activities and therefore exploratory activity **was** decreased following long-term ingestion of kola nut. The frequency and duration of grooming in the kola fed group **were** also significantly decreased at the end of 30 minutes. This implies decreased vertical locomotor (exploratory) activities following chronic ingestion of kola nut. Therefore long-term ingestion of 50%w/w kola nut diet decreased exploratory activity. It is most likely that consumption of large

quantities of kola nut will not serve the stimulant effect any longer but rather cause depression of the nervous system. These results however, do not support the report of previous researchers that kola nuts serve as a stimulant^{1,2}. The decrease in exploratory/vertical locomotor activity following long-term consumption of kola is in consonance with reports of Neil²² which showed that excessive consumption of caffeine caused mixed depressive states in psychiatric patients; and also the work of Greden²³ depressive syndrome as being associated with caffeine, which is one of the major constituents of kola nut.

The elevated plus maze has been proven as a model for assessing anxiety and fear^{16,17}. This test is based on the natural aversion of rodents for open space and heights. Mice fed 50%w/w kola diet when compared with control, spent more time in the open arm and less time in the closed arm. Since the open arm is the aversive arm, drugs that reduce anxiety would thus cause the animals to spend more time in the open arms of the maze. Therefore, that the kola-fed mice were less fearful compared to their control. To buttress these, there was a positive correlation between the duration in open arm and the frequency of downward head dipping, as mice which are less fearful would perform more head dips. Therefore, the kola diet reduced anxiety in the mice.

The light/dark transition box is also used as a model for assessing anxiety and fear. This light/dark test is based on the innate aversion of rodents for brightly illuminated areas and on the spontaneous exploratory behaviour of rodents in response to mild stressors that is novel environment and light²⁴. Mice which are less anxious would spend more time in the open space and brightly lit chamber. In this test, the kola fed group of mice spent more time in the illuminated (light) region of the box and also showed more activity (rearing and walling) in this region. This implies that long-term feeding of mice with kola diet produces an anxiolytic effect.

The kola fed group showed less grooming (non-exploratory behaviours associated with fear) in the dark region compared to their control. The frequency of the exploratory behaviours, rearing and walling, were higher in the test group. This is in agreement with findings of Costal *et al*²⁵ that increased exploratory behaviours was associated with an increase in the time spent in the light region of the light/ dark box. Grooming is a displacement reaction which happens when mice are anxious. Thus increased grooming behaviour indicate increased anxiety.

The mice in the test group were less fearful compared to control, which is in consonance with the test in the elevated plus maze. Therefore, long-term ingestion kola diet caused an anxiolytic effect. These results were contrary to earlier works which implicate caffeine (a major component of kola) as an anxiogenic agent^{26,27}.

In conclusion, long-term consumption of 50%w/w kola diet in mice caused decreased exploratory activity, instead of producing a stimulant effect, in mice. The kola diet also produced an anxiolytic effect, thereby reducing fear and anxiety in mice.

Ethical approval

Internationally acceptable ethics for laboratory animal used were strictly adhered to. Ethical approval was duly obtained from the Faculty of Basic Medical Sciences Animal Ethics Committee with Protocol number 014PHS015007

Consent: NA

REFERENCES

1. Ogutuga, D.B.A. (1975): Chemical composition and potential commercial uses of Kola nuts, *Cola nitida* vent cachott and Endlisher. Ghana J-Agric Sci. 8, 121-125
2. Martin, K.L; Morelli H. F, Schild, H.O; State land, B.E (1983): In Clinical Pharmacology Basic Principles in Therapeutics, 2nd (Ed.) Macmillan Publishing Co. Inc. New York p. 663
3. Beattle, G.B (1970): Soft drink flavours. Their History and Characteristics. 1 Cola or "Kola" flavours the flavour Industry pp. 390 – 394
4. Olounloyo, O.A (1979): Fungi Associated with deterioration of high kola nuts. Nigerian Journal of Agric Science (1) 52-59
5. Williams, S.O (1979): Project of kola chocolate processing and consumption CRIN, (Seminar paper 4).
6. Blades, M (2000): Functional foods or Nutraceutical Nutrition and food science vol. 30 Number 2 pp 73 – 75
7. Oladokun, M.A.O (1982): Morph-physiological aspect of germination, rooting and seedling growth in kola (*cola* spp) Ph. D thesis, the University of Ibadan pp. 230
8. Jayeal, C O (2001): Preliminary studies on the uses of kola nuts (*Cola nitida*) for soft drink production. The journal of food Technology in Africa, Vol. 6, No. 1, pp. 25 – 26
9. Egbe, N.E and Oladokun, M.A.O (1987): Factors limiting high yield in kola nut (*Cola nitida*) production Nigeria. Café cocoa, the vol. XXXL, 1004, Oct. - Dec.
10. Dewole EA, Dewumi DF, Alabi JY, Adegoke A. Proximate and phytochemical of *Cola nitida* and *Cola acuminata*. Pak J Biol Sci. 2013 Nov 15;16(22):1593-6.
11. Okoli, B.J, Abdullahi, K, Myina, O and Iwu .G (2012). Caffeine Content of Three Nigerian Cola. *Journal of Emerging Trends in Engineering and Applied Sciences (JETEAS)* 3(5):830-833
12. Botton PH, Pochmann D, Rocha AS, Nunes F, Almeida AS, Marques DM, Porciúncula LO. Aged mice receiving caffeine since adulthood show distinct patterns of anxiety-related behavior. *Physiol Behav.* 2017;170:47-53.
13. Sweeney P, Levack R, Watters J, Xu Z, Yang Y. Caffeine increases food intake while reducing anxiety-related behaviors. *Appetite.* 2016;101:171-7.
14. Richards G, Smith A. Caffeine consumption and self-assessed stress, anxiety, and depression in secondary school children. *J Psychopharmacol.* 2015;29(12):1236-47.
15. Osim, E.E and Udia, P.M (1993): The effect of consuming kola nut *Cola nitida* diet or mean arterial pressure in rats. Int. J. pharmacog. 31, No3, 193 - 197
16. Brown, R.E, Corey, S.C; Moore, A.K (1999): Differences in Measures of Exploration and Fear in MHC – Congenic C57BL/^ΔJ and B6 – H – 2k Mice. *Behaviour and Genetics*; 26, 263 – 271.
17. Lister, R.G. (1987): The Use of a Plus-Maze to Measure Anxiety in Mouse. *Psychopharmacology* 92: 180 – 185
18. Bisong S. A., U. E. Okon, E. A. Egbung, F. E. Abuo1 and O. A. Sanya (2017). Effect of Crude Ethanol Leaf-extract of *Murraya koenigii* on Anxiety in Mice. Asian Journal of Medicine and Health 7(1): 1-9; Article no.AJMAH.36294

- 371 19. Bisong S. A., C. O. Nku K. U. Nwoke and E. E. Osim (2018). Crude aqueous leave
372 extract of *Carica papaya* Linn (pawpaw) reduced anxiety and fear related
373 behaviour in CD1 mice. *European Journal of Pharmaceutical and Medical*
374 *Research*, 2018, 5(3), 488-493.
- 375 20. Hascoët, M; Bourin, M (1998): A New Approval to the Light/Dark Procedure in
376 Mice. *Pharmacological Biochem. Behav* 60, 645 – 653
- 377 21. Weiss, G and Greenberg, G. (1996). *Open field procedures*. In Greenberg, G. and
378 Haraway, M.H. (Eds) *Comparative psychology. A hand book*, Gaeland
379 New York. pp. 603-62.
- 380 22. Neil, J.F (1978): Caffeinism Complicating Hypersomnic Depressives Episodes.
381 *Comprehensive Psychiatry* 19, 377
- 382 23. Greden, J.F *et al* (1978): Anxiety and Depression Associated with Caffeine Among
383 Psychiatric patients. *Am. J* 131, 8, 963
- 384 24. Crawley, J. N., Godwin, F.K. (1980). Preliminary report of a simple
385 animal behaviour for the anxiolytic effects of benzodiazepines.
386 *Pharmacol. Biochem. Behave. B*, 167 – 170
- 387 25. Costall, B., Jones, B.J., Kelly, M.E., Naylor, R.J. & Tomkins, D.M. (1989).
388 Exploration of mice in a black and white test box: Validation as a model of
389 anxiety. *Pharmacology, Biochemistry and Behavior*, 32:777-785.
- 390 26. File, S.E; Hyde, J.R.G (1979): A Test of Anxiety that Distinguishes Between the
391 Actions of Benzodiazepines and those of other minor Tranquilizers and of
392 Stimulants. *Pharmacol. Biochem. Behav.* 11: 65 – 69
- 393 27. Charney, D.S; Galloway, M.P; Heninger, G.R (1984): The Effects of Caffeine of
394 Plasma MHPG; Subjective Anxiety, Automatic Symptoms and Blood
395 Pressure in Healthy Human. *Life Sci.* 35: 135 – 144
396