

**Species Composition and Diurnal Activity Rhythm of Tabanids
(Diptera: Tabanidae) at the Ivindo National Park and its Environs**

ABSTRACT

An entomological prospection to show the species composition and diurnal activity of tabanids was carried out using 15 Vavoua traps, during the rainy season (25th Sept-5th Oct and 21st Oct-9th Nov 2018), in the secondary forest [Ivindo National Park (INP)] and Village-Town sites in and around the biosphere reserve Ipassa-IRET Makokou in Gabon. In total, 839 tabanids were caught with 747 recorded at INP and regrouped under 11 species of the genus *Tabanus* [*T. taeniola* (57.76%), *T. ricardae* (26.32%), *T. par* (5.54%), *T. ruficrus* (3.74%), *T. socius* (3.74%), *T. disjunctus* (2.49%), *T. marmorosus* (0.28%) and *T. obscurehirtus* (0.14%)]. However, two species of the genus *Chrysops* [*C. dimidiatus* (70.83%) and *C. silaceus* (29.17%)] and one species of the genus *Haematopota* [*H. pluvialis* (100%)] were only identified at INP. While in the Village-Town sites, 92 tabanids were caught and regrouped under five species of the genus *Tabanus* [*T. ruficrus* (14.13%), *T. taeniola* (9.78%), *T. ricardae* (73.91%), *T. par* (1.09%), and *T. marmorosus* (1.09%)]. Tabanids were more abundant in the forest (7.12 tabanid/trap/day) as compared to the Village-Town (0.77 t/t/d) with a statistically significant difference ($P<0.05$). At INP, tabanids showed a unimodal activity pattern with peak attained between 12-14H and in the anthropized milieu (14-16H) and dominated by females.

Keywords: Hematophagous flies; park, reserve; Vavoua traps; biotope; rainy season; Gabon.

1. INTRODUCTION

1.1 Background

Tabanids represent a large group of dipterous insects with 4400 species regrouped under 144 genera [1]. Tabanids are a neglected subject of research but are vectors of pathogens of medical and veterinary importance [1,2]. Tabanids are known mechanical vectors of animal *Trypanosoma* spp., like *T. vivax*, *T. theileri* etc. [3] as

well as biological vectors (*Chrysops silacea* and *C. dimidiata*) of *Loa loa* filariasis [4,5,6].

In Gabon, there is no longitudinal study on tabanids, but the few existing reports on this group are cross sectional studies carried out in some protected areas and their environs [7, 8, 9, 10, 11].

The biting activity pattern of tabanids have been established for forest and savanna groups [5, 12]. The activity patterns vary with conditions of micro-environment across seasons with activity peaks ranging from unimodal, bimodal, trimodal etc [1, 5]. However, no information exists on the activity rhythm of tabanids with respect to sex at the Ivindo National Park (INP) and its environs. The present entomological prospection aims at determining the abundance and diurnal activity profile with respect to sex of tabanids at the Ivindo National Park and its surrounding.

2. MATERIALS AND METHODS

2.1 Study area

The study was carried out in Makokou, located in the Ogooué-Ivindo province in North-East Gabon. Trapping was carried out at the Institute of Research for Tropical Ecology (IRET) of Ipassa (0°.51'N; 12° 79'E) and its environs (0°.52'N; 12° 82'E) (Figure 1), elevated at an altitude of 500m [13]. The climate is of the equatorial and humid type with alternating rainy and dry seasons. The mean annual rainfall is 1600 to 1800mm while the mean annual temperatures are close to 24°C. The annual and daily thermal amplitudes are weak [7] and the main water body in the area is river Ivindo. The entomological prospection was carried out during the rainy season (25th Sept-5th Oct and 21st Oct-9th Nov 2018). The forest fauna of Gabon is rich and diversified and the Makokou region holds a significant share with one of the highest listed fauna in Gabon. It consists of 128 species of mammals, 424 species of birds, 65 species of reptiles, 47 species of amphibians among others [7].




Figure 1. Map of the study site showing the position of traps (red circles)

2.2 Capture of tabanids

Trapping was carried out using 15 Vavoua traps [14]. Vavoua traps have been reported to be efficient in the collection of tabanids [15]. Traps were set along a transect of about 17km, following the anthropogenic gradient from the secondary forest in the INP (non-anthropized environment), to the Village and Makokou town (highly anthropized environment). Trapping duration was three months (25th Sept-5th Oct and 21st Oct-9th Nov 2018). The 15 traps were divided into the two study areas, at a ratio of 7-8 traps per milieu. Trapping effort was: 15 traps × 30 days = 450 traps days. The traps were activated in the morning at 8 am and emptied at 6 pm. All trap cages were tagged with the trap number and date and returned to the laboratory. They were then placed in a freezer for 15 minutes to kill the insect prior identification. Identification was carried out using the dissecting microscope (LABOMED®, France) of the field station laboratory of IRET-Ipassa in Makokou.

2.3 Daily activity Rhythm

The diurnal activity pattern of tabanids in the study area was carried out using three traps in each of the prospection site for three days consecutively during prospection

94 days. The follow-up diurnal time ranges for this trial was 8-10H, 10-12H, 12-14H, 14-
95 16H and 16-18H. Trap-tags consisted of date, location and time interval.

96 **2.4 Fly identification**

97 The identification of tabanids was carried out using the identification key of Oldroyd
98 [16].

99 The abundance was defined by the Trap Apparent Density (ADT) known as the
100 number of tabanids caught per trap and day:

$$101 \text{ ADT} = \frac{\text{Number of tabanids flies captured}}{\text{Number of traps} \times \text{Number of trapping days}}$$

102 **2.5 Data analysis**

103 The statistical analysis was carried out using SPSS version 20. The one-way
104 ANOVA on ranks was used to compare the ADT in the two prospection biotopes.
105 The level significance was set at $P < 0.05$.

106 **3. RESULTS**

107 **3.1 The species composition of tabanids with respected to prospection site**

108 In total, 839 tabanids were caught with 747 collected at the INP which consisted of
109 11 species regrouped under three genera notably *Tabanus*, *Chrysops* and
110 *Haematopota*. At the INP, *Tabanus taeniola* was the most frequent species while
111 *Tabanus obscurehirtus* and *H. pluvialis* were very rare (Table 1). At the Village-Town
112 trap-sites, only 92 tabanids were caught and five species identified and all belonged
113 to the genus *Tabanus* (Table 1). In the Village-Town trap-sites, *T. ricardae* was the
114 most frequent while *T. par* and *T. marmorosus* were very rare (Table 1).

115

116

117

118

119

120

121 **Table 1. Percentage of tabanids fly species in prospected locations**

Biotope	Genus	Species	Number	%
INP	Tabanus (N=8)	Tabanus taeniola	417	57.76
		Tabanus ricardae	190	26.32
		Tabanus par	40	5.54
		Tabanus ruficrus	27	3.74
		Tabanus socius	27	3.74
		Tabanus marmorosus	2	0.28
		Tabanus obscurehirtus	1	0.14
		Tabanus disjunctus	18	2.49
	Total Tabanus		722	100
	Chrysops (N=2)	Chrysops dimidiatus	17	70.83
		C. silaceus	7	29.17
Total Chrysops		24	100	
Haematopota (N=1)		Haematopota pluvialis	1	100
Total Haematopota		1	100	
Village-Town	Tabanus (N=5)	Tabanus ruficrus	13	14.13
		Tabanus taeniola	9	9.78
		Tabanus ricardae	68	73.91
		Tabanus par	1	1.09
		Tabanus marmorosus	1	1.09
	Total Tabanus		92	100

122

123 3.2 Abundance of tabanids with respect to prospection sites

124 At the INP, *Tabanus taeniola* was the most abundant species of the genus *Tabanus*
 125 with ADT of 3.99. In *Chrysops*, *Chrysops dimidiata* (ADT = 0.16) was the most
 126 frequent species in the present collections. However, in the genus *Haematopota*,
 127 only one species was identified with a low apparent density (ADT= 0.01) (Figure 2A).

128 In the Village-Town sites, *Tabanus ricardae* was the most abundant species of the
 129 genus *Tabanus* with ADT of 0.57 (Figure 2B). The other species notably *Tabanus*

rifucrus and *Tabanus taeniola* were rarely captured. *Tabanus par* and *Tabanus marmoratus* were rarely caught (Figure 2B). Generally, the ADT of tabanids was higher in the forest (ADT=7.12) as compared to the Village-Town (ADT=0.77) sites with a statistically significant difference ($P<0.05$).




Figure 2. The ADT of tabanids in INP (A) and Village-Town (B)

3.3 Daily activity rhythm of tabanids with respect to prospection sites

At the INP, all the species of tabanids caught showed the same peaks of abundance between 12 and 14H (Figure 3A). In the Village-Town sites, *Tabanus taeniola*, *Tabanus socius* and *Tabanus ruficrus* showed a unimodal daily activity with peak noticed between 14 to 16H (Figure 3B). However, *Tabanus ricardae* was captured with the same abundance between 8 and 10H (Figure 3B).




Figure 3. Diurnal activity of tabanids with respect to site A(INP) and B (Village-Town)

4. DISCUSSION

The apparent density per trap and day of tabanids was higher than that reported by Mavoungou et al., [7]. This difference could be related to the discrepancies in the trapping efforts. In total, 16 species of tabanids were identified and represented 22.5% of known species of Central Africa [17]. This low species richness in the present study could be related to the type of trap used. Because Mihok [18]; Acapovi [19] rather revealed the Nzi trap was an efficient trapping system for tabanids. The Vavoua trap was used in this study for budgetary reasons i.e. it is cheap even though not efficient as compared to Nzi which is very expensive.

The tabanid species composition and ADT was higher at the INP trapping points as compared to the man-made areas. This difference in tabanid species composition

and number has been reported for these type-localities [7]. The high ADT of tabanids at the INP could be linked to the favourable environmental factors such as ambient temperature, relative humidity, rainfall and high host density [1] prevailing in the site that fostered the high development and survival of this group. It was interesting to find out that during the prospection period, *C. silacae* and *C. dimidiata* were only caught by traps set in the secondary forest of the INP and never caught by traps pitched at the Village-Town sites. The presence of the two species of Chrysops which are main vectors of *Loa loa* filariasis [5] indicates the risk of the transmission of the parasite to the human population frequenting the INP as well as the adjacent villages.

The peak of activity of tabanids followed a unimodal pattern and occurred at different diurnal time ranges in the INP (12-14H) and Village-Town (14-16H). This result is like that obtained by Acapovi [20] who showed a unimodal activity peak for tabanids in the mentioned periods. The high tabanid abundance during at time points of the day could be related to the prevailing meteorological factors (ambient temperature, wind speed and relative humidity) during such periods that favoured their activity in the prospected sites. The mentioned weather variables have been shown to be the main modulators of the activity of tabanids [1] and such conditions are optimal between 14 to 16H [20].

5. CONCLUSION

The genus *Tabanus* was highly represented at the INP trapping sites with 8 species (with *Tabanus taeniola* as the most abundant species) and less represented in anthropized medium with 4 species (with *T. ricardae* as the most abundant species). Three species notably *Chrysops dimidiata*, *Chrysops silacae* and *Haematopota pluvialis* were only identified in the secondary forest of the INP. The peak activity of tabanids in the Village-Town sites occurred between 14-16H while that at the INP occurred between 12-14H.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Baldacchino F, Desquesnes M, Mihok S, Foil LD, Duvallet G, Jittapalapong S. Tabanids: Neglected subjects of research, but important vectors of disease agents. *Infection, Genetics and Evolution*. 2014; 28:596-615.
2. Duvallet G, Boulanger N, Robert V. Arthropods: Definition and Medical Importance. Chapter 2 in *Skin and Arthropod Vectors*. Edited by Nathalie Boulangie; Elsevier Inc. 2018; 29-51.
3. Desquesnes M, Dia ML. Mechanical transmission of *Trypanosoma vivax* in cattle by the African tabanid *Atylotus fuscipes*. *Veterinary Parasitology*. 2004; 119: 9-19.
4. Toure FS, Mavoungou E, Deloron P, Egwang TG. Analyse comparative de deux méthodes de diagnostics de la loase humaine: sérologie IgG4 et PCR nichée. *Bulletin de la Société de Pathologie Exotique*. 1999; 92:167-170.
5. Wanji S, Tendongfor N, Esum M, Atanga SN, Enyong P. Heterogeneity in the prevalence and intensity of loiasis in five contrasting bioecological zones in Cameroon. *Trans. Royal Society of Tropical Medicine and Hygiene*. 2003; 97: 182-187.
6. Kelly-Hope LA, Paulo R, Thomas B, Brito M, Unnasch TR, Molyneux D. *Loa loa* vectors *Chrysops* spp.: perspectives on research, distribution, bionomics, and implications for elimination of lymphatic filariasis and onchocerciasis. *Parasite and Vectors*. 2017; 10:172.
7. Mavoungou JF, Makanga B, Acapovi YG, Desquesnes M, M'Batchi B. [Abundance and species diversity of tabanids (Diptera) in the biosphere reserve Ipassa-Makokou (Gabon) during the rainy season]. *Parasite*. 2012; 19: 165-171.
8. Zinga Koumba RC, Acapovi Yao GL, Mavoungou JF, Tongue KL, Mbang-Nguema OA, Obame OKP, Shango M. Influence de la saison sur l'écodistribution des glossines, tabanides, stomoxes du Baï de Momba Makokou, Gabon. *Agronomie Africaine*. 2013; 25: 149-158.
9. Bitome-Essono PY, Dechaume-Moncharmont FX, Mavoungou JF, Obiang Mba R, Duvallet G, Bretagnolle F. Distribution and abundance of hematophagous flies (Glossinidae, *Stomoxys*, and Tabanidae) in two national parks of Gabon. *Parasites*. 2015; 22: 23.

10. Mbang NOA, Mavoungou JF, Mawili-Mboumba DP, Zinga KRC, Bouyou-Akotet MK, M'batchi B. Inventory of potential vectors of *Trypanosoma* and infection rate of tsetse in the National Park of Ivindo, Gabon. *African Health Science*. 2015; 15: 762-767.
11. Mounioko F, Maganga GD, Mavoungou JF, Zinga KCR, Koumba AA, Sevidzem SL, Tamesse JL, Gustave S, M'batchi B. Molecular Screening of *Trypanosoma spp.* in *Glossina*, *Stomoxys* and Tabanids in the Moukalaba Doudou National Park (South-West, Gabon). *World Journal of Veterinary Science*. 2018; 6: 52-61.
12. Parrah-Henao G, Alarcon-Pineda EP, Lopez-Valencia G. Ecology and Parasitological analysis of Horseflies (Diptera: Tabanidae) in Antioquia, Columbia. *Caldasia*. 2008; 30 (1):179-188.
13. Wilks C. La conservation des écosystèmes forestiers du Gabon. Programme pour les forêts tropicales, UINC/CCE. 1990.
14. Laveissière C, Grébaut P. Recherche sur les pièges à glossines (Diptera: Glossinidae). Mise au point d'un modèle économique: le piège « Vavoua ». *Tropical Medicine and Parasitology*. 1990; 41:185-192.
15. Lendzele SS, Abdoulmoumini M, Timoleon T, Garabed R, Renz A, Dicmu S. Risk factors for the Contamination of *Stomoxys niger niger* Macquart 1851 (Diptera: Muscidae) With the Foot-and-Mouth Disease Virus. *Global Foot-and-Mouth Disease Research Alliance (GFRA) book of abstracts*. 2017; 149.
16. Oldroyd H. The horse-flies (Diptera: Tabanidae) of the Ethiopian region. II. *Tabanus* and related genera. London, UK, British Museum (Natural History). 1954; 341.
17. Taufflieb R, Finelle P. Étude écologique et biologique des tabanidés d'Afrique Équatoriale Française. *Bulletin Institut d'études centrafricaines, nouvelle série*, Brazzaville, Congo. 1956, 12: 209-251.
18. Mihok S (2002) The development of a multipurpose trap (the Nzi) for tsetse and other biting flies. *Bull Entomol Res* 92:385-403.
19. Acapovi GL, Yao Y, N'Goran E, Dia ML, Desquesnes M. Abondance relative des tabanidés dans la région des savanes de Côte d'Ivoire. *Revue d'élevage et de Médecine vétérinaire Pays tropicaux*. 2001; 54(2): 109-114.

268 20. Acapovi-Yao GL, Kohagne LT, Dieudonné TBT, Mavoungou JF. Dynamique et
269 dispersion spatiale des Tabanidae dans différents faciès écologiques de Korhogo en
270 Côte d'Ivoire. Entomologie Faunistique-Faunistic Entomology. 2017; 70:13-22.

271

272

273

UNDER PEER REVIEW