

Effect of the organic and NPK fertilizers on the growth and yield of sweet potato (*Ipomoea batatas* (L) Lam) in the centre of Côte d'Ivoire

Original research papers

ABSTRACT

In Côte d'Ivoire, the sweet potato is little cultivated but stays however a culture of pension and a food mattering in certain regions. This study was conducted to assess the effect of the application of mineral (NPK 15-15-15, NPK 12-22-22) and organic (poultry manure) fertilizers on yield components of sweet potato (*Ipomoea batatas* (L) Lam). The experiment was conducted over two years (2016 to 2017) in experimental station of National Center of Agronomic Research (CNRA) of Bouaké in the centre of Côte d'Ivoire. The experiment was conducted following a split plot with 2 factors (variety and fertilizers) and 3 replicates. The main factor was variety with 2 levels (variety Irene and variety TIB-440060) and the subplot was fertilizers application consisted 13 levels (200, 300, 400, 500 kg ha⁻¹ of NPK fertilizers combined 5, 10, 15 t ha⁻¹ of organic fertilizer). Thirteen treatments based on poultry manure (T6, T7), chemical fertilizer NPK 15 15 15 (T1, T3, T4, T5) and NPK 12 22 22 (T8, T10, T11, T12) and their combination (T2 and T9) were tested. The experimental results revealed that that the effects of chemical fertilizer treatments, manure and their combination with mineral fertilizers did not influence the measured parameters. The results also showed that fertilizers improved all the agronomic parameters of sweet potato compared to the control treatment during two years of experiment. However, the Irene variety with a mean weight of 180 g tuberous roots was significantly different from the TIB variety with a mean weight of 138.73 g.

In the conditions of our study and over the two-year period, producers of sweet potatoes would benefit from using organic fertilizers with the Irene variety.

Keywords: *Bouaké, Ipomoea batatas, mineral fertilizers, organic fertilizers, yield.*

1. INTRODUCTION

Sweet potato (*Ipomoea batatas* (L.) Lam) is a perennial herbaceous vine cultivated as an annual. It is suited to the humid tropics but can be grown in southern temperate regions. The tubers and sometimes the leaves are eaten [1]. She can grow in numerous different ecological conditions [1]. Her world production was 106 million of tons in 2016 among which more than 67% coming from Asia, particularly, China with 71 million of tons [2]. In Africa, the sweet potato is an important economic culture because of its potential of struggle against poverty and improvement of the nutritional state of the rural populations in a little expensive and long-lasting way [3]. She is widely cultivated as a basic food mattering in a number of country African and its nutritional value exceeds by far the yam, the

cassava and the taro [4]. In Côte d'Ivoire, the sweet potato is little cultivated but stays however a culture of pension and a food mattering in certain regions [5]. The national production was estimated at approximately 49 357 tons in 2016 [2]. In spite of its nutritional and economic assets, her production is relatively low because confronted with enormous difficulties. Sweet potato is cultivated on poor or marginal lands. Furthermore, the majority of the cultivars receive only few or no fertilizer. In the face of these constraints, it is thus necessary to have the good cultural technique with an adapted regime of fertilization which offers better yields in the ivorian conditions. According to [6] and Palm and al. [7], it is generally admitted that organic and chemical fertilizers are necessary to increase

the agricultural production in west Africa. Several hypotheses were formulated concerning a possible application or a combination of organic and chemical fertilizers [8], so improving the yields of the cultures, the fertility of grounds and production costs [7]. In spite of the numerous hypotheses which were formulated, little information is brought back on the optimal level of application of organic and chemical fertilizers for the culture of the sweet

potato, in particular in Côte d'Ivoire. Consequently, it is necessary to study and to recommend the best application of organic and chemical fertilizers to the farmers committed in the production of sweet potato. Consequently, it is necessary to study and to recommend the best application of organic and chemical fertilizers to the farmers committed in the production of the sweet potato.

2. MATERIALS AND METHODS

2.1 Site description

The present study was led to the Station of Research on Subsistence crops (SRCV) of the National Center of Agronomic Research (CNRA) of Bouaké situated in the centre of Côte d'Ivoire (7°46 'N, 5°06 'W, 376 m) [9]. Bouaké is in the zone of transition between the forest climate of the South and the climate of the savanna of the North [10]. The climate is of wet tropical type with four seasons, of which a big dry season (November in February), a big rainy season (March in June), a small dry season (July in August) and a small rainy season (September in October). These periods are less and less marked these last years [11]. The hydrography mainly consists of the river

Bandama and the N'Zi and its ponds hillsides which drain Bouaké [12]. The vegetation is constituted by savanna raised with several species of Poaceae [13]. Soils are ferrallitic, averagely saturated, reshaped, shallow and stemming from a material of granitic change with a sandy-clayey texture [14]. The mean annual precipitation are 1200 mm of rains [15], with a mean annual temperature of 25.73 °C [10].

2.2 Chemical characteristics of the used fertilizer

The manure used has very high levels of carbon, nitrogen and CEC with high levels of organic matter. The chemical characteristics of the manure used are shown in Table I below.

Table I: Initial chemical characteristics of manure used

pH	Organic matter				P	Adsorbing complex (cmol.kg ⁻¹)				
	C (%)	N (%)	C/N	MO (%)		P.ass (g.kg ⁻¹)	CEC	Ca ²⁺	Mg ²⁺	K ⁺
7,3	7,05	0,65	10,84	12,15	0,056	11,24	8,77	2,29	0,90	0,16

2.3 Plant material

The plant material was constituted of two (2) varieties of sweet potato with orange flesh : Irene and TIB-440060. The cuttings of these varieties were removed on plots of land of multiplication of cuttings of the Station of Research on Subsistence crops (SRCV) of the CNRA of Bouaké (Fig. 1).



Fig. 1. Tunnel of multiplication of cuttings of sweet potato.

2.4 Fertilizing material

The fertilizing materials used for the try were constituted by two types of fertilizer organic manure and an artificial fertilizer. Chemical

fertilizers consisted of the granular N-P-K 15-15-15 and the N-P-K 12-22-22 + 2SO₃ + 1MgO + 5CaO. The organic fertilizer was a mixture of dung of cow and droppings of chicken.

2.5 Experimental design

The try was led during two consecutive years (2016 and 2017) following to a design in Split-Plot to three repetitions. The main factor was constituted by the variety with 2 modalities (variety Irene and variety TIB-440060) and the second factor by fertilizers. So, the chemical fertilizer contained four levels : 200; 300; 400 and 500 kg.ha⁻¹ of N-P-K and the organic manure included three (3) levels, worth knowing, 5, 10 and 15 t.ha⁻¹ and their

combinations distributed in a random way within the elementary plots of land. Treatments are presented in table II.

The total surface of the try was 702 m² with 39 elementary plots of land corresponding to treatments. Every elementary plot was constituted of sixteen (16) mounds on a surface of 3 m x 3 m with a space between the mounds of 1 m on 1 m, that is a density of 10 000 mounds in the hectare. Blocks were separated from each other by paths of 2 m and the main plots of land and main plots of land, by a spacing of 1.5 m.

Table II : List of the various applied treatments

Identification of treatments	Composition of treatments
Te	Control without fertilizer
T ₁	200 kg.ha ⁻¹ NPK 15 15 15
T ₂	200 kg.ha ⁻¹ NPK 15 15 15 + 5 t.ha ⁻¹ of manure
T ₃	300 kg.ha ⁻¹ NPK 15 15 15
T ₄	400 kg.ha ⁻¹ NPK 15 15 15
T ₅	500 kg.ha ⁻¹ NPK 15 15 15
T ₆	10 t.ha ⁻¹ of manure
T ₇	15 t.ha ⁻¹ of manure
T ₈	200 kg.ha ⁻¹ NPK 12 22 22
T ₉	200 kg.ha ⁻¹ NPK 12 22 22 + 5 t.ha ⁻¹ of manure
T ₁₀	300 kg.ha ⁻¹ NPK 12 22 22
T ₁₁	400 kg.ha ⁻¹ NPK 12 22 22
T ₁₂	500 kg.ha ⁻¹ NPK 12 22 22

2.6 Preparing land

The parcel was cleaned with a machete and the crop residue was removed. These crops residues were removed from the plot and the ground was furnished, followed by the construction of humps at a height of at least 60 cm to prevent the phenomena of settlement due to rains.

2.7 Mound making and planting

The sweet potatoes were planted with stripped cuttings between 15 and 30 cm, bearing 3 to 5 knots. Three (3) cuttings were planted at equidistance (approximately 30 cm apart) at the top of each mound with a density of 30,000 plants per hectare. The height of the log at planting was at least 60 cm to prevent the phenomena of settlement due to rains (Fig. 2).



Mound of plantation

Cuttings

Fig. 2. Hump cuttings planting (Dibi and *al*, 2015)

2.8 Application of fertilizers

The various fertilizers were manually applied to the mounds in the planting holes before the cuttings were planted. The fertilizers were buried by mixing the fertilizers with the soil in the cuttings planting holes.

2.9 Maintaining of the parcels

Dead plant replacements were performed one week after cuttings were planted. Three manual weeds (weeding) were performed during the test. They consisted in ridding the hummocks and intermediates of the weeds. It is then advisable to trace the logs to the first weeding (between the 30th and the 40th day after planting, depending on the case). The first weeding was performed two weeks after transplanting, the second at six weeks and the third at ten weeks. The plots were irrigated regularly during the dry season.

The water intake was done weekly using an irrigation system.

2.10 Harvesting of tubers

The harvesting was done by hand at the ripening of the tubers using tools with great care in order to avoid the various alterations of the tuberous roots to have satisfactory yields which are kept well and long. Harvest occurred about 4 to 6 months after planting. Ripening was also appreciated by observing yellowing of the leaves, followed by thinning or by the clear colour of the latex that comes out when cutting the tubers.

2.11 Collect of the agronomic parameters

2.11.1 Number of tuberous roots of large and small size

The average number of tuberous roots of large and small size was determined at harvest,

classifying tuberous roots according to their

2.11.2 Number of tuberous roots per plant and weight of tuberous roots

The number of tuberous roots for each parcel was determined by counting. Next, the total number of tuberous roots was divided by the number of productive plants. The number of tuberous roots per plant is given by the following formula:

$$N_{tub} = \frac{N_{ttub}}{N_{pp}}$$

2.11.3 Yield in fresh tubers

The yield ($t \cdot ha^{-1}$) of fresh tubers was determined by the ratio of the total weight in tuberous roots (P_{ttub}) to the area (Sup) of the

2.11.4 Crop index

The crop index (I) which represents the efficiency of the production of tuberous roots was determined by the ratio between the total weight of the tuberous roots (P_{tub}) and the total weight of the plant ($P_{ttub} + P_{tig}$).

2.11.5 Dry matter levels

Dry matter levels were determined by selecting tubers at random by repetition and treatment for laboratory analysis. Samples were taken at three different levels: in the middle and close to the two ends of each selected tuberous root. All of these samples were then merged to have a representative sample per treatment. The resulting pieces were peeled and a fraction of 200g was collected to give the fresh weight (PF) per sample. These 200g fractions were

2.11.6 Increase level

The increase level or increase in yield (increase level) relative to the control, which shows the effect of fertilizers on yields, was calculated according to the following formula :

marketable or non-marketable size.

(N_{tub}): Number of tuberous roots for each parcel;

(N_{ttub}): Total number of tuberous roots;

(N_{pp}): Number of productive plants.

For the mean weight (P_{tub}), the tuberous roots of each treatment were weighed and the mean was determined for two test years. It is given by the formula below :

$$P_{tub} (g) = \frac{P_{ttub}}{N_{pp}}$$

parcel. It is determined at harvest and is given by the following formula

$$Rendement (t/ha) = \frac{P_{ttub}}{Sup}$$

$$I R = \frac{P_{tub}}{P_{tub} + P_{tig}}$$

then cut into small strips and then put into aluminum paper depending on the treatment. Aluminum paper containing these sweet potato slats was put in the oven at 100°C for 24 hours. Dry and cooled samples were weighed to determine dry weight (PS). This allowed the following formula to determine the dry matter level (TMS):

$$TMS (\%) = \frac{PS}{PF} \times 100$$

$$Taux accr. (\%) = \frac{Rend (F) - Rend (T)}{Rend T} \times 100$$

Rend (T) = yield of the control parcel ; Rend (F) = yield of the fertilized parcel.

2.11.7 Coefficient of efficiency of fertilizers

The efficiency of the fertiliser which is determined by the increase in yield due to the intake of 1 kg of fertiliser has been given by the relationship :

$$Ce = \frac{\text{Rend (F)} - \text{Rend (T)}}{\text{Dose engrais}}$$

Ce : Efficiency coefficient ; Rend (T) = control plot yield; Rend (F) = fertilized plot yield.

2.12 Statistical analysis

At the end of the experiment, the data collected were subjected to data analysis (ANOVA) using the software Statistica version 7.1. This analysis was used to assess the effect of fertilizers on yield and its components on both sweet potato varieties and significant difference was determined. The Fisher 5% test was used to classify averages into homogeneous groups. A Mann Whitney test compared mean two to two.

3. RESULTS

3.1 Effect of treatments on number of marketable and non-marketable tuberous roots

For the number of non-marketable tuberous roots, the results showed a significant difference ($P = .05$) between the two varieties (Table III). Irene (234 roots) recorded the highest number of non-marketable tuberous roots compared to TIB-440060 (151 roots).

For treatments, T1 (NPK 15-15-15) with a number of 168 tuberous roots was not significant from T11 (NPK 12-22-22) which

obtained the highest number of non-marketable tuberous roots with an average of 215.

The effect of the control treatment was not significantly different from that of T1 (200 kg.ha⁻¹ NPK 15 15 15). Other treatments had mean numbers that ranged from 183 to 209 non-marketable tuberous roots. The interaction between the variety and the fertilizer, was not significant ($P = .05$).

As regards the average numbers of marketable tuberous roots, they oscillated between 82 and 123 for all treatments applied. T12 (500 kg.ha⁻¹ NPK 12 22 22) and T8 (200 kg.ha⁻¹ NPK 12 22 22) obtained the highest mean numbers of tuberous roots, with 123 and 114 marketable roots respectively, an increase of 19% and 10% compared to the witch with an average of 103 marketable roots (Table III). Chemical fertilizer treatments were not significantly different from the control and manure treatments. At variety level, Irene recorded the highest mean number (108 marketable roots), while TIB recorded 98 marketable roots. However, the variance analysis of the mean number of marketable tuberous roots did not show a significant difference for varieties and their interaction with varieties (Table III).

Table III: Effect of fertilizers on average number of marketable and non-marketable tuberous roots

Variables	Levels	Number of small tuberous roots	increase level (%)	Number of big tuberous roots	increase level (%)
Variety	Irene	234 ^a		108 ^a	
	TIB	151 ^b		90 ^a	
<i>P</i>		0.001		0.313	
Treatments	T _e	201	0	103	0
	T ₁	168	-17	84	-19
	T ₂	194	-3	95	-8
	T ₃	198	-2	94	-9
	T ₄	185	-8	113	9
	T ₅	188	-7	102	-2
	T ₆	192	-5	91	-12
	T ₇	183	-9	97	-7
	T ₈	202	0	114	10
	T ₉	183	-9	82	-20
	T ₁₀	185	-8	88	-15
	T ₁₁	215	7	99	-4
T ₁₂	209	4	123	19	
<i>p</i>		1.00		1.00	
<i>p</i> (interaction)		1.00		1.00	

*Assigned column averages of the same letter do not differ from the 5% threshold (Fisher test)

3.2 Effects of varieties and fertilizers on average number of tuberous roots per plant

The results showed that the varieties had highly significant effects on the mean number of tuberous roots per plant ($P < .001$). Indeed, the TIB variety obtained the highest mean number with 3 roots per plant. The Irene variety recorded the low mean number, with 2 roots per plant (Table IV).

3.3 Effects of fertilizers on the average weight of tuberous roots

Analysis of the results showed that the variety factor had a highly significant effect ($P < .001$) on the mean weight of the tuberous roots (Table V). Irene obtained the highest mean number of tuberous roots (0.18 kg) and TIB-440060 with the lowest mean number (0.13 kg).

As regards the effect of fertilizers, the mean numbers of tuberous roots varied between 2 and 3 tuberous roots per plant for all treatments applied. The mean numbers of tuberous roots obtained from treatments involving chemical fertilizers and treatments consisting of the combination of chemical fertilizers and organic fertilizers were not statistically different from the control. The same was true for the variety fertilizer interaction (Table IV).

For the fertilizer factor, the T8 treatment (200 kg.ha⁻¹ NPK 12 22 22) obtained the highest average number of tuberose roots (0.19 kg) or an increase of 31%, followed by treatment T10 (300 kg.ha⁻¹ NPK 12 22 22) with a mean number of 0.17 kg. The smallest mean number of tuberous roots (0.12 kg) was obtained with the T7 treatment (15 t.ha⁻¹ of manure).

Compared to the witness, the increases recorded were in the range of 7 to 31%. However, the variance analysis of the mean

3.4 Effect of Fertilizers on crop index

Results for the evolution of the crop index are presented in Table VI. Results obtained indicated that the variety TIB-440060 had the best crop index with an average number of 0.60, significantly different from the Irene

3.5 Effect of fertilizers on yield in fresh tuberous roots

The highest mean yield in fresh tuberose roots (9.4 t.ha⁻¹) was obtained with the T8 treatment (200 kg.ha⁻¹ NPK 12-22-22) with a 40% growth rate compared to the untreated control which recorded a mean yield of 6.8 t.ha⁻¹ and a zero increase rate (Table VI). This mean yield was significantly different from the effect of manure treatment ($P = 1.00$). On the other hand, the lowest mean yield (5.7 t.ha⁻¹) was recorded with

tuberous root weights did not show significant differences between fertilizers and their interaction with varieties.

variety, which recorded a mean harvest index of 0.50 ($P = .05$).

Concerning the fertilisers, the results obtained did not show a significant difference between crop indices in the different treatments. Harvest index values ranged from 0.53 to 0.58 for all applied fertilizers.

the T6 treatment (10 t.ha⁻¹ of manure. The T12 and T4 treatments also achieved high yields in fresh tubers, respectively, 8.1 t.ha⁻¹ and 7.7 t.ha⁻¹. Compared to the control, T9, T2, and T11 showed low yields with 5.9 t.ha⁻¹, 6.5 and 6.6 t.ha⁻¹, respectively. The smallest increase was achieved by T3 treatment (7.1 t.ha⁻¹) with at 5%.

For varieties, the results showed that there is no significant difference between yields.

Table IV: Effect of fertilizers on mean number of tuberous roots and mean weight

Variables	Levels	Number of tuberous roots/plant	Increase level (%)	Mean weight (g)	Increase level (%)
Variety	Irene	2 ^b		180.17 ^a	
	TIB	3 ^a		138.73 ^b	
<i>P</i>		0.001		0.012	
Treatments	T _e	3	0	144.63	0
	T ₁	2	-9	158.24	9
	T ₂	2	-6	154.89	7
	T ₃	3	-3	169.91	17
	T ₄	3	0	165.13	14
	T ₅	2	-6	175.37	21
	T ₆	3	-3	144.32	0
	T ₇	3	-4	126.48	-13
	T ₈	3	-2	189.34	31
	T ₉	2	-11	135.9	-6
	T ₁₀	2	-12	177.43	23
	T ₁₁	3	5	155.25	7
T ₁₂	3	6	175.93	22	
<i>p</i>		0.653		0.10	
<i>p</i> (interaction)		0.657		0.10	

* Assigned column averages of the same letter do not differ from the 5% threshold (Fisher test)

Table V: Effect of Fertilizers on crop index

Variables	Levels	Crop index	Increase level (%)
Variety	Irene	0.50 ^b	
	TIB	0.60 ^a	
<i>P</i>		0.032	
Treatments	T _e	0.58	0
	T ₁	0.53	-9
	T ₂	0.52	-10
	T ₃	0.57	-2
	T ₄	0.54	-6
	T ₅	0.54	-7
	T ₆	0.54	-8
	T ₇	0.54	-6
	T ₈	0.56	-3
	T ₉	0.58	-1
	T ₁₀	0.53	-9
	T ₁₁	0.54	-7
T ₁₂	0.54	-7	
<i>p</i>		1.00	
<i>p</i> (interaction)		0.10	

* Assigned column averages of the same letter do not differ from the 5% threshold (Fisher test)

Table VI: Effect of treatments on the mean yield in fresh tubers

Variables	Levels	Mean yield in fresh tubers (t.ha ⁻¹)	Increase level (%)
Variety	Irene	7.78 ^a	
	TIB	6.31 ^a	
<i>P</i>		0.339	
Treatments	T _e	6.8	0
	T ₁	6.5	-3
	T ₂	6.5	-4
	T ₃	7.1	5
	T ₄	7.7	14
	T ₅	7.3	8
	T ₆	5.7	-16
	T ₇	6.5	-3
	T ₈	9.4	40
	T ₉	5.9	-13
	T ₁₀	7.4	10
	T ₁₁	6.6	-2
T ₁₂	8.1	20	
<i>p</i>		0.10	
<i>p</i> (interaction)		0.10	

* Assigned column averages of the same letter do not differ from the 5% threshold (Fisher test)

3.6 Coefficient of efficiency of organic and mineral fertilizers

The results of the analysis showed that the T8 treatment (200 kg.ha⁻¹ NPK 12 22 22) with the highest efficiency also had the highest

efficiency coefficient (13). It was followed by T12, T10 and T4 treatments with efficacy coefficient 3 and 2, respectively. T2, T6, T7, T9 and T11 treatments recorded a zero efficiency coefficient (Fig. 2).

3.7 Effect of fertilizers on dry matter content of fresh tuberous roots

The mean dry matter content of fresh tuberose roots ranged from 34.12% to 36.48% for all fertilizer levels applied to the plots, compared to 35.40% for the control (Table VII). Fertilizers

and varieties did not significantly affect the mean dry matter content of tuberous roots. Indeed, they showed mean rates of statistically similar dry matter, regardless of the nature of fertilisation and variety ($P = .05$).




Fig. 2. Evolution of the efficiency coefficient according to fertilizers

Table VII: Effect of fertilizers on the rate of fresh dry matter of tuberous roots

Variables	Levels	Dry matter content (%)
Variety	Irene	35.19 ^a
	TIB	35.10 ^a
<i>P</i>		0.10
Treatments	T _e	35.40 ^a
	T ₁	36.48 ^a
	T ₂	34.12 ^a
	T ₃	35.38 ^a
	T ₄	36.20 ^a
	T ₅	35.22 ^a
	T ₆	34.75 ^a
	T ₇	34.57 ^a
	T ₈	34.05 ^a
	T ₉	35.19 ^a
	T ₁₀	34.80 ^a
	T ₁₁	35.29 ^a
T ₁₂	35.49 ^a	
<i>P</i>		0.10
<i>p</i> (interaction)		0.623

* Assigned column averages of the same letter do not differ from the 5% threshold (Fisher test)

4. DISCUSSION

4.1 Effect of organic fertilizers on agronomic parameters of sweet potato

Organic fertilizers did not affect the mean number of marketable tuberous roots, the number and weight of tuberous roots per plant, and yield despite organic fertilizer inputs of 10 and 15 t.ha⁻¹. This result could be explained by the fact that nutrients such as nitrogen and potassium from organic fertiliser are not in sufficient quantity [16] to produce large caliber tuberous roots. Nitrogen and potassium play a well-known role in the production of large calibers [17]. In addition, [16] reported that potassium contributes to the improvement of calibers, while a deficiency in K results in a higher proportion of small calibers. In addition, work by [18] has also confirmed that the input of N and K is essential for the production of sweet potatoes.

The decline in yield from organic fertilizers could be explained by the slow process of manure decomposition. Nutrients from organic matter are progressively released through mineralization and their action is much slower [19]. The results were in full agreement with those of [20] who also found no significant differences in performance and components with the application of organic fumire.

4.2 Effect of chemical fertilizers on agronomic parameters of sweet potato

The application of chemical fertilizers did not positively affect the numbers of marketable and plant tuberous roots. However, the application of mineral fertilizers allowed for an increase in the weight and yield of fresh tubers compared to the control without fertilizers. This would result in the direct accessibility of N, P and K nutrients brought to the soil by compound fertilizer [21]. It is therefore likely that increases of 31% and 40% between the control (Te) and T8 treatment are due to the conjugated effects of the three elements N, P and K. Similar results were obtained in Côte d'Ivoire on cassava [22] and in other countries ([23 ; 24]). These results showed that the highest yields

were recorded by fertilizer inputs with N, P and K. Our results are consistent with those of Lompo and Belem (2012) who obtained the best yields and fresh tuberous root weights with the mineral formula of N₆₀P₃₀K₁₀₀ on sweet potato.

In addition, other authors had previously reported that N and K were essential to the production of sweet potatoes. Thus, the application of chemical fertilizers that provide nutrients such as potassium and nitrogen significantly increases root yield ([18] ; [26]). In addition, the yield of the tuberous roots of the sweet potato may depend on the photosynthetic capacity of the foliar cover, the ability of the crop to translocate the assimilates of the leaf to the root and their ability to assimilate them. It is therefore possible that assimilation of photosynthesis translocated to the roots may have contributed to the high yields of the mineral fertilizer treatment [3].

4.3 Effect of the combination of chemical fertilizers and organic fertilizers on the agronomic parameters of sweet potato

The application of 200 kg.ha⁻¹ of NPK + 5 t.ha⁻¹ of manure did not affect the number, weight, and average yield of tuberous roots, as well as the dry matter content of tubers and the crop index. The low yields obtained with organic mineral fertilization would be related to the application dose of the fertilizer. [27] obtained the best yields in tuberose roots with the application of 150 kg.ha⁻¹ NPK + 2.4 t.ha⁻¹ manure. Indeed, the amount of nutrients such as nitrogen and potassium contained in fertilizers would be the basis for this result. Nitrogen and potassium are essential in the production of sweet potato, but when they are brought in excess or absorbed too late, they prolong vegetation at the expense of tubers. Thus the high concentration of nitrogen in the fertilizer could have slowed the activity of potassium. [28] reported that tuberous root magnification occurs when the nitrogen/potassium ratio is low (1:3 ratio), because a high nitrogen concentration reduces the potassium concentration, affecting the storage roots.

4.4 Efficiencies of mineral and organic fertilizers

The low efficiency coefficient value of fertilizer elements observed for organic fertilizers would likely be due to the slow and progressive mineralization of manure to provide mineral elements to soils and crops. Similar results were obtained by [29] with household waste composts on a Kinshasa Ferralsol, on peanut, soy and sorrel crops. Organic matter plays an important role in soil, thus promoting the growth of micro-organisms that induce activation of nutrient solubilization. Nutrients made sufficiently available over time in soil are used effectively by crops ([30]). In the case of our study, inputs of 10 and 15 t.ha⁻¹ of organic inputs yields and efficiency coefficient were the lowest, compared to mineral fertilizers that were more effective. This could be explained by the fact that the nutrients contained in mineral fertilizers are available and readily absorbable by crops as demonstrated by the work of [31] on the cultivation of gumbo and cabbage using mineral and organic fertilizers. For example, the 200 kg.ha⁻¹ dose of NPK 12-22-22 recorded the best average weights and yields compared to NPK 300, 400 and 500 kg.ha⁻¹. In fact, for any

4.5 Effect of varieties on agronomic parameters of sweet potato

The mean number of tuberose roots varied according to varieties as was the case in the 4 sweet potato varieties Mugandé, Spk013, Kemb 10 and SPK004 grown in western Kenya, with values of 4.2, respectively; 3.1; 3.5 and 2.8 [36]. However, in the Yan Shu1 and White Delite varieties grown in eastern Congo, the number of tubers is higher at 15.5 and 14 roots/plant, respectively [37]. This large number obtained in these varieties grown in the Congo relative to the yields of our study would be related to soil type, rainfall and especially genetic variability. Thus, the differences in the weight and yield of tuberose roots in our study compared to those reported by [38] should be within the different varieties used in these different trials. Work by [39] reported a marketable root yield of 0.72 kg per plant, which is higher than the 0.14 kg/plant

fertiliser added, the efficiency coefficient of fertilisers for different treatments decreases with increasing doses ([32]). These results show the interest and need in low availability of mineral fertilizers to use low doses. This confirms the studies of [33], which obtained similar results in yam culture.

This decline in yield, due to the combination of mineral and organic fertiliser, would also be due to rainfall, which is one of the factors determining the decline in yields of fresh tuberous roots. The effect of fertilisers could have been better if the rains had been regular throughout the cycle. Drought pockets were noted at certain periods of fertilizer intake, resulting in low solubilization and volatilization of the elements, making them less effective. Indeed, our work was conducted on horseback during the small dry season and the small rainy season during which there was not enough precipitation. Treated plants had to suffer from water stress, especially during the first two months after planting and application of fertilizers. Yields of sweet potato are reduced if severe water stress occurs during tuberization [34]. [35] reported that soil moisture status affects plant growth and yield.

and 0.18 kg/plant found for fertilizer application in this study. This same result was observed in Nyagezi marsh conditions by [40], who found that the weight and yield of tuberose roots varied with varieties. This uniformity of the results would be due to the ecophysiology of sweet potato, which requires very sunny exposures for photosynthesis to be done under better conditions; This would allow starch to form in large quantities and subsequently to obtain large tuberous roots. Physiologically, the conversion of CO₂ to carbon hydrate was well established.

The crop index showed significant differences between varieties. Current results indicate that the crop index is low. This could be due to poor separation or poor translocation of dry matter to storage roots, as claimed by [41].

Results suggest that the increase in the fresh weight of aerial biomass at harvest may have

contributed to the decline in root yield and, in turn, to the decline in the crop index, as the higher fresh weight of aerial biomass could be influenced by a high translocation of vegetative biomass assimilates to the detriment of tuberous roots. Some authors, including [42], have shown a correlation between yield and dry

5. CONCLUSION

The results showed that the effects of chemical fertilizer treatments, manure and their combination with mineral fertilizers did not influence the measured parameters. The results also showed that fertilizers improved all the agronomic parameters of sweet potato compared to the control treatment during two years of experiment. However, the Irene variety with a mean weight of 180 g tuberous roots was

CONFLICT OF INTERESTS

ACKNOWLEDGEMENTS

The authors thank the researchers from Jean Lorougnon Guédé University for their collaboration and the National Center of Agronomic Research (CNRA) for their financial

REFERENCES

1. Vernier P. et Varin D. La culture de la patate douce. Agriculture et Développement ; 1994.
2. FAOSTAT. <http://faostat.fao.org/site/567/default.aspx#ancor>; 2017 (accédé le 29/10 2017).
3. Brobbrey A. Growth, yield and quality factors of sweetpotato (*Ipomoea batatas* (L) lam) as affected by seedbed type and fertilizer application and utilization of root and tuber crops in africa. Master of philosophy in agronomy department of crop and soil sciences Kwame Nkrumah University of Science and Technology (Kumasi); 2015.

matter in aerial biomass. These differences between varieties could be attributed to variations in the genetic composition of varieties and, therefore, to their photosynthetic sharing habits.

significantly different from the TIB variety with a mean weight of 138.73 g.

These results are based only on a two-year study and for a single site, for a better precision of the effects of each nutrient composing fertilizers, it would be advisable for our results to extend the duration of the test to benefit from the back effects of using these organic fertilizers on soil fertility and sweet potato productivity.

The authors have not declared any conflict of interests.

support. We thank the managers of the experimental station of CNRA of Bouaké, where the experiments were set up. We also thank Dr DIBI Konan Evrard Brice (CNRA of Bouaké) for the statistical analysis and Dr BAKAYOKO Sidiky for the valuable remarks and criticisms.

4. Katherine L. A. La production biologique de la patate douce ; 2005.
5. Dibi KEB., Essis B., N'Zué B., Kouakou AM. & Zohouri GP. Appui à la promotion de la patate douce à chair orange en Côte d'Ivoire. Rapport final avenant du Projet / Agriculture familiale pour la nutrition et l'équité de genre (CHANGE). CNRA Bouaké ; 2016.
6. Buresh IJ., Sanchez PA. & Calhoun F. Replenishing soil fertility in Africa: *SSSA Special Publication 51*, Soil Science Society of America American Society of Agronomy Madison, Wisconsin, USA; 1997.
7. Palm CA., Myers RJK. & Nandwa SM. Combined use of organic and inorganic

- nutrient sources for soil fertility maintenance and replenishment. In: Buresh R.J, Sanchez P.A. and Calhoun (Eds.), Replenishing soil fertility in Africa. ASA, CCSA, SSSA Special Republication 51, Madison, WI, USA; 1997: 193-217.
8. **Giller E., Cadisch G. & Mugwira LM.** Potential benefits from interaction between organic and inorganic nutrient sources. *In: Fertility Research for maize-based farming systems in Malawi and Zimbabwe.* (Ed) S.R. Waddington ; 1998 : 155-158.
 9. **N'zi JC., Kouamé C., N'guetta ASP., Fondio L., Djidji AH. & Sangaré A.** Evolution des populations de *Bemissia tabaci* Genn selon les variétés de tomate (*Solanum lycopersicum* L.) au Centre de la Côte d'Ivoire. *Scien. & Nat.*, 2010 ; 7 (1) : 31 – 40.
 10. **Traoré K., Sorho F., Dramane DD. & Sylla M.** Adventices hôtes alternatifs de virus en culture de Solanaceae en Côte d'Ivoire. *Agro. Afric.*, 2013 ; 25 (3) : 231 – 237.
 11. **Huber R.** La fumure. Edit. LMZ. Centrale des moyens d'enseignement agricole. Ch. 3052 Zollikofen ; 1988.
 12. **Konan J.** Compétition entre bâti et agriculture dans la conquête des bas-fonds de la ville de Bouaké : le savoir-faire ou les actions stratégiques des citadins agriculteurs pour préserver les espaces agricoles. *Vertigo. La revue électronique en sciences de l'environnement* ; 2017 ; <http://vertigo.revues.org/18302> ; DOI : 10.4000/vertigo.18302.
 13. **Séka K., Diallo AH., Kouassi NK. & Aké S.** Incidence du *Yam mosaic virus* (YMV) et du *Cucumber mosaic virus* (CMV) sur des variétés de *Dioscorea* spp. Cultivées dans les régions de Bouaké et de Toumodi en Côte d'Ivoire. *Int. J. Biol. Chem. Sci.*, 2009; 3 (4): 694 – 703.
 14. **Ettien DJB.** Intensification de la production d'igname (*Dioscorea* spp.) par la fertilisation minérale et l'identification de nouvelles variétés en zone forestière et savanicole de Côte d'Ivoire. Thèse de Doctorat Unique en Sciences de la Terre, Option : Agro-Pédologie. Université de Cocody (Abidjan, Côte d'Ivoire) ; 2004.
 15. **Akassimadou EF. & Yao-Kouamé A.** Caractéristiques morpho-pédologiques et potentiels d'un sol de bas-fonds secondaire développé sur granito-gneiss en région de savane guinéenne (Centre de la Côte d'Ivoire). *Journal of Applied Biosciences* ; 2014 ; 79 : 6968 – 6982 ISSN 1997-5902, 15p. <http://dx.doi.org/10.4314/jab.v79i1.17>.
 16. **Loe A.** Fertilisation et nutrition minérale de la pomme de terre Mulhouse : S.C.P.A (direction technique) ; 1977.
 17. **Kebdani B. & Missat L.** Etude de l'influence de type de la fertilisation et l'apport de fumure sur la culture de pomme de terre (*Solanum tuberosum* L) Mémoire de Master II En Agronomie Option production et amélioration végétales Université Abou Bekr Belkaid (Tlemcen) ; 2014.
 18. **Njoku JC., Okpara DA. & Asiegbu JE.** Growth and yield response of sweet potato to inorganic nitrogen and potassium in a tropical ultisol. *Nigerian Agricultural Journal* ; 2001 ; 32: 30-41.
 19. **Rishirmuhirwa T. & Roose E.** Effets des matières organiques et minérales sur la réhabilitation des sols acides de montagne du Burundi. ORSTOM, Montpellier (FRA). *16è Congrès Mondial de la Science du Sol*, 20-26 août 1998, Montpellier (France) ; 1998.
 20. **Mukhtar AA., Tanimu B., Arunah UL. & Babaji BA.** Evaluation of the agronomic characters of sweet potato varieties grown at varying levels of organic and inorganic fertilizer. *World journal of agricultural sciences* ; 2010 ; 6 (4): 370-373.
 21. **Kaboré B.** Les contributions en azote des légumineuses, des amendements

- organo- minéraux dans les systèmes de culture : Impact sur les rendements des céréales et sur la fertilité des sols à long terme. Mémoire d'Ingénieur du Développement Rural/Option Agronomie IDR/UFB (Burkina Faso) ; 2004.
22. **Bakayoko S., Tschannen A., Nindjin C., Dao D., Girardin O. & Assa A.** Impact of water stress on fresh tuber yield and dry matter content of cassava (*Manihot esculenta Crantz*) in Côte d'Ivoire. *African Journal of Agricultural Research*; 2009; 4: 21-27.
 23. **Borah KK., Bhuyan B. & Sarma H.** Assessment of Soil Fertility Status in and Around the Tea Gardens of Undivided Darrang District, Assam. *International Journal of Applied Environmental Sciences* ; 2010.
 24. **Zoundi SJ.** Agriculture vivrière : Les Africains confrontés à des choix controversés de modèles agricoles. *Cahiers Agricultures*; 2012; 21 : 366-373.
 25. **Lompo F. & Belem J.** Promotion of orange-fleshed sweetpotato to control vitamin A and antioxidant deficiencies in Burkina Faso, Annual report Year 3 Oct. 11-Sept. 12; 2012.
 26. **Saraswati P. & Antonius S. (2015).** Water Use Efficiency and Yield of Sweetpotato as Affected by Nitrogen and Potassium Application. *Journal of Agricultural Science*; 2015; Vol. 7, No.7. doi:10.5539/jas.v7n7p128
 27. **Onunka NA., Chukwu LI., Mbanasor EO. & Ebeniro CN.** Effect of organic and inorganic manures and time of application on soil properties and yield of sweet potato in a tropical ultisol. *Journal of Agriculture & Social Reseach*; 2012; 12(1): 182-193.
 28. **AVRDC.** Annual report for 1974. Asian Vegetable Research and Development Center, Shanhua, Taiwan; 1975.
 29. **Mulaji KC.** Utilisation des composts de biodéchets ménagers pour l'amélioration de la fertilité des sols acides de la province de Kinshasa. Thèse de doctorat, Gembloux Agro bio tech (République Démocratique du Congo) ; 2011.
 30. **Ojetayo AE., Olaniyi JO., Akanbi WB. & Olabiyi TI.** Effect of fertilizer types on nutritional quality of two cabbage varieties before and after storage. *Journal of Applied Biosciences*; 2011; 48: 3322– 3330.
 31. **Bhardwaj ML., Raj H. & Koul BL.** Yield response and economics of Organics sources and inorganic source in tomato (*Lycopersicon esculentum*), okra (*Hibiscus esculentus*), cabbage (*Brassica oleracea var. botrytis*). *Indian Journal of Agricultural Science*; 2000; 70 (10): 653 – 656.
 32. **Kimuni LN., Mwali MK., Mulembo TM., Lwalaba JWL., Kanyenga LA., Katombe, JL., Mpundu BN., Mubemba M. & Longanza BL.** Effets de doses croissantes des composts de fumiers de poules sur le rendement de chou de Chine installé sur un sol acide de Lubumbashi. *Journal of Applied Biosciences*; 2014; 77 :6509– 6522. <http://dx.doi.org/10.4314/jab.v77i1.4>
 33. **Soro D., Dao D., Carsky R., Asiedu R., Tra TB., Assa A. & Girardin O.** Amélioration de la production de l'igname à travers la fertilisation minérale en zone de savane de Côte d'Ivoire. *Agronomie Africaine* (numéro spécial, Atelier National sur l'igname d'octobre 2001) ; 2003 ; 4 : 39 – 46.
 34. **Degras L.** Sweetpotato (English Eds by Simon Charter) *Macmillan Publishers Ltd.* Oxford, English ; 2003.
 35. **Singer JM. & Munns DN.** *Soils: An introduction.* Macmillan Publishing Company. New York, P 492. Republication 51, Madison, WI, USA; 1987 : 193-217.
 36. **Ndolo PJ., Mcharo T., Carey EE., Gichuki ST., Ndinya C. & Maling'a J.** Participatory on-Farm selection of sweetpotato varieties in western Kenya. *Afr. Crop.Sci.J.*, 2001; 9(1): 41-48.
 37. **Phemba P., Mutombo T., Lutaladio NB. & Carey EE.** Performance et

- stabilité de rendement des génotypes de patate douce dans divers environnements à l'Est du Congo. *Afr. Crop. Sci. J.*, 1998; 6(2) :109-118.
38. **Mulundu M.** Relationship between vine propagation method, and growth and yield in sweetpotato [*Ipomoea batatas* (L.) Lam.]. Master of philosophy in agronomy, Department of plant sciences, University of Zambia (Lusaka); 2014.
39. **Belehu T.** Agronomical and physiological factors affecting growth, development and yield of sweet potato in Ethiopia. PhD Thesis, University of Pretoria (South Africa), 2003.
40. **Shaba M.** Etude comparative de rendement des clones et variétés de patate douce (*Ipomoea batatas*) dans les conditions de marais à Nyangezi, TFE, UEA inédit Bukavu ; 2007.
41. **Yeng SB., Agyarko K., Dapaah HK., Adomako WJ. & Asare E.** Growth and yield of sweetpotato (*Ipomoea batatas* L.) as influenced by integrated application of chicken manure and inorganic fertilizer. *African Journal of Agricultural Research*; 2012; Vol. 7(39):5387-5395. DOI: 10.5897/AJAR12.1447
42. **Sajjapongse A. & Roan YC.** Les facteurs physiques affectant le rendement en racine de la patate douce. In *Patate Douce. CTA / ACCT*. Shanhua ; 1982 : 217-229.

UNDER PEER REVIEW