

1
2
3
4
5
6

Original Research Article Effects of storage and priming on seed germination emergence in soil and embryo culture of *Musa acuminata* Calcutta 4

Formatted: Highlight
Formatted: Highlight
Formatted: Font: Arial, 18 pt, Highlight

ABSTRACT:

Aims: Effects of 3 storage durations, 3 hydro priming protocols and 6 chemical priming protocols on germination emergence in soil (*in vivo*) and embryo culture (*in vitro*) of *Musa acuminata* Calcutta 4 were investigated.

Study design: The experimental design was a completely randomised with three replicates. Analysis of variance was used (P=.05) to test treatment effects in a Completely Randomised design. mean comparison was by LSD.

Place and Duration of Study: This study was carried out for a period of 10 months at the International Institute of Tropical Agriculture High Rainfall Station, Onne, in Rivers State, Nigeria.

Methodology: Seed pre-sowing treatments consisted of 3 storage protocols, 3 hydro priming and 6 chemical treatment protocols. After which treated seeds were divided into two sets. One set was sown directly in soil and the other set subjected to embryo culture technique.

Results: Seeds sown in soil immediately they were extracted had significantly higher germination emergence than stored seeds. Emergence Germination declined by 20% and 23% after 2 weeks and 4 weeks of storage respectively. For embryo culture, seeds stored for 2 weeks had significantly higher germination (40%) than seeds that were not stored or seeds stored for 4 weeks (38%). Emergence Germination *in vivo* was significantly higher for seeds that were not hydro primed than for seeds hydro primed for 4 days or 8 days. Emergence Germination declined by 33% and 38% in seeds hydro primed for 4 days and 8 days respectively. Hydro priming for embryo culture for 4 days increased germination significantly by 60% compared to those without hydro priming. All the chemicals reduced emergence germination in both soil and *in vitro* procedures except that of Copper oxychloride in embryo culture which increased germination by 18%, compared to the control achieving 47% germination.

Conclusion: Higher germination was recorded with *in vitro* than *in vivo* procedures irrespective of the treatments applied. Perhaps inherent factors in the seed coat and possible interactions in soil may account for the poor emergence germination exhibited *in vivo* and will require further investigation.

Keywords: [*Musa acuminata* Hydro-priming chemical-priming *in vivo*; *in vitro*].

Formatted: Highlight

7
8
9
10
11
12
13
14
15
16

1. INTRODUCTION

Seed production is required in plantain and banana (*Musa* spp.) mainly for breeding purposes. At maturity, *Musa* seeds are black or dark brown stony bodies. The seed has a rough seed coat [1]. It contains an embryo, which is embedded in a copious endosperm and chalazal mass [2]. Seeds vary in size (about 4-6 mm), colour (brown or black) and shape (angular or globose). Seed shape varies due to compression between neighbouring seeds [1]. The structure of the *Musa* seed is complex, hence making germination very difficult [1,3, 4]. It was found that seed viability was also affected by moisture content, oxygen and

17 temperature [5]. The presence of a semi-permeable inner membranous seed coat restricts
18 the movement of moisture and oxygen into the embryo. In addition, while a hard seed coat
19 provides effective protection during maturation, dispersal and dormancy, it hampers
20 germination because the embryo requires extra energy to rupture the seed coat.

21 Seed set in *Musa* spp. varies greatly among seed-fertile cultivars. This limitation in variable
22 seed set is further compounded by an extremely low rate, slow and non-uniform
23 emergencegermination in soil thus making creation of new cultivars and other breeding
24 activities of plantains and bananas difficult [1,6]. In fact seed emergencegermination
25 especially of hybrid seeds in soil is reported to be less than 1% [7]. While seeds of *Musa*
26 *balbisiana* (with the B genome) readily germinate in culture and soil [8], seeds of *M.*
27 *acuminata* (with the A genome) and most interspecific hybrids have poor germination and
28 are not viable especially if the fingers are left to over ripen (blackened or rotten) before
29 extraction [8]. However, a major source of pollen in plantain & banana breeding is the wild
30 diploid accession, *Musa acuminata* Calcutta 4, which though agronomically poor, produces
31 abundant and viable pollen [9]. It is resistant to black Sigatoka disease, but produces non-
32 parthenocarpic fruits due to the presence of two complementary recessive genes for
33 parthenocarpy [10]. It is important in germplasm enhancement because it serves as a source
34 of plantain alleles and resistance to black Sigatoka disease [11].

35 The parental differences between seeds from *M. balbisiana* and other accessions could be
36 histological, physiological or genetic in nature. Another study have identified single sequence
37 repeats (SSRs) that could help in understanding the divergence between *M. acuminata* and
38 *M. balbisiana* [12]. For example, in *Vicia* spp., germination ability has been linked to
39 permeability of the seed coat, a condition that was found to be controlled by a two-gene
40 system [13]. Similarly seed coat permeability in cotton *Gossypium hirsutum* L. was found to
41 be controlled by a single gene [14].

42 Due to their triploid nature, plantains and bananas are almost completely female sterile,
43 resulting in low seed set upon pollination and poor seed quality. If seeds from Tetraploid (4x)
44 and Diploid (2x) crosses can be made to germinate at relatively high frequency when planted
45 in soil, they can be grown in environments other than those under which they were produced
46 provided that an efficient method for seed germination is available. Perhaps seeds obtained
47 from the 4x - 2x crosses, if treated with some chemicals and sown in the soil could have
48 relatively high emergencegermination.

49 In order to increase germination, seeds are scarified by physical or chemical means to
50 permit imbibition and improve the rate of germination or shorten the time required for
51 germination [15]. Unfortunately, some pre-sowing treatments such as chipping of testa,
52 scorching and the application of temperature shocks are usually deleterious and often lethal

53 to *Musa* seeds [16]. Other methods used to overcome dormancy in *Musa* seeds include
54 treatment with different concentrations of potassium hydroxide, sulphuric acid and carbon
55 dioxide [17]. The use of potassium hydroxide has also been found to improve germination
56 and emergence of several other crop species [18]. This was demonstrated in oat (*Avena*
57 *fatua* L.) as dormancy was broken and germination was significantly enhanced. Other
58 studies have pointed out that seed germination and seedling vigour can be improved as a
59 result of various hydro priming protocols [19,20,21,22,23,24].

60 Since emergencegermination of plantain and banana seeds in soil is abysmally low, hybrid
61 seed propagation in *Musa* is usually difficult [4,7]. Therefore, the regeneration of hybrid
62 seedlings has relied more on *in vitro* culture of excised embryos [25], a technical and
63 relatively more expensive procedure than planting in soil. Improving *Musa* seed
64 emergencegermination in the soil being the natural medium of plant growth (designated as *in*
65 *vivo*) could accelerate hybrid development, selection, and evaluation of several cultivars in
66 *Musa* breeding efforts that meet the production and consumption requirements of target
67 populations as advocated by [26]. In addition enhanced seed germination would encourage
68 seed storage in gene banks [4] for germplasm preservation. Moreover, it would facilitate the
69 production of large number of segregating planting materials and decentralize hybrid
70 distribution for research and production under various agro-ecologies at a relatively low cost.

71 The main objective of this study was to investigate how to enhance emergencegermination
72 of *Musa acumunata* Calcutta 4 (AA genome) seeds when planted in soil (*in vivo*) and by
73 embryo culture technique (*in vitro*). Specifically:

- 74 I. whether varying storage durations will affect emergencegermination of seeds planted in
75 soil (*in vivo*) and embryo culture (*in vitro*) differently;
- 76 II. determine how hydro priming protocols will affect emergencegermination of seeds
77 planted in soil (*in vivo*) and embryo culture (*in vitro*); and
- 78 III. find out how chemical priming with various chemicals will affect emergencegermination
79 of seeds planted in soil (*in vivo*) and embryo culture (*in vitro*).

80

81 **2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY**

82 **2.1 Experimental Site and Planting Material**

83 This study was carried out at the International Institute of Tropical Agriculture (IITA) High
84 Rainfall Station, Onne (4°51'N, 7° 03'E, 10 m above sea level), in Rivers State, south-eastern
85 Nigeria. The rainfall pattern is monomodal, distributed over a 10 month period from February
86 through December, with an annual average of 2400 mm. Relative humidity remains high all

87 year round with mean values of 78% in February, increasing to 89% in the months of July
88 and September. The mean annual minimum and maximum temperatures are 25°C and 27°C,
89 respectively, while solar radiation/sunshine lasts an average of 4 hours daily [27].
90 Seeds of the wild banana *M. acumunata*, Calcutta 4 (diploid AA) which is resistant to black
91 Sigatoka disease were used to evaluate the effect of different seed treatments on direct seed
92 emergencegermination in soil (*in vivo*) and on embryo culture (*in vitro*). Bunches were
93 harvested when the fruits of the proximal nodal cluster (first hand) had reached physiological
94 maturity. Harvested bunches were ripened with ethylene for four 4 days, after which the
95 seeds were extracted mechanically, washed and air-dried.

Formatted: Highlight

96 **2.2 Treatments and Experimental Details**

97 Seed pre-sowing treatments consisted of 3 storage protocols, 3 hydro priming and 6
98 chemical treatment protocols. Seed pre-sowing treatments consisted of the under listed
99 protocols, after which treated seeds were divided into 2 sets. One set was sown directly in
100 soil (*in vivo*) and the other set subjected to embryo culture technique (*in vitro*) after embryo
101 rescue. Each set of treatments was replicated 3 times, in a completely randomized design.

102 **Treatment protocols**

- 104 1. Three storage protocols of seed in transparent air-tight plastic jars at ambient
105 temperature for 0 (sowing immediately on extraction), 2 weeks and 4 weeks after
106 extraction of seeds;
- 107 2. Three hydro priming protocols, i.e., soaking of seeds in water for 0 (no soaking in water
108 before sowing), 4 days and 8 days before sowing. Seeds were soaked in water with 2
109 drops of Tween 80 (Sorbitan), agitated initially and allowed to stand for 24 hrs, with
110 changes of solution (at 24 hrs intervals) for the different soaking durations.
- 111 3. Six chemical treatment protocols, with copper-oxychloride (0.052 M), 25% sulphuric acid
112 (0.23 M), silver nitrate (0.06 M) plus streptomycin sulphate (0.0002 M), hydrogen
113 peroxide (0.1 M), potassium nitrate (0.01 M) and water (control). Seeds were soaked in
114 chemical solution with 2 drops of Tween 80 (Sorbitan), agitated initially and allowed to
115 stand for 24 hrs.

116 **2.3 Crop Conduction**

117 **2.3.1 Planting in soil – (*in vivo*)**

118 On completion of treatment protocols, seeds were immediately washed with tap water and
119 sown at a rate of 11 seeds per pot, in perforated plastic pots (16 cm x 13 cm x 4.9 cm), three-

120 quarters filled with soil (soil, dried palm fibre and dried poultry manure in a 7:3:1 ratio).
121 Watering of sown seeds was carried out as required. Germination was considered to have
122 occurred when the plumule emerged about 1cm above the soil level.

123 **2.3.1 Embryo culture – (*in vitro*)**

124 Treated seeds were subjected to *in vitro* culture [28]. Seeds were surface sterilized (with 70%
125 methylated spirit for 2 minutes) and transferred to a 1% solution of silver nitrate plus Tween
126 80 for 20 minutes and rinsed in sterilized distilled water. Embryos were excised from seeds
127 using forceps and a scalpel under a stereoscopic microscope in a laminar flow cabinet. The
128 excised embryos were inoculated in culture tubes, each containing 20 ml of modified MS
129 (Murashige and Skoog) medium [29]. The medium was half the standard concentration of
130 MS, supplemented with 3% sucrose, 2 mg 1-1 glycerine, 0.5 mg 1-1 nicotinic acid, 0.5 mg 1-
131 1 pyridoxine, 0.4 mg 1-1 thiamine and 20 mg 1-1 ascorbic acid. Gelrite (Sigma, USA) was
132 used to solidify the medium. Cultures were incubated under continuous light at a temperature
133 of 10°C and examined daily. Germination was recorded when shoots emerged to about 1 cm
134 above the medium.

135

136 **Data Collection and Statistical Analyses**

137 The number of **emergegerminated** seeds in both *in vivo* and *in vitro* procedures was recorded
138 weekly until no further **emergencegermination** occurred. **The experimental design was a**
139 **completely randomised design with treatments replicated 3 times.** Analysis of variance was
140 used to test treatment effects. All data were analysed using the GLM procedure of Statistical
141 Analyses Software and any effects found to be significant have been tested at a significance
142 level of 5% while means were compared using the LSD test at $P \geq 0.05$. Graphs of means
143 with associated standard errors were drawn to show treatment effects.

144

145 **3. RESULTS AND DISCUSSION**


146

147 **3.1. Seed Storage Duration**

148 Best **emergencegermination** of 31% (*in vivo*) was achieved when seeds were not stored.
149 Seeds sown in soil (*in vivo*) that were not stored had significantly higher
150 **emergencegermination** compared to seeds that were stored for **2 weeks** or **4 weeks** (Fig. 1).
151 **EmergenceGermination** declined by 20% and 23% after **2 weeks** and **4 weeks** of storage
152 respectively. However, seeds stored for **2 weeks** did not significantly differ in
153 **emergencegermination** from those stored for **4 weeks**. For embryo culture (*in vitro*), seeds
154 stored for **2 weeks** had 42% germination; significantly higher germination than seeds that
155 were not stored or seeds that were stored for **4 weeks** (Fig. 1). Germination increased by

Formatted: Highlight

156 40% at 2 weeks of storage but declined by 38% beyond 2 weeks at 4 weeks of storage.
157 There was no significant difference in germination between seeds that were not stored and
158 those that were stored for 4 weeks. On the average germination in embryo culture (*in vitro*)
159 was 29% higher than emergence germination in soil (*in vivo*) and was significantly better.
160 Planting immediately after seed extraction was best for soil (*in vivo*) and planting at 2 weeks
161 of storage was best for embryo culture (*in vitro*). This was significantly better and 36% higher
162 than the best soil (*in vivo*) emergence germination.
163 Emergence Germination in soil (*in vivo*) was significantly higher in seeds that were not stored
164 than for seeds stored for 2 weeks or 4 weeks (Fig. 1). Storage of seeds of pearl millet
165 (*Pennisetum glaucum*, Slapf & Habbnd) for 10 days reduced soil germination, and declined
166 further after 14 days of storage [30]. A major limitation of stored seeds may result from the
167 seed coat. The seed coat contains ferulic acid and polyphenolic compounds that affect soil
168 germination by restricting the embryo development [1]. Germination in soil (*in vivo*) was
169 reported to be as erratic as <5%-23% over 20 days [31]. In this study, *in vivo*
170 emergence germination ranged from 24%-31% for storage duration. For embryo culture (*in*
171 *vitro*) germination was significantly higher in seeds stored for 2 weeks, and declined
172 thereafter. Perhaps lower moisture content could have played a role in the higher level of
173 emergence germination observed after 2 weeks of storage and the decline beyond 2 weeks
174 to 4 weeks [31]. While storage for 2 weeks increased *in vitro* germination, it reduced the rate
175 of *in vivo* emergence germination, although the data suggests that the seeds in both
176 instances retained viability for as long as 4 weeks. For best results, seeds to be planted in
177 soil (*in vivo*) should be planted as soon as they are excised from the fruits while seeds for
178 embryo culture (*in vitro*) should be stored for 2 weeks before use.
179




180
181
182 Fig. 1. Effect of storage duration on *in vivo* and *in vitro* percentage germination
183 germination of *Musa acuminata* Calcutta 4
184

3.2. Hydro Priming Protocol

185 When seeds were sown in soil (*in vivo*), emergence germination was significantly higher for
186 seeds that were not hydro primed than for seeds that were hydro primed for 4 days or 8 days
187 (Fig. 2). Emergence Germination declined by 33% and 38% in seeds hydro primed for 4 days
188 and 8 days respectively. Emergence Germination of seeds that were hydro primed for 4 days
189 did not significantly differ from those that were hydro primed for 8 days. In embryo culture (*in*
190 *vitro*) germination of those not hydro primed was significantly lower than those hydro primed
191 for 4 days and 8 days (Fig. 2). Hydro priming for 4 days increased germination significantly
192 by 60% compared to those without hydro priming but reduced germination by 3% when
193 hydro primed for 8 days compared to that hydro primed for 4 days. There was no significant
194 difference in germination between seeds hydro primed for 4 days and that hydro primed for 8
195 days (Fig. 2). Comparing both procedures, again on the average, germination in embryo
196 culture (*in vitro*) was 29% higher than in soil (*in vivo*) and was significantly better. Planting
197 without hydro priming was best for soil (*in vivo*). Planting after 4 days hydro priming was best

198 for embryo culture (*in vitro*) resulting in a 15% higher germination than the best soil (*in vivo*)
199 emergencegermination.
200 Seeds that were not hydro primed gave significantly higher *in vivo* emergencegermination
201 compared to seeds hydro primed for 4 days and 8 days (Fig. 2). Sowing of hydro primed
202 seeds in the soil led to decline in the rate of emergencegermination. It is likely that hydro
203 priming reduced the protection level provided by the seed coat under normal circumstances.
204 Hence, the seeds became exposed to microbial attack and other detrimental soil factors [31]
205 or the moisture content of the seeds could have exceeded the optimum required for
206 germination over the 4 days - 8 days period [31].
207 In embryo culture (*in vitro*) seeds hydro primed for 4 days gave significantly higher
208 germination than seeds hydro primed for 8 days. Hydro priming for 5 days was reported to
209 have increased *in vitro* germination in *M. balbisiana* more than hydro priming for 3 days and
210 9 days [32]. They found germination after hydro priming for 5 days was 94% (*in vitro*) within 7
211 days compared to 50% after 54 days for greenhouse-sown seeds (*in vivo*). Similarly, in this
212 study, hydro priming only increased *in vitro* but not *in vivo* emergencegermination. Hydro
213 priming is also thought to increase free-radical scavenging enzyme activity, counteracting the
214 effects of lipid peroxidation and reducing leakage of metabolites [33, 34,35]. It is likely that by
215 the fourth day of soaking, the embryos could have become metabolically active for rapid
216 germination under aseptic conditions. The reasons for the difficulty in achieving high
217 emergencegermination of *Musa* under natural conditions (*in vivo*) need to be further
218 investigated.




219 Fig. 2: Effect of hydro-priming on *in vivo* and *in vitro* germination of
 220 *Musa acuminata* Calcutta 4
 221
 222

223 3.3. Chemical Priming Protocol

224 Emergence Germination of seeds primed with sulphuric acid was significantly lower than
 225 those of seeds primed with other chemicals or the control in both *in vivo* and *in vitro*
 226 procedures (Fig. 3). Other than this, for soil, (*in vivo*) there was no significant difference in
 227 emergence germination of seeds primed with other chemicals and the control (water). In fact,
 228 all the chemicals reduced emergence germination by 6% (KNO₃), 8% (CuOCl), 11% (AgNO₃
 229 + Streptomycin), 21% (H₂O₂) and significantly by as much as 63% (H₂SO₄) compared to the
 230 control. However, in embryo culture (*in vitro*), copper oxychloride increased germination by
 231 18% compared to the control achieving 47% germination. All other chemicals reduced
 232 germination by 11% (AgNO₃ + Streptomycin), 21% (KNO₃), 23% (H₂O₂) and significantly by
 233 50% (H₂SO₄) compared to the control. Emergence Germination in all chemical priming
 234 treatments was lower for *in vivo* than for *in vitro* procedures. The best *in vitro* germination

235 was with copper oxychloride priming which was significantly better and 46% higher than the
236 best *in vivo* emergence germination, the control -priming with water (32% germination).
237 Chemical priming of seeds did not improve germination of *M. acuminata* Calcutta 4, although
238 their efficacy has been reported in several other crop species [36]. This study has shown
239 that sulphuric acid priming at the concentration used, significantly reduced both *in vivo* and
240 *in vitro* germinations (Fig. 3). Emergence Germination *in vivo* did not significantly differ
241 between chemically primed seeds and the control. This suggests that the chemical priming
242 at the concentrations used, did not improve *in vivo* emergence germination. However,
243 significantly higher germination was reported with application of copper oxychloride in hybrid
244 plantain seeds when applied at low concentrations as a fungicide to soil [6] rather than as a
245 seed primer. This perhaps indicates that soil treatment rather than seed treatment could be
246 an avenue for further exploration. They also identified the average weather conditions such
247 as air temperature and sunshine at the time of seed treatment as a significant factor
248 influencing the germination of the seeds.
249



250

251 **Fig. 3: Effect of chemical priming on *in vivo* and *in vitro* germination of *Musa***
252 ***acuminata* Calcutta 4**

253
254 For *in vitro* germination, priming with copper oxychloride produced a significantly higher
255 germination than other chemicals implicating perhaps its anti-fungal properties relative to the
256 other chemicals in embryo culture.

257 Consistently higher germination was recorded with *in vitro* than *in vivo* procedures
258 irrespective of the treatments applied. Almost all studies conducted have reported the same
259 trend [32,37,38,39].

260 **4. CONCLUSION**

261 Consistently higher germination was recorded with *in vitro* than *in vivo* procedures
262 irrespective of the treatments applied. Perhaps inherent factors in the seed coat and
263 possible interactions in soil may account for the poor emergence germination exhibited *in*
264 *vivo* and will require further investigation. In this study, sowing seeds extracted immediately
265 without hydro priming was best for (*in vivo*), while for embryo culture (*in vitro*), storage for 2
266 weeks and hydro priming for 4 days or priming with copper oxychloride gave the best results.

267

268

269

REFERENCES

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

1. Graven P, De Koster CG, Boon JJ, Bouman F. Structure and macromolecular composition of seed coat of the *Musaceae*. *Annals of Botany*. 1996. 77:105-122.
2. Cronquist A. An integrated system of classification of flowering plants. Columbia University Press, New York. 1981.
3. Kiew R. Notes on the natural history of the Johore banana, *Musa gracilis* Holtum. *Malayan Nature Journal*. 1987. 41:239-248.
4. Chin, HF Germination and storage of bananas seeds. In: frison, ea., horry jp. de waele d. (eds) new frontiers in resistance breeding for nematode, fusarium and Sigatoka. INIBAP Proceedings of workshop held in Kuala Lumpur, Malaysia. 1995. 218-227.
5. Owen EB. The storage of seeds for maintenance of viability. Commonwealth Agricultural Bureaux. *Field Crops Bulletin*. 1956. 43:81.
6. Dumpe BD, Wokoma ECW. Factors influencing germination of hybrid plantain seeds in soil. *Plant Foods for Human Nutrition* 2003. 58: 1-11,
7. Ortiz R, Vuylsteke D. Factors influencing seed set in triploid *Musa* spp. L. and production of euploid hybrids. *Annals of Botany*. 1995. 75:151-155.
8. Simmonds NW. The germination of banana seeds. *Tropical Agriculture Trinidad*. 1952. 29:2-16.
9. Swennen R, Vuylsteke D. Breeding black Sigatoka resistant plantains with a wild banana. *Tropical Agriculture Trinidad*. 1993. 70 (1):74-77.
10. Simmonds NW. Segregations in some diploid bananas. *Journal of Genetics*. 1953. 51: 458-469.
11. Vuylsteke D, Ortiz R. Plantain-derived diploid hybrids (TMP2x) with black Sigatoka resistance. *American Journal of Horticultural Science*. 1995. 30:147-149.
12. Ravishankar KV, Sampangi-Ramaiah M.H, Ajitha R, Khadke GN, Chellama V. Insights into *Musa balbisiana* and *Musa acuminata* species divergence and

- 295 development of genic microsatellites by transcriptomics approach. *Plant Gene* 2015.
296 4:78-82
- 297 13. Donnelly ED, Watson JE, McGuire, JA. Inheritance of hard seed in *Vicia*. *Journal of*
298 *Hereditary*. 1972. 63:361-365.
- 299 14. Lee JA. Inheritance of hard seed in cotton. *Crop Science*. 1975. 15:149-152.
- 300 15. Lopez JH, Aviles RB. The pre-treatment of seeds of four Chilean prosopis to improve
301 their germination response. *Seed Science and Technology*. 1988. 16:239-246.
- 302 16. Simmonds NW. *The evolution of the bananas*. Longmans, London. 1962.
- 303 17. Simmonds NW. Experiments on the germination of banana seeds. *Tropical Agriculture,*
304 *Trinidad*. 1959. 36 (4):259-273.
- 305 18. Gao YP, Zheng GH, Gusta LV. Potassium hydroxide improves seed germination and
306 emergence in five native plant species. *American Journal of Horticultural Science*.
307 1998. 32 (2):274-276
- 308 19. McDonald MB. Seed priming. In: black m, bewley jd. (eds.). *Seed technology and its*
309 *biological basis*. Sheffield Academic Press, England, 2000. 287–325.
- 310 20. Halmer P. Methods to improve seed performance in the field. In: benech-arnold rl,
311 sanchez r. (eds.). *Handbook of seed physiology*. Food Product Press, New York.
312 2004. 125–156
- 313 21. Ghassemi-Golezani K, Sheikhzadeh-Mosaddegh P, Valizadeh M. Effects of hydro-
314 priming duration and limited irrigation on field performance of chickpea. *Res. J. Seed*
315 *Sci*. 2008. 1:34–40
- 316 22. Ghassemi-Golezani K, Chadordooz-Jeddi A, Nasrullahzadeh S, Moghaddam M.
317 Effects of hydro-priming duration on seedling vigor and grain yield of pinto bean
318 (*Phaseolus vulgaris* L.) cultivars. *Not. Bot. Hor. Agro. ClujNap*. 2010. 38:109–113.
- 319 23. Ghassemi-Golezani K., Hosseinzadeh-Mahootchy A, Zehtab-Salmasi S, Tourchi M.
320 Improving field performance of aged chickpea seeds by hydro-priming under water
321 stress. *Int. J. Plant Animal Environ. Sci*. 2012. 2:168–176.
- 322 24. Ghassemi-Golezani K, Hosseinzadeh-Mahootchy A. Influence of hydro-priming on
323 reserve utilization of differentially aged chickpea seeds. *Seed Technology*, 2013.
324 135: (1):117-124
- 325 25. Vuylsteke D, Swennen R, De Langhe E. Tissue culture technology for the improvement
326 of African plantains. In: fullerton ra, stover rh. (eds.). *Sigatoka leaf spot diseases of*
327 *bananas INIBAP Proceedings*, San José, Costa Rica. 1990. 316-337.
- 328 26. Coffman, WR, Smith ME. Role of public, industry and international research centre
329 breeding programs in developing germplasm for sustainable agriculture. In: sleeper,
330 da, barker, tc, bramel-cox, pj. ((eds) *plant breeding and sustainable agriculture,*
331 *considerations for objectives and methods*, CSSA Special Publication. 1991. 18:1-9.
- 332 27. Ortiz R, Austin PD, Vuylsteke D. IITA High Rainfall Station African humid forest.
333 *American Journal of Horticultural Science*. 1997. 32: 969-972.
- 334 28. Vuylsteke D, Swennen R, De Langhe E. Tissue culture technology for the improvement
335 of African plantains. In: fullerton ra, stover rh (eds.). *Sigatoka leaf spot diseases of*
336 *bananas. INIBAP Proceedings*, San José, Costa Rica. 1990. 16-337.
- 337 29. Vuylsteke D. Shoot-tip culture for the production, conservation and exchange of *Musa*
338 germplasm. *Practical manuals for handling crop germplasm in vitro* 2. International
339 Board for Plant Genetic Resources, Rome. 1989. 56p
- 340 30. Singh J, Govila OP, Agrawal PK. Preliminary results from a study of seed germinability
341 of pearl millet (*Pennisetum typhoides* L.) F₁ hybrids and their parents during accelerated
342 ageing test. *Seed Science and Technology*. 1988. 16:685-692
- 343 31. Vineesh PS, Skaria R, Mukunthakumar S, Padmesh P, Decruse SW. Seed germination
344 and cryostorage of *Musa acuminata* subsp. *Burmannica* from Western Ghats. *South*
345 *African Journal of Botany*. 2015. 100:158–163

- 346 32. Afele JC, De Langhe E. Increasing *in vitro* germination of *Musa balbisiana* seed. Plant
347 Cell, Tissue and Organ Culture. Kluwer Academic Publishers, Netherlands. 1991. 27:
348 33-36.
- 349 33. McDonald MB. Seed deterioration: physiology, repair and assessment. Seed Sci.
350 Technol. 1999. 27: 177-237
- 351 34. Hsu CC, Chen CL, Chen JJ, Sung JM. Accelerated aging-enhanced lipid peroxidation in
352 bitter gourd seeds and effects of priming and hot water soaking treatments. Sci. Hortic.
353 2003. 98: 201-212.
- 354 35. Wang HY, Chen CL, Sung JM. Both warm water soaking and solid priming treatments
355 enhance anti-oxidation of bitter gourd seeds germinated at sub-optimal temperature.
356 Seed Sci. Technol. 2003. 31 (1): 47-56. DOI: <https://doi.org/10.15258/sst.2003.31.1.06>
- 357 36. Copeland LO, McDonald MB. Principles of seed science and technology. 3rd edition.
358 Macmillan, New York. 1995. 409p.
- 359 37. Chin HF. Germination and storage of banana seeds. In: Frison AE, Horry JP, De Waele
360 D. (eds.). New Frontiers in Resistance Breeding for Nematodes, Fusarium and
361 Sigatoka. INIBAP, Montpellier, France. 1996. 218-227.
- 362 38. Uma S, Lakshmi S, Saraswathi MS, Akbar A, Mustaffa MM. Embryo rescue and plant
363 regeneration in banana (*Musa* spp.). Plant Cell Tissue Organ Cult. 2011. 105-111.
- 364 39. Uma S, Lakshmi S, Saraswathi MS, Akbar A, Mustaffa MM. Plant regeneration through
365 somatic embryogenesis from immature and mature zygotic embryos of *Musa acuminata*
366 ssp. burmannica. In Vitro Cell Dev. Biol. Plant. 2012. 48, 539-545.

UNDER REER REVIEW