

PREVALENCE OF GEOHELMINTH PARASITES OF HORSES IN DUTSINMA METROPOLIS, KATSINA STATE NIGERIA

ABSTRACT

This study was carried out to investigate the prevalence of geohelminth parasites of horses in Dutsinma metropolis between May to August, 2016. Fresh fecal sample were obtained from the rectum of 48 randomly selected horses. A floatation technique was used for the lab work experiment. From the result, out of the total sample collected 6 (12.50%) were positive with eggs of two parasites *Strongyles* and *Oxyuris spp* with the prevalence of (6.25%) for each. Among the three points of sample collection, the highest prevalence was recorded at Kadangaru (20.00%) followed by Unguwa Yandaka (18.18%) and no case recorded at Hayingada. X^2 cal = 36.84 at $P=0.05$ and 2df. The highest prevalence was recorded among the males (18.52%) while only 1 female was affected with the prevalence of (4.76%) X^2 cal = 36.57 at $P=0.05$ and 2df. Prevalence was recorded in adult of 14.29% while young horses recorded a prevalence of 10.00%, X^2 cal = 36.84 at $P=0.05$ *Strongyles* and *Oxyuris spp* were recognized as important geohelminth parasites in Horses in Dutsinma Local Government Area of Katsina State with various rates of infection (12.50%). Managerial control tools should be integrated to improve prevention of geohelminth parasites by communication, information and health education.

Key words: Prevalence, Fecal samples, Geohelminth, Parasites, Horses

Introduction

Geohelminth parasites are soil-transmitted helminths which are groups of parasitic worms causing human and other Animals infection through contact with parasite eggs or larvae that thrive in the warm and moist soil of the world's tropical and subtropical countries. As adult worms, the soil-transmitted helminths live for years in the human gastrointestinal tract. More than a billion people are infected with at least one species of geohelminth parasites worldwide. They are considered together because it is common for a single individual, especially a child living in a less developed country, to be chronically infected with all three worms. Such children have malnutrition, growth stunting, intellectual retardation, and cognitive and educational deficits.

Parasitic diseases are the major obstacle in the growth and development of animal health all over the world. Horses, among most domestic animals are reported to be more susceptible to large number of parasites and may harbour different species at any time. An apparently healthy horse can harbour over one half million geohelminth parasites, trematodes, cestodes and nematodes (Martins *et al.*, 2009). This is because, the gastrointestinal tract provides favorable environment for the survival and proliferation of many of these parasites.

Horses (*Equus ferus caballus*) are one of the two extant sub species of horse *Equus ferus* (Getachew, *et al.*, 2010). They evolve over 45-55 million years. Humans began to domesticate horse around 4000 B C. Horses and humans interact in a wide variety of sport competition and man competitive activities.


Intestinal parasites such as helminths usually produce insidious diseases in animals. Infected horse may show signs of weakness, emaciation, restlessness, unthriftiness, diarrhea, anemia and sometimes intestinal obstruction or perforation.

The aim of this Study is to determine the Prevalence of geohelminth Parasites of Horses in Dutsinma Metropolis, Katsina State with the following objectives:


- To determine the prevalence of geohelminth parasites of horses in Dutsinma metropolis
- To identify the geohelminth parasites among horses in the study area
- To isolate the geohelminth parasites species in the Horses

STUDIED AREA

52 Dutsinma (Lat 12° 27' 01.18" N/ Long 7° 29'.29" E), Katsina State, Nigeria Dutsinma Local
 53 Government covers an area of 527 km squares and has a population of 16971, at the 2006 census.
 54 Total annual rainfall around Dutsinma is about 800mm Retrieved 10th July 2011 from
 55 www.albarkablog.com. The inhabitants of the local government are predominantly Hausa and Fulani
 56 by tribe. Their main occupations are farming and animal rearing.
 57



58 **Fig 1:** Map of Katsina locating Dutsinma at the centre.
 59
 60



61 **Fig.2: Map of the Study Area showing the**
 62 **Sampling Points.**
 63 **Source:- Cnes / Spot Image 2014.**

64 **Fig. 2 Map of Dutsinma**

65 **Sample Population**

66 A total of 48 faecal samples of horses were collected from different places within Dutsinma
 67 metropolis including Unguwar Yandaka, Hayin gada, and Kadangaru areas

68 **Sample Collection**

Samples were collected between the periods of May to August, 2016. Faecal samples were randomly collected from the rectum of horses in each place of the study area during the early hours of the day while owners restrained the animals (Ayele, 2006). Aseptic measures were maintained during collection. Each sample was collected into a clean polythene bag while wearing gloves, labeled, noting the sex and age of the animal. These samples were immediately conveyed to the Biology Laboratory of Department of Biological Science, Federal University Dutsinma for examination.

Sample Examination

All faecal samples collected from the rectum of horses were screened eggs/oocysts of the parasite using the floatation technique. A technique with a principle where the floatation medium (Saturated salt NaCl solution) aids the rise of eggs and/or oocysts of geohelminth parasites by virtue of the medium's high specific gravity, allowing the faecal material to settle to the bottom.

Preparation of Floatation Medium

To prepare the floatation medium, 10grams of NaCl was dissolved in 500mL distilled water contained in a beaker. A stirring rod was used to stir the mixture until the NaCl dissolved completely. The solution was made up to 1½ liter by addition of distilled water (Jorgen and Brain, 1994). For each of the faecal sample collected, 3 grams was weighed into a crucible add 50mL of the floatation fluid was added. This was gently emulsified using a pestle and the resultant faecal suspension was poured into a 60ml universal bottle through a strainer (sieve), more of the solution was added until a convex meniscus was formed. A clean, dry and grease-free slide was placed over the mouth of the universal bottle, ensuring contact with the filtrate. This was allowed to stand for 5 minutes to allow the parasite eggs/oocysts to float. The slide was carefully removed and a cover slip was placed on the slide suspended sample and examined for geohelminth parasite under low (10×) and high (40×) power objectives of a microscope (Yagoob, 2015).

Statistical analysis

The data were analyzed using the chi-square test according to the species, age and body condition of animals. In all cases, the SPSS version 16 was used and the confidence level was held at 95% and the results were considered significant when $P \leq 0.05$.

Results

Prevalence of Geohelminth Parasites

A total of 48 Horses were screened for the research, in which 6(12.50%) were harboring one or more Geohelminth parasites. Among the three point of sample collection, the highest prevalence was recorded at Kadangaru (20.00%) followed Unguwa Yandaka (18.18%) and no case recorded at Hayingada.

Table1. Prevalence of Geohelminth Parasites in the study area

Location	No. screened	No. positive	Prevalence	E	X ²
Hayin Gada	16	0	0.00	0.00	0.00
Unguwar Yandaka	22	4	18.18	0.50	24.50
Kadangaru	10	2	20.00	0.25	12.25
Total	48	6	12.50	0.75	36.75

$X^2_{cal} = 36.75 > X^2_{tab} = 10.60$, therefore significant difference exist at $P \leq 0.05$ and 2df.

KEY

X² = Chi-square, and E = expected value,

Prevalence of Geohelminth Parasites in Relation to Sex

Out of the 48 horses examined, 27 were males and 21 were females. The highest prevalence was recorded among the males (18.52%) while only 1 female was affected with the prevalence of (4.76%).

115 Table2. Prevalence of Geohelminth Parasites in relation to the sex of horses

116

Location	MALE					FEMALE				
	No. scre.	Positive	%	E	X ²	No. scre.	Positive	%	E	X ²
Unguwar Yndaka	12	4	33.33	0.74	14.17	10	0	0.00	0.00	0.00
Hayin gada	9	0	0.00	0.00	3.45	7	0	0.00	0.00	18.05
Kadangaru	6	1	16.67	0.19	0.00	4	1	25.00	0.05	0.00
Total	27	5	18.52		17.62	21	1	4.76	0.05	18.05

117 X² cal = 36.57 > X² tab = 10.60, therefore significant difference exist at P ≤ 0.05 and 2df.

118 Key

119 X²=Chi-square, E = expected value, and % = prevalence and No = number screened

120 **Prevalence of Geohelminth Parasites in Relation to Age**

121 28 (58.33%) adult and 20 (41.67%) young Horses were examined. Slightly highest prevalence
 122 was recorded at adult with the prevalence of 14.29% while low prevalence was founded at
 123 young horses with the prevalence of 10.00%.

124
 125 Table3. Prevalence of Geohelminth Parasites in relation to the age of horses

126

Location	ADULT					YOUNG				
	Adult	Positive	%	E	X ²	Young	Positive	%	E	X ²
Unguwar Yandaka	13	3	23.08	0.43	15.36	9	1	11.11	0.10	8.10
Hayin gada	9	0	0.00	0.00	0.00	7	0	0.00	0.00	0.00
Kadangaru	6	1	16.67	0.14	5.28	4	1	25.00	0.10	8.10
Total	28	4	14.29	0.57	20.64	20	2	10.00	0.20	16.20

127 X² cal = 36.84 > X² tab = 10.60, therefore significant difference exist at P ≤ 0.05 and 2df.

128 Key

129 X²=Chi-square, E = expected value, and % = prevalence

130
 131
 132 **Geohelminth Parasites Identified in the Study Area**

133 Eggs of the parasites as identified in the study area including those of *Strongyles* and *Oxyuris spp*
 134 recorded the same prevalence of (6.25%) each.

135 Table4. Quantitative prevalence of geohelminth parasite identified in the study area

136

Location	<i>Strongyles</i>					<i>Oxyuris spp</i>				
	No. screened	Positive	%	E	X ²	No. screened	Positive	%	E	X ²
Unguwar Yandaka	22	1	4.55	0.06	14.73	22	2	9.09	0.13	26.90
Hayin gada	16	0	0.00	0.00	0.00	16	0	0.00	0.00	0.00
Kadangaru	10	2	20.0	0.13	26.90	10	1	10.00	0.06	14.73
Total	48	3	6.25	0.19	41.63	48	3	6.25	0.19	41.63

137 X² cal = 83.26 is significant difference exist at P ≤ 0.05 and 2df.

138 Key

139 X²=Chi-square, E = expected value, and % = prevalence

140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193

Discussion

This study was designed to investigate the Prevalence of Geohelminth Parasites of Horses in Dutsinma. A total of 48 Horses from Unguwar Yandaka, Hayingada and Kadangaru were screened for the research, eggs of two species of geohelminth parasites were recorded 12.50% (*Strongyles* and *Oxyuris spp*). This agrees with previous studies by (Saeed *et al.*, 2010, and Wannas *et al.*, 2012) which further explained the predomination of Strongyle eggs with prevalence rates of 58.5%, 57.14%, and 66.67%, respectively. The overall geohelminth prevalence appears to be low, in which only 6(12.50%) were harboring one or more Geohelminth parasites. Among the three point of sample collection, the highest prevalence was recorded at Kadangaru (20.00%) followed Unguwa Yandaka (18.18%) and no case recorded at Hayingada, table 1. This is low when compared to that of studies in Ethiopia (Fikru and Bizunesh, 2005) estimating prevalence in horses to be 91%, and similar estimates of 80% (Burden *et al.*, 2010) and 81% (du Toit *et al.*, 2008) in Mexico. Potential variation arising from management differences and differences with respect to topography, climate, animal working conditions and seasonality make direct comparisons difficult. The differences in intensity of infection could be due to variations in parasite biology relating to climatic conditions, pasture infection intensity relating to grazing practices and/or differences in use of anthelmintics; detailed information on such variables would be required to determine their importance

Out of the 48 horses examined, 27 were males and 21 were female horses. The highest prevalence was recorded among the males (18.52%) while only 1 female was affected with the prevalence of (4.76%). This difference in the prevalence is statistically significant ($P < 0.05$), table 2. Which is in line with the work of (Hinney *et al.*, 2011), who reported a higher prevalence of (22.09%) male than that of the female Horses (13.56%).

28 (58.33%) adult and 20 (41.67%) young Horses were examined. Slightly highest prevalence was recorded at adult with the prevalence of 14.29% while low prevalence was founded at young horses with the prevalence of 10.00%, table 3. This is not in line the work of (Marariu *et al.*, 2012) who recorded a slightly higher rate among young (100%) horses than adult (97.8%) This might be associated with apparent inability of the adult younger age groups to develop adequate acquired immunity predisposing them to high risks of severe infection with geohelminth parasites when compared with adult donkeys. Higher infection rates and more severe infections reflect lack of immunity in younger population (Urquhart *et al.*, 1996).

The low prevalence we found (12.50% of horses) is similar to the findings of the 2009 survey from Greece, that found Strongyle eggs in 25.6% of horses (Papazahariadou *et al.*, 2009). But this finding is less than the 78% prevalence that was found in the 1986 survey (Tolliver *et al.*, 1987). It might be due to routine deworming programs that might were introduced to most of the horse population in this study area. This decline in prevalence was statistically significant ($P < 0.05$), it may suggest a trend. This concurs with another study from Louisiana conducted over similar time scale, which demonstrated a decrease in the infestation of most geohelminth species, (Crane *et al.*, 2008).

However, the reasons why the parasites vary between age and sex groups is require further investigation, base on my knowledge this is the first attempt to investigate the geohelminth parasites in Dutsinma metropolis. Such further studies could encourage more specific needs for owner education that could reduce endo-parasites burdens and also improve horse welfare.

Oxyuris equi with prevalence rate of 6.25 % was very low when compared with the work of Tolossa and Fikru *et al.*

Conclusion

It has been deduced in this research that *Strongyles* and *Oxyuris spp* were identified as important geohelminth parasites in Horses in Dutsinma Local Government Area of Katsina State.

Recommendation

- ❖ Based on the results obtained and in order to avoid frequent infection, it is recommended that routine faecal samples go through Parasitological diagnostic techniques to determine cases of geohelminth parasites.
- ❖ Communication information and health education be integrated as preventive tool for horse owners.

194 ❖ Owners of horses should routinely take them to Veterinary Clinics for accurate and
195 appropriate diagnosis.

196

197

198

199 **COMPETING INTEREST**

200 Because many people come in contact with horses, the likelihood of being contaminated with the
201 horses' parasite and spread to other members of the community is relatively high and very significant.
202 Therefore, the attempt to study these parasites is important to the community members in general by
203 bringing to bare the ways of infection or contamination by these parasites.

204 **ACKNOWLEDGEMENT**

205 The authors would like to acknowledge all staff of Federal University Dutsinma and all the reviewers
206 of this paper.

207 **REFERENCES**

208 Abbott EM. Larval Cyathostomosis, In: Part 1 Equine Practice.2008; 20, (3): 56 – 67.

209

210 Ayele CF. Principal health problems of donkeys in Dugda Bora district of Ethiopia. In: Proceedings of
211 the 5th International Colloquium on Working Equines. Sidmouth, 2006 pp. 162- 168.

212

213 Burden FA, du Toit N, Hernandez-Gil N, Prado-Ortiz O, Trawford AF. Selected health and
214 management issues for working donkeys in rural Mexico, Tropical Animal Health and Production.
215 2010 42:597-605.

216

217 Crane M, Khallaayoune K, Christley R, Scantlebury C, Faoud M. The effect of an
218 anthelmintic programme for working equids in Morocco: a comparison of treated and placebo control
219 Populations. In: Proceedings of the 9th International Equine Colic Research, Liverpool, UK, 2008 171-
220 172.

221

222 duToit N, Crewe W. Haemoparasites. In: Blacklock and Southwell, A guide to human
223 parasitology, Laboratory Techniques, 10th edition, H.K. Lewis and Co Ltd, ELBS series; 2008 Pp.
224 177 – 186.

225

226 Fikru R, Bizunesh T. Prevalence of equine gastrointestinal parasites in western highlands of Oromia,
227 Ethiopia, Bulletin of Animal Health and Production in Africa 2005 53: 161-166.

228

229 Getachew AM, Trawford AF, Feseha G , Reid SWJ. Gastrointestinal parasites of
230 working donkeys in Ethiopia, Tropical Animal Health and Production 2010 42: 27-33.

231

232 Hinney B, Wirtherle NC, Kyule M, Miethe N, Zessin KH, Klausen PH. Prevalence of Helminths in
233 Horses in the state of Brandenburg, Germany, Parasitol Res 2011 108: 1083-1091.

234

235 Jorgen S, Brain R. Isolation of Infection. Dictyocaulus larvae from herbage", Vet.
236 Parasitology.1994 16, (18) 16.

237

238 Morariu S, Bogdan AT, Dărăbuș G. Heminth Parasites in Horses from Ten Locations of Timiș
239 County, Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca,
240 2012 69(1/2): 381-384.

241

242 Martins S, Krecek RC, Milne SA. Prevalence and biodiversity of helminth parasites in
243 donkeys from South Africa, Journal of Parasitology 2009 86, 756-762.

244

245 Papazahariadou M, Papadopoulos E, Diakou A, Ptochos S. Gastrointestinal Parasites of Stabled and
246 Grazing Horses in Central and Northern Greece, Journal of Equine Veterinary Science, 2009 29 (4):
247 233-236.

248

- 249 Saeed K, Qadir Z, Ashraf K, Ahmad N. Role of intrinsic and extrinsic epidemiological factors on
250 strongylosis in horses, the Journal of Animal and Plant Sciences,2010 20: 277-280 ISSN: 1018-7081.
251
- 252 Tolliver SC, Lyons ET, Drudge JH. Prevalence of internal parasites in horses in critical tests of
253 activity of parasiticides over a 28- year period (1956-1983) in Kentucky. Vet Parasitol,
254 1987 23, 273-284.
- 255 Urquhart GM, Aremour JV , Duncan AM, Dunn FW, Jennis FW. Veterinary parasitology 2 edition.
256 The University of Glasgw, Black Well Science, Scotland, 1996 pp: 3-137.
257
- 258 Wannas HY, Dawood KA, Gassem GA. Prevalence of Gastro-intestinal Parasites in Horses
259 Condition of Horses and Donkeys in Al Diwaniyah Governorate, AL-Qadisiya Journal of Vet. Med.
260 Sci.2012, 11(1), 2012.
261
- 262 Yagoob G. Seroepidemiology of Neospora Sp. In horses in East Azerbaijan Province of Iran,
263 Research journal of biological sciences,2015 6(5); 224-226.