

## Original Research Article

### Soil fertility as influenced by incorporation of K enriched Azolla

#### Abstract

Use of chemical fertilizer in judiciously without organic manure has created many soil health problems. Biological Nitrogen Fixation (BNF) is a major source of fixed N for plant life and to sustain production and productivity of agricultural crops. All other biofertilizers simply solubilize or mobilize the nutrients that are already present in soils. Whereas the *Azolla* is unique in the sense that it acts as host to the N-fixing cyanobacteria after which it is used virtually as a green manure. An incubation experiment was conducted by growing *Azollae filiculoides* with 2 agriculturally important potassic fertilizers (Potassium Chloride, Potassium shulphate) as main plot in seven concentrations (0, 5, 10, 20, 30, 40 and 50 ppm of K) as sub-plots laid down in split plot design replicated three times. The collected azolla was incorporated with soil at 10 t/ha and maintained at two moisture condition such as 60 and 100 percent and assessed soil fertility by estimating various available plant nutrients and organic carbon status..The mean organic carbon content of the soil was 0.657 and 0.525% by K enriched azolla at 60 and 100% moisture contents respectively..The available N content ranged from 216.2 to 327.3 and 191.1 to 285.3 kg/ha from 0 to 40 ppm of K concentration at 60 and 100 % moisture respectively because the Azolla had a high N content released into the soil after decomposition. Azolla also contributed to the supply of phosphorus, potassium, sulfur, zinc, iron and molybdenum in sufficient amounts in addition to other micronutrients besides addition of nitrogen. Among the various concentration, 40 and 50 ppm K were significantly maintained higher and equal soil available P status of 75.17 and 77.33 kg / ha respectively. The fertilizer,  $K_2SO_4$  with 30, 40 and 50 ppm at 60 % moisture content and 40 and 50 ppm of  $K_2SO_4$  and 40 ppm of KCl produced statistically higher and equal available K in azolla incorporated soil maintained at 100 % moisture. The soil biological health,

mineralization and consequent increase in nutrient status by the application of K enriched Azolla was more under 60 % soil moisture content than fully saturated soil.

**Keywords: Biofertilizer, Azolla, K-enrichment, Soil fertility, Organic carbon and Available NPK.**

## **1. Introduction**

Intensive crop production is the demand of time to feed the vast growing population in India. This has created a pressure to use more chemical fertilizer. Use of chemical fertilizer injudiciously without organic manure has created many soil health problems like low fertilizer use efficiency, poor soil physical condition, reduced water holding capacity, degraded rhizospheric properties, and low fertility (Awodun, 2008). Rice crops remove around 16-17 kg N for the production of each ton of paddy cultivation (Sahrawat, 2000). Most of the rice soils of the world are deficient in N, so, fertilizer N applications are required to meet its N demand (Datta and Buresh, 1989). Generally, urea is applied as the N source for rice production. But the efficiency of added urea-N is very low, due to denitrification, NH<sub>3</sub> volatilization and leaching (De Datta and Buresh 1989). Therefore, alternate sources of N has to be evolved to supply crop demanded N with less or no environmental pollution. The demand of fertilizers and manures is increasing at the present days scenarios to maximize the crop production. Biological Nitrogen Fixation (BNF) is a major source of fixed N for plant life and to sustain production and productivity of agricultural crops. Estimate of global terrestrial showed that, the BNF ranged from 100 to 290 million tonnes of N/year. Of this, 40–48 million tonnes is estimated to be biologically fixed in agricultural crops and fields. BNF is one of the natural sources of nitrogen for rice and *Azolla*-cyanobacteria biomass has been identified as potential source of nitrogen. The integrated nutrient management is to maintain or adjust plant nutrient supply to achieve a given level of crop production by optimizing the benefits from all possible sources of plant nutrients (Subba Rao, 2005).

Organic manures are considered to play a significant role in nutrient contribution. The use of organic fertilizer is a way to improve soil fertility. Azolla can be used as organic fertilizer (Syamsiyah *et al.*, 2015). Subedi and Shrestha (2015) explained that, Azolla does not only increase the productivity of rice but also improve the long-term soil fertility. All other biofertilizers simply solubilize or mobilize the nutrients that are already present in soils. Whereas the *Azolla* is unique in the sense that it acts as host to the N-fixing cyanobacteria after which it is used virtually as a green manure. In the process, it adds not only the biologically fixed N but also the other nutrients absorbed from the soil and present in its biomass. Against the total anticipated biofertilizers demand of 1 million tonne in the country, the current supply position is very low (<10 000 tonnes). The present investigation studied soil fertility improvement by azolla grown under varied K fertilizer commonly used in agriculture.

63

## 2. Materials and Methods

### 2.1. Study site

An incubation experiment was conducted by growing *Azolla filiculoides* with 2 agriculturally important potassic fertilizers (Potassium Chloride, Potassium sulphate) as main plot in seven concentrations (0, 5, 10, 20, 30, 40 and 50 ppm of K) as sub-plots laid down in split plot design replicated three times. The experimental site was Rice Research Station, Ambasamudram located between 8° 42' N and 77° 28' E with an altitude of 64.8 M above mean sea level. The mean annual rainfall received was 913 mm. The experimental soil was acidic in reaction (pH 5.84) and free from salinity (0.08 dS m<sup>-1</sup>) with sandy clay in texture. The organic carbon content was 0.56 %. The soil was low in available nitrogen (210 kg ha<sup>-1</sup>), high in available P (24.5 kg ha<sup>-1</sup>) and medium in available K (150 kg ha<sup>-1</sup>). One gram of Azolla fern was grown in a tray with a dimension of 23 x 15x 6 cm<sup>3</sup> filled with 1.5 litres of potassic solutions (Plate 1) and the fern was collected on 7<sup>th</sup>, 15<sup>th</sup>, 30<sup>th</sup>, 60<sup>th</sup>, 90<sup>th</sup> and 120<sup>th</sup> day after incubation / culturing, rinsed with distilled water and analysed for various

biometric and biochemical parameters. The collected Azolla was incorporated with soil at 10 t/ha and maintained at two moisture condition such as 60 and 100 percent. The Azolla incorporated soil was collected after 30 days of incubation, processed and analyzed for various available plant nutrients and organic carbon status. Organic carbon present in soil /was oxidized/oxidized by chromic acid ( $K_2Cr_2O_7$ ) in the presence of conc.  $H_2SO_4$ . Potassium dichromate on reaction with  $H_2SO_4$  provided nascent oxygen which combined with carbon to form  $CO_2$ . The excess chromic acid left unused by the organic matter /was determined by back titration with 0.5 N ferrous sulphate or ferrous ammonium sulphate using diphenylamine indicator (Walkley and Black, 1934). Available nitrogen in the soil was estimated by alkaline permanganate method (Subbiah and Asija, 1956). Available phosphorus extracted with 0.03 N  $NH_4F$  and 0.025 N  $HCl$ . The amount of P extracted was treated with ammonium molybdate and antimony potassium tartarate and developed colour with ascorbic acid. The intensity of blue colour /was determined colorimetrically at 660nm (Bray *et al.*, 1945). The soil was leached with neutral normal ammonium acetate and the  $K^+$  ions in the exchange sites were replaced by  $NH_4^+$  ions. The  $K^+$  ions in solution was then determined with the flame photometer (Standford and English (1949).

**Data Collection** The soil samples were collected from the incubation bottle after 90 days of incubation were shade dried, processed and sieved through 0.5 and 2 mm sieve for estimating organic carbon and available NPK respectively. **Statistical Analysis** The data were analysed statistically using computer software IRRISTAT (IRRI, 1993). Differences among the mean values of the treatment were compared by the LSD test when the F test from the analysis of variance was significant at the  $p=0.005$  level.

### **3. Results**

### 3.1.Organic carbon

The effect of K fertilizer, its concentration and their interaction on the organic carbon content of Azolla incorporated into the soil at 60 %and 100 % moisture content is presented in Table 1 and 2. The concentration of K fertilizer alone influenced the organic carbon content of the Azolla incorporated at both 60 % and 100 % moisture content.The mean organic carbon content of the soil was 0.657 and 0.525% by K enriched Azolla at 60 and 100% moisture contents respectively. Azolla enriched with 40ppm K solution recorded significantly higher soil organic carbon content of 0.763 and 0.623% respectively at 60 and 100 % soil moisture content (Table 1) which was 42 and 56% more than the Azolla grown under 0ppm K. However, it was on par with 50 and 30ppm of K solution in both the moisture-maintained soil.The 20ppm K registered 0.697 and 0.567% of organic carbon content in soil maintained at 60 and 100% moisture content which were 9 to 10% less than the highest organic carbon maintained by 40ppm of K.

### 3.2.Available N

The main effect of Azolla grown under K fertilizer and its concentration alone significantly influenced the available N content at both 60 and 100% moisture content. On an average 291.9 and 256.9 kg ha<sup>-1</sup> of available N was maintained by the incorporation of K enriched Azolla in soil maintained at 60 and 100% moisture content respectively (Table 1). Among the K fertilizer, K<sub>2</sub>SO<sub>4</sub> significantly maintained higher available N status of 297.2 and 262.2 kg ha<sup>-1</sup> which was 4 to 5% higher than the KCl which registered 286.6 and 251.6 kg ha<sup>-1</sup> of soil available nitrogen content at 60 and 100% moisture content respectively. The available N content ranged from 216.2 to 327.3 and 191.1 to 285.3 kg ha<sup>-1</sup> from 0 to 40ppm of K concentration at 60 and 100% moisture content respectively. Though, the 40 and 50ppm of K were on par with each other, they however, maintained significantly higher available N status

of 327 and 285kg/ha  $\text{kg ha}^{-1}$  at 60 and 100% moisture content respectively and it was 4.0% higher than 30ppm of K enriched Azolla.

128

### 129 3.3.Available P

130 The main and interaction of fertilizer and their concentration significantly influenced the  
131 available P status of Azolla incorporated into the soil maintained at 100% moisture content  
132 where as the main effect was alone significantly influenced the available P content at 60%  
133 moisture condition. On an average, the K enriched Azolla registered 59.63 and 42.13  $\text{kg ha}^{-1}$   
134 of available P at 60 and 100% soil moisture respectively. Among the fertilizers used, the  
135  $\text{K}_2\text{SO}_4$  maintained higher available P content of 61.71 and 44.21  $\text{kg ha}^{-1}$  in the soil  
136 incorporated with K enriched Azolla at 60 and 100 % moisture content respectively followed  
137 by KCl which registered the available P content of 57.54 and 40.04  $\text{kg ha}^{-1}$  (Table 1). Among  
138 the various concentration, 40 and 50ppm K was significantly maintained higher and equal  
139 soil available P status of 75.17 and 77.33  $\text{kg ha}^{-1}$  respectively which was 8.6% more than the  
140 30ppm K (71.17  $\text{kg ha}^{-1}$ ) at 60% moisture content.

141 Table 1. Soil fertility of K enriched Azolla as influenced by main effect of fertilizer and  
142 concentration at 60 and 100% moisture content

Soil Fertility Parameters	Organic Carbon (%)		Available N ( $\text{kg ha}^{-1}$ )		Available P ( $\text{kg ha}^{-1}$ )		Available K ( $\text{kg ha}^{-1}$ )	
Moisture	60%	100%	60%	100%	60%	100%	60%	100%
Fertilizer								
F <sub>1</sub> (KCl)	0.644	0.513	286.6 <sup>b</sup>	251.6 <sup>b</sup>	57.54 <sup>b</sup>	40.04 <sup>b</sup>	205.1 <sup>b</sup>	160.9 <sup>b</sup>
F <sub>2</sub> ( $\text{K}_2\text{SO}_4$ )	0.670	0.538	297.2 <sup>a</sup>	262.2 <sup>a</sup>	61.71 <sup>a</sup>	44.21 <sup>a</sup>	215.5 <sup>a</sup>	171.3 <sup>a</sup>
Mean	<b>0.657</b>	<b>0.525</b>	<b>291.9</b>	<b>256.9</b>	<b>59.63</b>	<b>42.13</b>	<b>210.3</b>	<b>166.1</b>
SEd	0.006	0.006	1.17	1.16	0.85	0.85	1.02	1.02
CD (0.05)	NS	NS	5.02	5.0	3.68	3.68	4.40	4.4
Concentration								
C1 (0 ppm)	0.518 <sup>f</sup>	0.398 <sup>e</sup>	216.2 <sup>g</sup>	191.2 <sup>g</sup>	34.50 <sup>g</sup>	22.50 <sup>f</sup>	157.8 <sup>g</sup>	125.8 <sup>f</sup>
C2 (2 ppm)	0.552 <sup>e</sup>	0.422 <sup>e</sup>	266.8 <sup>f</sup>	236.8 <sup>f</sup>	44.50 <sup>f</sup>	29.50 <sup>e</sup>	172.5 <sup>f</sup>	135.5 <sup>e</sup>
C3 (5 ppm)	0.587 <sup>d</sup>	0.457 <sup>d</sup>	280.0 <sup>e</sup>	250.0 <sup>e</sup>	51.67 <sup>e</sup>	36.67 <sup>d</sup>	190.2 <sup>e</sup>	153.2 <sup>d</sup>
C4 (10 ppm)	0.632 <sup>c</sup>	0.502 <sup>c</sup>	296.2 <sup>d</sup>	259.2 <sup>d</sup>	57.83 <sup>d</sup>	39.83 <sup>c</sup>	204.5 <sup>d</sup>	162.5 <sup>c</sup>

C5 (20 ppm)	0.697 <sup>b</sup>	0.567 <sup>b</sup>	307.5 <sup>c</sup>	270.5 <sup>c</sup>	64.83 <sup>c</sup>	46.83 <sup>b</sup>	219.3 <sup>c</sup>	177.3 <sup>b</sup>
C6 (30 ppm)	0.755 <sup>a</sup>	0.622 <sup>a</sup>	314.0 <sup>b</sup>	277.0 <sup>b</sup>	71.17 <sup>b</sup>	53.17 <sup>a</sup>	238.7 <sup>b</sup>	189.7 <sup>a</sup>
C7 (40 ppm)	0.763 <sup>a</sup>	0.623 <sup>a</sup>	327.3 <sup>a</sup>	285.3 <sup>a</sup>	75.17 <sup>a</sup>	53.17 <sup>a</sup>	250.0 <sup>a</sup>	195.0 <sup>a</sup>
C8 (50 ppm)	0.753 <sup>a</sup>	0.613 <sup>a</sup>	327.2 <sup>a</sup>	285.2 <sup>a</sup>	77.33 <sup>a</sup>	55.33 <sup>a</sup>	249.5 <sup>a</sup>	189.5 <sup>a</sup>
Mean	<b>0.657</b>	<b>0.525</b>	<b>291.9</b>	<b>256.9</b>	<b>59.63</b>	<b>42.13</b>	<b>210.3</b>	<b>166.1</b>
SEd	0.011	0.013	2.96	3.0	1.08	1.07	2.89	2.89
CD (0.05)	0.023	0.235	6.06	6.1	2.20	2.19	5.93	5.92

**Table 2. Soil fertility of K enriched Azolla as influenced by interaction effect between fertilizer Vs. concentration at 60 and 100% moisture content**

Soil Fertility Parameters	Organic Carbon (%)		Available N ( kg ha <sup>-1</sup> )		Available P ( kg ha <sup>-1</sup> )		Available K ( kg ha <sup>-1</sup> )		
	Moisture	60%	100%	60%	100%	60%	100%	60%	100%
F <sub>1</sub> C <sub>1</sub>		0.510	0.390	214.3	189.3	33.33 <sup>j</sup>	21.33 <sup>j</sup>	156.3 <sup>f</sup>	124.3 <sup>h</sup>
F <sub>1</sub> C <sub>2</sub>		0.543	0.413	263.7	233.7	44.00 <sup>i</sup>	29.00 <sup>i</sup>	171.3 <sup>e</sup>	134.3 <sup>f</sup>
F <sub>1</sub> C <sub>3</sub>		0.577	0.447	278.0	248.0	51.67 <sup>h</sup>	36.67 <sup>h</sup>	187.3 <sup>d</sup>	150.3 <sup>e</sup>
F <sub>1</sub> C <sub>4</sub>		0.617	0.487	287.0	250.0	56.67 <sup>g</sup>	38.67 <sup>gh</sup>	195.3 <sup>d</sup>	153.3 <sup>e</sup>
F <sub>1</sub> C <sub>5</sub>		0.670	0.540	302.7	265.7	61.00 <sup>f</sup>	43.00 <sup>f</sup>	207.0 <sup>c</sup>	165.0 <sup>d</sup>
F <sub>1</sub> C <sub>6</sub>		0.727	0.593	307.3	270.3	65.00 <sup>e</sup>	47.00 <sup>e</sup>	226.0 <sup>b</sup>	177.0 <sup>c</sup>
F <sub>1</sub> C <sub>7</sub>		0.757	0.617	319.7	277.7	73.00 <sup>c</sup>	51.00 <sup>d</sup>	249.3 <sup>a</sup>	194.3 <sup>ab</sup>
F <sub>1</sub> C <sub>8</sub>		0.753	0.613	320.3	278.3	75.67 <sup>bc</sup>	53.67 <sup>cd</sup>	248.3 <sup>a</sup>	188.3 <sup>b</sup>
F <sub>2</sub> C <sub>1</sub>		0.527	0.407	218.0	193.0	35.67 <sup>j</sup>	23.67 <sup>j</sup>	159.3 <sup>f</sup>	127.3 <sup>h</sup>
F <sub>2</sub> C <sub>2</sub>		0.560	0.430	270.0	240.0	45.00 <sup>i</sup>	30.00 <sup>i</sup>	173.7 <sup>e</sup>	136.7 <sup>f</sup>
F <sub>2</sub> C <sub>3</sub>		0.597	0.467	282.0	252.0	51.67 <sup>h</sup>	36.67 <sup>h</sup>	193.0 <sup>d</sup>	156.0 <sup>e</sup>
F <sub>2</sub> C <sub>4</sub>		0.647	0.517	305.3	268.3	59.00 <sup>fg</sup>	41.00 <sup>fg</sup>	213.7 <sup>c</sup>	171.7 <sup>cd</sup>
F <sub>2</sub> C <sub>5</sub>		0.723	0.593	312.3	275.3	68.67 <sup>d</sup>	50.67 <sup>d</sup>	231.7 <sup>b</sup>	189.7 <sup>b</sup>
F <sub>2</sub> C <sub>6</sub>		0.783	0.650	320.7	283.7	77.33 <sup>ab</sup>	59.33 <sup>a</sup>	251.3 <sup>a</sup>	202.3 <sup>a</sup>
F <sub>2</sub> C <sub>7</sub>		0.770	0.630	335.0	293.0	77.33 <sup>ab</sup>	55.33 <sup>bc</sup>	250.7 <sup>a</sup>	195.7 <sup>ab</sup>
F <sub>2</sub> C <sub>8</sub>		0.753	0.613	334.0	292.0	79.00 <sup>a</sup>	57.00 <sup>ab</sup>	250.7 <sup>a</sup>	190.7 <sup>b</sup>
Mean		0.657	0.525	291.9	256.9	59.63	42.13	210.3	166.1
SEd									
F at C		0.016	0.017	4.08	4.09	1.66	1.65	3.96	3.96
C at F		0.016	0.016	4.18	4.19	1.52	1.51	4.09	4.09
CD (0.05)									
F at C		NS	NS	NS	NS	4.39	4.39	8.71	3.9
C at F		NS	NS	NS	NS	3.11	3.11	8.38	4.1

But in 100% soil moisture content at 30, 40 and 50ppm, K was significantly maintained higher and on par soil available P status followed by 20ppm K. The lowest available P of

34.5 and 22.5 kg ha<sup>-1</sup> was registered by soil incorporated with Azolla enriched with 0 ppm of K at 60 and 100% moisture content. Under the interaction between K fertilizer and its concentration, K<sub>2</sub>SO<sub>4</sub> at 50 and 30 ppm were significantly superior in maintaining higher available P status at 100 % moisture content followed by 40 ppm K<sub>2</sub>SO<sub>4</sub> and it was on par with 50 ppm of KCl (Table 2). However, the 0 ppm of both the fertilizer produced lowest available P content (21.33 and 23.67 kg ha<sup>-1</sup>) of Azolla incorporated soil.

### 3.4. Available K

The available K content of soil incorporated with K enriched Azolla was significantly influenced by the main and interaction effect of K fertilizer and its concentration. Irrespective of the fertilizer and their concentration about 210.3 and 166.1 kg ha<sup>-1</sup> of available K was maintained by the incorporation of K enriched Azolla in soil at 60 and 100 % moisture respectively (Table 1). Among the K fertilizer, K<sub>2</sub>SO<sub>4</sub> was superior in maintaining available K content (215.5 and 171.3 kg ha<sup>-1</sup>) in Azolla incorporated soil which was 5-7 % more than the KCl (205.1 and 160.9 kg ha<sup>-1</sup>). With respect to concentration of K, 40 and 50 ppm of K at 60 % moisture and 30.40 and 50 ppm of K at 100 % moisture registered significantly higher and equal available K in Azolla incorporated soil followed by 30 and 20 ppm of K at 60 and 100 % moisture respectively. The Azolla grown under 0 ppm of K registered the lowest available K content of 157.8 and 125.8 kg ha<sup>-1</sup> at 60 and 100 % moisture content respectively. Under the interaction between K fertilizer and its concentration, K<sub>2</sub>SO<sub>4</sub> with 30, 40 and 50 ppm at 60 % moisture content and 40 and 50 ppm of K<sub>2</sub>SO<sub>4</sub> and 40 ppm of KCl produced statistically higher and equal available K in Azolla incorporated soil maintained at 60 and 100 % moisture respectively (Table 2). The 0 ppm of both the fertilizer registered the lowest available K content (159.3 and 156.3 kg ha<sup>-1</sup>) in soil incorporated with Azolla grown under these concentrations at both moisture content.

#### 4. Discussion

Soil fertility is influenced by the humic substances formed during the decomposition of *Azolla* (Bhardwaj and Gaur 1970). Incorporation of *Azolla* enhanced the soil nutrients availability by their biological activity. The decomposed organic matter from *Azolla* biomass played an active role in the development of microbial population. Similarly, Kannaiyan and Subramani (1992) showed the increased of cellulolytic and urea hydrolyzing activities in addition to significant increase in the population of heterotrophic bacteria by the added *Azolla*. Soil incorporation of *Azolla* also increased urease and phosphatase activity (Thanikachalam *et al.* 1984). *Azolla* contains macro, secondary and micronutrients that is important for quality rice production (Kumar and Shahi, 2016). Sutanto (2002) stated that, the use of 7.5 ton ha<sup>-1</sup> *Azolla* to paddy field increased soil organic matter (C-organic) 0.09 times of control (without *Azolla*). Syamsiyah *et al.* (2015) proved that, application of *Azolla* at 2 ton ha<sup>-1</sup> could increase the organic matter up to 3.69% compare to the field without *Azolla*. The increasing of organic C is caused by the high content of organic C in *Azolla*. The incorporated *Azolla* into soil would soon be mineralized. Watanabe *et al.*, (1991) stated that 90% of *Azolla* was decomposed in 4 weeks and releases humic substances in to the soil. The increase in grain yield might be due to build up of soil organic carbon and more nitrogen through the integrated use of NPK and green manuring with *Azolla*.

Biological Nitrogen Fixation (BNF) is a major source of fixed N for plant life and to sustain production and productivity of agricultural crops. Estimate of global terrestrial showed that the BNF ranged from 100 to 290 million tonnes of N/ year. Of this, 40–48 million tonnes is estimated to be biologically fixed in agricultural crops and fields. BNF is one of the natural sources of nitrogen for rice and *Azolla*-cyanobacteria biomass has been identified as potential source of nitrogen. The glutamate synthase enzyme dominated in ammonia assimilation

202 followed by glutamine synthetase and glutamate dehydrogenase in Azolla (Fig. 1). The activity  
 203 of all the three enzymes were more at the 40 ppm of K followed by 30 and 50 ppm of K.  
 204 Incorporation of 40 ppm K either as KCl and K<sub>2</sub>SO<sub>4</sub> incubated Azolla enhanced ammonia  
 205 assimilation and improved soil fertility (Fig. 2) on 30 days which may help to reduce nitrogen  
 206 demand for rice crop (Muruganayagi, 2017).




Figure 1. Nitrogen assimilating enzymes as influenced by K fertilizer




Figure 2. Nitrogen fixing enzyme as influenced by concentration of fertilizers

207  
 208 According to Roy, (1981), incorporation of 6 t of *Azolla* ha<sup>-1</sup>, equivalent to 36 kg of N ha<sup>-1</sup>  
 209 before planting and incorporation of 1 ton ha<sup>-1</sup> *Azolla*, equivalent to 24 kgN ha<sup>-1</sup> after 3-4  
 210 days of planting. Fogg *et al.* (1973) have found that *Azolla* and cyanobacteria bio-fertilizer can  
 211 add as much as 30-120 kg N ha<sup>-1</sup> per crop. Many researchers have considered cyanobacteria  
 212 as a promising source of nitrogen in tropical rice soils. (Valiente *et al.*, 1998) investigated the  
 213 potential contribution of N<sub>2</sub> fixation by indigenous cyanobacteria to rice soil with increasing  
 214 amount of fertilizers. Ventura *et al.* (2012) concluded that about 50% of the N in *Azolla* was  
 215 mineralized after 2 weeks of incubation with more than 3% N content. Use of *Azolla* as green  
 216 manuring and as intercrop proved beneficial and significant result over control in respect of P  
 217 content. The highest available phosphorus (29.6 kg ha<sup>-1</sup>) was recorded with 100% NPK +

green manuring of Azolla (Kumar and Shahi, 2016). Azolla also contributes to the supply of Phosphorus, Potassium, Sulfur, Zinc, Iron and Molybdenum in sufficient amounts in addition to other micronutrients besides addition of Nitrogen. Similarly, the soil biological health due to application of Azolla has resulted in improving mineralization and consequent increase in the soil microbial status (Yadav *et al.* 2014).

## 5. Conclusion

Soil biological health, mineralization and consequent increase in nutrient status by the application of K enriched Azolla was more under 60 % soil moisture content than fully saturated soil. Further, enrichment of Azolla with 40 ppm of K<sub>2</sub>SO<sub>4</sub> enhanced nutrient content in Azolla and in soil.

## References

- AwodunMA. Effect of Azolla (*Azolla species*) on physicochemical properties of the soil. *World J. Agril. Sci.* 2008; **4** (2): 157-160.
- Bhardwaj KKR, Gaur AC. Effect of humic and pulvic acid on growth and efficiency of nitrogen fixation by *Azotobacterchroococcum*, *Folia Microbiology* 1970; **15**: 364.
- Bray RA, Kurt LT. Determination of Total Organic and Available Form of Phosphorus in Soil. *Soil Sci.*, 1945; 59: 34-35.
- Corazon MRamirez. Relationship between soil phosphorus availability and azolla growth, *Soil Sci. Plant Nut.* 2012; **30** (4) : 595-598.
- Debjani Halder, Shyamal Khoroar. Mineralization and Availability of *Azolla* and Cyanobacteria Biomass Nutrients in Rice Soil. *J. Agril. Sci. and Tech.* 2013; **3**: 782-789.
- De DattaSK, Buresh RJ. Integrated nitrogen management in irrigated rice, *Adv. Soil Sci.* 1989; **10** : 143-169.
- Fogg GE, Steward WDP, FayP Walsky AE. The Blue Green Algae, Academic Press, New York; 1973
- Halder;D, ; KhoroarS. Mineralization and Availability of *Azolla* and Cyanobacteria Biomass Nutrients in Rice Soil. *J. Agril. Sci. Tech.* 2013; **3** : 782-789
- IRRI, IRRISTAT, Biometric unit, IRRI, Los Banos, Philippines; 1993

249 Kannaiyan S, Subramani S. Use of Azolla as biofertilizer for rice crop, cyanobacterial  
250 nitrogen fixation, (Ed: Kaushik B. D.), 1992; 281-289, Indian Agricultural Research  
251 Institute, New Delhi

252 Kumar B ShahiDK.. Effect of azolla as green manure on soil properties and grain yield of  
253 rice in acid soil of Jharkhand. *Annals plant soil Res.*, 2016; **18** (3): 214-218.

254 MichihikoY, NaokiT, Noriyo. HDA KatsunoriNNitrogen fixation in azolla-anabaena  
255 symbiosis as affected by mineral nutrient status. *Soil Sci. Plant Nut.* 2012; **26** (3) :  
256 415-426.

257 MuruganayakiSJeyachitra A. Nitrogen assimilating enzyme of azolla as influenced by  
258 potassic fertilizer, *Inter. J, Curr. Adv. Res.* 2017; **6**(3), pp. 3012-3016.

259 Roy B. Manuring of rice with Azolla, *Oryza*, 1981; **21** : 238-241.

260 SahrawatKL. Macro and micronutrients removed by upland and lowland rice cultivars in  
261 West Africa, *Comm. Soil Sci. Plant Analysis*, 2000; **31**:717-723.

262 Singh A L, Singh PK. "Influence of Azolla management on the growth, yield of rice and soil  
263 fertility": II. N and P contents of plant and soil. *Plant Soil*. 1987; **102** (1) : 49-54.

264 Singh PK. "Nitrogen Economy of Rice Soils in Relation to Nitrogen Fixation by Blue-Green  
265 Algae and Azolla. In Increasing Rice Yield in Kharif" Central Rice Research  
266 Institute: Cuttack, India. 1987; 221–239.

267 SinghPK, Subudhi BP. Utilize Azolla in poultry feed, in: Subjected Matter Training-Cum-  
268 Discussion Seminar on Use of Bio-fertilizer with Special Reference to Azolla, Held  
269 from 16th to 21st Feb., CRRI, Orissa, 1981.

270 StanfordS. English. Use of flame photometer in rapid soils tests for Potassium and calcium.  
271 *Agron. J*, 1949; **41**: 446-447.

272 Subba Rao A, Sammi Reddy K. Integrated nutrient management vis-à-vis crop  
273 production/productivity, nutrient balance, farmer livelihood and environment: India.  
274 Proc. Regional Workshop Beijing, China, 2005.

275 Subbiah BV, Asija. A rapid procedure for estimation of available nitrogen in soils. *Curr. Sci.*,  
276 1956; **25**: 259-260.

277 Subedi P, Shrestha J. "Improving soil fertility through Azolla application in low land rice": A  
278 review. *Azarian J. Agri.*, 2015; **2** (2): 35-39.

279 SutantoR. Penerapanpertanianorganik. PenerbitKanisius. Yogyakarta 2002.

280 SyamsiyahJ, SunarmintoBH. Mujiyo. Changes in soil chemical properties of organic paddy  
281 field with azolla application. *J. Soil Sci. Agroclim.*, 2015; **13** (1): 68-73

282 Thanikachalam A, Rajakannu K, Kannaiyan S. Effect of Neem cake, Carbofuran and Azolla  
283 application on phosphatase activity in soil. 25th Annual Conference of Association  
284 of Microbiologists of India, GB Pant University of Agriculture and Technology, Pant  
285 Nagar, India. 1984.

286 VenturaW, WatanabeI, Mascariña GB. Mineralization of Azolla N and its availability to  
287 wetland rice. *Soil Sci. Plant Nutr.*, 2012; **38**(3): 505-516

- 288 Valiente EF, Ucha A, Quesada A, LeganesF, Carreres R. Contribution of N<sub>2</sub>-fixing  
 289 cyanobacteria to rice production, 5th International Symposium, Inorganic N  
 290 Assimilation, Luso, Portugal, *Plant Soil*, 1998; **1** :107-112.
- 291 Walkley A, Black CA. An estimation of methods for determining organic carbon and  
 292 nitrogen in the soils. *J. Agric. Sci.*, 1934; **25**: 598-609.
- 293 Watanabe I, Benjamin Padre Jr, Corazon Ramirez. “Mineralization of Azolla N and Its  
 294 Availability to Wetland Rice”. *Soil Sci. Plant Nutr.*, 1991; **37** (4): 679-688.
- 295 Yadav RK, Abraham G, Singh Y V, Singh PK. “Advancement in the utilization of Azolla-  
 296 Anabaena system in relation to sustainable agricultural practices”. Proc. Indian  
 297 National Science Academy, 2014; **80**:(2): 301-316.

298



299

300