

Clonal propagation of Strawberry (*Fragaria x ananassa* Duch.)
through *in vitro* runner tip culture through incorporation of
growth hormones

Md. Nasir Uddin Badal^{1,3*}, Md. Shoyeb¹, Md. Abdur Rauf Sarkar^{1,2}, Md. Shahedur
Rahman¹ and Shaikh Mizanur Rahman¹

¹Department of Genetic Engineering and Biotechnology, Jessore University of Science
and Technology, Jessore 7408, Bangladesh

²Laboratory of Plant Genetics and Breeding, Faculty of Agriculture, Saga University,
Saga 840-8502, Japan

³Protein Dynamics Research Group, Faculty of Medicine and Life Science, University
of Tampere, Tampere 33520, Finland.

*Corresponding author: abdullahsyum1992@gmail.com

Abstract

Runner tips explants of strawberry give rise to multiple shoots when cultured on MS medium supplemented with different concentrations and combinations of BAP with KIN or NAA or GA₃. The highest response of shoot multiplication was obtained on MS containing 2.5 mg l⁻¹ BAP + 0.5 mg l⁻¹ Kin + 0.5 mg l⁻¹ GA₃. The maximum frequency of rooting (83%) and highest number of roots (3.49) was produced in medium containing 1.0 mg l⁻¹ IBA. The well grown rooted plantlets were acclimatized and successfully established in autoclaved vermiculate soil and as well as natural condition. Using our established protocol, it is also possible to provide large numbers of micropropagated

plantlets of this cultivars to produce high quality strawberry fruit for commercial cultivation practices.

Key words: Strawberry, in vitro, regeneration, multiple shoots and roots, acclimatization

1. Introduction

Strawberry is a member of the Rosaceae family which is genetically octoploid ($2n=8x=56$) in nature, is the most important soft fruit worldwide ^[1]. The cultivated strawberry (*Fragaria X ananassa* Duch.) is a natural hybrid between the Scarlet or Virginia strawberry (*F. virginiana* Duch.) and pistillate South American (*F. chiloensis* L. Duch), ^[2]. It is a perennial, stoloniferous herb. Strawberries have traditionally been a popular delicious fruit for its flavor and taste. It can be consumed fresh, frozen or processed. It is produced in 71 countries worldwide on 506000 acres ^[3].

Strawberry is one of the most fascinating fruits of the world. People of all spheres like this fruit because of some unavoidable reason. Firstly, it's bright red color, juicy texture and tantalizing aroma attracts all very easily. Secondly, it contains comparatively high percentage of minerals including manganese, calcium, potassium, iron, copper, numerous dietary fibers, carbohydrates, unsaturated fatty acid, proteins and most of the nutritious elements essential for human being ^[5]. It is also a rich source of vitamins like B1, B2, C. Especially the fruit contains higher vitamin C concentration than orange or lemon. Finally, it has relatively high quantities of ellagic acid, which has a wide range of biological activity including anti-carcinogenic impact ^[4]. This dilactone of hexahydroxy diphenic acid is a natural phenol antioxidant, gradually getting polarity in

mental function, viral and bacterial infection and cancer treatment [3]. It is consumed in large quantities, either fresh or in prepared foods, such as preserved fruit juice, pies, ice creams, and milk shakes. Artificial strawberry flavor is also widely used in many industrialized food products. It is produced in 73 countries worldwide on 200,000 hectares and produced 31 lac metric tons strawberry [6].

In conventional cultivation, the strawberry (*Fragaria X ananassa* Duch.) is a species vegetatively propagated via runner seedlings. Nevertheless, such seedlings due to their weakness and susceptibility to pathological agents are not always suitable for this cultivation. Moreover, most of the strawberry cultivars are grown in the temperate parts of the world, although a few day-neutral variety can be grown in the sub-tropical regions. However, commercial cultivation is not popular in Bangladesh due to lack of proper cultivation knowledge, profitability and lack of available season friendly variety [7]. For better strawberry production, 10 to 20h photoperiod, 12 to 30°C day temperature and 12 to 24 short days are essential [8]. Bangladesh is a sub-tropical country with 15 to 25°C winter average day temperature, where almost two months include short photoperiod of 12 to 16 h [9]. Therefore, from the beginning through the whole winter season strawberry can be grown without that much excessive nursing and care. In present time, gradually framers are becoming interested in strawberry culture due to high demands and more profits. Since last few years, strawberries were cultivating but the main constrain of its cultivation was to maintain plant materials in hot summer of Bangladesh [10].

To solve the problems of natural seedlings and make it available nowadays, the advantageous alternative to this conventional method seems to be the use of micro-propagated plants for cultivation [14]. Micro-propagation of strawberry has been applied

on a large scale in commercial production since mid 1970s [15]. First applied to strawberry meristems *in vitro* techniques have been amplified to an efficient method of mass propagation [14][16]. Strawberries can be propagated *in vitro* condition by tissue culture methods whereas micro-propagation is a very useful technique of improvement of plant. Micro-propagated strawberry plants were comparatively better in different characters (crown size, number of runners, flowering time and yield of berries) than conventionally propagated runner plants [11][12]. Although production of propagules through runner has been reported to contribute 90% of total Dutch strawberry production, the product in Elsanta cultivars was found to be susceptible to several fungal diseases [13].

However, strawberry was introduced in Bangladesh two decades ago and in becoming popular in recent years. Bangladesh Agricultural Research Institute (BARI) recently released a strawberry variety known as BARI Strawberry-1 [17]. Conventional propagation methods are slow, laborious, and expensive with many limitations and may not be recommended for effective and commercial multiplication [18]. Commercial multiplication of the released variety in this country is essential for rapid extension. The advantages of *in vitro* propagation are that it offers fast multiplication rates [19]. The significant advantages of micro-propagation by which many plants can be produced from a single individual in a relatively short span of time and space [20].

Moreover, the conventional way of production is not adequate to meet the commercial demand. Micropropagated strawberry plant has been introduced to prevent most of the plant and soil transmissible diseases [9]. Large-scale commercial propagation by *in vitro* techniques has been used widely in the strawberry industry [21].

Therefore, in our investigation to considering the view of the potential commercial value and market demand we established an efficient and cost-effective method of *in vitro* propagation of strawberry to ensure availability of planting materials for commercial cultivation practice.

2. Materials and Methods

2.1 Collection of explants and surface sterilization

Juvenile runner tips from strawberry mature and healthy plants (Fig. 1A) were collected from the strawberry garden of Jessore University of Science and Technology, Jessore 7408, Bangladesh. From the strawberry field we had collected around 5 cm runner tip explant with sterile scalpel and immediately dropped into distil water to avoid the cell damage and contamination. Collected explants were washed in a running tap water for 30 min and then thoroughly washed again by adding a few drops of Savlon (ACI Ltd. Bangladesh) and Tween-20 (UNI Chem, China) for 5 minutes followed by rinsing with three times autoclaved distilled water. Again, surface sterilized in a 0.1% mercuric chloride (MERCK, India) for 5 min followed by rinsing them five times with autoclaved dH₂O inside the Laminar Air Flow Cabinet.

2.2 Inoculation

Small runner tips (0.5-1.0 cm) were inoculated on MS medium (Murashige, and Skoog, 1962) supplemented with different concentration of growth regulators (BA, NAA, KIN and GA₃). In addition to that, 30 g l⁻¹ sugar were used as carbon source and 0.6% agar have used to solidify the medium. The pH of the medium was adjusted to 5.7 by using 1M KOH before autoclaving at 0.15 MPa in 121°C for 20 min. Being prepared the sterilized explants and solidified cooled medium we have inoculated them under laminar

air flow hood. Sterilized scalpel has used to inoculate the explants in a fashion that half of the explants should put inside the medium.

2.3 Incubation

The cultures were maintained in a growth chamber at $25 \pm 2^\circ\text{C}$ under photoperiod of 16 h/day and a light intensity of 2000-3000 lux provided by white fluorescence tube (Philips) including 55-60% relative humidity.

2.4 Sub-culture and multiple shoot regeneration

Subcultures were done every 2-3 weeks interval depending on the explants growth conditions. Nodal segments from the proliferated shoots were sub-cultured again for further multiple shoots regeneration.

2.5 Rooting and acclimatization

Regenerated multiple shoots were cut (approximately 2-3 cm long) and individual shoots were placed on MS solidified medium containing different single concentrations of IBA and IAA for root induction. Well grown *in vitro* rooted plantlets were taken out from culture tubes and washed thoroughly with tap water to remove the culture medium from the roots. Washed plantlets were planted to sterilized vermiculate soil in small pot and after acclimatization grown in a natural condition.

2.6 Data analysis

Surface sterilization, shoot regeneration, multiple shooting and rooting data were recorded during the experiments. In the data table we have presented those experimental data which have gotten from 15 explants inoculation at a time. After getting results we

have repeated the same hormone concentration three times to be sure about the homogeneity and appropriate hormone ratio. We have used SPSS software to measure the standard deviation of the data.

3. Results and Discussion

Runner tips explants of strawberry cultured on MS medium with different concentration of BAP (0.5-2.5 mg l⁻¹), KIN (1.0-2.0 mg l⁻¹), and GA₃ (0.1-1.0 mg l⁻¹) (Table 1). Within 4-5 weeks of culture multiple shoots were directly initiated from the explants. The maximum number of shoots (7) in the medium with BAP (2.0 mg l⁻¹) were greater than those observed in the medium supplemented with KIN and GA₃ singly, where 82% explants showed shoot proliferation (Table 1 and Fig. 1, B). But when BAP concentration had lowered than 2.5 mg l⁻¹ and increased more than 2.5 mg l⁻¹, the shoot proliferation rate was decreased. To be more accurate we have applied lower and higher hormone concentration than this optimum BAP (0.5-2.5 mg l⁻¹), every time we have experienced the same results.

On the other hand, runner tips explants were cultured on MS medium supplemented with different concentration and combination of BAP (1.0-2.5 mg l⁻¹) with KIN (0.1-0.5 mg l⁻¹) or BAP (1.0-2.5 mg l⁻¹) with GA₃ (0.05-0.5 mg l⁻¹) or BAP (1.0-2.0 mg l⁻¹) with NAA (0.1-0.5 mg l⁻¹). And cultured on MS medium with BAP (1.0-2.5 mg l⁻¹) with KIN (0.5-1.0 mg l⁻¹) and GA₃ 0.5 mg l⁻¹. The numbers of shoots in medium with BAP+KIN+GA₃ combinations were greater than those observed in the medium supplemented with BAP + KIN or BAP + GA₃ or BAP + NAA treatments. The highest rate of response was obtained at 2.5 mg l⁻¹ BAP + 0.5 mg l⁻¹ KIN combination (Table 2) where 92% explants showed shoot proliferation and 7 ± 0.23 shoots were developed. When BA concentration was decreased below 2.0 mg l⁻¹, the rate of shoot multiplication

reduced. In our present investigation, maximum number of shoots per explant (20 ± 0.23) and highest average length (3.5 ± 0.26) were recorded at BAP 2.5 mg l^{-1} + KIN 0.5 mg l^{-1} + GA₃ 0.5 mg l^{-1} , which was found to be the best combination for high frequency of multiple shoot induction of strawberry (Table 2, Fig. 1, C). The observed rate of shoot proliferation 98%. When BAP was supplemented with KIN or GA₃ or NAA the rate of shoot proliferation and the shoot length were increased but number of shoots were decreased. The high concentration of cytokinin reduced the number of micropropagated shoots [23]. Similar results have already been reported in *Fragaria indica* Andr. [24]. Also, this result is in consistent with the findings of in papaya [25] as well as in *Eucalyptus grandis* [26]. The shoot organogenesis varied from strawberry genotypes to genotypes [27]. Though some workers reported high concentration of BAP is the best for strawberry micro propagation [28] while other authors suggested 1.0 mg/l IAA + 1.0 mg/l BAP + 0.05 mg/l GA₃; 0.5 mg/l BA + 0.1 mg/l GA₃ + 0.1 mg/l IBA [29][30] and 0.5 mg/l BA + 0.1 mg/l IBA for strawberry micropropagation. In a nutshell we can state that hormone concentration BAP 2.5 mg l^{-1} + KIN 0.5 mg l^{-1} + GA₃ 0.5 mg l^{-1} is the best combination for shoot proliferation than other studied combination. This combination gives the best outcomes and not that much higher concentration, which indicate the cost-effective shoot proliferation combination for strawberry micropropagation.

The developing shoots were elongated by sub-culturing on the same combinations of growth regulators. Later, elongated shoots were excised and used for root induction. Well grown shooted plantlets were cultured on different concentrations of IBA (0.5 - 2.0 mg l^{-1}) and IAA (0.5 - 2.0 mg l^{-1}) singly on $\frac{1}{2}$ MS solidified medium. Among the different treatments 1.0 mg l^{-1} IBA proved to be the most suitable for root induction with 83% of

rooting and 3.49 ± 0.05 roots per explant and the average root length being 3.18 cm (Table 3; Fig. 1, F). Also, were observed similar effects of IBA in case of rooting of strawberry plantlets [3].

We have seen that higher concentration of hormone is not the best way of micropropagation, rather than, precise combination of different hormone is also very important. In all above described hormone combinations where lower or higher rate than the concentrations we found fail to give high amount of shoot proliferation and it is also true in case of root induction and proliferation. Thoroughly analysis of all hormones combination both in shoot and root proliferation we found the best amount is BAP 2.5 mg l^{-1} + KIN 0.5 mg l^{-1} + GA₃ 0.5 mg l^{-1} (Table 2, Fig. 1, C) for shoot and 1.0 mg l^{-1} IBA (Table 3; Fig. 1, F) for root proliferation.

Table 1: Effects of different concentration of BAP, KIN and GA₃ on multiple shoot regeneration from runner tips explants of strawberry.

Growth regulators conc.(mg l ⁻¹)	No. of explants inoculated	% of explants responded	Days to shoot formation	No. of shoots per culture (M±S.E.)	Highest length of shoots in cm (M±S.E.)
BAP					
0.5	15	20	18-22	3±0.57	1.90±0.05
1.0	15	40	15-20	5±0.58	2.10±0.19
1.5	15	70	12-15	5±0.45	2.51±0.57
2.0	15	82	8-15	7±0.73	2.10±0.15
2.5	15	71	12-15	5±0.66	2.03±0.03
KIN					
1.0	15	40	10-15	2±0.43	2.17±0.08
1.5	15	65	10-16	4±0.45	2.06±0.03
2.0	15	73	8-15	5±0.21	2.20±0.06
GA₃					
0.1	15	25	10-15	3±0.57	2.04±0.04
0.5	15	45	8-15	4±0.22	2.25±0.02
1.0	15	33	10-15	5±0.65	2.03±0.10

Note: M = Mean, S.E. =Standard Error

Table 2: Effects of different concentration and combination of auxin with cytokinin and GA₃ in MS medium on multiple shoot regeneration from runner tips explants of strawberry.

Growth regulators (mg l ⁻¹)	No. of explants inoculated	% of explants responded	Days to shoot formation	No. of shoots per culture (M±S.E.)	Highest length of shoots in cm (M±S.E.)
BAP+KIN					
1.5+0.5	15	65	10-15	3±0.58	2.20±.11
2.0+0.1	15	75	9-12	3±0.40	3.0±0.16
2.0+0.5	15	85	8-12	5±0.34	3.4±0.21
2.5+0.5	15	92	8-12	7±0.23	3.5±0.66
BAP+GA₃					
1.0+0.05	15	28	12-16	2±0.21	3.2±0.11
1.5+0.1	15	60	10-15	3±0.76	2.9±0.42
1.5+0.5	15	87	12-16	5±0.24	3.06±0.05
2.0+0.5	15	90	12-16	8±0.58	3.3±0.24
2.5+0.5	15	82	12-16	7±0.33	3.47±0.63
BAP+NAA					
1.0+0.5	15	28	12-20	2±0.55	2.0±0.26
2.0+0.1	15	57	12-20	4±0.32	2.3±0.27
2.0+0.5	15	60	10-15	5±0.35	2.56±0.38
BAP+KIN+GA₃					
1.0+0.5+0.5	15	80	7-12	12±.33	3.2±0.51
1.5+0.5+0.5	15	85	7-14	12±.66	2.5±0.88
1.5+1.0+0.5	15	72	6-15	14±.57	2.5±0.58
2.0+0.5+0.5	15	90	6-12	17±.39	3.4±0.32
2.5+0.5+0.5	15	98	5-10	20±.23	3.5±0.26

Note: M = Mean, S.E. =Standard Error

Table 3: Effects of different concentration of auxins (IBA, IAA) in MS medium on *in vitro*

root induction from regenerated shoots of strawberry after 7-8 weeks of culture.

Growth regulators conc. (mg l ⁻¹)	No. of shoot sub-cultured	Shoot derived from the explants of mature plants			
		% of rooting	Days to root generation	Average no. of roots	Average root length (cm)
IBA					
0.5	15	26	16-22	2.26±0.23	2.86±0.09
1.0	15	83	12-20	3.49±0.05	3.18±0.04
1.5	15	64	14-20	2.50±0.04	2.90±0.65
2.0	15	71	16-22	3.36±0.03	3.18±0.56
IAA					
0.5	15	39	18-26	2.04±0.35	3.18±0.30
1.0	15	56	16-22	3.45±0.33	3.50±0.50
1.5	15	64	16-26	3.20±0.20	3.22±0.39
2.0	15	53	21-26	2.92±0.32	3.08±0.52

Note: M = Mean, S.E. =Standard Error




Figure 1: *In vitro* propagation of Strawberry plantlets from runner tips explants; A. Runner tips used as explants. B. Shoot proliferation on MS supplemented with 2.0 mg l^{-1} BAP after 3 weeks of culture. C. Proliferated of multiple shoots on MS+ 2.0 mg l^{-1} BAP+ 0.5 mg l^{-1} KIN+ 0.5 mg l^{-1} GA₃ after 4 weeks of culture. D and E. Establishment of shoot after 7 weeks of culture. E. Rooted shoots on 1.0 mg/l IBA after 7-8 weeks of culture.

4. Conclusion

From our investigation, it can be concluded that enormous strawberry plantlets even in off season can be produced from an explant within a short period of time by using our established protocol of micropropagation. Also, our investigation, provides a reliable and economical method of maintaining pathogen free plants. Although, very recently strawberry plants have been introduced in Bangladesh but within a short time this fruit has become very popular to whole country. There for requirements of healthy, disease free and high productive and cost-effective plantlets are increasing day by day. So, applying our above described micropropagation protocol we can easily fulfill our demand of strawberry plantlets for growing in commercial cultivation purposes.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Ethical issues:

There are no ethical issues in the whole research work and in this manuscript. Our findings and this protocol is out of conflict.

Acknowledgment

We thank to Department of Genetic Engineering and Biotechnology, Jessore University of Science and Technology, Jessore 7408, Bangladesh for providing us all kinds of laboratory facilities. We also thank to Jessore University of Science and Technology, Jessore 7408, Bangladesh for providing the partial financial support for this research.

References

1. Jiajun L, Yuhua L, Guodong D, Hanping D, Mingqin D (2005). A natural pentaploid strawberry genotype from the Changbai Mountains in Northeast China. *Horticultural Science* 40: 1194-1195.
2. Debnath S C, Teixeira da Silva J A (2007). Strawberry culture in vitro: application in genetic transformation and biotechnology. *Fruit, Vegetable and cereal Science and Biotechnology* 1(1): 1-12.
3. Sakila S, Ahmed M B, Roy U K, Biswas M K, Karim R, Razvy M A, Hossain M, Islam R, and Hoque A (2007). Micropropagation of Strawberry (*Fragaria X ananassa* Duch.) A Newly Introduced Crop in Bangladesh. *American-Eurasian Journal of Scientific Research* 2 (2): 151-154.
4. ICAR News (2005). Indian council of agriculture research. 11(4).
5. Nehra N S, Kartha K K, Stushnoff, Giles K L (1994). Effect of in vitro propagation methods on field performance of two strawberry cultivars. *Euphytica* 76: 107-115.
6. FAO (2008). FAOSTAT Agricultural Statistics Database. <http://www.Fao.org>
7. Biswas M K, Dutta M, Roy U K, Islam R, and Hossain M (2009). Development and evaluation of in vitro somaclonal variation in strawberry for improved horticultural traits. *Scientia Horticulture* 122: 409-416.
8. Michel J V, Anita S, and Svein O G (2006). Interactions of photoperiod, temperature, duration of short-day treatment and plant age on flowering of *Fragaria X ananassa* Duch. Cv. Korona. *Scientia Horticulture* 107: 64-170.

9. Biswas M K, Islam R, Hossain M Micropropagation and field evaluation of strawberry in Bangladesh,” *Journal of Agricultural Technology*, vol. 4, no. 1, pp. 167-182, 2008.
10. Ara T, Karim M A, Aziz A A, Islam R, and Hossain M (2013). Micropropagation and field evaluation of seven strawberry genotypes suitable for agro-climatic condition of Bangladesh. *African Journal of Agricultural Research* 8(13): 1194-1199.
11. Karhu S, Hakala K (2002). Micropropagated strawberries on the field. *Acta Horticulture* 2: 182-186.
12. Karhu S, Hakala K (2007) Micropropagated Strawberries on the field. *Acta Horticulture* 567: 321-324.
13. Dijkstra J (1993). Research on strawberries focusses on healthy plant material. Expensive cultural method requires excellent material. *Fruittelt-Den-Hang* 83(34): 14-15.
14. Jadwiga Z, Elżbieta K, Jacek G (2015). Comparative studies on the agronomic value of in vitro and conventionally propagated strawberry (*Fragaria × ananassa* Duch.) plants *Acta Sci. Pol. Hortorum Cultus* 14(3): 25-35.
15. Boxus P, Quoirin M, Laine M J (1977). Large scale propagation of strawberry plants from tissue culture. In: *Applied and fundamental aspects of plant cell, tissue and organ culture*. Reinert, J., Bajaj, Y.P.S. (eds.). New York, Springer-Verlag, 130-143.
16. Boxus P (1992). Mass production of strawberry and new alternatives for some horticultural crops. In: *Proceedings of the International Symposium on*

Transplant Production Systems. Kurata, K., Kozai, T. (eds). Yokohama, Japan
151-62, 21-26.

17. Ashrafuzzaman M, Faisal S M, Yadav D, Khanam D, and Raihan F (2013).
Micropropagation of strawberry (*fragaria ananassa*) through runner culture.
Bangladesh Journal of Agricultural Research 38(3): 467-472.
18. Dhar, M (1998). Techniques of vegetative and in vitro propagation of Jackfruit.
Ph.D. Thesis, Banggabandhu Shaikh Mujibar Rahman Agricultural University,
Salna, Gazipur, Bangladesh 120.
19. Mott R L (1981). Trees, In: Conger, B.V. (ed.), Cloning Agricultural Plants via
in vitro Techniques. CRC Press, Boca Ratan 217-254.
20. Chawla H S (2002). Introduction to Plant Biotechnology. Oxford & IBH
publishing Co. Pvt. Ltd., 66 Janapath, New Delhi 110001, India.
21. Abdullah G R, Al-Khateeb A A, Layous L N (2013). Response of the
Strawberry Cv. "Elsanta" Micro Propagation in vitro to Different Carbon
Sources and Concentrations. Jordan Journal of Agricultural Sciences, 9: 1.
22. Murashige T, Skoog F (1962). A revised medium for rapid growth and bioassays
with tobacco tissue cultures. Physiologia Plantarum, 15: 473-497.
23. Hu C Y, and Wang P J (1983). Meristem shoot tip and bud culture. *In*: Evans
DA, Sharp WR. Ammirato PV and Yamada Y (Eds) Hand book of Plant Tissue
Culture, Macmillan, New York I: 177-227.
24. Bhatt I D, Dhar U (2000). Micropropagation of Indian wild strawberry. Plant
Cell, Tissue and Organ Culture 60: 83-88.

25. Cononer R A, Litz R E (1978). In vitro propagation of papaya. Horticultural Science, 13: 241-242.
26. Teixetra S L, L. L. D. Silva L L D (1990). In vitro propagation of adults *Eucalyptus grandis* Hill Ex. Maiden from epicormic shoots. VII Intl. Cong. On Plant, Tissue and Cell Cult. (IAPTC), Amsterdam, 218.
27. Singh A K, Pandey S N (2004). Genotypic variation among strawberry cultivars for shoot organogenesis. Acta Horticulture, 662: 277-280.
28. Morozova T (2002). Genetic stability of Pure lines of *Fragaria vesca* L in micropropagation and long-term storage in vitro. Acta Horticulture 567: 85-87.
29. Boxus P (1999). Micropropagation of strawberry via axillary shoot proliferation. In: Plant Cell Culture Protocols. Methods in Molecular Biology. Part III. Plant Propagation In Vitro. Hall R. D. (ed.) Humana Press Inc., Totowa NJ 111: 103-114.
30. Litwińczuk W (2004). Field performance of 'senga sengana' strawberry plants (*Fragaria* × *ananassa* Duch.) Obtained by runners and in vitro through auxiliary and adventitious shoots. Electronic Journal of Polish Agricultural Universities, Horticulture, 7: 1.