

WATER REQUIREMENT OF SOLANACEOUS CROPS GROWN IN KASHMIR VALLEY BY FAO-56 PENMEN MONTEITH METHOD

Latief Ahmad, Sameera Qayoom, Baseerat Afroza, Owais A.Bhat and Nighat Mushtaq

Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir-Shalimar (J&K) 190025

Email id of corresponding author: drlatief@skuastkashmir.ac.in

ABSTRACT

Studies on water balance and irrigation at agricultural areas require accurate values of reference evapotranspiration. The values obtained can help farmers to determine the water requirement of these vegetable crops so that proper irrigation can be applied according to its growth stage and weather condition. The objective of this study was to calculate water requirements for four Solanaceous vegetable crops: Tomato, Potato, Chilli and Brinjal grown in temperate conditions of Kashmir Valley. Water requirements were calculated using the FAO-56 Penman-Monteith Method. The total water requirement for tomato, potato, chilli and brinjal was found 458 mm, 286 mm, 343 mm and 410 mm respectively. The maximum water was required by all crops during the mid-season stage comprising nearly 45-60% of the total water requirement of the crop.

Keywords: *Tomato; potato; chilli; brinjal; crop coefficient; evapotranspiration, Penman-Monteith Method*

INTRODUCTION

The family Solanaceae includes potato, tomato, eggplant, capsicum and chillies. These comprise of some of the most popular vegetables grown and consumed in Kashmir Valley. A large quantity of water is used by these crops for their growth and development. Nearly 90% of the water absorbed by roots is transpired back to the atmosphere. This increases the water requirement of these crops (Grace and Williams, 2004). Hence adequate water management is essential for proper growth and yield of these crops. Water stress affects the crop yields especially if it occurs at the key stages of growth (Salter and Good, 1967). Insufficient water

supply inhibits plant growth in terms of leaf area and plant height (Thompson and Chase; 1992 and Abdrabbo *et al.*,2007).

Crop water use is a function of evaporation (E) and transpiration (T) that fluctuates daily. Evapotranspiration (ET) is one of the most difficult components to be determined in the water balance as compared to the other components like precipitation or irrigation (Fisher *et al.*, 2005; Xu and Singh, 2005).The crop evapotranspiration (ET_c) is defined as “the ET rate of crop under standard conditions where there is no stress by water quality constraints, pests, or inadequate soil fertility” (Allen *et al.*, 1998). Correct estimation of Etc and thus the crop water requirement is essential for determination of water availability, crop water balance and crop water requirements (Pereira *et al.*, 1999).

Crop water requirement is defined as the “depth of water required by plants to compensate the water loss via ET so that plants are able to grow optimally while soil water balance shows amount of soil water added, removed or stored in volume of soil during a time period” (Allen *et al.*, 1998).ET_c can help farmers in order to decide irrigation period and irrigation frequency to increase crop yields and profits while minimizing input costs, energy and environmental impacts.

A number of models for the estimation of ET₀ are available but the results are inconsistent with the values of ET₀. This is due to the differences in modeling assumptions and input data requirements. Another reason for inconsistency is that the models are developed for specific regions (Lu *et al.*, 2005).FAO-56 Penman-Monteith due to close approximation to lysimeter observations is considered to be the best approach for estimating ET₀ (Droogers and Allen, 2002; Popova *et al.*,2006).

STUDY AREA AND DATA USED

The state of Jammu and Kashmir falls politically within the Indian union and is located in the north western extremity of the Himalayas between 32°10' and 37°10'N latitudes and 72°30' and 80°30'E longitudes. Covering a total area of about 2, 22,236 km², the state is divided into three provinces (Figure 1) viz. Jammu (26293 km²), Ladakh (59146 km²) and Kashmir Valley(15948 km²), with each province differing markedly from one another in its climate, vegetation and culture. The altitude of the irrigated temperate region of Kashmir Valley ranges from 1524–2100 m AMSL .The weather data for the present study was obtained from

Regional Meteorological Centre, Srinagar. The data set consisted of daily values of maximum temperature, minimum temperature, morning relative humidity, afternoon relative humidity, sunshine hours and wind speed for the period 1992-2017.



Figure 1. Study Area

METHODOLOGY

The reference evapotranspiration and the crop water requirement of the solanaceous crops were estimated by means of FAO-56 Penman-Monteith Method (Allen et al, 1998). The FAO-56 Penman-Monteith Method has been established as a standard method for calculation of reference evapotranspiration (Jensen et al, 1990; Itenfisu et al, 2003; Lopez-Urrea et al, 2006). The method has also been verified to be more suitable for Indian conditions than FAO-24 Penman method (Kashyap and Panda, 2001). The irrigation water use efficiency can be enhanced by the application of FAO-56 PM Method and thus the distribution of water at different levels can be improved (Nandagiri and Koor, 2006). The method has been applied to determine the water requirement of number of crops grown in Kashmir valley like cereal crops, pulses apple and saffron (Ahmad et al., 2017a, 2017b, 2017c).

FAO-56 Penman-Monteith Method

The penman method was developed to calculate the evaporation from open water surfaces (Penman, 1948). The model was then modified by Monteith for its application to cropped surfaces as well (Monteith, 1965). The equation of modified Penman-Monteith method also known as FAO-56 Penman-Monteith method is given as:

$$ET_0 = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T_{mean} + 273} U_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34U_2)}$$

Where,

- ET_0 = reference evapotranspiration [mm day^{-1}],
 R_n = net radiation at the crop surface [$\text{MJ m}^{-2} \text{day}^{-1}$],
 G = soil heat flux density [$\text{MJ m}^{-2} \text{day}^{-1}$],
 T_{mean} = mean daily air temperature at 2 m height [$^{\circ}\text{C}$],
 U_2 = wind speed at 2 m height [m s^{-1}],
 e_s = saturation vapor pressure [kPa],
 e_a = actual vapor pressure [kPa],
 $e_s - e_a$ = saturation vapor pressure deficit [kPa],
 Δ = slope vapor pressure curve [$\text{kPa } ^{\circ}\text{C}^{-1}$],
 γ = psychrometric constant [$\text{kPa } ^{\circ}\text{C}^{-1}$].

CROP COEFFICIENT (K_c)

Crop coefficients (K_c) are used with reference evapotranspiration (ET_0) to estimate specific crop evapotranspiration rates (ET_c) and are defined as the ratio of ET_c to ET_0 . Crop coefficients vary with crop type, the growth stage of the crop, weather and irrigation method, and with some cultural practices.

Crop coefficient is an important parameter for indirect estimation of crop evapotranspiration. It gives information on water requirement of crop and the impact of other parameters on it, for instance, biological characteristics of crops, level of crop yield, conditions of soil tillage etc. (Wright, 1982; Liu et al., 1998; Haiyanand Xiyun, 2006). The values of crop coefficients for the crops under study at different stages of growth were obtained from FAO.

Crop Water Requirement

The canopy properties and aerodynamic resistance of the crop results in difference between the reference evapotranspiration and crop water requirement. The water requirement of the crops is also different for different stages of growth. Crop Water Requirement is thus computed from reference evapotranspiration and crop coefficient as (Doorenbos and Pruitt, 1977):

$$ET_c = K_c \times ET_0$$

Where,

ET_c = crop evapotranspiration/Crop water requirement (mm day^{-1}),

ET_o = reference evapotranspiration (mm day^{-1}),

K_c = crop coefficient.

Solanaceous Vegetables

Tomato (*Solanum lycopersicum* L.)

Tomato is the most important fruit vegetable grown throughout the world and ranking second in importance to potato in many countries. It is widely accepted as 'Protective Food' and is grown extensively in the state of Jammu and Kashmir. Tomato is a warm season crop and cannot withstand severe frosts. A warm sunny weather is most suited for its proper ripening, color development, quality and high yields. The optimum temperature for germination ranges from 18°C to 26°C . The irrigation should be so arranged that the soil remains moderately moist. Excessive irrigation induces the plant to vine and dropping off the blossoms. During summer season, irrigation at every 3 to 4 days interval is necessary, whereas for spring season crop 10 to 15 days interval is sufficient. Subsequent irrigations are given according to need of the crop. Irrigation at flowering and fruiting stage is imperative for higher quality yields.

Brinjal / Eggplant (*Solanum melongena* L.)

Brinjal is the most common vegetable crop grown widely in India for its varied shape, size and colour of fruits. It is a popular vegetable due to its high productivity and wide adaptation. It is a warm season crop and requires a relatively long growing season with plenty of sunshine and moderate day temperature of 21° - 27°C . Brinjal being moderately deep rooted crop needs irrigation at frequent intervals. During summer, irrigation is applied at 4-5 days interval and during cold weather irrigation is applied at 10-15 days interval. Furrow method of irrigation is the most common method.

Chilli (*Capsicum annuum* L.)

Chilli or hot pepper is a tropical vegetable crop commonly used throughout the world as a spice for its pungency and red ripe color of dried fruits. It is also cultivated for its green immature fruits. It performs well in warm humid tropical and subtropical regions. Optimum temperature for its successful cultivation is 20 - 33°C . Extreme temperature (below 15°C and above 35°C) significantly reduces the reproductive growth and pollen viability. Humidity favors

vegetative growth but frost is injurious. Low temperature at fruit ripening stage is likely to delay the color development in fruits. It performs well in areas with annual rainfall ranging from 600 - 1200 mm spread over a period of 4- 5 months.

Chilli needs judicious irrigation for proper growth and yield. Frequent and heavy irrigations induce lanky growth and cause flower shedding. Critical stages of irrigation are flowering, fruit setting and fruit development and after periodical harvests. During summer, irrigation is applied at 4-5 days interval and during cold weather irrigation is applied at 10-12 days interval. Furrow method is the most common method of irrigation.

Potato (*Solanum tuberosum* L.)

Potato is the fourth major food crops of the world after rice, wheat and maize. Being a major source of carbohydrates, it is often used as a substitute for cereals and is grown in almost every country. Potato is a cool season crop; however at low temperature vegetative growth is restricted and at freezing point irrecoverable frost injury occurs to the plant. Long days coupled with high temperature conditions promote vegetative growth without formation of tubers and short day with low temperature induce tuberization. Optimum temperature for good plant growth is 15-20°C. Most of the cultivars do not tuberize when the prevailing night temperature is above 21 °C.

Adequate irrigation is essential for proper growth and yield of potato. Stolon formation, tuber initiation and tuber development stages are the critical stages for irrigation. Irrigating the crop at an interval of 7-10 days after earthing up is essential. The level of irrigation water must not exceed 3/4th part of ridges. Total water requirement for the crop varies between 350-550 mm.

RESULTS AND DISCUSSION

The total water requirement for the different crops of Solanaceous family under Kashmir valley conditions was determined using FAO-56 Penman-Monteith Method. The water requirement was low during initial stages of growth in all the crops. This is due to low canopy cover of the crop during this stage. With the advancement of vegetative stage and development of crop, the percent ground cover, leaf area and crop height increased resulting in increase in water requirement. The water requirement was maximum during the mid-season stage i.e., vegetative

growth, flowering and fruiting period. This comprised nearly 40-60% of the total water requirement of the crops. The water requirement again depleted with the decrease in canopy cover during the late season stage. In this stage, the leaves and roots begin to dry and finally the crops die. The relative water requirement during different stages is shown in Figure 2.

The water requirement of the four crops namely tomato, potato, chilli and brinjal during different stages of growth is given in Table 1.

Table 1 Crop water requirement of different crops of Solanaceae family in Kashmir Valley

	Tomato	Brinjal	Potato	Chilli
Initial	49	38	22	20
Development	112	112	37	58
Mid-Season	195	191	131	198
Late-season	102	68	97	68
Total	457	410	286	343

The water requirement is represented graphically in Figure 3. The water requirement was lowest for the potato crop being 286 mm. while the highest water requirement was obtained for the tomato crop being 457 mm. the water requirement for brinjal and chilly was found to be 410 and 340 mm respectively.




Figure 2 Relative water requirement during different growth stages




Figure 3 Water Requirement (mm) of different Solanaceous crops for different stages of growth.

CONCLUSION

In the present study, the crop water requirement of four Solanaceous vegetable crops: Tomato, Potato, Chilli and Brinjal grown in temperate conditions of Kashmir Valley was determined based on crop coefficient (K_c) and reference evapotranspiration (ET_0). Results showed that these crops required moderate amount of water throughout their period of growth (286-457 mm). The maximum water requirement was during the vegetative growth stage for all the crops. Efficient irrigation practices, especially at critical stages of crop growth, are necessary to ensure good yield particularly in absence or failure of rainfall during the growing season. The data on potential evapotranspiration and the Crop Water Requirement is essential for determination of irrigation water requirements. This in turn is helpful for efficient scheduling of irrigation for harvesting maximum benefits.

REFERENCES

- Abdrabbo, M. A. A., M. k. Hassanein and M. A. Medany. 2007. Effect of Irrigation Regime and Compost Level on Potato Production in Northern Delta, Egypt. *Proceeding of the 7th African Potato Association Conference /Exhibition, Alexandria, Egypt*.185-197.
- Ahmad, L., Parvaze, S. and Kanth, R.H., 2017a. Crop water requirement of major crops of Srinagar, Kashmir (J&K). *J. Exp. Agr*, 15(2), pp.1-9.
- Ahmad, L., Parvaze, S., Parvaze, S. and Kanth, R.H., 2017b. Reference evapotranspiration and crop water requirement of apple (*Malus Pumila*) in Kashmir Valley. *Journal of Agrometeorology*, 19(3), pp.262-264.
- Ahmad, L., Parvaze, S., Parvaze, S. and Kanth, R.H., 2017c. Crop water requirement of saffron (*Crocus sativus*) in Kashmir valley. *Journal of Agrometeorology*, 19(4), pp.380-381.
- Allen, R.G., Pereira, L.S., Raes, D. and Smith, M., 1998. Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. *FAO, Rome*, 300(9), p.D05109.
- Doorenbos J, Pruitt WO. Guidelines for predicting crop water requirements. *FAO, UN, Irrigation and Drainage. Paper No.24*.(2nd Ed.) FAO, Rome; 1977.
- Droogers, P., Allen, R.G., 2002. Estimating reference evapotranspiration under inaccurate data conditions. *Irrig. Drain. Syst.* 16, 33-45.
- Fisher, J.B., DeBiase, T.A., Qi, Y., Xu, M., Goldstein, A.H., 2005. Evapotranspiration models compared on a Sierra Nevada forest ecosystem. *Environ. Model. Softw.* 20, 783-796.
- Grace, J., Williams, M., 2004. Understanding and parameterizing the soil-water-atmosphere transfer through vegetation. In: Feddes et al. (Org.) *Unsaturated-Zone Modeling: Progress, Challenges and Applications*. Springer, Wageningen, Netherlands, 73-94.
- Haiyan, M. and Xiyun, J., 2006. Research progress of the crop water demand calculation. *Water Sciences and Engineering Technology*, 5, pp.5-7.

- Itenfisu, D., Elliott, R.L., Allen, R.G. and Walter, I.A., 2003. Comparison of reference evapotranspiration calculations as part of the ASCE standardization effort. *Journal of Irrigation and Drainage Engineering*, 129(6), pp.440-448.
- Jensen, M.E., Burman, R.D. and Allen, R.G., 1990. Evapotranspiration and irrigation water requirements. ASCE.
- Kashyap, P.S. and Panda, R.K., 2001. Evaluation of evapotranspiration estimation methods and development of crop-coefficients for potato crop in a sub-humid region. *Agricultural water management*, 50(1), pp.9-25.
- Liu, Y., Teixeira, J.L., Zhang, H.J. and Pereira, L.S., 1998. Model validation and crop coefficients for irrigation scheduling in the North China Plain. *Agricultural Water Management*, 36(3), pp.233-246.
- López-Urrea, R., de Santa Olalla, F.M., Fabeiro, C. and Moratalla, A., 2006. Testing evapotranspiration equations using lysimeter observations in a semiarid climate. *Agricultural water management*, 85(1), pp.15-26.
- Lu, J., Sun, G., McNulty, S.G., Amatya, D.M., 2005. A comparison of six potential evapotranspiration methods for regional use in the southeastern United States. *J. Am. Water Resour. Assoc.* 41, 621–633.
- Monteith J.L. 1965; Evaporation and environment. Rages, Inc., New York in GF Fogg, ed. The state and movement of water in living organisms. Academic Press. 205-234.
- Nandagiri, L. and Kovoov, G.M., 2006. Performance evaluation of reference evapotranspiration equations across a range of Indian climates. *Journal of irrigation and drainage engineering*, 132(3), pp.238-249.
- Penman H.L. 1948. Natural evaporation from open water, bare soil and grass. In Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences. *The Royal Society*, 3(1032):120-145.
- Pereira, L.S., Perrier, A., Allen, R.G., Alves, I., 1999. Evapotranspiration: concepts and future trends. *J. Irrig. Drain. Eng.* 125, 45–51.
- Popova, Z., Kercheva, M., Pereira, L.S., 2006. Validation of the FAO methodology for computing ETo with limited data: Application to south Bulgaria. *Irrig. Drain.* 55, 201–215.
- Salter, P.J. and J.E Good (1967): Crop response to water at different stage of growth. *Common Wealth Agric.* Bur. Farham Royal, Bucks England, 246.
- Thompson J.A. and D. L. Chase (1992): Effect of limited irrigation as growth and yield of a semidwarf wheat in southern New South Wales Australian *J. of Exp. Agric.*, 32:725–730.
- Xu, C.-Y., Singh, V., 2005. Evaluation of three complementary relationships evapotranspiration models by water balance approach to estimate actual regional evapotranspiration in different climatic regions. *J. Hydrol.* 308, 105–121.
- Mohammadian, M., Arfania, R., & Sahour, H. (2017). Evaluation of SEBS algorithm for estimation of daily evapotranspiration using landsat-8 dataset in a Semi-Arid Region of Central Iran. *Open Journal of Geology*, 7(03), 335.
- Tehrani, E. N., Sahour, H., & Booij, M. J. (2018). Trend analysis of hydro-climatic variables in the north of Iran. *Theoretical and Applied Climatology*, 1-13.