

Assessment of Protein C and Protein S of Pregnancy Loss Victims

Igwe, Chioma Ada¹ Adias Teddy Charles² Eze Evelyn Mgbeoma³ Nwachuku Edna Ogechi³

1 Department of Haematology and Blood Group Serology, Federal Medical Centre, Umuahia, Abia State, Nigeria.

2 Federal University, Otueke, Bayelsa State, Nigeria.

3 Department of Medical Laboratory Science, Rivers State University, Port Harcourt, Nigeria.

ABSTRACT

Background: The haemostatic changes that result in thrombophilia during the pregnant state have been linked to pregnancy loss. **Objective:** Assessment of Protein S, and Protein C assays in pregnancy loss in Abia State, South East, Nigeria. **Materials and Methods:** This was a cross-sectional study involving women in their reproductive years. Study population was stratified into 3 groups and the Protein C and Protein S concentrations measured and compared among the three groups. **Results:** A total of 130 apparently healthy Nigerian women of child-bearing age were enrolled in the study. The study groups consisted of 70 women who had just lost a pregnancy, 30 women with normally progressing pregnancy and 30 nonpregnant women. The protein C concentration for the pregnancy-loss subjects was significantly lower than that of the normal pregnancy at $p \leq .01$ while that of Protein S showed non-significance ($p > 0.05$). **Conclusion:** Protein C deficiency is associated with increase in pregnancy loss. **Key words:** Pregnancy loss, Protein C, Protein S.

INTRODUCTION

Human reproduction is accompanied by physiological changes that occur primarily to nurture pregnancy as they ensure the proper development, growth and ultimate survival of the embryo/foetus. These changes include (but are not limited to) cardiovascular, respiratory, metabolic/endocrinal, renal, haematologic and immunologic changes (Priya *et al.*, 2016). Some of these changes influence normal biochemical values while others may mimic symptoms of medical disease (Priya *et al.*, 2016). The pregnant state is not failure-proof. Pregnancy loss (also termed miscarriage or abortion) is the spontaneous loss of an intra-uterine pregnancy without outside intervention, and occurring before 20 weeks of gestation (Abdullah and Mahdi, 2013). Pregnancy is associated with significant changes in the haemostatic profile (Chandra *et al.*, 2012) creating a hypercoagulable state, and which has been attributed to alterations of the proteins that govern blood clotting (Gersh *et al.*, 2009), thus enforcing a thrombophilic state (Walker, 2000). There are varied reports regarding the levels of Protein C and Protein S during pregnancy. Oruc *et al.*, (2000), reported reduced levels of both Protein C and Protein S in pregnancy (most probably dilutional). Said *et al.*, (2010) and Petrozza *et al.*, (2014) reported of reduced levels of Protein S during pregnancy. While Petrozza *et al.*, (2014) noted no change in Protein C levels in pregnancy when compared with the nonpregnancy, Said *et al.*, (2010) observed a significant increase in Protein C activity during pregnancy.

Women with a history of recurrent pregnancy loss (RPL) are in a procoagulant state even when they are not pregnant (Petrozza and Berin, 2014). This thrombophilic state has been linked to pregnancy loss (Mekaj *et al.*, 2015). In this study we investigated the relationship between pregnancy loss and deficiencies of both Protein C and Protein S in women of Abia State, Nigeria.

MATERIALS AND METHODS

A total of 130 apparently healthy women of childbearing age (18-45 years) were enrolled in this study. It was designed as a cross-sectional study. The study population was divided into three groups as follows: pregnancy-loss (70 subjects), normal pregnancy (30 subjects) and non-pregnant control (30 subjects). Study lasted the period between December 2017 and March 2018.

Study Setting

The study took place at the following hospitals: Federal Medical Centre, Umuahia, Nazareth Specialist Hospital Aba, (a specialist gynaecological clinic), General Hospital, Aba and General Hospital, Ohafia, all in Abia state, Nigeria.

Blood collection

3mls of the blood were collected via venepuncture and put into plain vacuum containers. For the pregnancy loss subjects, this collection was done within 48 hours of pregnancy loss. The serum was retracted after 30 minutes of clotting. The serum was used for the estimation of Protein C and Protein S.

Laboratory Analysis

The microtitre plate provided in this kit has been pre-coated with antibody. Add standard, samples and HRP conjugated antibody to wells according to the manufacturer, Melsin Laboratories, China. After incubation and washing to remove the uncombined enzyme, chromogen Solution A and B was added. The colour of the liquid changed to blue. At the effect of acid, the colour finally became yellow. The colour change is measured spectrophotometrically

at a wavelength of 450nm. The concentration of Protein C and Protein S in the sample was then determined by comparing the O.D of the samples to the standard curve.

Normal values

Normal values of Protein S for Caucasians is 15-25 μ /ml while that for Protein C is 3.9-5.4 μ /ml. But the respective levels are lower in the pregnant state as well as in Black Africans (Jerrard-Dune *et al*, 2003). Aside, Momodu and Buseri (2015) attributed inconsistencies in the reference values of these anticoagulants to differences in the sensitivities and specificities of the reagents used as well as to assay techniques. In this study assay was done with ELISA technique and assays below 2.5 μ /ml for Protein S and 0.7 μ /ml for Protein C were excluded from the study.

Exclusion Criteria

Those excluded from this study included subjects who were reactive to HIV 1 and II, HbsAg, HCV as well as smokers, patients with co-morbidities like hypertension, pelvic inflammatory disease, known thrombotic illness and patients with abnormal reproductive hormones. Also excluded are subjects with assays below 2.5 μ /ml for Protein S and 0.7 μ /ml for Protein C.

Ethical Approval

Ethical approval was obtained from the Ethics Committees of the Federal Medical Centre, Umuahia, the Hospitals Management Board, Umuahia and the Nazareth Specialist Hospital, Aba.

Statistical Analysis

Statistical analysis was done using SPSS windows version 20 (IBM Corporation, 2011). Data was grouped into pregnancy-loss, normal pregnancy and non-pregnant control subjects. The Kolmogorov-Smirnov test was used to determine normality. Parametric data were expressed as $\pm$ SD. Parameters were skewed. Hence, they were expressed as median, 2.5th (p2.5) and 97.5th (p97.5) percentile, and were log-transformed prior to analysis. ANOVA and Games-Howell post hoc procedures were used to test for group differences. Pearson's correlation was used to determine relationship between parameters. The predictive analysis of parameters on pregnancy loss was explored via binary logistic regression model. Data was considered significant at $P < .05$.

RESULTS

A total of 130 apparently healthy women of childbearing age (18-45 years) were recruited for this study. Subjects consisted of 70 (53.85%) pregnancy loss subjects, 30 (23.08%) normal pregnancy subjects, and 30 (23.08%) non-pregnant control subjects. The mean $\pm$ standard deviation of age for the test subjects (Pregnancy loss subjects) was 32.46 ± 5.41 years, while those for normal pregnancy and nonpregnant controls were 31.67 ± 4.25 years and $28.30 \pm$

3.72 years respectively. The majority of this group of subjects (58.57%) were primigravida (gravida I) patients, 31.43% were gravida II, while 10% were gravida III patients (Table 1). Results from Table 1 also show that among the pregnancy-loss group, 41 women (58.57%), were experiencing their miscarriage in their very first pregnancy in life, 22 (31.43%), in their second pregnancy and 7 women (10%) in their third pregnancy. Results show that 65 (92.86%) miscarriages in this group occurred during the first trimester and 5 (7.14%) occurred during the second trimester. Among this group too, 54 (77.14%) women have had one pregnancy loss, 14 (20%) have experienced two pregnancy losses while 2 (2.86%) have experienced three pregnancy losses (in all, the present miscarriage inclusive) (Table 1).

Table 1: Essential Statistical Details of Pregnancy Loss Subjects

Parameter	Freq.	%	Mean $\pm$ SD
Pregnancy Loss Subjects			
• Total	70	53.85	NA
Pregnancy Status (Gravidity)			
• Primigravida (Gravida I)	41	58.57	NA
• Gravida II	22	31.43	NA
• Gravida III	7	10.00	NA
Trimester of Pregnancy Loss			
• 1 st Trimester	65	92.86	NA
• 2 nd Trimester	5	7.14	NA
• 3 rd Trimester	NIL	NIL	NA

**Number of Pregnancy Loss(es)
experienced by subject**

• 1 pregnancy loss	54	77.14	NA
• 2 pregnancy losses	14	20.00	NA
• 3 and above	2	2.86	NA

Age (y)

32.46 ± 5.41

Key: Freq. = frequency, y = years, NA = not applicable, NIL = nil, % = percentage, SD = standard deviation, primigravida (gravida I) = pregnant for the first time, gravida II= pregnant for the second time, gravida III = pregnant for the third time

Table 2: Comparison of Protein C and Protein S concentrations between Pregnancy-Loss, Normal Pregnancy, and Non-pregnant Subjects.

Parameter		Pregnancy Loss (n=70)	Normal Pregnancy (n=30)	Non-pregnant (n=30)	F	P
Protein S (µg/ml)	<i>Mdn</i>	9.00	9.30	8.60	0.27	0.77
	P2.5 – P97.5	6.04 – 11.50	4.0 – 9.30	2.60 – 8.60		
Protein C (µg/ml)	<i>Mdn</i>	0.90	1.20	1.20	11.74	0.00
	P2.5 – P97.5	0.70 – 3.89	0.80 – 1.20	0.70 – 1.20		

Key: n = number of subjects, *Mdn* = median, P2.5 = 2.5th percentile, P97.5 = 97.5th percentile, *F* = F-test statistic, *p* = error probability, ** Significant difference observed at *p* < .01, using ANOVA.

Table 3: Post hoc Testing of Protein C and Protein S concentrations between Pregnancy-Loss, Normal Pregnancy, and Non-pregnant Subjects.

Immunological Parameter	Post hoc Pair	95% CI	P
Protein S (µg/ml)	PL vs Norm P.	-0.08 – 0.06	0.91
	Norm P. vs Non-P.	-0.08 – 0.12	0.86
	PL vs Non-P	-0.06 – 0.08	0.95

Protein C (µg/ml)	PL vs Norm P.	-0.25 – -0.05	0.00**
	Norm P. vs Non-P.	-0.17 – 0.12	0.92
	PL vs Non-P	-0.30 – -0.05	0.00**

Key: PL = Pregnancy Loss, Norm P. = Normal Pregnancy, Non-P. = Non-pregnant, p = error probability, 95% CI = 95% Confidence Interval of the Difference, ** Significant difference observed at $p \leq .01$, * Significant difference observed at $p \leq .05$. All *post hoc* testing were done using Turkey HSD and Games-Howell methods as applicable.

Table 4: Logistic Regression on the Relationship between Protein S/Protein C and Pregnancy Loss

	95% CI for exp b			
	$B(SE)$	Lower	exp b	Upper
Included				
Constant	-3.57(1.14)**		0.02	
Protein S	0.07(0.06)	0.96	1.07	1.19
Protein C	0.31(0.17)	0.98	1.37	1.91

Key: Key: exp b = change in odds per unit change in predictor, B = coefficients of predictors, SE = standard error, 95% CI = 95% Confidence Interval $R^2 = .09$ (Cox & Snell), .13 (Nagelkerke), Model $\chi^2(3) = 9.28$, $p = .03$. ** $p < .01$.

DISCUSSION

Heritable thrombotic states have been linked to pregnancy loss, most especially recurrent pregnancy loss (RPL) (Rey *et al.*, 2003; Van Dreden *et al.*, 2012; Yilmaz *et al.*, 2015). RPL is thought to have multiple aetiologies including chromosomal abnormalities, immune dysfunction, thrombophilic disorders and hormonal disturbances (Larsen *et al.*, 2013). Thrombophilia is a predisposition to thrombosis and describes a tendency of increased blood clotting (Walker, 2000). It creates a hypercoagulable state that leads to arterial and/or venous thrombosis at the site of implantation or in the placental blood vessels (Abu-Heija, 2014). It has been reported that defects are noticed in the haemostatic variables just before miscarriages take place (Petrozza and Berin, 2014).

Protein C and Protein S are natural anticoagulants (Parand *et al.*, 2013). When activated, Protein C inhibits clotting by proteolytic cleavage (and thus deactivation) of factors Va and VIIIa, using protein S as a co-factor (Esmon, 2001). Factors Va and VIIIa are important in the coagulation cascade and their inhibition helps to prevent thrombosis, thereby helping to keep blood in a fluid state. Thus deficiencies of Protein C and Protein S results in the development of a procoagulant state (Singla and Jain, 2018), thus worsening the procoagulant state already existing in pregnancy. Protein C equally enhances the viability and growth of trophoblast cells (Iserman *et al.*, 2003), thus ensuring foetal survival. This shift to a procoagulant state is evidenced by a reported shift in the thromboxane/prostacycline ratio in favour of thromboxane which is a known prothrombotic agent, resulting in vasospasm and platelet

aggregation in the trophoblasts, eventually leading to the development of microthrombi and placental necrosis (Singla and Jain, 2018).

In this study, Protein S and Protein C assays were compared between pregnancy-loss, normal pregnancy, and non-pregnant control subjects as shown in Table 2 and Table 3. Protein C concentration for the pregnancy-loss subjects (*Mdn*: 0.90 µg/ml, *p*_{2.5} – *p*_{97.5}: 0.70 – 3.89 µg/ml) was significantly lower than that of the normal pregnancy (*Mdn*: 1.20 µg/ml, *p*_{2.5} – *p*_{97.5}: 0.80 – 1.20 µg/ml) and non-pregnant control subjects (*Mdn*: 1.20 µg/ml, *p*_{2.5} – *p*_{97.5}: 0.70 – 1.20 µg/ml) (*p* < .01, respectively). However Protein C assay showed no significant difference between the normal pregnancy and non-pregnant control groups (*p* > .05). However Protein S showed non-significance (*p* > .05) when compared across all groups.

Gene polymorphism occurring in either Protein C or Protein S can lead to abnormalities in either output or functional expressivity of these anticoagulant proteins (Chumpa *et al.*, 2006). This may partly account for any discordant genetic activity expressed by Protein S or Protein C in our study. In fact a subset of subjects in the pregnancy loss group showed increased Protein C activity.

Several studies have reported a positive correlation between RPL and heritable thrombotic states (Van Dreden *et al.*, 2012; Yilmaz *et al.*, 2015). However the existence of conflicting data in literature regarding any association between miscarriage on one hand and Protein C as well as Protein S on the other hand is acknowledged (Mekaj *et al.*, (2015). In their work, Mekaj *et al.*, (2015) found that Protein C and Protein S are causal factors for miscarriage. In contrast with our work, Parand *et al.*, (2013), found a higher frequency of Protein S deficiency in patients with RPL

compared with controls but found that the frequency of Protein C, was not significantly different between patients with RPL and healthy women. Protein C and Protein S deficiencies are inherited independently (Chaudhari *et al.*, 2017). Combined deficiencies are rare and come with increased and earlier onset of risk of thrombosis (Chaudhari *et al.*, 2017). In their own work, Ogasarawa *et al.*, (2001) found no differences in the subsequent miscarriage rates between patients with abnormal and normal values of Protein C and Protein S. In the majority of women with inherited thrombophilia, pregnancy is uneventful (Parand *et al.*, 2013). However the risks for miscarriage and some other complications of pregnancy are increased in carriers of thrombophilia. Why certain women with thrombophilia present with gestational vascular complications is still unknown but may be related to a combined effect with another inherited or acquired prothrombotic risk either systemic or localised at the level of the placenta (Altintas *et al.*, 2007) which leads to compromised foetal viability and loss.

Result also showed (from Table 4) that neither Protein C nor Protein S showed significant likelihood of predicting pregnancy loss ($P > 0.05$). However, the overall model was significant with small effect sizes (Model $X^2(3) = 9.28$, $p = .03$, $R^2 = .09$ [Cox & Snell], .13 [Nagelkerke]).

Conclusion

The study assessed serum concentrations of the natural anticoagulants Protein C and Protein S in pregnancy loss, normal pregnancy and nonpregnant controls. Results showed significant association between pregnancy loss and Protein C deficiency. But this association lacks predictive value. However a subset of the pregnancy loss victims showed heightened activity of Protein C activity. In contrast Protein S levels remained stable during the period of pregnancy and showed non-significant variation across all groups.

Reference

- Priya, S., Nelson-Piercy, C., Heli, T. & Alexandra, M. (2016). Physiological changes in pregnancy. *Cardiovascular Journal of Africa*, 27(2), 89-94.
- Abdullah, GA. and Mahdi, N. (2013.) The role of cytokines among women with spontaneous abortion. *Medical Journal of Islamic World Academy of Sciences*, 21(3): 119-124
- Chandra, S., Tripathi, AK., Mishra, S., Amzarul, M. And Vaish, AK. (2012). Physiological changes in haematological parameters during pregnancy. *Indian Journal of Hematology and Blood Tranfusion*; 28(3): 144-146
- Petrozza, JC. & Berin, I. (2014). Recurrent early pregnancy loss. *emedicine.medscape.com*. Accessed 20/9/2016.
- Gersh, KC., Nagawami, C. and Wiessel, JM. (2009). Fibrin network structure and clot mechanical properties are altered by incorporation of erythrocytes. *Thrombosis and Hemostasis*, 102:1169-1175.
- Walker, I. D. (2000). Thrombophilia in pregnancy. *Journal of Clinical Pathology*, 53: 573-580.
- Mekaj, Y., Lulaj, S., Daci, F., Rafuna, N., Miftari, E., Hoxha, H., Silamniku, X. & Mekaji, A. (2015). Prevalence and role of antithrombin III, protein C and protein C resistance in Kosovo women with recurrent pregnancy loss during the first trimester of pregnancy. *Journal of Human Reproductive Sciences*, 8(4): 224-229.
- Rey, E., Khan, SR., Shrier, I. and David, M. (2003). Thrombophilic disorders and foetal loss. *Lancet*, 361:901-8.
- Van Dreden, P., Woodhams, B., Rousseau, A., Favier, M. and Favier, R. (2012). Comparative evaluation of tissue factor and thrombomodulin activity changes during normal and idiopathic early and late foetal loss. The cause of hypercoagulability. *Thrombosis Research*, 129:1787-792.
- Yilmaz, M., Delibas, I.B., Isaoglu, U., Ingec, M., Borekci, B. and Ulug, P. (2015). Relationship between mean platelet volume and current miscarriage: a preliminary study. *Archives of Medical Science*, 11(15):989-993.
- Larsen, E.C., Christiansen, O.B, Kolte, A.M. & Macklon, N. (2013). New insights into mechanism behind miscarriage. *BMC Medicine*, 11: 54.

Abu-Heija, A. (2014). Thrombophilia and recurrent pregnancy loss: is heparin still the drug of choice? *Sultan Qaboos University Medical Journal*, 14(1): 26-36.

Parand, A., Zolghadri, J., Nezam, M., Afrasiabi, A., Haghpanah, S. & Karimi, M. (2013). Inherited thrombophilia and pregnancy loss. *Iran Red Crescent Medical Journal*, 15(12): e13708

Chaudhari, HK., Shah, PK., Pai, KR. & D'Souza, BD. (2017). Combined protein C and protein S deficiency with pregnancy. *International Journal of Reproduction, Contraception, Obstetrics and Gynaecology*, 5(7): 2450-2452.

Ogasawara, MS., Aoki, K., Katano, K., Ozaki, Y. and Suzumori, K. (2001). Factor XII but not protein C, protein S, antithrombin III, or factor XIII is a predictor of recurrent miscarriage. *Fertility and Sterility*, 75:5.

Altintas, A., Pasa, S., Akdeniz, N., Cil, T., Yurt, M. & Ayyildiz, O. (2007). Factor V Leiden and G20210 A prothrombin mutations in patients with recurrent pregnancy loss: data from the southeast of Turkey. *Annals of Haematology*, 86(10): 727-731.

Iserman, B., Sood, R., Pawlinski, R., Zogg, M., Kalloway, S., Degen J. L Mackman, N. & Wevler, H: The Thrombomodulin Protein C System is Essential for the Maintenance of Pregnancy. *Nat Medicine* 9(3): 163-80

Esmon, C. (2001). Protein C Antiway Last Pathway and Its Role in Controlling Microvascular Thrombosis and inflammation. *Critical Care medicine* 29 (Suppl 7): 48

Jerrard-Dune, P., Evans, A., McGovern, R., Hajat, C., Kalra, L., Rudd, AG., Wolfe, C. D. and Markus, H. S. (2003), Ethnic Differences in Risk of Thrombophilia: Implications for Stroke in a Multiethnic population: the South London Ethnicity and Strokes Study. *Stroke*, 34(8), 1821-6

Chumpa, w., Peerapittayamongkol, C., Angchaisuksiri, P., Komansin, N., Muta, K., Kuaha, K., Lida, H., Innove, S., wada, Y., Kurihara, M, Hamasaki, N., & Fucharoen, S. (2006). Single nucleotide Polymorphisms and Haplotypes of Protein C and Protein S Genes in the Thai Population. *Blood Coagulation & Fibrinolysis*, 17(1), 13-18.

Oruc, S., Saruc, M., Koyuncu, F. M. & Ozdemir, E. (2000). Changes in the Plasma Activities of Protein C and Protein S During Pregnancy. *Australian N Z Journal of Obstetrics and Gynaecology* 40(4), 448-50

Petrozza, J. C & Berin, I. (2014). Recurrent Early Pregnancy Loss . eMedicine.medscape.com. Accessed 20/9/2016.

Momodu, I & Buseri, F. I. (2015). Protein C and Antithrombin III Activities in Neatly Nigerian Women. (2015). International Journal of Haematology, Research, (1), 20-23.