

Original Research Article

ASSESSMENT OF THE PREVALENCE OF *Chlamydia trachomatis* INFECTION IN DONATED BLOOD AT REGIONAL BLOOD TRANSFUSION CENTRE, ELDORET, KENYA.

ABSTRACT

Aim: To detect *Chlamydia trachomatis* in donated blood at Regional Blood Transfusion center, Eldoret.

Methodology: A random selection blood sample was done for this study for a period of one month. Blood bags were numbered 1-87 and picked amongst unnumbered blood bags by each donor in order of donation. The samples obtained were subjected to centrifugation and serum used in the immunoassay test by using the *Chlamydia trachomatis* IgG/IgM Elisa test kit.

Result: This study showed that *C. trachomatis* was present in 8% (7 samples) of the blood samples analyzed and 92% were found to be negative.

Conclusion: With 8% of the samples testing positive for *Chlamydia trachomatis* this indicated that some of the blood donated to the Regional Transfusion Center is contaminated.

Keywords: Blood, *Chlamydia trachomatis*, Detection, Kenya, Transfusion

1. INTRODUCTION

Chlamydia is a genus of pathogenic bacteria that are obligate intracellular parasites. *Chlamydia* infections are the most common bacterial sexually transmitted diseases in humans and are the leading cause of infectious blindness worldwide [1]. *Chlamydia trachomatis*, commonly known as chlamydia, is one of four bacterial species in the genus *Chlamydia*.

C. trachomatis is globally a major cause for sexually transmitted diseases (STDs) including, urogenital and Chlamydia. Chlamydia is one of the most common sexually transmitted infections worldwide affecting about 4.2% of women and 2.7% of men [2]. *C. trachomatis* cause various infections that include trachoma (conjunctivitis which is the most reported reason for blindness globally), pneumonic infections, and genital infections in people of both sexes and conjunctivitis in newborns. It is estimated that there are over 50 million new cases of *C. trachomatis* infection annually [3]. However another study done in 2015 showed that about 61 million new cases occurred globally [4].

C. trachomatis causes various contaminations among among which is donated blood. A study estimated that about 92 million blood donations are collected per year in different parts of the world [5]. In addition, WHO published that 39 countries in the world still do not routinely screen for transfusion-transmissible infections (TTIs) including HIV, Hepatitis B, Hepatitis C and syphilis [6]. About 47% of blood donations in low-income countries are screened in laboratories which do not have capacity for *C. trachomatis* contamination [7]. Blood transfusion centers are expected to provide blood

37 that is free of contaminants. However, without this there is likeliness of transmitting fatal agents like *C.*
38 *trachomatis* that may result in deadly diseases that may be found in blood [8].

39 Weström [9] asserted that one of the factors that may contribute to unnoticed *C. trachomatis*
40 contamination of donated blood is that its infections are to a bigger percentage asymptomatic. This
41 has commonly been reported in both males and females with about 50% males and 75% females of
42 those affected having no symptoms.

43 **2. MATERIALS AND METHODS**

44 **2.1. Sample Collection**

45 Blood samples used in the study were collected aseptically using venipuncture techniques by
46 qualified personnel of the center following the Pendergraph procedure of phlebotomy [10].

47 Use of aseptic techniques were observed to preserve the integrity of the specimen. Blood samples
48 obtained were labelled with the donors code for *C. trachomatis* test. The sample blood obtained was
49 subjected to centrifugation to separate the serum from the other blood components. The serum
50 samples were then refrigerated at 2^o-8^oC in RBTC Hematology Laboratory to be used in the
51 immunoassay test by using the *Chlamydia trachomatis* IgG/IgM Elisa test kit manufactured by Vircell
52 Company Spain to find out the presence of *C. trachomatis* in donated blood using the ELISA
53 procedure as shown below.

54 **2.2. Sample Processing**

55 10ml of blood sample from each donor was centrifuged at 1600rpm for 15 minutes. This was done
56 when clotting had occurred and within 1 hour after blood sample was drawn. Serum was then used for
57 the indirect immunoassay test.

58 **2.2.1. Indirect Immunoassay**

59 **Preparations and Assay Procedure**

60 The washing solution was prepared in advance. This was done by using distilled water to fill 50 ml of
61 20x washing solution up to 1 liter after which the solution was warmed up to 37°C in the washing
62 concentrate during storage before diluting. Once diluted, it was stored at 4°C.

63 **The Indirect Immunoassay Procedure**

64 The incubator was set at 37°C and all the reagents brought to room temperature before use
65 (approximately 1hour), without removing the plate from the bag. The components were then shaken
66 well.

67 The plate was removed from the package and the numbers of wells to be employed determined that
68 is: Four wells for the controls; two for the cut off serum and one each for the negative and positive
69 sera. Wells not required for the test were returned to the pouch and sealed.

70 For IgG test, 100 µl of serum diluent **2** was added to all wells followed by 5 µl of each sample, 5 µl of
71 positive control **3G**, 5 µl of cut off control **4G** (in duplicate) and 5 µl of negative control **5G** into the
72 corresponding wells. The plates were then shaken in a plate shaker for 2 min in order to achieve a
73 homogenous mixture of the reagents. A sealing sheet was then used to cover the wells and incubated
74 at 37°C for 45 min.

75 For IgM test, 25 µl of VIRCELL IgG sorbent was added to each of the required wells, except for the
76 wells where controls will be dispensed. 5 µl of sample was added and then 75 µl of the serum diluent
77 **2** to each well. Control wells were prepared by adding first 100 µl of the serum diluent **2** to each well
78 and then 5 µl of the positive control **3M**, 5 µl of the cut off control **4M** (in duplicate) and 5 µl of the
79 negative control **5M** to the corresponding wells. This was followed by plate shaking in a plate shaker
80 for 2 min in order to achieve a homogenous mixture of the reagents. A sealing sheet was then used to
81 cover the wells and incubated at 37°C for 45 min.

82 After incubation, the seal was removed, liquid aspirated from all wells and washed five times with 0.3
 83 ml of washing solution **9** per well and any remaining liquid was drained off. 100 µl of IgG conjugate
 84 solution **6G** or IgM conjugate solution **6M** was immediately added into each well and a sealing sheet
 85 used to cover followed by incubation at 37°C for 30 minutes.

86 After 30mins the seal was removed, liquid aspirated from all wells and washed five times with 0.3 ml
 87 of washing solution **9** per well. Any remaining liquid was drained off and 100 µl of substrate solution **7**
 88 immediately added into each well followed by incubation at room temperature for 20 minutes
 89 protected from light. After incubation 50 µl of stopping solution **8** was immediately added into all wells
 90 and finally readings were carried out within 1 hour of stopping.

91 **3. RESULTS**

92 In this study, a greater percentage of the samples were from the donors between 20-29 years of age
 93 (60.9%), while the least were below 20 years old group. There was a higher proportion of females
 94 (55.2%) than males (44.8%). Samples from single donors were 64.4% while divorced/separated and
 95 widowed were both 3.4%. 75% of the samples were from donors who were currently at tertiary
 96 education level while none was from those who had never attended attained any form of education.
 97 The highest proportion of the samples was from unemployed donors (73.6%), 11.5% from self-
 98 employed and 14.5% from employed persons (Table 1).

99 The largest percentage of samples were from donors who felt they were currently in good health
 100 (97.7%), had no sexual encounters with persons whom they did not know their background (96.6%),
 101 had irregular sexual partners (85.1).

102 **Table 1. Overall characteristics of blood donors whose samples were analyzed at RBTC,**
 103 **Eldoret.**

104

Characteristics	Categories	Total (N=87)
		No (%)
Age groups	Below 20 years	4 (4.6)
	20-29 years	53(60.9)
	30-39 years	25(28.7)
	40 and above	5(5.7)
Gender	Male	39(44.8)
	Female	48(55.2)
Marital status	Single	56(64.4)
	Married	25(28.7)
	Divorced/Separated	3(3.4)
	Widowed	3(3.4)
Education level	None	0(0)
	Primary	7(8.0)
	Secondary	14(16.1)
	Tertiary	66(75.9)
Employment status	Unemployed	64(73.6)
	Self-employed	10(11.5)
	Employed	13(14.9)

Good health	Yes	85(97.7)
	No	2(2.3)
Sexual activity	Yes	3(3.4)
	No	84(96.6)
Irregular sexual partners	No	13(14.9)
	Yes	74(85.1)

105 **Detection of *Chlamydia trachomatis* in donated blood using IgG/IgM Elisa kit**


106 Samples with indexes below 9 were considered as not having IgG or IgM specific antibodies against
 107 *C. trachomatis* while samples with indexes above 11 were considered as having IgG or IgM specific
 108 antibodies against *C. trachomatis*. Therefore this study showed that *C. trachomatis* in blood donors'
 109 samples at RBTC was found to be present in 8% (7 samples) of the blood samples analyzed at RBTC
 110 and 92% were found to be negative. (Table 2).

111 **Table 2: Antibodies Present in donor samples indicating *Chlamydia trachomatis* infection**

Antibodies against <i>C. trachomatis</i>			
Week	Donor No.	IgM Antibodies	IgG Antibodies
1	1-25	-	-
2	32	-	+
	37	-	+
	41	+	+
	48	+	-
3	53	+	-
	59	-	+
4	86	-	+

112 - =Absence of antibodies, + = Presence of antibodies

113 **Figure 1: Proportion of donors' samples with *Chlamydia trachomatis* at RBTC**



114

115

116 **Associations Between *C. trachomatis* and Selected Variables**

117 With the significance level at 0.05 the statistics revealed that having irregular sexual partners
 118 dependency with occurrence of *C. trachomatis* $\chi^2 (1) = 4.667$, $p=0.031$ but not significant. (Table 3)

119

120 **Table 3: Detection of *C. trachomatis* infection in donated blood with respect to risk factors at**
 121 **RBTC, Kenya.**

Characteristics			<i>C. trachomatis</i> status		Chi Square tests	
			Positive N=7 Out of 87 samples	Pearson Chi-Square	Valid cases	Asymp. Sig. (2-sided)
Age groups	Below 30 years	4	0.236	87	0.627	
	30 years and above	3				
Gender	Male	2	0.706	87	0.401	
	Female	5				
Marital status	Single	4	0.173	87	0.677	
	Non-singles	3				
Education level	Primary and below	1	0.401	87	0.527	
	Secondary and above	6				
Employment status	Unemployed	4	1.055	87	0.304	
	Self-employed and/ Employed	3				
Irregular sexual partners	No	3	4.667	87	0.031*	
	Yes	4				

122 **Note:** Non-singles include married, divorced, separated or widowed

123 * Significant at $p<0.05$.

124 **4. DISCUSSION**

125 The presence of *C. trachomatis* in blood samples was determined based upon the reaction of
 126 antibodies in the donor sample tested with the antigen adsorbed on the polystyrene surface. Unbound
 127 immunoglobulins were washed off, this was then followed by an enzyme anti-human globulin binding

128 the antigen-antibody complex in a second step. After a new washing step, bound conjugate was
129 developed with the aid of a substrate solution (TMB) to render a blue colored soluble product which
130 turned into yellow after adding the acid stopping solution [11]. IgG/IgM antibody detection are markers
131 for a Chlamydia-positive immune response, either for current, chronic or past infections. The assay
132 used in this study was the COMP (Complexes of Outer Membrane Proteins) of *C. trachomatis*, free
133 from LPS which is responsible for most cross-reaction with other Chlamydia species. ELISA yields
134 very good agreement with the target results (IgM100%, IgG 98%) in quality assessment schemes
135 [12]. It provides good agreement with characterized samples from quality assessment schemes of
136 IgM 100%, IgG 98% [12].Therefore ELISA technique is paramount for detection of *C. trachomatis*
137 antibodies.

138

139 Occurrence of IgM antibodies without IgG being present is an indication of primary infection while
140 presence of IgG antibodies without IgM antibodies in the blood sample is an indication of re-infection
141 or presence of previous infections since IgG antibodies persists in the body for a long period of time.
142 Presence of both IgG and IgM antibodies in the sample shows an ongoing infection.

143 In a study from Makkah Saudi Arabia, 8.7% of the women were positive for IgG antibodies to *C.*
144 *trachomatis* with antibody indexes of 1.4–2.0 [13]. Low prevalence in Saudi patient population may be
145 due to the adherence of strict moral principles and code of ethics in Saudi Arabia. In comparison to
146 this study, 71.4% of women were positive for both IgG/IgM antibodies which is much higher.

147 Similarly, it has been shown in the USA and Europe that demographic factors which increase the risk
148 of chlamydial infection include youth, single marital status and multiple sexual partners [14].
149 Therefore, a closer attempt should be made to correlate risk factors and disease entity when
150 screening for *C. trachomatis*.

151 According to a study done by CDC,[15] on the prevalence of *C. trachomatis* among the adults, the
152 highest rates is between 20 to 24 years old, however the age-specific rates for women was as low as
153 to 16 to 26 years old. This current study however slightly differs from the statistics by CDC. The age
154 between 20 to 29 years of age showed the highest prevalence.

155 Multiple partnerships may increase the likelihood of encountering a sexually transmitted pathogen
156 through the increased probability of choosing a partner with infection, while having new or casual
157 sexual contacts may be related to increased risk because of a reduced familiarity between partners
158 [16]. Statistics from this study revealed that having irregular sexual partners might have had
159 dependency with occurrence of *C. trachomatis* $X^2(1) = 4.667, p=0.031$.Therefore study concurs with
160 the later however it is not conclusive as per the statistics on the *chi square* table.

161 5. CONCLUSION

162 The findings of this study therefore indicates that some blood donated to the Regional Transfusion
163 Center was contaminated with *Chlamydia trachomatis* with 8% of the total samples indicating positive
164 for *C. trachomatis* infection while 92 % were negative. Moreover the age group of between 20 to 29
165 years old had the greatest percentage of those with *C. trachomatis*. Having multiple irregular number
166 of sexual partners increases the chances of having *Chlamydia trachomatis* infection. However this is
167 not be conclusive according to the statistics of this study.

168 The sex partners of donors with Chlamydial infection should therefore be referred for medical care
169 through the efforts of individual donors. A further study is also recommended on the exact level of *C.*
170 *trachomatis* by use of absorbance ranges.

171 6. COMPETING INTERESTS

172 "The authors declares that they have no competing interests"

173 7. ACKNOWLEDGMENTS

174 We thank the Blood transfusion services Eldoret for their technical support and Professor Elizabeth
175 Role for her advice, encouragement and her role in interpretation of data.

176 **8. AUTHORS' CONTRIBUTIONS**

177 The authors' responsibilities were as follows;

178 **REFERENCES**

- 179 [1]. Ryan KJ. & Ray CG. Sherris Medical Microbiology (4th Ed.). McGraw Hill. pp. 463–70;2004.
- 180 [2]. Newman L, Rowley J, Vander Hoorn S, Wijesooriya NS, Unemo M, Low N, & Temmerman M.
181 Global estimates of the prevalence and incidence of four curable sexually transmitted infections in
182 2012 based on systematic review and global reporting. *PloS one*, (2015)10(12).
- 183 [3]. Centers for Disease Control and Prevention. *Sexually Transmitted Disease Treatment Guidelines*.
184 2003; 51(6):1-78.
- 185 [4]. Global Burden of Disease. "Global, regional, and national incidence, prevalence, and years lived
186 with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden
187 of Disease Study 2015". *Lancet*. 2015; 388 (10053).
- 188 [5]. World Health Organization. WHO Global Data Base on Blood Safety, Summary Report. Geneva.
189 2011.
- 190 [6]. World Health Organization. Aide-memoire for national blood programmes. Geneva,
191 Switzerland:http://www.who.int/bloodsafety/transfusion_services/en/Blood_Safety_Eng.pdf World
192 Health Organization. Global database on blood safety: report 2001--2002. Available at
193 http://www.who.int/bloodsafety/GDBS_Report_2001-2002.pdf. 2002.
- 194 [7]. World Health Organization. Prevalence and incidence of selected sexually transmitted infections;
195 *Chlamydia trachomatis, Neisseria gonorrhoeae, syphilis and Trichomonas vaginalis*. 2008.
- 196 [8]. Mudasaar M. Transfusion-transmitted diseases. Department of Internal Medicine, Division of
197 Geriatrics, Duke University Health System: 2008;10.1046/j.1537-2995.2000.40030335.
- 198 [9]. Weström LARS, Joesoef, Reynolds GLADYS, Hagdu A, Thompson SE. Pelvic inflammatory
199 disease and fertility. *Sexually transmitted diseases*, 1999; 19(4), 185-192.
- 200 [10]. Pendergraph GE & Pendergraph CB. *Handbook of phlebotomy and patient service techniques*.
201 Lippincott Williams & Wilkins.1998
- 202 [11]. Bas S, Muzzin P, Ninet B, Bornand JE, Scieux C, Vischer TL. Chlamydial serology: comparative
203 diagnostic value of immunoblotting, microimmunofluorescence test, and immunoassays using
204 different recombinant proteins as antigens. *Journal of clinical microbiology*, 2001; 39(4), 1368-1377.
- 205 [12]. Gosink, J. Serological Diagnostic for Chlamydia antibodies Infections. *International Journal of*
206 *Medical Microbiology*.2015; 293 (148).
- 207 [13]. Ghazi HO, Daghestani MH, Mohamed MF. Seropositivity of *chlamydia trachomatis* among Saudi
208 pregnant women in Makkah, *J Family Community Med* 2006; 13 (2):61–4.
- 209 [14]. Pramanik –Mania J, Kerkar S, Sonawane S, Mehta P, Salvi, V. Current Chlamydia trachomatis
210 infection, a major cause of infertility. *Journal of reproduction & infertility*. 2012; 13(4) 204.

- 211 [15]. Centers for Disease Control and Prevention. Atlanta, GA: US Department of Health and Human
212 Services.2005.
- 213 [16]. Vranic SM. Chlamydia trachomatis Infections of the Adults. In *Sexually Transmitted Infections*.
214 InTech.2012.

UNDER PEER REVIEW