

1  
2 **EVALUTION OF ANTI-DIABETIC POTENTIAL OF AQUEOUS EXTRACT OF**  
3 **“LUFFA CYLINDRICA” (NATIVE SPONGE/SPONGE GOURD) LEAF AND SEED**  
4 **ON ALLOXAN INDUCED DIABETIC WISTAR RATS.**  
5

---

6  
7  
8  
9 **ABSTRACT**  
10

The study was carried out to evaluate the anti-diabetic effect of *Luffa cylindrical* (native sponge /sponge gourd) seed and leaf extracts in alloxan- induced diabetic rats. Sixteen experimental rats were divided into four groups of four rats each: a, diabetic control; b, normal control; c, diabetic rats treated with seed extract (400mg/kg) and d, diabetic rats treated with leaf extract (400mg/kg). The groups **A, C** and **D** rats were induced with diabetes intraperitoneally with alloxan (150mg/kg bw). Phytochemical screening was carried out on the plant seed and leaf extracts and the following biochemical tests were carried out: blood glucose, serum lipid profile, serum alanine aminotransferase, serum aspartate aminotransferase, serum alkaline phosphatase, total protein, albumin, creatinine, urea, uric acid and some electrolytes like **Na<sup>+</sup>, K<sup>+</sup>, HCO<sub>3</sub><sup>-</sup>**, and **Cl<sup>-</sup>** the administration of alloxan to experimental rats resulted in an increased level of most biochemical parameters; blood glucose, serum alanine aminotransferase, serum aspartate aminotransferase and serum alkaline phosphatase, serum total cholesterol, triglyceride, low density lipoprotein, creatinine, urea and uric acid. *Luffa cylindrical* seed and leaf extracts was administered to groups c and d diabetic rats respectively for two weeks, results were compared with normal control and diabetic control rats these parameters were found to be significantly ( $p < 0.05$ ) high in the diabetic groups than in the normal control groups. Treatment with the plant extract significantly ( $p < 0.05$ ) reduced elevated blood levels of glucose, cholesterol, triglyceride, alkaline phosphatase, amylase, aspartate aminotransferase, alanine aminotransferase, creatinine, urea, uric acid associated with alloxan-induced diabetic rats. The plant tested positive for alkaloids, flavonoids, saponins and tannins, negative for cardiac glycosides, phenols, resins, terpenes and steroids. Extracts of *Luffa cylindrical* seed and leaf has shown to have anti-diabetic and anti-lipidemic effects generally on alloxan induced diabetic rats. The study's findings has shown that the plant possess hypoglycaemic and hypolipidaemic property and has supported the traditional use of *Luffa cylindrical* plant in the management of diabetes and its complications.

11  
12 **Keywords:** *Luffa cylindrical*, alloxan, leaf, seed, diabetes

13 **1. INTRODUCTION**

14 A number of plants have been used in traditional medicine for many years. Some do seem to  
15 work although there may not be sufficient scientific data (double-blind trials, for example) to  
16 confirm their efficacy. Such plants should qualify as medicinal plants. The term 'crude drugs  
17 of natural or biological origin' is used by pharmacists and pharmacologists to describe whole  
18 plants or parts of plants which have medicinal properties [1]. In view of the fact that at the  
19 time there was not sufficient information either concerning the reasons for the illnesses or  
20 concerning which plant and how it could be utilized as a cure, everything was based on  
21 experience. In time, the reasons for the usage of specific medicinal plants for treatment of  
22 certain diseases were being discovered; thus, the medicinal plants' usage gradually  
23 abandoned the empiric framework and became founded on explicatory facts. Until the  
24 advent of iatrochemistry, plants had been the source of treatment and prophylaxis [2].  
25 Nonetheless, the decreasing usefulness of synthetic drugs and the increasing  
26 contraindications of their usage make the usage of natural drugs topical again. Traditional  
27 medicine has been used by the majority of the world population for thousands of year [3].



28  
29 Fig. 1: pictorial representation of the *Luffa cylindrica* plant

30 Amongst all the medicinal plants used in Nigeria for management and treatment of various  
31 types of ailments is the native sponge, scientific name is *Luffa cylindrica* in figure 1 above. It  
32 has other common names as smooth Luffa, sponge Luffa, vegetable sponge  
33 gourd, climbing okra, dishcloth gourd, chinese okra, it belongs to the family cucurbitaceae.  
34 *Luffa cylindrica* is native to India. Locations within which *Luffa cylindrica*  
35 is naturalized include: eastern africa and some pacific islands. *Luffa*  
36 *cylindrica* is naturalized in parts of Nigeria, Kenya and Tanzania and invasive in parts of  
37 Uganda.

38 *Luffa cylindrica* (sponge gourd) belonging to family *cucurbitaceae* is widely used across the  
39 globe as a vegetable. *L. Cylindrica* roem fruit extract (Ice) has been found to be an excellent  
40 antidiabetic and antioxidant[4][5].

41 *Luffa cylindrica* as a medicinal plant has been widely active in treatment of many diseases  
42 and used in proffering solutions to clinical problems relating to child birth. Although too many  
43 communities where this plant is used have little idea about the secret of its potency.  
44 Scientific research has shown the presence of some chemical components and proteins in  
45 *Luffa cylindrica*, and many others, which made it possible for *Luffa cylindrica* to be used as  
46 potentially effective chemical agent in health care delivery. Thus, possibility of transforming  
47 the chemical agents implicated in the plant of study into synthetic drugs to combat endemic  
48 diseases such as cancer and HIV should be the next focus of the clinical scientists [1].

49 Diabetes is a disease in which the body's ability to produce or respond to the hormone  
50 insulin is impaired, resulting in abnormal metabolism of carbohydrates and elevated levels of  
51 glucose in the blood. Diabetes mellitus is a clinically and genetically heterogeneous group of  
52 disorders that has a common feature of abnormally high levels of glucose in the blood due  
53 either to insulin deficiency or to resistance of the body's cells to the action of insulin.  
54 Diabetes mellitus or commonly diabetes is considered to be one of most serious, endocrine  
55 syndrome. It is a metabolic disorder characterized by hyperglycemia, glycosuria,  
56 hyperlipidemia, negative nitrogen balance, and sometimes ketonemia. Type 1 diabetes is  
57 caused by deficiency of insulin secretion from  $\beta$ -pancreatic cells [5]. On the other hand, type  
58 2 diabetes is characterized by initial phases of progressive insulin resistance.

59

## 60 **2. MATERIAL AND METHODS**

61 Plant material

62 *Luffa cylindrica* leaves were obtained from a growing tendril *Luffa cylindrica* plant, from  
63 kafachang kaduna state of Nigeria while the seeds were also obtained from a growing tendril  
64 *Luffa cylindrica* plant, From Jos, Plateau State of Nigeria and they were both identified in the  
65 Biochemistry Laboratory Of Bingham University, Karu Nassarawa State Nigeria.

66

### 67 **Preparation and Administration of The *Luffa Cylindrica* Leaves and Seeds Extract**

68 *Luffa cylindrica* leaves and seed extract was prepared by drying of the leaves and seeds  
69 collected. The *Luffa cylindrica* leaves and seed were pounded and matched to powder, 100g

70 and 50g of the samples were weighed respectively and was soaked in 1000ml and 500ml of  
 71 distilled water (100mg/ml and 50mg/ml respectively) respectively and then stirred and  
 72 heated on a hot plate for 15mins. The extracts were filtered with cloth sieve and then heated  
 73 in the water bath to dry at 60°C till samples completely dry. 400mg of *Luffa cylindrica* plant  
 74 extracts per kg body weight of rats was administered to each rat of each group of extract  
 75 once a day.

76 **Experimental Specimen**

77 **Albino** rats of 150-200g weight were purchased from Plateau State Nigeria. The experiment  
 78 was **approved** by the HOD Biochemistry Bingham University Karu, Nassarawa State and  
 79 HOD Animal Farm Unit of University Of Jos, Plateau State, Nigeria. The rats were housed in  
 80 metal cages with steel net covers and kept at room temperature (24-28°C) under 12hours  
 81 dark-light cycles. All rats were fed appropriately with their respective diet feed, water and  
 82 were acclimatized for 2 weeks in the animal house in University Of Jos, Plateau State.

83 **Induction of Diabetes**

84 Alloxan was induced in experimental rats after 12hours of fasting (overnight) by  
 85 intraperitoneally administration of 150mg/kg body weight of alloxan. After the above  
 86 observations, the fasting blood glucose concentration of all experimental rat were  
 87 determined with the aid of a glucometer (blood was taken from the respective rat's tail) for  
 88 concentration greater than 120mg/dl.

89 **Experimental Design**

The control and experimental rats divided into different groups and treated accordingly;	
Group 1:	normal control: non-diabetic group
Group 2:	diabetic control: diabetic group
Group 3:	<i>Luffa cylindrica</i> leaves extract group: diabetic rats receiving 400mg <i>Luffa cylindrica</i> leaves extract per kg body weight once daily.
Group 4:	<i>Luffa cylindrica</i> seeds extract group: diabetic rats receiving 400mg <i>Luffa cylindrica</i> seeds extract per kg body weight once daily.

90  
91

## 92 **Determination Of Biochemical Parameters**

93 Determination of uric acid

94 Determination of blood glucose

95 The fasting blood glucose concentration was determined by o'toluidine method.

96 O'toluidine solution

97 Determination of serum total protein

98 Biuret reagent

99 Determination of serum total cholesterol

## 100 **3. RESULT AN DISCUSSION**

### 101 **Statistical analysis**

102 The data were expressed as Mean  $\pm$  standard error of Mean. Statistical analysis was  
103 performed using analysis of variance (anova) at 5% level of confidence ( $p < 0.05$ ). Using spss  
104 analytical software.

105 Results

106 Table 1: percentage extraction of plant samples extracted and used on the experimental  
107 rats.

Sample	Weight of raw plant (g)	Weight of plant extract (g)	Percentage of plant extraction (%)
Leaves	100	13.35	13.35
Seeds	50	7.60	15.2

108

109 Table 2: phytochemicals present in the plant samples (leaves and seeds) used in the  
110 analysis and administered to experimental rats.

S/n	Phytochemicals	Leaf	Seed
-----	----------------	------	------

1.	Alkaloids	+	+
2.	Flavonoids	+	+
3.	Tannins	+	-
4.	Saponins	+	+
5.	Terpenes	-	-
6.	Steroids	-	-
7.	Cardiac glycosides	-	-
8.	Balsam	+	-
9.	Carbohydrates	+	-
10.	Phenol	+	-
11.	Resins	-	-

111 Key

112 + = detected                      - = not detected

113 Table 3: effect of administration of *Luffa cylindrica* seeds and leaves extract on biochemical  
114 parameters (such as glucose level, total protein and albumin).

Group	Treatment	Glucose (mmol/l)	Total protein (g/l)	Albumin (g/l)
A	Diabetic control	12.46±0.022	68.50±0.426	30.42±0.510
B	Normal control	3.48±0.029 <sup>a</sup>	76.47±0.442 <sup>a</sup>	38.50±0.430 <sup>a</sup>
C	Diabetic + seed	8.34±0.029 <sup>ab</sup>	70.47±0.521 <sup>ab</sup>	33.44±0.464 <sup>ab</sup>
D	Diabetic + leaf	6.70±0.037 <sup>ab</sup>	73.39±0.447 <sup>ab</sup>	31.58±0.457 <sup>ab</sup>

115 Values are expressed as Mean ± SD, n= 4 for each group.

116 <sup>A</sup>values are significantly different when compared with diabetic control (p<0.05).

117 <sup>B</sup>values are significantly different when compared with normal control (p<0).

118

119 Table 4. Effect of administration of *Luffa cylindrica* seeds and leaves extract on lipid profile  
 120 parameters.

Group	Treatment	Total cholesterol (mmol/l)	Triglyceride (tg) (mmol/l)	High density lipoprotein (hdl) (mmol/l)	Low density lipoprotein (ldl) (mmol/l)
A	Diabetic control	5.35±0.103	1.98±0.166	0.52±0.089	2.26±0.317
B	Normal control	3.14±0.025 <sup>a</sup>	0.80±0.138 <sup>a</sup>	1.31± 0.257 <sup>a</sup>	1.44±0.178 <sup>a</sup>
C	Diabetic + seed	4.45±0.029 <sup>ab</sup>	1.64±0.173 <sup>ab</sup>	0.78±0.129 <sup>ab</sup>	1.88±0.245 <sup>ab</sup>
D	Diabetic + leaf	3.46±0.033 <sup>ab</sup>	1.21±0.166 <sup>a</sup>	0.88±0.141 <sup>ab</sup>	1.63±0.171 <sup>ab</sup>

121

122 Values are expressed as Mean ± SD, n= 4 for each group.

123 <sup>A</sup>values are significantly different when compared with diabetic control (p<0.05).

124 <sup>B</sup>values are significantly different when compared with normal control (p<0.05).

125 Table 5: effect of administration of *Luffa cylindrica* seeds and leaves extract on some liver  
 126 function test parameters {ALT (Alanine Aminotransferase), AST (Aspartate  
 127 Aminotransferase), and ALP (Alkaline Phosphatase)}

Group	Treatment	ALT (u/l)	AST (u/l)	ALP (u/l)
A	Diabetic control	24.53±0.830	31.49±0.843	359.35±0.520
B	Normal control	12.58±0.822 <sup>a</sup>	16.50±0.684 <sup>a</sup>	168.41±0.724 <sup>a</sup>
C	Diabetic + seed	18.74±0.827 <sup>ab</sup>	23.52±0.874 <sup>ab</sup>	284.31±0.602 <sup>ab</sup>
D	Diabetic + leaf	15.76±0.708 <sup>ab</sup>	19.59±0.852 <sup>ab</sup>	251.55±0.731 <sup>ab</sup>

128 Values are expressed as Mean ± SD, n= 4 for each group.

129 <sup>A</sup>values are significantly different when compared with diabetic control (p<0.05).

130 <sup>B</sup>values are significantly different when compared with normal control (p<0.05).

131 Table 6: effect of administration of *Luffa cylindrica* seeds and leaves extract on other  
 132 biochemical parameters (such as creatinine, urea and uric acid).

Group	Treatment	Creatinine ( $\mu\text{mol/l}$ )	Urea ( $\text{mmol/l}$ )	Uric acid ( $\mu\text{mol/l}$ )
A	Diabetic control	11.54 $\pm$ 0.420	197.53 $\pm$ 0.501	408.30 $\pm$ 0.580
B	Normal control	4.34 $\pm$ 0.477 <sup>a</sup>	104.48 $\pm$ 0.437 <sup>a</sup>	268.30 $\pm$ 0.658 <sup>a</sup>
C	Diabetic + seed	9.44 $\pm$ 0.435 <sup>ab</sup>	159.70 $\pm$ 0.480 <sup>ab</sup>	314.25 $\pm$ 0.645 <sup>ab</sup>
D	Diabetic + leaf	7.35 $\pm$ 0.505 <sup>ab</sup>	124.44 $\pm$ 0.433 <sup>ab</sup>	351.55 $\pm$ 0.701 <sup>ab</sup>

133

134 Values are expressed as Mean  $\pm$  SD, n= 4 for each group.

135 <sup>A</sup>values are significantly different when compared with normal control (p<0.05).

136 <sup>B</sup>values are significantly different when compared with diabetic control (p<0.05).

137

Table 7 Effect of administration of *Luffa Cylindrica* seeds and leaves extract on biochemical electrolytes parameter (Such as Sodium, Potassium, Chloride and Biocarbonate)

140

Group	Treatment	Sodium Na <sup>+</sup> ( $\text{mmol/L}$ )	Potassium K <sup>+</sup> ( $\text{mmol/L}$ )	Chloride Cl ( $\text{mmol/L}$ )	Bicarbonate HCO <sub>3</sub> <sup>-</sup> ( $\text{mmol/L}$ )
A	Diabetic Control	135.29 $\pm$ 0.766	5.90 $\pm$ 0.026	106.26 $\pm$ 0.501	18.26 $\pm$ 0.522
B	Normal control	141.29 $\pm$ 1.056 <sup>a</sup>	3.70 $\pm$ 0.022 <sup>a</sup>	113.28 $\pm$ 0.684 <sup>a</sup>	26.25 $\pm$ 0.510 <sup>a</sup>
C	Diabetic + Seed	137.00 $\pm$ 0.816 <sup>ab</sup>	5.20 $\pm$ 0.026 <sup>ab</sup>	108.28 $\pm$ 0.643 <sup>ab</sup>	21.25 $\pm$ 0.506 <sup>ab</sup>
D	Diabetic + Leaf	139.01 $\pm$ 0.816 <sup>ab</sup>	4.30 $\pm$ 0.050 <sup>ab</sup>	110.28 $\pm$ 0.597 <sup>ab</sup>	24.25 $\pm$ 0.507 <sup>ab</sup>

158

## 159 DISCUSSION

160

161 Treatment with *Luffa cylindrica* plant extracts (seed and leaf) produced a time dependent  
 162 decreased concentration in blood glucose level and other biochemical parameters: total  
 163 protein, total cholesterol and liver enzymes (alanine aminotransferase, aspartate  
 164 aminotransferase and alkaline phosphatase).



165

166 The high level of glucose observed in blood of induced experimental rats by the  
167 administration of alloxan, in our case, which is cytotoxic (toxic to living cells) specifically for  
168 the  $\beta$ -cells of the islets of langerhans in the pancreas which function in regulation of insulin  
169 secretion [6]

170

171 It has been identified that the liver is necrotized in alloxan induced diabetic rats which leads  
172 to release or increase activities of liver enzymes (alanine aminotransferase, aspartate  
173 aminotransferase and alkaline phosphatase) as they leak due to cirrhosis from the liver to  
174 the bloodstream and this is an indicator of the hepatotoxicity caused by the induction of  
175 alloxan shown clearly in the diabetic group [7]. The time dependent decrease of these liver  
176 enzymes in the blood stream maybe due to the administration of the plant extracts which  
177 may have helped in retrogressing the hepatocellular damage caused by alloxan  
178 administration initially, thereby helping in refurbishing and mending the hepatocyte  
179 membrane integrity.

180 The table 3 above shows the result of the analysis of biochemical parameters (glucose, total  
181 protein and albumin) on experimental rats. In consideration of groups of the experimental  
182 rats, the glucose concentration of the normal control had no significant change in the  
183 concentrations, considering the diabetic + seed and diabetic + leaf treatment group, which  
184 was shown to be significantly different when compared respectively to the diabetic control  
185 group ( $p < 0.05$ ). As compared generally, the results shows that administration of *Luffa*  
186 *cylindrica* plant extracts (seed and leaf) were effective in reducing blood glucose level after  
187 14days of treatment, as earlier observed in the same research carried out utilizing *Luffa*  
188 *cylindrica* fruits which tested for antihyperglycemic activity in alloxan induced hyperglycemic  
189 rats [8].

190

191 Secondly, the total protein and albumin tests may be ordered in a variety of settings to help  
192 diagnose disease, to monitor changes in health status, total protein measurements can  
193 reflect nutritional status and may be used to screen for and help diagnose kidney  
194 disease or liver disease. A low total protein such as the diabetic group level can suggest  
195 a liver disorder, kidney disorder, or a disorder in which protein is not digested or absorbed  
196 properly. Low levels may be seen in severe malnutrition and with conditions that cause mal-  
197 absorption, such as celiac disease or inflammatory bowel disease [9]. The destruction of the  
198 pancreas results in the utilization of non-carbohydrate precursors such as protein for the  
199 synthesis of glucose to form energy need in the cells, generally leads to increased lipolysis  
200 and increased synthesis of ketone bodies results in severe decrease in the total protein level

201 observed in diabetic group. The table 3 in the results above shows the result of the analysis  
202 of biochemical parameters (total protein and albumin) on experimental rats. After 14days of  
203 treatment and induction of diabetes in experimental rats it was observed that the  
204 concentration of the total protein comparing the diabetic control group of concentration which  
205 was shown to increase. Compared with that of the diabetic + seed and diabetic + leaf  
206 groups, which shows that all groups maintained the normal range of total protein which is  
207 between 60g/l to 80g/l. However, results indicate the ability of *Luffa cylindrica* plant extracts  
208 to be effective in enhancing the level of total protein observed in the extraction groups of  
209 seed and leaf when compared with the diabetic group, as also seen in the table ( $p < 0.05$ ).  
210 Also, after 14days of treatment and induction of diabetes in experimental rats it was  
211 observed that the concentration of serum albumin which measures the amount of liver  
212 protein contained in the clear liquid protein of the blood, when compared with the normal  
213 control group seemed to be significantly different from the diabetic + seed and diabetic + leaf  
214 groups ( $p < 0.05$ ). Also, in the same research, venous blood samples was used for  
215 estimation of plasma glucose, total proteins, albumin, fibrinogen which tested in the study of  
216 type 2 diabetics, plasma albumin levels were decreased compared to controls and plasma  
217 fibrinogen, total protein levels were statistically significantly increased compared to controls  
218 [10].

219

220 The increase in total cholesterol level of the diabetic group was due to the hyperglycemia  
221 confirmed in the diabetic group. *Luffa cylindrica* plant extracts were able to improve lipid  
222 metabolites generally including the correction of the high density lipoproteins known as good  
223 cholesterol which aids as carriers for the removal of low density lipoproteins and triglyceride  
224 from the blood to prevent the blockage of arteries, results indicates the ability of *Luffa*  
225 *cylindrica* plant extracts to be effective in correction of these levels of metabolites in  
226 experimental rats diabetic and treated respectively.

227

228 The table 4 in the results above shows the result of the analysis of biochemical lipid profile  
229 parameters (total cholesterol, triglyceride, high density lipoproteins and low density  
230 lipoproteins) on experimental rats. In consideration of the groups of the experimental rats  
231 (experimental design) the result of the serum total cholesterol, shows that *Luffa cylindrica*  
232 seed and leaf extracts is effective in significantly reducing the level of serum total cholesterol  
233 for diabetic control group ( $p < 0.05$ ). Comparing the diabetic + seed and diabetic + leaf extract  
234 groups, the leaf of the plant reflects the ability to reduce serum total cholesterol more than  
235 the seed of the *Luffa cylindrica* plant.

236

237 After 14days of treatment and induction of diabetes in experimental rats it was observed that  
238 the concentration of the triglyceride and low density lipoproteins when compared with the  
239 diabetic control groups, which showed a significant decrease ( $p<0.05$ ) on treatment with  
240 seed in the diabetic + seed group, which reflects that the *Luffa cylindrica* plant is effective in  
241 reducing the concentration of triglyceride and low density lipoproteins. Also, the result of the  
242 high density lipoproteins concentration after 14days of treatment with the *Luffa cylindrica*  
243 plant extracts, in the comparison with the diabetic group also shows that there is an increase  
244 in the level of the concentration of the high density lipoproteins in the diabetic + seed and  
245 diabetic + leaf groups respectively ( $p<0.05$ ).

246

247 The table 5 in the results above shows the result of the analysis of some biochemical liver  
248 function test parameters (alanine aminotransferase, aspartate aminotransferase and alkaline  
249 phosphatase) on experimental rats. Considering the groups of experimental rats it was  
250 observed that the concentration of the liver enzyme (serum ALT, AST and ALP) in normal  
251 control group increased significantly when compared to that of the diabetic control group  
252 ( $p<0.05$ ).

253 After 14days of the administration of the seed and leaf extracts, there was a decrease in the  
254 diabetic + seed treatment group respectively of each of the enzymes and the diabetic + leaf  
255 treatment group also ( $p<0.05$ ). The result shows that when comparing the diabetic + seed  
256 with diabetic + leaf the leaf treatment is more effective in reducing liver damage. But  
257 generally *Luffa cylindrica* plant extracts were effective in reduction of liver damage which  
258 result in high level serum alanine aminotransferase, serum aspartate aminotransferase and  
259 serum alkaline phosphatase.

260

261 The table 6 in the results above shows the result of creatinine, urea and uric acid on  
262 experimental rats, it was observed that the concentration of the creatinine, urea and uric acid  
263 in diabetic control group increased significantly when compared to the normal control group  
264 ( $p<0.05$ ).

265

266 After 14days of the administration of the seed and leaf extracts, there was a decrease in the  
267 diabetic + seed treatment group and the diabetic + leaf treatment group ( $p<0.05$ ). The result  
268 shows that when comparing the diabetic + seed with diabetic + leaf the leaf treatment was  
269 more effective and generally shows that *Luffa cylindrica* plant extracts were effective in  
270 reducing creatinine, urea and uric acid levels in the blood which aids in the reduction of  
271 kidney disease and dysfunction [12].

272

273 Table 7 shows the result of the analysis of sodium, potassium, chloride and bicarbonate on  
274 experimental rats. They are generally chemicals in the blood stream that regulate important  
275 functions in the body, when dissolved in water electrolytes separates into positively and  
276 negatively charged ions, it was observed that the concentration of all the electrolytes except  
277 potassium in normal control group reduced significantly when compared with that of the  
278 diabetic control group ( $p < 0.05$ ) but that of potassium rather increased for normal and  
279 diabetic respectively. The result shows that when the diabetic + seed and diabetic + leaf  
280 treatments were both effective in reducing electrolyte imbalance which may have occurred  
281 due to hormonal or endocrine disorders, kidney disease and dysfunction.

282

283 Also, methanolic extract of *Luffa cylindrica* fruits on oral glucose tolerance and its effect on  
284 normoglycemic rats were studied [8]. The same was tested for antihyperglycemic activity in  
285 alloxan induced hyperglycemic rats at the two dose levels 200 and 400mg/kg body weight.  
286 The serum biochemical parameters were also assessed in the alloxan induced experimental  
287 animals. The methanolic extract of *Luffa cylindrica* exhibited remarkable antihyperglycemic  
288 activity [8]. The treatment of diabetic rats with methanolic extract of the test plant improved  
289 the serum biochemical parameters and the activities were found to be dose dependent, the  
290 respective effects were basically observed on fasted normal, alloxanised hyperglycemic and  
291 glucose tolerance rats. Diabetes is associated with profound alterations in lipid and  
292 lipoproteins profiles, early detection and treatment of hyperlipidemia in diabetic patients  
293 reduces the risk for cardiovascular and cerebrovascular diseases [13]. Therefore lowering of  
294 plasma or tissue lipids levels generally may lead to decrease in the risk of micro and macro  
295 vascular disease related complications [14]. It therefore can be suggested that *Luffa*  
296 *cylindrica* plant extracts may improve lipid profiles as shown in the results above either  
297 directly or indirectly through reduction of blood glucose level generally in experimental rats  
298 diabetic or treated respectively as observed in this research study.

299

#### 300 4. CONCLUSION

301 In conclusion, *Luffa cylindrica* seed and leaf extracts were able to reduce elevated level of  
302 blood glucose, lipid profile and serum enzymes. The result confirms antidiabetic potential of  
303 *Luffa cylindrica* plant in alloxan induced diabetic wistar rats. The results suggest that *Luffa*  
304 *cylindrica* plant extracts' have the possibilities to improve and enhance treatment of diabetes  
305 complications.

306

307 **REFERENCES**

- 308 1. Sofowora A. Medicinal Plants and Traditional Medicine in Africa'. 3rd edn. Ibadan:  
309 Spectrum Books; 2008.
- 310 2. Evans WC. Trease and Evans' Pharmacognosy. 16th Edition. London: WB  
311 Saunders Company Ltd; 2008.
- 312 3. Kelly k. History of medicine. New york: facts on file; 2009. Pp. 29–50.
- 313 4. Seifu t. Ethnobotanical and ethnopharmaceutical studies on medicinal plants of  
314 chifra district, afar region, north eastern ethiopia. Phd thesis. Addis ababa  
315 university: school of graduate studies; 2004.
- 316 5. Du q, xu y, li l, zhao y, jerz g, winterhalter p. Antioxidant constituents in the fruits of  
317 *Luffa cylindrica* (L.) Roem. J agric food chem. 2006; 54:4186–90.
- 318 6. Karasawa h, takaishi k, kumagae y. Obesity-induced diabetes in mouse strains  
319 treated with gold thioglucose: a novel animal model for studying  $\beta$ -cell  
320 dysfunction. Obesity (silver spring, md.) 2011; 19(3):514-21.
- 321 7. Adiga s, bairy kl, meharban a, punita isr. Hypoglycemic effect of aqueous extract of  
322 *trichosanthes dioica* in normal and diabetic rats. Intl j diab dev cties 2010;  
323 30: 38-42.
- 324 8. Johnston d .e and lee, mary (2009-03-10). "*special considerations in interpreting*  
325 *liver function tests*". Am fam physician. 59 (8): 2223–30.
- 326 9. American association for clinical chemistry (aacc),  
327 <https://labtestsonline.org/understanding/analytes/tp/tab/test/> article was last  
328 reviewed on april 10, 2016
- 329 10. American association for clinical chemistry (aacc),  
330 <https://labtestsonline.org/understanding/analytes/tp/tab/test/> article was last  
331 reviewed on april 10, 2016
- 332 11. Manjusha hazra, sriparna kundusen, sanjib bhattacharya, pallab k haldar, malaya  
333 gupta, upal k mazumder, (2011). Journal of advanced pharmacy education &  
334 research. "*evaluation of hypoglycemic and antihyperglycemic effects of Luffa*  
335 *cylindrica fruit extract in rats.*" Vol. 2. (2011). Pg. 138-146  
336
- 337 12. Venkataramana g., indira p., d.v.m. rao. (2013) indian journal of basic & applied  
338 medical research. "*changes of plasma total proteins, albumin and fibrinogen*  
339 *in type 2 diabetes mellitus- a pilot study*". Vol. 2, pg. 679-685  
340

- 341 13. Razafindraibe M, Kuhlman AR, Rabarison H, Rakotoarimanana V, Rajeriarison C,  
342 Rakotoarivelo N, et al. Medicinal plants used by women from Agnalazaha littoral  
343 forest (Southeastern Madagascar). J Ethnobiol Ethnomed. 2013;9:73.
- 344 14. Arjola z, klodiana s, gentian v et al (2014).lipid profile in diabetes mellitus type 2  
345 patients in albania and the correlation with bmi, htn and hepatosteatosi. J  
346 family med community health; 1(4):1018.
- 347 15. Marinou, k. A., georgopoulou, k., agrogiannis, g., karatzas , t., iliopoulos, d.,  
348 papalois, a., dontas, i. A. (2010). Differential effect of *pistacia vera* extracts on  
349 experimental atherosclerosis in the rabbit animal model: an experimental  
350 study. *Lipids in health and disease*, 9, 73.

UNDER PEER REVIEW