

## **Original Research Article**

### **PHYTOCHEMICAL COMPOSITION AND IN VITRO ANTIOXIDANT ACTIVITY OF GOLDEN MELON (*CUCUMIS MELO L*) SEEDS FOR FUNCTIONAL FOOD APPLICATION**

Aim: Golden melon (*Cucumis melo*) is an annual herbaceous plant belonging to the family of *Cucurbitaceae* (*Cucurbit*). This study was carried out to evaluate the phytochemical composition and in vitro antioxidant activity of golden melon seed extract.

Place and Duration of study: The study was carried out between a period of July and August 2017 at Baking Milling Division, Federal Institute of Industrial Research Oshodi Nigeria.

Methodology: The crude methanolic extracts of the seed was tested for phytochemical and antioxidant activities according to standard analytical procedure. The antioxidant potential of the seed extracts was examined using different assays by determining total phenolic content, total flavonoid content, total antioxidant capacity. The free radical scavenging activities of the extract such as 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity, in vitro lipid peroxidation, and nitric oxide (NO) scavenging assay were determined spectrophotometrically.

Results: The phytochemical screening of the seed extracts revealed the presence of some secondary metabolites such as alkaloids, phenolic, steroids, flavonoids, terpenoids and cardiac glycosides. The total phenolic content of extract was found to be 29.39mg/100g while the amount of total flavonoid content was 20.67mg/100g. Scavenging ability was observed to increase in proportion to concentration for all the scavenging assays and at the highest concentration. Total antioxidant capacity assay showed 19.44mg per 100 g. This high scavenging ability in the seed extracts may be attributed to the presence of phenolic

and flavonoids compounds in the extract. The DPPH free radical scavenging activity of 100µg/ml *Cucumis melo* extract was 75.20% ± 0.72 while the reference standard (Ascorbic acid) was 83.24% ± 0.31. Lipid peroxidation inhibition ability of 100µg/ml *Cucumis melo* extract was 87.18% ± 0.16 while the standard (ascorbic acid) was 94.96% ± 0.16 at the same concentration. Results obtained from this study showed that the Nitric Oxide scavenging ability of the extract was 80.50%±0.63 while the standard antioxidant was 85.94% ± 0.54.

Conclusion: In all the assays, *Cucumis melo* extract showed maximum percentage of antioxidant potentials at 100µg/ml. Additionally, golden melon seed possess appreciable amount of phenols and high antioxidant properties which could be explored and incorporated in functional food applications particularly in baked products.

***Golden melon seed, phytochemical composition, antioxidant activity and free radical scavenging activity***

## 1. INTRODUCTION

Fruits and vegetables are vital components of human diet and their consumption has been shown to confer a great deal of nutritional and health benefits on humans including preventing and reducing the risk of certain chronic degenerative diseases [1]. They contain some biological active components known as phytochemicals that promote health through the prevention of specific degenerative diseases such as cardiovascular diseases, diabetes, obesity, cancers and gastrointestinal tract disorders. Golden melon (*Cucumis melo*) is an annual, drooping herbaceous plant that is distinguished by a short angular-stem woody rootstock with bristly hairs large belonging to the family of *Cucurbitaceae* (*Cucurbit*), one of the

most genetically diverse groups of food plants that are drought-tolerant [2]. It originated in Europe and Africa before spreading to other parts of the world. It is a bright-yellow melon with a pale green to white inner flesh which is succulent and juicy grown in the Sahel and Sudan savanna of Nigeria. Nutritionally, golden melon is a great constituent of essential minerals and nutrients such as vitamin C, pantothenic acid, calcium, zinc, vitamin B6, fibre, magnesium, iron, potassium, vitamin A and omega-3 & 6 [3].

In Nigeria, there are various types of fruit and vegetables commonly consumed according to their geographical locations and seasons. Among these fruits are citrus, oranges, mangoes, watermelon, apple, cabbage, carrot, cucumbers, pumpkin, grapes, plantain /bananas among many others. However, only the fleshy parts of these fruits and vegetables are eaten leaving the peels, rinds and/or the seeds as waste materials. There are numbers of studies that have been done by researchers exploring the potentials source of antioxidant in plant waste materials. Fruits have high vitamin, mineral, fibre, phytochemical and antioxidant contents in their pulps, seeds and rinds and *Cucumis melo* is no exception.

Reports showed that besides dietary sources, antioxidant can be gained from food processing industries especially agricultural by products [4]. These respective by products are peel, rind, seed, mill wastes and trimming wastes. Reports showed that these by products are promising sources of important bioactive substances called phytochemicals (carotenoids, phenolics, and flavonoids), antioxidants vitamins that possess beneficial physiological properties [5, 6]. Previous studies have revealed that the seeds are concentrated sources of many health-benefiting vitamins, minerals, antioxidants, and essential amino acids such as tryptophan, and glutamate [7]. These phytochemicals and antioxidants can be used or incorporated into foods as ingredients for functional foods in order to promote health and prevent the risk of some degenerative diseases. In recent times, there has been an increasing consumer interests in functional foods as a result of their nutraceutical and health benefits. Functional foods are foods that provide additional health benefits beyond the basic nutritional functions.

The use of fruits and vegetables and their by-products as ingredient in food processing as well as products requiring hydration, viscosity development, and freshness preservation, such as baked foods has been well documented [8, 9, 10, 11, and 12]. Except for these reported studies, there is a dearth of information on the phytochemical composition and antioxidant activity of *Cucumis melo* L seeds. This

study, therefore, investigates the phytochemical composition and in vitro antioxidant activity of golden melon (*Cucumis melo* L) seed.

## 2. MATERIALS AND METHODS

### 2.1 Sourcing and preparation of Golden melon (*Cucumis melo*) seeds

Golden melon (*Cucumis melo* L) fruits were obtained from local market, Lagos state, Nigeria. The seeds were removed, washed and dried in a cabinet dryer at 50°C for 4 hours before being milled. The dried seed was milled into flour which was packaged in a polyethylene for further analyses.

The seed extract was prepared according to the decoction method described by [8]. Milled *Cucumis melo* L seeds were transferred into a round bottom flask and submerged in methanol in ratio 1:10 (w/v). The flask was stoppered and left to stand for 72 hours. The extract was then filtered using Whatman No 1 filter paper. The filtrate was concentrated using a rotary evaporator at 40°C. The resultant residue were weighed and stored at 4°C.

### 2.2 Phytochemical screening of crude extract

The crude methanolic extracts of the seeds were tested for the presence of alkaloids, steroids, tannins, saponins and glycosides according to the methods described by [13, 14, 15, 16, and 17]. The qualitative results are expressed as (+) for the presence and (–) for the absence of phytochemicals.

**Test for flavonoids:** The presence of flavonoids in the plant samples was determined as follows: (a) Five (5) ml of dilute ammonia solution was added to 5 ml of the extract solutions followed by the addition of concentrated H<sub>2</sub>SO<sub>4</sub>. A yellow colouration observed in each extract indicated the presence of flavonoids. (b) Few drops of 1% Aluminum solution were added to 2 ml of each extract. A yellow colouration was observed indicating the presence of flavonoids. (c) Five (5) ml of the extract solution was in each case heated with 10 ml of ethyl acetate over a steam bath for 3 minutes. The mixture was filtered and 4 ml of filtrate was shaken with 1 ml dilute ammonia solution. A yellow colouration is an indication of flavonoids presence.

**Test for terpenoids (Salkowski test):** A 5 ml sample of the extract was mixed in 2 ml of chloroform, and 3 ml concentrated H<sub>2</sub>SO<sub>4</sub> was carefully added along the sides of the test tube to form a layer. The formation of a reddish brown colouration at the interface indicated the presence of terpenoids.

**Test for steroids:** A 5 ml sample of the extract was added to 2 ml acetic anhydride and 2 ml  $\text{H}_2\text{SO}_4$ . The colour changed from violet to blue or green indicated the presence of steroids.

**Test for alkaloid:** A 5 mg sample of the extract dissolved in 3 ml of acidified ethanol was warmed slightly and then filtered. Few drops of Mayer's reagent and 1 ml of Dragendroff's reagent were added to 1 ml of the filtrate and turbidity was observed.

**Test for tannins:** A few drops of 0.1% ferric chloride was added to the extract solutions and observed brownish-green or a blue-black colouration, which signified the presence of tannins.

**Test for saponins:** Five (5) ml of the extract solution was shaken vigorously for a stable persistent froth. The frothing was mixed with olive oil and shaken vigorously. The formation of emulsion indicated the presence of saponins in the samples.

**Anthraquinone glycoside (Borntrager's test):** To the extract solution (1 ml), 5%  $\text{H}_2\text{SO}_4$  (1 ml) was added. The mixture was boiled in a water bath and then filtered. Filtrate was then shaken with equal volume of chloroform and kept to stand for 5 min. Then lower layer of chloroform was shaken with half of its volume with dilute ammonia. The formation of rose pink to red color of the ammonical layer gives indication of anthraquinone glycosides [18].

**b) Cardiac glycoside (Keller-Killiani test):** Extract (0.5 g) was shaken with distilled water (5 ml). To this, glacial acetic acid (2 ml) containing a few drops of ferric chloride was added, followed by  $\text{H}_2\text{SO}_4$  (1 ml) along the side of the test tube. The formation of brown ring at the interface gives positive indication for cardiac glycoside and a violet ring may appear below the brown ring [19]

## **2.3 Antioxidant Activity Assay**

### **2.3.1 DPPH radical scavenging activity assay**

The free radical scavenging activity of the extract was determined spectrophotometrically. It was based on the scavenging of the stable 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical which was estimated according to the procedure described by [20, 21]. An aliquot of 0.5 ml of sample extract in ethanol (95%) at different concentrations (20, 40, 60, 80, 100  $\mu\text{g}/\text{ml}$ ) was mixed with 2.0 ml of reagent solution (0.004 g of DPPH in 100 ml methanol). The control contained only DPPH solution in place of the sample while methanol was used as the blank. The mixture was vigorously shaken and left to stand at room

temperature. After 30 minutes the decrease in absorbance of test mixture (due to quenching of DPPH free radicals) was read at 517 nm on a UV- visible spectrophotometer (Milton Roy Spectronic 601, USA). The scavenging effect was calculated using the expression:

$$\% \text{ inhibition} = [A_0 - A_1] / A_0 \times 100$$

Where  $A_0$  is the absorption of the blank sample and  $A_1$  is the absorption of the extract

Ascorbic acid was used as standard.

### 2.3.2 Nitric oxide scavenging activity assay

The compound sodium nitroprusside is known to decompose in aqueous solution at physiological pH (7.2) producing nitric ions ( $\text{NO}^\bullet$ ). Under aerobic condition,  $\text{NO}^\bullet$  reacts with oxygen to produce stable products (nitrate and nitrite), which can be determined using Griess reagent. The absorbance of the chromophore that formed during diazotization of the nitrite with sulphanilamide and subsequent coupling with Naphthylethylenediamine dihydrochloride can be immediately read at 550 nm.

A 4 ml sample of flour extract or standard solution of different concentrations (20, 40, 60, 80, 100  $\mu\text{g/ml}$ ) were taken in different test tubes and 1 ml of Sodium nitroprusside (5 Mm in phosphate buffered saline) solution was added into the test tubes. They were incubated for 2 hours at 30 °C to complete the reaction. A 2 ml sample was withdrawn from the mixture and mixed with 1.2 ml of Griess reagent (1% Sulphanilamide, 0.1% naphthylethylene diamine dihydrochloride in 2%  $\text{H}_3\text{PO}_4$ ). The absorbance of the chromophore formed during diazotization of nitrite with sulphanilamide and its subsequent coupling with naphthylethylene diamine was measured at 550 nm on a UV- visible spectrophotometer (Milton Roy Spectronic 601, USA) as modified by [22]. Ascorbic acid was used as standard and inhibition of nitric oxide radical was calculated using the expression:

$$(\%) \text{ inhibition activity} = [(A_0 - A_1)/A_0] \times 100.$$

Where,  $A_0$  is the absorbance of the Control and  $A_1$  is the absorbance of the extract or standard.

### 2.3.3 Lipid peroxidation assay

Lipid peroxidation was induced by  $\text{Fe}^{2+}$ -ascorbate system in liver homogenate and estimated as thiobarbituric acid reacting substances (TBARS) by the method of [23] as described by [24]. In this assay, the end product of lipid peroxidation using liver homogenate as lipid- rich media [25] was quantified by determining the formed adduct of malonyldialdehyde (MDA) reaction with thiobarbituric acid (TBA) under

acidic condition. The pink colored product was measured at 532 nm on a UV- visible spectrophotometer (Milton Roy Spectronic 601, USA). Into 0.5ml of a 0.1mg/ml liver homogenate was added 0.1 ml of varying concentrations of the extract (20, 40, 60, 80, 100µg/ml) in a test tube followed by the addition of 1 ml distilled water. Then 50µl of FeSO<sub>4</sub> (0.07 M) was added to the reaction mixture. The reaction mixture was vortexed and allowed to stand for 30 minutes at ambient temperature after which 1.5ml of 20% (v/v) acetic acid and 1.5 ml of 0.8% (w/v) thiobarbituric acid in 1.1% (w/v) sodium dodecyl sulphate were added. The resulting mixture was then incubated in a water bath at 95°C for 1 hr. After cooling, 4.0 ml of butan-1-ol was added to each tube, shaken vigorously and centrifuged at 3000 rpm for 10 min. the absorbance of the organic upper layer was measured at 532nm. The TBARS values were calculated using the extinction coefficient  $1.56 \times 10^{-5}$  M/cm.

Inhibition of lipid peroxidation (%) by the extract was calculated using the formula

$$(1-E/C) \times 100$$

Where

C= absorbance value of the fully oxidized control

E= absorbance in the presence of extract as  $(A_{532} + \text{TBA}) - (A_{532} - \text{TBA})$

#### 2.3.4 Reducing Power Assay

The ability of methanolic extracts to reduce iron (III) to iron (II) was assessed by the method of [26]. The dried extract (20–100 µg) in 1 ml of the corresponding solvent was mixed with 2.5 ml of phosphate buffer (0.2 M, pH 6.6) and 2.5 ml of potassium ferricyanide (K<sub>3</sub>Fe(CN)<sub>6</sub>; 10 g/l), and then the mixture was incubated at 50 °C for 30 min. After incubation, 2.5 ml of TCA (100 g/l) was added and the mixture was centrifuged at 1650 rpm for 10 min. Finally, 2.5 ml of the supernatant solution were mixed with 2.5 ml of distilled water and 0.5 ml of FeCl<sub>3</sub> (1 g/l) and the absorbance was measured at 700 nm on a UV- visible spectrophotometer (Milton Roy Spectronic 601, USA). High absorbance indicates high reducing power.

#### 2.4 Estimation of total phenolic content

The amount of total phenol content was determined by Folin-Ciocalteu colorimetric method [27] using gallic acid as a standard following the method as described by [28]. A 0.5 ml sample of extract and 0.1 ml of Folin- Ciocalteu reagent (0.5 N) were mixed and incubated at room temperature for 15 minutes. After

this, 2.5 ml sodium carbonate solution (7.5% w/v) was added and further incubated for 30 minutes at room temperature. The absorbance of the solution was measured on a UV- visible spectrophotometer (Milton Roy Spectronic 601, USA) at 760 nm. The concentration of total phenol was expressed as gallic acid equivalent (GAE) (mg/g of dry mass) which is a commonly used reference value.

## **2.5 Total flavonoid content estimation**

Total soluble flavonoid of the extract was determined using aluminium chloride colorimetric method using quercetin as standard [29]. 1 ml of sample solution (100µg/ ml) was mixed with 3 ml of methanol, 0.2 ml of 10% Aluminum chloride, 0.2 ml of 1 M potassium acetate and 5.6 ml of distilled water. The resulting mixture was incubated at room temperature for 30 minutes and the absorbance of the reaction mixture was measured on a UV- visible spectrophotometer (Milton Roy Spectronic 601, USA) at 415 nm. The calibration curve was prepared by preparing quercetin solutions at various concentrations in methanol.

## **2.6 Total antioxidant capacity determination**

The total antioxidant capacity of the extracts was determined using the method of [30]. A sample of the extract (0.3 ml) was mixed with 3 ml of reagent solution (0.6 M sulphuric acid, 28 mM sodium phosphate and 4 mM ammonium molybdate). The tubes were capped and incubated in a boiling water bath at 95<sup>0</sup> C for 90 minutes. After the samples had cooled to room temperature, the absorbance of the aqueous solution of each was measured at 695 nm. The total antioxidant capacity was expressed as equivalent of ascorbic acid.

## **Statistical analysis**

Statistical significance was established using student t test paired comparison and data were reported as mean ± of standard error (SEM) of three determinations using Microsoft Excel 2010. The p values of  $p < 0.05$  were considered statistically significant for differences in mean between the standard Ascorbic acid and the seed extract. The IC<sub>50</sub> values were estimated from the % inhibition versus concentration plot using linear regression analysis.

# **3. RESULTS AND DISCUSSION**

## **3.1 Phytochemical constituents**



The phytochemical screening of the seed extracts revealed the presence of different secondary metabolites like alkaloids, flavonoids, steroids, phenols, cardiac glycosides and terpenoids as presented in Table 1. The other phyto-constituents such as saponin, tannins and phlobatannins were totally absent.

This present study showed that the predominant phytoconstituent of the seed extract is phenol, which was followed by steroid, then flavonoids, terpenoids and alkaloids while cardiac glycoside was the least compound.

Presence of flavonoids, tannins, and alkaloids has been observed to possess various biological properties related to antioxidant mechanisms [31]. They are effective hydrogen donors that inhibit the lipid oxidation and chelating metal ions, making them good anti-oxidants [32]. The cardiac glycosides are basically steroids with an inherent ability to afford a very specific and powerful action mainly on the cardiac muscle when administered through injection into man or animal. Cardiac glycosides have been reported to be used to treat congestive heart failure and cardiac arrhythmia [33]. Their mode of action starts by inhibiting  $\text{Na}^+/\text{K}^+$  pump which then increases the level of calcium ion, so more  $\text{Ca}^{+2}$  would be available for the contraction of heart muscles which recover cardiac output and reduce the distension of heart [34, 35].

Alkaloids are the largest group of secondary chemical constituents made largely of ammonia compounds comprising basically of nitrogen bases synthesized from amino acid building blocks with various radicals replacing one or more of the hydrogen atoms in the peptide ring, most containing oxygen.

Steroids (anabolic steroids) have been observed to promote nitrogen retention in osteoporosis and in animals with wasting illness [36, 37]. Similarly, steroids derived from plants are known to have cardiotonic effect and also possess antibacterial and insecticidal properties [38]. They are very often used in medicines due to their well-known biological activities

The total phenolic content expressed as mg per 100gram of the seed extract was found to be 29.39mg/100g while the total flavonoid content was 20.67mg/100g. The phenolic content can be used as

an important indicator of antioxidant capacity which can be used as a primary screening for any product when intended for use as a natural source of antioxidants in functional foods.

Plants contain a wide variety of free radicals scavenging molecules including phenols, flavonoids, vitamins, terpenoids that are rich in antioxidant activity [39, 40]. Antioxidants protect cells against the damaging effects of reactive oxygen species otherwise called, free radicals such as singlet oxygen, super oxide, peroxy radicals, hydroxyl radicals and peroxynite which results in oxidative stress leading to cellular damage [41]. Natural antioxidants play a key role in health maintenance and prevention of the chronic and degenerative diseases, such as atherosclerosis, cardiac and cerebral ischemia, carcinogenesis, neurodegenerative disorders, diabetic pregnancy, rheumatic disorder, DNA damage and ageing [42]. Plant polyphenols are the significant group of compounds acting as free radical scavenging or primary antioxidants; therefore, it is justifiable to determine phenolic content in plant extract.

Phenolics essentially represent a host of natural antioxidants, used as nutraceuticals, and found in apples, green-tea, and red-wine for their enormous ability to combat cancer and are also thought to prevent heart ailments to an appreciable degree and sometimes are anti-inflammatory agents. The most common group of plant phenolics are the flavonoids, the structures of which are based on that of flavone, consisting of two benzene rings linked through a three-carbon  $\gamma$ -pyrone ring. Common classes of flavonoids include flavones, flavonols, isoflavones, anthocyanins, catechins (flavanols) and flavanones. Flavonoids are a group of phenolic compounds with antioxidant activity that have been identified in fruits, vegetables, and other plant foods and that have been linked to reducing the risk of major chronic diseases.

The health-related properties of phenolic compounds, particularly flavonoids, are believed to be based on their antioxidant activity as hydrogen donating free radical scavengers [43, 44].

Table 1: Qualitative and Quantitative Phytochemical Screenings of Aqueous Extract of Golden melon (*Cucumis melo L*) Seed

Compound	Qualitative	Quantitative (mg/100g)
Flavonoid	+	20.67±0.19
Alkaloid	+	18.85±0.12
Phenol	+	29.39±0.11
Saponin	-	ND
Cardiac glycoside	+	11.34±0.18
Phlobatanin	-	ND
Tannin	-	ND
Steroid	+	22.63±0.22
Terpenoid	+	19.49±0.14

Values are means ± SDs of triplicate determinations. ND- Not Detected

### 3.2 Free Radical Scavenging Activity

The antioxidant potential of the *Cucumis melo* seed extracts was examined by determining 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging activity, in vitro lipid peroxidation, and nitric oxide (NO) scavenging assay. The free radical scavenging capacity was observed at 20 µg/ml, 40 µg/ml, 60 µg/ml, 80 µg/ml and 100 µg/ml of methanol extracts. The results showed that the seed extracts of test fruit had radical scavenging activity. A number of assays were conducted to analyze antioxidant activities of the extract since antioxidant activities of different types of substances involve different mechanisms.

Scavenging ability of the methanolic extract was observed to increase in proportion to concentration for the sample and at the highest concentration of 100 µg/ml.

The DPPH free radical scavenging activity of *Cucumis melo* extract at different concentrations from 20 to 100 µg/ml differed significantly ( $p < 0.05$ ) compared to the reference standard which is ascorbic acid. It ranged from  $37.86 \% \pm 0.64$  to  $75.20 \% \pm 0.72$  while the reference standard (Ascorbic acid) ranged from  $44.72 \% \pm 0.23$  to  $83.24 \% \pm 0.31$  as was depicted in fig 1. *Cucumis melo* seed showed maximum inhibition (75.20%) of DPPH free radical at 100 µg/ml while the  $IC_{50}$  values for the extract and the reference standard (ascorbic acid) was found to be  $53.28 \pm 0.56$  µg/ml and  $46.77 \pm 0.34$  µg/ml respectively.  $IC_{50}$  value is defined as the concentration of the sample require to scavenge DPPH radical by 50%, was obtained from a calibration curve for the extract. The lower the  $IC_{50}$  value the higher the antioxidant activity of a sample. It was clear that the free radical scavenging activities of the extract was in a concentration dependent manner. This high scavenging ability in the seed extracts maybe attributed to the presence of phenolic and flavonoids compounds in the extract as in-vitro studies have shown that flavonoids compounds may have positive effects against inhibition oxidation of biomolecules by its anti-oxidative potential, anti-inflammatory, anti-allergic, and antifungal effect [45, 46, and 47]. This study is in agreement with the result obtained by [48] who worked on the free radical scavenging activity of *Bergenia stracheyi* extracts.

Free radicals are known to be scavenged by synthetic antioxidants, but due to their adverse side effects leading to carcinogenicity, search for effective and natural antioxidants has become crucial [49]. Natural antioxidants are believed to be safer and bioactive [50]. DPPH assay was carried out to measure the primary antioxidant activity of the sample in the present study. The ability of a substance to remove or scavenge free radicals is classified as a primary antioxidant [51]. DPPH assay reaction depends on the

ability of the samples to scavenge free radicals which is visually noticeable as the colour change from purple to yellow due to hydrogen donating ability [52]. The more rapid the absorbance decreases, the more potent the primary antioxidant activity [53].

Figure 2 presents lipid peroxidation inhibition ability of *Cucumis melo* seed extract at different concentrations of 20 to 100 µg/ml. It ranged from  $47.18 \pm 0.13$  -  $87.18\% \pm 0.16$  while the standard (ascorbic acid) ranged from  $45.55\% \pm 0.12$  -  $94.96\% \pm 0.16$ . The  $IC_{50}$  was found to be  $50.34 \pm 0.36$  µg/ml for the extract, and ascorbic acid which is the standard reference antioxidant as  $45.75 \pm 0.42$  µg/ml. Lipid peroxidation scavenging activity of *Cucumis melo* seed showed maximum inhibition (94.96%) at 100 µg/ml in a concentration dependent manner and differed significantly ( $p < 0.05$ ) from the standard antioxidant. Extensive research studies have shown that lipid peroxidation is probably the most investigated free radical induced process [54, 55, 56, and 57]. The production of malondialdehyde (MDA) becomes elevated after oxidative stress has initiated lipid peroxidation [58]. MDA is expressed when the oxidative insult exceeds the threshold value [59]. Antioxidants are important inhibitors of lipid per-oxidation not only as a defense mechanism of living cells against oxidative damage but also for food preservation [48].

Reducing power of the methanolic of *Cucumis melo* seed extract was determined and the results are shown in Figure 3. Significant reduction ( $p < 0.05$ ) was observed in the reducing power of the seed extract when compared with the standard reference ascorbic acid. In the same manner, the methanolic extract displayed the highest reducing power at 100 µg/ml in a concentration dependent manner. In this assay, the presence of reducers (i.e., antioxidants) causes the reduction of the  $Fe^{3+}$ /ferricyanide complex to the ferrous form. Therefore, measuring the formation of Perl's Prussian blue at 700 nm can monitor the  $Fe^{2+}$  concentration [41]. The mechanisms of antioxidant properties of plant extracts are assumed to be based on their ability to trap positively charged electrophilic species, to scavenge oxygen radicals, to have reducing power, and to chelate metals to form inactive complexes [60] and these properties are majorly dependent on phenolic compounds found in the extracts. Some studies have however suggested that a proper and reasonable heat treatment application to food materials such as citrus peel could be used to enhance its antioxidant properties. In addition, polyphenols, including flavonoids that are soluble in less polar solvents, may be able to chelate metal ions such as iron and copper because of the large

number of hydroxyl groups (OH) of their chemical structure, which are responsible for conferring the chelating ability [61].

In order to evaluate the antioxidant potency through NO scavenging by the test samples, the change of optical density of NO was monitored. Results obtained from this study showed that the Nitric Oxide scavenging ability of the extract ranged from 33.19- 80.50%±0.63 while the standard antioxidant ranged from 44.97% ± 0.74 - 85.94% ± 0.54. There was a significance difference ( $p<0.05$ ) between the seed extract and the standard antioxidant at the different concentrations studied. The  $IC_{50}$  value was estimated as  $58.36 \pm 0.34$  µg/ml for the extract and  $51.66 \pm 0.22$  µg/ml for ascorbic acid which was the standard reference antioxidant.

Nitric oxide scavenging activity of *Cucumis melo* seed showed maximum inhibition (80.50%) at 100µg/ml in a concentration dependent manner as well (Figure 4). These demonstrated the ability of the seed in arresting the chain of reactions initiated by excess generation of reactive nitric oxide species (NO.) that are deleterious to the human health [62]. Nitric oxide is classified as a free radical because of its unpaired electron and displays important reactivity with certain types of proteins and other free radicals. This result was in agreement with study conducted by [22].

In all the assays, *Cucumis melo* extract showed maximum percentage of antioxidant potentials at 100µg/ml with the  $IC_{50}$  values well comparable to that of the ascorbic acid, the standard reference antioxidant used in the study.




Figure1. DPPH radical scavenging activity of *Cucumis melo* L seed extract at different concentrations.

Ascorbic acid was used as positive control. Bars represent means  $\pm$  standard error (SEM) of three replications. \*Statistically significant at  $p < 0.05$ .




Figure 2: Lipid peroxidation inhibition by *Cucumis melo L* seed extract at different concentrations.

Ascorbic acid was used as positive control. Bars represent means  $\pm$  standard error (SEM) of three replications. \*Statistically significant at  $p < 0.05$ .






Figure 3: Reducing power of different concentration of *Cucumis melo L* seed extract which was estimated by potassium ferricyanide method. Ascorbic acid was used as positive control. Bars represent means  $\pm$  standard error (SEM) of three replications. \*Statistically significant at  $p < 0.05$ .




Figure 4: Nitric oxide scavenging activity of *Cucumis melo* L seed extract at different concentrations.

Ascorbic acid was used as positive control. Bars represent means  $\pm$  standard error (SEM) of three replications. \*Statistically significant at  $p < 0.05$ .

#### 4. CONCLUSION

The results of phytochemical screening of methanolic extracts of *Cucumis melo L* seed reveals the presence of alkaloids, flavonoids, phenolics, steroids, cardiac glycosides and terpenoids. The free radicals scavenging activity results show that *Cucumis melo L* seed has compounds of radical scavenging potential. Antioxidant activity of plant material is one of the key indicators of the health beneficial effects in prevention and reduction of risk of diseases which is becoming increasingly rampant even in developing countries. The phenol, flavonoid contents and radical scavenging properties of the seed are associated with its antioxidant properties. These findings further demonstrate that the seed extract is not toxic and safe for human consumption which could probably be considered as functional ingredient. Hence, this study had revealed that golden melon seed possess appreciable amount of phenols and flavonoids and high antioxidant properties which could be explored and incorporated in functional food applications particularly in baked products.

#### REFERENCES

- 1 Genkinger JM, Platz EA, Hoffman SC, Comstock GW, Helzlsouer KM (2004): Fruit, vegetable and antioxidant intake and all-cause cancer, and cardio vascular disease mortality in a community dwelling population in Washington country, Maryland . Am J Epidemiol; 160: 1223-33.
- 2 Ajuru, M. G. and Okoli, B. E. (2013), The Morphological Characterization of the Melon Species in the Family Cucurbitaceae Juss., and their Utilization in Nigeria, International Journal of Modern Botany 2013, 3(2): pp. 15-19. DOI: 10.5923/j.ijmb.20130302.01.
- 3 Raji, O. H. and Orelaja, O. T. (2014) Nutritional Composition and Oil Characteristics of Golden Melon (*Cucumis melo*) Seeds, *Food Science and Quality Management*, pp. 18-20.
- 4 Duda-Chodak Aleksandra, Tomasz Tarko (2007): Antioxidant Properties of different fruit seeds and peels. Tiarum Polonorum Acta Acta Sci. Pol., Technol. Aliment. 6(3), 29-36

- 5 Lachman, J.; Hamouz, K.; Musilova, J.; Hejtmankova, K.; Kotikova, Z.; Pazderu, K.; Domkarova, J.; Pivec, V.; Cimr, J. (2013): Effect of peeling and three cooking methods on the content of selected phytochemicals in potato tubers with various colour of flesh. *Food Chem.* 138, 1189–1197.
- 6 Yildirim, A.; Mavi, A.; Kara, A.A. (2001) Determination of antioxidant and antimicrobial activities of *Rumex crispus* L. extracts. *J. Agric. Food Chem.* 49, 4083–4089.
- 7 Arora, R., Kaur, M. and Gill, N. S. (2011), Antioxidant activity and Pharmacological Evaluation of *Cucumis melo* var. *agrestis* Methanolic seed Extract, *Research Journal of Phytochemistry*, 5(3), 146-155
- 8 Brewer M.S., Potter S.M., Sprouls G. and M. Reinhard (1992): Effect of soy protein isolate and soy fiber on color, physical and sensory characteristics of baked products. *Journal of Food Quality* 15, 245-262
- 9 Fernández-López J. (; 2008): Development of functional ingredients: fruits fibre. In: Fernández-López J, Pérez-Alvarez JA, editors. *Technological strategies for functional meat products development. Kerala: Transworld Research.* pp. 41–57.
- 10 Moazzezi, S., S.M. Seyedain and L. Nateghi, (2012): Rheological properties of barbari bread containing apple pomace and carboxy methyl cellulose. *J. Life Sci.*, 9(3): 1318-1325.
- 11 Bhise, S.R., & Kaur.A (2014): Incorporation of Oat, Psyllium and Barley Fibers: Effect on Baking Quality, Sensory Properties and Shelf Life of Bread. *International Journal of Engineering Practical Research (IJEPR)* Volume 3 Issue 3, 5258, August 2014. Doi:10.14355/ijepr.2014.0303.02
- 12 Bhol Soumya, Lanka Divyajyoti, and John Don Bosco Sowriappan (2016): Quality characteristics and antioxidant properties of breads incorporated with pomegranate whole fruit bagasse. *J Food Sci Technol.* ; 53(3): 1717–1721. doi: 10.1007/s13197-015-2085-8

- 13 Pavia, D.L.L., Lampman, G M and Kriz G.S, (1976): Introduction to Organic laboratory techniques : a contemporary approach . Second edition Saunders Golden Sanburst Series,W.B Saunders Company, Philadelphia, London, Toronto; pp599-614.
- 14 Ojiako, O.A and Smith H.A (1991): Phytochemical screening of Nigerian medicinal plants. *Nature*; 1: 260-264
- 15 Handa ,SS,. Khanuja, S.P. S., Longo, G and Rakesh, D.D(2008): Extraction Technologies for Medicinal and Aromatic Plants. *International Centre for Science and High Technology, Trieste*, 21-25
- 16 Magadula, J.J and Tewtrakul, S (2010) Anti-HIV-1 protease activities of crude extracts of some *Garcinia* species growing in Tanzania. *African Journal of Biotechnology*, 9 (12): 1848-1852.
- 17 Obasi N.L, Egbuonu, A.C.C., Ukoha, P.O and Ejikeme P.M (2010) Comparative phytochemical and antimicrobial screening of some solvent extracts of *Samanea saman* pods. *African journal of pure and applied chemistry*; 4(9): 206-212.
- 18 Harborne JB (1973); *Phytochemical Methods: A guide to modern techniques of plant analysis*. Chapman and Hall, New York, pp. 279, 3rd Edition,
- 19 Joshi A, Bhobe, M; Saatarkar, A (2013): Phytochemical investigation of the roots of *Grevia microcos*. *Linn. J. Chem. Pharm.Res.* 5, 80-87.
- 20 Ayoola. GA, Coker HB, Adesegun SA, Adepoju-Bello, AA; Obaweya K. Ezennia, EC., Attangbayila TO (2008): Phytochemical screening and antioxidant activities of some selected medicinal plants used for malarial therapy in Southwestern Nigeria, *Trop.J. Pharm Res.* 7, 1019-1024.
- 21 Burits M, and Bucar F (2000) Antioxidant activity of *Nigella sativa* essential oil", *Phytother Research* 14: 323–328.

- 22 Ojewunmi Oyesola, Tope Oshodi, Omobolanle Ogundele, Chijioke Micah and Sunday Adenekan (2013) Toxicity screening and in vitro antioxidant activities of aqueous extracts of *Morinda lucida* and *Saccharum officinarum* leaves *Biokemistri Vol. 25 (2)* 72–78 31
- 23 Alisi CS and Onyeze GOC (2008) Nitric oxide scavenging ability of ethyl acetate fraction of methanolic leaf extracts of *Chromolaena odorata* (Linn.). *African Journal of Biochemistry Research* 2: 145–50.
- 24 Buege J and Aust DS (1978) Microsomal lipid peroxidation. In: Fleisscher S, Packer L, eds. *Methods in enzymology*. New York: Academic Press, 52: 302–311.
- 25 Nagmoti, D. M., Khatri, D. K Juvekar P. R., and. Juvekar A. R (2011): Antioxidant activity and free radical-scavenging potential of *Pithecellobium dulce* Benth seed extracts *Free Radical and Antioxidants*, vol. 2, no. 2, pp. 37–43.
- 26 Gutteridge, J.M. and Halliwell, B. (1990). The measurement and mechanism of lipid peroxidation in biological systems. *Trends Biochemical Science*; 15:129-135.
- 27 McDonald S, Prenzler PD, Antolovich M and Robards K (2001) Phenolic content and antioxidant activity of olive extracts. *Food Chemistry* 73:73–84.
- 28 Slinkard K and Singleton VL (1977) Total phenol analysis: automation and comparison with manual methods. *American Journal of Enology and Viticulture* 28: 49–55.
- 29 Chang C, Yang M, Wen H (2002): Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of Food and Drug Analysis* 10:178–182.
- 30 Prieto P, Pineda M and Aguilar M (1999): Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Analytical Biochemistry* 269: 337–341.
- 31 Latha PG, Suja SR, Abraham A, Rajasekharan S and Panikkar KR (2003) Hepatoprotective effects of *Ixora coccinea* flower extract in rats. *Journal of Tropical Medicinal Plants* 41: 33–38.

- 32 Shahidi F (1997) Natural antioxidants: An overview. In: Natural antioxidants, Shahidi F. (ed.), AOCS press, Champaign, IL, 1–11.
- 33 Vladimir, K, Ludmila, M (2001): Glycosides in Medicine: the role of glycosidic residue in biological activity. *Curr. Med. Chem.* 8, 1303-1328.
- 34 Banso A., Adeyemo S (2006): Phytochemical screening and Antimalarial assessment of *Abutilon mauritianum*, *Bacopa monnifera* and *Datura stramonium*. *Biokemistri* 18, 39-44.
- 35 Aiyelaagbe, OO., Osamudiamen, PM (2009): Phytochemical screening for active compounds in *Magnifera indica* leaves from Ibadan, Oyo state. *Plant Sci. Res.* 2: 11-13
- 36 Maurya, R.; Singh G. & Yadav, P.P. (2008). Antiosteoporotic agents from Natural sources. In: Atta-ur-Rahman (Ed.) *Studies in Natural Products Chemistry*, Vol. 35. Elsevier. pp 517-545.
- 37 Madziga H.A, Sanni S and Sandabe U.K. (2010). Phytochemical and Elemental Analysis of *Acalypha wilkesiana* Leaf. *Journal of American Science*. 6(11): 510-514.
38. Alexei Y. Bagrov, Joseph I. Shapiro, and Olga V. Fedorova (2009): Endogenous Cardiotonic Steroids: Physiology, Pharmacology, and Novel Therapeutic Targets. *Pharmacological Reviews*, 61 (1) 9- 38, doi: <https://doi.org/10.1124/pr.108.000711>.
- 39 Madsen, H.L. & Bertelsen G (1995): Spices as antioxidants. *Trends Food Science and Technology*. 6: 271-277.
- 40 Cai, Y.Z. & Sun, M. (2003): Antioxidant activity of betalins from plants of the Amaranthacea. *Journal of Agriculture and Food Chemistry*. 51: 2288-2294.
- 41 Mattson, M.P. & Cheng A (2006): Neurohormetic phytochemicals: lowdose toxins that induce adaptive neuronal stress responses. *Trends in Neurosciences*. 29(11): 632-639.
- 42 Uddin, S.N.; Akond, M.A., Mubassara, S. & Yesmin MN (2008). Antioxidant and Antibacterial activities of *Trema cannabina*. *Middle-East Journal of Scientific Research*. 3: 105-108

- 43 Rice-Evans CA, Miller NJ Paganga G (1996). Structure antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic. Biol. Med.*, 20: 933-956.
- 44 Prior RL, Cao G (2000) Antioxidant phytochemicals in fruits and vegetables: Diet and health implications. *Hort Sci.* 35:588-92.
- 45 Friedman, M. (2007): Overview of antibacterial, antitoxin, antiviral, and antifungal activities of tea flavonoids and teas. *Mol. Nutr. Food Res.*, 51, 116–134.
- 46 Giovannini, C.; Filesi, C.; D'Archivio, M.; Scazzocchio, B.; Santangelo, C.; Masella, R. (2006). Polyphenols and endogenous antioxidant defences: Effects on glutathione and glutathione related enzymes. *Ann. Ist. Super. Sanita.* 42, 336–347.
- 47 Shapiro, H.; Singer, P.; Halpern, Z.; Bruck, R. (2007): Polyphenols in the treatment of inflammatory bowel disease and acute pancreatitis. *Gut.* 56, 426–435.
- 48 Kumar Vineshand Devendra Tyagi. (2013): Phytochemical Screening and Free-Radical Scavenging Activity of *Bergenia stracheyi*. *Journal of Pharmacognosy and Phytochemistry* Vol. 2 No. 2 pp 175-180.
- 49 Ebrahimzadeh, M. A., Nabavi S. M, Nabavi S. F., Bahramian, F. and Bekhradnia, A. R. (2010) Antioxidant and free radical scavenging activity of *H. officinalis* L. var. *angustifolius*, *V. odorata*, *B. hyrcana* and *C. speciosum* *Pakistan Journal of Pharmaceutical Sciences*, vol. 23, no. 1, pp. 29–34, 2010.
- 50 Adedapo, F. O. Jimoh, S. Koduru, P. J. Masika, and A. J. Afolayan, (2009): Assessment of the medicinal potentials of the methanol extracts of the leaves and stems of *Buddleja saligna*,” *BMC Complementary and Alternative Medicine*, vol. 9, article 21
- 51 Wang, H., Gao, X. D., Zhou, G. C., Cai, L. and Yao, W. B. (2008): In vitro and in vivo antioxidant activity of aqueous extract from *Choerospondias axillaries* fruit. *Food Chemistry* 106: 888-895



- 52 Ajila, C. M., Naidu, K. A., Bhat, S. G. and Prasada R. U. J. S. 2007. Bioactive compounds and antioxidant potential of mango peel extract. *Food Chemistry* 105: 982-988.
- 53 Siddhuraju, P., Mohan, P. S. and Becker, K. 2002. Studies on the antioxidant activity of Indian Laburnum (*Cassia fistula* L.): a preliminary assessment of crude extracts from stem bark, leaves, flowers and fruit pulp. *Food Chemistry* 79: 61-67.
- 54 De Zwart, L.L., Meerman, J.H., Commandeur, J.N. and Vermeulen, N.P. 1999. Biomarkers of free radical damage applications in experimental animals and in humans. *Free Radical Biological Medicine*. 26:202-226.
- 55 Moore, K. and Roberts, L.J. (1998): Measurement of lipid peroxidation. *Free Radical Research*; 28:659-671.
- 56 Gutteridge, J.M. (1995) : Lipid peroxidation and antioxidants as biomarkers of tissue damage. *Clinical Chemistry*; 41:1819-1828.
- 57 Halliwell, B. and Gutteridge, J.M. (1990). Role of free radicals and catalytic metal ions in human disease: an overview. *Methods of Enzymology*;186:1-85
- 58 Catala, A. (2011): Lipid peroxidation of membrane phospholipids in the vertebrate retina. *Front Bioscience (Schol Ed)* 3: 52–60.
- 59 Jakovljevic B, Novakov-Mikic A, Brkic S, Bogavac MA, Tomic S, Miler V. (2012). Lipid peroxidation in the first trimester of pregnancy. *Journal of Maternal, Fetal and Neonatal Medicine*;25:1316–1318.
- 60 Echavarría AP, Pagán J, Ibarz A. (2012): Melanoidins formed by Maillard reaction in food and their biological activity. *Food Eng Rev* 4(4):203–23.
- 61 Nandhakumar E, Indumathi P. (2013): In vitro antioxidant activities of methanol and aqueous extract of *Annona squamosa* (L.) fruit pulp. *Journal of Acupuncture and Meridian Studies* 6: 142-148.

- 62 Hasan SMR, Hossain MM, Akter R, Jamila M, Mazumder MEH and Islam I (2009) Antioxidant, antidiarrhoeal and cytotoxic properties of Punica granatum Linn. *Latin American Journal of Pharmacy* 28: 783–788.

UNDER PEER REVIEW