

Effect of Indole Acetic Acid And Boron on the Nutrient Contents of Summer Tomato Cultivar Bina Tomato-3

ABSTRACT

A pot experiment was carried out with tomato variety BINA Tomato-3 in the net house of the Department of Agricultural Chemistry, Bangladesh Agricultural University (BAU), Mymensingh from March to August, 2010 to study the effect of Indole Acetic Acid (IAA) and Boron (B) on the nutrient contents of BINA Tomato-3. There were four doses of IAA viz. 0, 20, 40 and 60 ppm and B viz. 0, 1.5, 2.0 and 2.5 kg ha⁻¹. The experiment was laid out in a Completely Randomized Design (CRD) with 3 replications. The result revealed that nutrient contents of tomato fruits and plants such as N, P, Ca, Mg and B were significantly influenced by the application of IAA and B. The highest N (0.70%), Ca (0.69%), Mg (0.80%) in tomato fruits and highest N (0.40%), Ca (0.93%), Mg (0.90%) in tomato plants were obtained in 60 ppm IAA along with 2.0 kg B ha⁻¹ and the lowest was obtained in the control treatment. In tomato fruits the highest P (0.41%) was found in IAA₄₀B_{2.0} treatment, while the lowest P (0.18%) was recorded in IAA₀B_{0.0}. The lowest B (4.55 μg g⁻¹) was recorded in control, while the highest B (15.72 μg g⁻¹) was found in IAA₂₀B_{2.0} in tomato fruits. In tomato plants the highest P (0.34%) was recorded in IAA₆₀B_{1.5} and the lowest P (0.20%) was found in control. The lowest B (4.15 μg g⁻¹) was recorded in IAA₀B_{0.0}, while the highest B (13.33 μg g⁻¹) was found in IAA₄₀B_{2.0} in tomato plants. The overall results suggest that treatment IAA₆₀B_{2.0} was the best from other treatments.

Keywords: Indole Acetic Acid, Boron, Nutrient, summer tomato

1. INTRODUCTION

Tomato (*Lycopersicon esculentum*, mill.), belongs to the family solanaceae. It is one of the most popular, nutritious, palatable and versatile cultivated vegetable all over the world including Bangladesh. It was originated in tropical America [1]. In Bangladesh, it ranks 2nd next to potato [2] and top of the list of canned vegetables. At present China, India, Egypt, Turkey, Iran, Italy, Mexico and Brazil are the leading tomato producing countries in the world [3]. The chemical composition of tomato includes total sugar 2.50-4.50%, vitamin C 15-20 mg/100g, calcium 0.25-0.50 g/100g, magnesium 0.10-1.50 g/100g, phosphorus 0.20-0.80 g/100g, iron 40-500 ppm, zinc 10-50 ppm, lycopene 20-50 g/100g (dry fruit weight basis) [4].

In Bangladesh, it is widely cultivated due to its adaptability to wide range of soil and climate [5]. Soil and climatic conditions of all the districts of Bangladesh, especially Rajshahi, Natore, Chapainawabganj, Naogaon, Pabna, Bogra, Sirajgonj, Joypurhat, Rangpur and Gaibandha districts are very suitable for tomato cultivation. Tomato is a very nutritious vegetable containing considerable amounts of vitamins and minerals [6]. Tomato production during the summer season is restricted due to high temperature, high humidity, and rainfall. Adverse climatic condition during summer causes severe flower dropping in tomato. High

temperature dries up stigmatic fluid and thus resulted in poor fruit setting and yield of tomato. However, scientists have been trying to develop heat tolerant varieties and Bangladesh Institute of Nuclear Agriculture (BINA) has developed some heat tolerant varieties like BINA Tomato-2, BINA Tomato-3 and BINA Tomato-5 [7]. Plant growth regulator has been reported to enhance fruit set under both normal and adverse weather conditions. The study on IAA for tomato production in Bangladesh is very rare and limited at the Research Institutes. IAA not only helps in fruit setting but also improves the biochemical and nutritional composition of tomato. This crop differs in their sensitivity to Boron (B) deficiency and the absorption of B varies with the types of soil. The reduced level of water in the soil also causes a proportionate to decrease the rate of B diffusion to root [8]. Visible symptoms of B deficiency (0.3 μm) initiated on young leaves as internal chlorosis later leading to necrosis [9]. Keeping these points in view the experiment was undertaken to determine the effect of IAA and B on the nutrient contents of BINATomato-3.

2. MATERIAL AND METHODS

A pot experiment was conducted at the net house of the department of Agricultural Chemistry, Bangladesh Agricultural University (BAU), Mymensingh. The research work was accomplished in earthen pot and the different nutrients of the fruit and plant samples were analyzed in the laboratory of the Department of Agricultural Chemistry, BAU, Mymensingh.

The climate of the experiment area was under the sub-tropical climatic zone, which is characterized by moderate to high temperature, heavy rainfall, high humidity and relatively long day during Kharif season (March to August). Before starting the experiment initial soil sample of the experimental plots was analyzed in the laboratory for determination of its physical and chemical properties. The treatments included in the experiment were as follows:

Factor A : Four levels of IAA viz.

1. No IAA (IAA_0)
2. 20 ppm IAA (IAA_{20})
3. 40 ppm IAA (IAA_{40})
4. 60 ppm IAA (IAA_{60})

Factor B: Four levels of Boron viz.

1. No B ($\text{B}_{0,0}$)
2. 1.5 kg B ha^{-1} ($\text{B}_{1,5}$)
3. 2.0 kg B ha^{-1} ($\text{B}_{2,0}$)
4. 2.5 kg B ha^{-1} ($\text{B}_{2,5}$)

2.1 Chemical analysis of fruit

2.1.1 Total nitrogen determination

Total N was estimated by semi-micro kjeldahl method [10] and [12]. In this method, exactly 0.5 g oven dried and the ground sample was digested with 3 mL of H_2O_2 and 6 mL concentration H_2SO_4 in presence of 1 g catalyst mixture in the digestion tube. After the completion of digestion, the flask was cooled at room temperature and about 25 mL of distilled water was added. For performing distillation, 10 mL of the digested solution was taken in a distillation unit with 10 mL of 40% NaOH. The distillate was collected in 10 mL 2% boric acid containing a mixed indicator. The collected distillate in boric acid was titrated with standardized 0.1N H_2SO_4 . The N percentage was calculated by the following formula-

$$\% N = \frac{(T-B) \times N \times 1.4}{S}$$

where,

T = Sample titration (mL) value

B = Blank titration (mL) value

N = Strength of H₂SO₄ (N)

S = Sample weight (g)

2.1.2 Preparation of fruit extract for the determination of different nutrient contents

Exactly 1g of dried and finely ground fruit material was taken into a 250 mL conical flask and 10mL of di-acid mixture (HNO₃: HClO₄ = 2:1) was added to it. Then it was placed on an electric hot plate for heating at 180-200°C until the solid particles disappeared and white fumes were evolved from the flask. Then, it was cooled at room temperature, washed with distilled water and filtered through filter paper (Whatman No. 42) making the volume up to the mark with distilled water following wet oxidation methods as described by [11]. The solution was used for the analysis of P, Ca, Mg and B.

2.1.3 Phosphorus determination

The phosphorus content of the fruit extract was determined by developing blue colour with stannous chloride (SnCl₂.2H₂O) reduction of phosphomolybdate complex and measuring the colour with the help of a spectrophotometer (Model-LT-31) at 660 nm **wavelengths** [12]. Stannous chloride was used as a reducing agent to form molybdophosphoric blue complex with sulphomolybdate. Exactly 2 mL aliquot was taken in a 100 mL volumetric flask followed by the addition of 4 mL of sulphomolybdic acid and 6 drops of stannous chloride solution. The volume was made up to the mark with distilled water and was shaken thoroughly. Finally, the absorbance of light at 660 nm wave length was measured with the help of a spectrophotometer.

2.1.4 Calcium determination

Calcium content of the fruit extract was determined by complexometric method of titration using Na₂EDTA as a complexing agent at pH 12 where calcon was taken followed by 40 mL water, 3 mL 10% NaOH solution, 10 drops each of the hydroxylamine hydrochloride (NH₂OH.HCl), potassium ferrocyanide [K₄Fe(CN)₆.3H₂O] and TEA (triethanol amine; C₆H₁₅NO₃), as masking agent. After the addition of calcon indicator solution, the test sample was titrated against Na₂-EDTA (0.01M) solution from a burette until the pink colour of the solution turned into pure blue.

2.1.5 Magnesium determination

Magnesium content of the fruit extract was determined by the complexometric method of titration using Na₂EDTA as a complexing agent at pH 10 where EBT (Eriochrome Black T) was used as an indicator [12]. Exactly 5 mL of the aliquot was taken following the addition of 40 mL distilled water, 5 mL NH₃-NH₄ buffer solution, 10 drops each of the hydroxyl amine hydrochloride (NH₂OH.HCl), potassium ferrocyanide **[K₄Fe(CN)₆.3H₂O]**, sodium tungstate (Na₂WO₄.2H₂O) and TEA.

2.1.6 Boron determination

The content of boron in fruit extract was determined by a spectrophotometric method using azomethine-H reagent. Exactly 5mL of extract was taken in a 250 mL volumetric flask. Then 4mL buffer masking solution and 4mL azomethine-H was added. Then the volume was made upto the mark. After one hour, absorbance was measured at 420 nm wavelength with help of a spectrophotometer following the analytical technique outlined by [13]. (Triethanol amine; $C_6H_{15}NO_3$), as a masking agent, after the addition of EBT indicator solution, the aliquot was titrated against Na_2EDTA (0.01M) solution from a burette until the pink colour of the solution turned into pure blue.

2.1.7 Plant analysis after harvesting

For the determination of N, Ca, Mg, P and B in plant, 1g dry ground plant sample was taken. Preparation of the stock solution and analyses of the elements were the same as the previous methods of fruit analysis.

2.2 Experimental design and statistical analysis

The experiment was laid out in Completely Randomized Design (CRD) with three replications. The recorded data were subjected to statistical analysis. All the data were analyzed for ANOVA with the help of a computer package program of MSTAT-C (Mathematical and Statistical calculation) [14].

3. RESULTS AND DISCUSSION

3.1 The effect of IAA on different nutrients content in fruits of BINA tomato-3

The nutrient contents of tomato fruits were found highest for N (0.64%), P (0.37%), Ca (0.64%), Mg (0.73%) and B ($12.63 \mu g g^{-1}$) in treatment IAA_{60} and all these fine nutrient elements were lowest in the treatment where no IAA was used (Table.1). The contents of N, P, Ca, Mg and B in tomato fruits significantly increased due to IAA application.

Table 1. The effect of different levels of IAA on nutrients content in fruits of BINA tomato-3

Treatments	N (%)	P (%)	Ca (%)	Mg (%)	B ($\mu\text{g g}^{-1}$)
	IAA ₀	0.54c	0.24c	0.43	0.56b
IAA ₂₀	0.56bc	0.33b	0.36	0.67a	11.13ab
IAA ₄₀	0.62ab	0.34ab	0.53	0.56b	12.13a
IAA ₆₀	0.64a	0.37a	0.64	0.73a	12.63a
Level of significance	**	**	**	**	**
LSD	0.01	0.00	0.02	0.01	0.46

NS = Not significant, **indicates significant at 1% level of probability. Figures in a column having same or no letter(s) do not differ significantly.

3.2 The effect of B on different nutrients content in fruits of BINA tomato-3

BINA tomato-3 fruits contained the highest amount of nutrients viz. N (0.64%), P (0.36%), Mg (0.71%) and B $14.13\mu\text{g g}^{-1}$ at B_{2.0} but the content of Ca (0.52%) was highest in B_{1.5} treatments. All these nutrients were lowest in control (B_{0.0}) treatment (Table.2). A significant variation in N, P, Mg and B contents of tomato fruits were observed due to different levels of B fertilizer. Boron did not show a significant effect on Ca content in tomato fruits.

Table 2. The effect of different levels of B on nutrients content in fruits of BINA tomato-3

Treatments	N (%)	P (%)	Ca (%)	Mg (%)	B ($\mu\text{g g}^{-1}$)
	B _{0.0}	0.54c	0.28c	0.47	0.56b
B _{1.5}	0.61a	0.31bc	0.52	0.57b	10.73b
B _{2.0}	0.64a	0.36a	0.49	0.71a	14.13a
B _{2.5}	0.57ab	0.33ab	0.49	0.68a	12.63a
Level of significance	**	**	NS	**	**
LSD	0.01	0.00	0.02	0.01	0.46

NS = Not significant, **indicates significant at 1% level of probability. Figures in a column having same or no letter(s) do not differ significantly.

3.3 Interaction effect of different levels of IAA and B on different nutrients content in fruits of BINA tomato-3

Effect of interaction on different nutrients content in fruits of BINA tomato-3 is presented in Table 3. The effects of IAA with different doses of boron on N, P, Ca, Mg and B contents in tomato fruits were statistically significant. The nutrients contents of tomato fruits were found highest for N (0.70%) in treatment IAA₆₀B_{2.0} and lowest value (0.49%) was obtained in treatment IAA₀B_{0.0}. The P content of tomato fruits was highest (0.41%) in treatment IAA₄₀B_{2.0} and it was lowest (0.18%) in treatment IAA₀B_{0.0}. The Ca content of tomato fruits was highest (0.69%) in treatment IAA₆₀B_{2.0} and its lowest value (0.32%) was found in IAA₂₀B_{2.0} and IAA₂₀B_{2.5} treatments. The Mg content of tomato fruits was highest (0.80%) in treatment IAA₆₀B_{2.0} and its lowest value (0.30%) was obtained in control. The B content of tomato fruits was highest (15.72 µg g⁻¹) in treatment IAA₂₀B_{2.0} and its lowest value (4.55 µg g⁻¹) was found in treatment control. The results clearly indicated that N, P, Ca, Mg and B contents increased in tomato fruits due to the application of IAA and B.

Table 3. Interaction effect of IAA and B on nutrients content in fruits of BINA Tomato-3

Treatments	N (%)	P (%)	Ca (%)	Mg (%)	B (µg g ⁻¹)
IAA ₀ B _{0.0}	0.49c	0.18h	0.40 cde	0.30f	4.55f
IAA ₀ B _{1.5}	0.57abc	0.23gh	0.45 b-d	0.55de	9.34de
IAA ₀ B _{2.0}	0.59abc	0.28fg	0.43 b-d	0.71ab	11.73bcd
IAA ₀ B _{2.5}	0.50bc	0.26g	0.45 b-d	0.66bcd	12.13a-d
IAA ₂₀ B _{0.0}	0.51bc	0.29efg	0.43 b-d	0.75ab	7.34ef
IAA ₂₀ B _{1.5}	0.62abc	0.30def	0.37de	0.50e	10.14cde
IAA ₂₀ B _{2.0}	0.63abc	0.37abc	0.32e	0.68abc	15.72a
IAA ₂₀ B _{2.5}	0.49c	0.34cd	0.32e	0.76ab	11.33bcd
IAA ₄₀ B _{0.0}	0.56abc	0.29efg	0.45b-d	0.56cde	8.54de
IAA ₄₀ B _{1.5}	0.66a	0.33bcd	0.61abc	0.48e	12.13a-d
IAA ₄₀ B _{2.0}	0.64ab	0.41a	0.51a-d	0.66bcd	14.53ab
IAA ₄₀ B _{2.5}	0.60abc	0.35bc	0.56a-d	0.55de	13.33abc
IAA ₆₀ B _{0.0}	0.59abc	0.34bc	0.61abc	0.63bcd	10.93cde
IAA ₆₀ B _{1.5}	0.60abc	0.37abc	0.64ab	0.74ab	11.33bcd
IAA ₆₀ B _{2.0}	0.70a	0.39ab	0.69a	0.80a	14.52ab
IAA ₆₀ B _{2.5}	0.68a	0.36bc	0.61abc	0.76ab	13.73abc
Level of significance	*	**	**	**	**

B _{2.0}	0.33	0.31a	0.79	0.68	11.63a
B _{2.5}	0.33	0.30a	0.80	0.64	10.53a
Level of significance	NS	**	NS	NS	**
LSD	0.01	0.00	0.02	0.02	0.50

NS = Not significant, **indicates significant at 1% level of probability. Figures in a column having same or no letter(s) do not differ significantly.

3.6 The interaction effect of different levels of IAA and B on different nutrients content in a plant of BINA tomato-3

Effect of interaction on different nutrients content in a plant of BINA tomato-3 is presented in Table 6. All the treatments significantly increased N, P, Ca, Mg and B contents in tomato plants as compared to the control treatment (IAA₀B_{0.0}). The interaction effect of different levels of IAA and B on nutrients composition of tomato plants was found highest for N (0.40%) in treatment IAA₆₀B_{2.0}. P (0.34%) was found highest in treatment IAA₆₀ B_{1.5} and Ca was highest (0.93%) in treatments IAA₆₀B_{1.5} and IAA₆₀ B_{2.0}. Mg was found highest (0.90%) in treatment IAA₆₀B_{2.0}. The maximum B content (13.33 $\mu\text{g g}^{-1}$) in treatment IAA₄₀ B_{2.0} and all these nutrients were lowest in treatment IAA₀ B_{0.0} in control. The results showed that N, P, Ca, Mg and B contents increased in tomato plants due to the application of IAA and B.

It was further observed that the nutrients content in both tomato fruits and plants increased with the increasing level of IAA and a minimum level of B but better nutrients content in tomato fruits and plants were possible by using 60 ppm IAA and 2 kg B ha⁻¹. Applying of both natural and synthetic auxin helps the farmer in cultivating tomato in an adverse climatic condition which can give good fruit yield by increasing vegetative and reproductive growth and reducing the flower and fruit drop [15, 16].

Table 6. Interaction effect of different levels of IAA and B on plant nutrients content of BINA tomato-3

Treatments	N (%)	P (%)	Ca (%)	Mg (%)	B ($\mu\text{g g}^{-1}$)
IAA ₀ B _{0.0}	0.17f	0.20g	0.51d	0.36f	4.15e
IAA ₀ B _{1.5}	0.22ef	0.26ef	0.59cd	0.36f	6.94de
IAA ₀ B _{2.0}	0.26de	0.28cde	0.64bc	0.39f	8.54bcd
IAA ₀ B _{2.5}	0.23def	0.26ef	0.61cd	0.39f	9.34a-d
IAA ₂₀ B _{0.0}	0.35ab	0.25f	0.61cd	0.42f	8.54bcd
IAA ₂₀ B _{1.5}	0.30bcd	0.28cde	0.69bc	0.56de	11.33abc
IAA ₂₀ B _{2.0}	0.31bc	0.29cd	0.75abc	0.68bcd	12.13ab
IAA ₂₀ B _{2.5}	0.35ab	0.31bc	0.83ab	0.60cd	10.93a-d
IAA ₄₀ B _{0.0}	0.35ab	0.28cde	0.72bc	0.56de	7.34cde

IAA ₄₀ B _{1.5}	0.38ab	0.30bc	0.85ab	0.68bcd	11.33abc
IAA ₄₀ B _{2.0}	0.35ab	0.33ab	0.83ab	0.76abc	13.33a
IAA ₄₀ B _{2.5}	0.37ab	0.31bc	0.88ab	0.74a-d	10.93a-d
IAA ₆₀ B _{0.0}	0.32bc	0.29cd	0.83ab	0.76abc	9.74a-d
IAA ₆₀ B _{1.5}	0.38ab	0.34a	0.93a	0.86ab	11.33abc
IAA ₆₀ B _{2.0}	0.40a	0.33ab	0.93a	0.90a	12.53ab
IAA ₆₀ B _{2.5}	0.36ab	0.30	0.88ab	0.83ab	10.93a-d
Level of significance	**	**	**	**	**
LSD	0.04	0.01	0.09	0.09	2.00

NS = Not significant, **indicates significant at 1% level of probability. Figures in a column having same or no letter(s) do not differ significantly.

4. CONCLUSION

When IAA and B were applied together, it gave a remarkably higher response in terms of nutrient contents of a summer tomato. The highest Nitrogen, Calcium and Magnesium of BINA Tomato-3 in both cases of fruits and plants were obtained from the application of IAA at the rate of 60 ppm along with B at the rate of 2.0 kg ha⁻¹.

REFERENCES

1. Solunkhe DK, Desai BB and Bhat NR. Vegetables and flower seed production. 1st Edn. Agricola publishing Academy, New Delhi, India; 1987.
2. BBS (Bangladesh Bureau of Statistics). The hand book of Agricultural Statistics, Ministry of planning, Govt. People's Republic of Bangladesh, Dhaka. 2009; 14.
3. FAO (Food and Agriculture Organization). FAO Production Year Book. Basic Data Branch, Statistics Division, FAO; 2002.
4. Frusciantè L, Carli P, Ercolano MR, Pernice R and Matteo AD. Antioxidant nutritional quality of tomato. Molecular Nutrient and Food Research. 2007; 51 : 609-617.
5. Ahmed KU. Phul phal O Shak Shabji (in Bangla). 2nd Edition. Banglow No. 2, Farm Gate, Dhaka, Bangladesh. 1995;470.
6. Bose TK and Some MG. Vegetable Crops. Naya Prokash, 206 Bidhan Sarani, Kolkata, India. 2002;35.
7. Begum SM, Hayder RM and Pasha NUM. New varieties of summer tomato. Genetics and Plant Breeding Division, Bangladesh Institute of Nuclear Agriculture (BINA). [A leaflet in Bangla entitled "Grishma Kalin Tomator Natun Jat"]. 1998.
8. Barber SA. Soil Nutrient Bioavailability. A mechanistic Approach. John Wiley and Sons, New York, USA. 1995;23.
9. Sinha P, Dabe BK and Chatterjee C. Influence of boron stress on biomass yield, metabolisms and quality of groundnut. Indian J. Plant physiol. 2002;7(2): 131-134.

10. Jackson, ML. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd., New Delhi. 1973;151-154.
11. Singh D, Chhonkar PK and Pandey RN. Soil Plant Water Analysis. A Method Manual. IARI, New Delhi, India. 1999;72-86.
12. Page AL, Miller RH and Keeney DR. Methods of Soil Analysis, Part-2. 2nd Edn. Amer. Soc. Agron. Inc. Madison, Washington, USA. 1982;98-765.
13. Tandon HLS. Methods of Analysis of Soils, Plants, waters and Fertilizers. Fertilizer Development and Consultation Organization, New Delhi, India. 1995;84-90.
14. Russel DF. MSTAT-C Package Programme. Dept. of Crop and Soil Science, Michigan State University, USA;1986.
15. Pramanik, K., Pradhan, J., & Sahoo, S. K. (2018). Role of auxin and gibberellins growth, yield and quality of tomato: A review. *International Journal of Current Microbiology and Applied Sciences*, ISSN: 2319-7706, 6 (11), 1624-1636.
16. Verma P. S., Meena M. L. And Meena S. K. 2014. Influence Of Plant Growth Regulators On Growth, Flowering And Quality Of Tomato (*Lycopersicon Esculentum* Mill), Cv. H-86. *Indian Journal Of Hill Farming* 27(2):19-22.

UNDER PEER REVIEW