

# Occurrence of Bacterial leaf Blight of Cocoyam and Characterisation of the Causal Organism in Northern Nigeria

Abdullahi, A.<sup>1</sup>, Alao, S.E.L.<sup>1</sup>, Banwo, O. O.<sup>1</sup>, Alabi, O.<sup>1</sup>, Lu, S.<sup>2</sup>, Bandyopadhyay, R.<sup>3</sup>

<sup>1</sup>Department of Crop Protection, Ahmadu Bello University, Zaria, Nigeria;

<sup>2</sup>Department of Biochemistry, Entomology, Molecular Biology and Plant Pathology, Mississippi State University, USA.

<sup>3</sup>International Institute of Tropical Agriculture, Ibadan.

Corresponding author's email: aabdullahi2015@gmail.com

## ABSTRACT

**Aim:** To determine the occurrence of bacterial leaf blight of cocoyam and to characterize isolates of the causal organism in Northern Nigeria.

**Place and Duration of Study:** The study was conducted in cocoyam farmers' fields in Kaduna and Kano States, Department of Crop Protection, Ahmadu Bello University, Zaria, Nigeria and the Mississippi State University, USA in 2015.

**Methodology:** Field surveys were conducted in Kaduna and Kano States of northern Nigeria to determine occurrence of bacterial leaf blight of cocoyam. Sampling was done using quadrat. Incidence and severity of the disease was assessed on the field. Bacterial isolates obtained from the diseased samples were used for pathogenicity and hypersensitive reaction (HR) tests. Biochemical tests such as Gram, catalase, oxidase, pectolytic, amylolytic, and production of acid from glycerol were carried out on the isolates. A nearly full length of the 16S rDNA gene of selected isolates was PCR amplified using 16S rRNA primers. The resulting amplicons were sequenced using Sanger sequencing. The 16S rDNA gene sequences were aligned along with other *Xanthomonas* sequences imported from the NCBI database using muscle tool from MEGA6.

**Results:** The results showed Kaduna State had higher incidence and severity (50.2 %, 13.5 %) than Kano State (20.5%, 5.1%). and that bacterial isolates induced blight symptoms on cocoyam and elicited HR reactions on tobacco. Isolates were Gram negative, catalase positive, oxidase negative, amylolytic, pectolytic and produced no acids from glycerol. BLAST search of sequenced genes showed 98-100% homology to *Xad*. Maximum likelihood phylogenetic trees constructed for the 16S rDNA gene sequences revealed isolates were identical to the *Xad* reference strains KP247494, KM576803 and EU203153. Generally, all the isolates obtained were *Xad*.

**Conclusion:** Bacterial leaf blight of cocoyam occurred in Kaduna and Kano States and the bacterial isolates were identical to *Xanthomonas axonopodis* pv. *dieffenbachiae*.

**Keywords:** Incidence; Severity; Kaduna; Kano; Biochemical; Molecular; 16S rDNA

## **1.0 INTRODUCTION**

Cocoyam (*Colocasia* and *Xanthosoma* species) which belongs to the family Araceae is one of the oldest crops grown mostly in the tropics, for its edible corms and leaves and as an ornamental plant (1). Cocoyam is one of the basic food crops of major economic importance and ranks third in importance after cassava and yam among the root and tuber crops cultivated and consumed in Nigeria (2). It is superior to yam and cassava nutritionally, with higher protein, mineral and vitamin contents and the starch more readily digested (3). Nigeria is the world's leading producer of cocoyam, with annual production of 3.27 million metric tonnes, accounting for about 36 percent of total world output of cocoyam (4). The yield and quality of this crop are threatened by various abiotic and biotic factors. Diseases caused by fungi (5; 6; 7) and bacteria (8; 9) are among biotic factors militating against the production of cocoyam in Nigeria.

Bacterial leaf blight (BLB) was first reported as an important disease on cocoyam in Nigeria (9). The causal agent was identified as *Xanthomonas axonopodis* pv. *dieffenbachiae* using pathogenicity and biochemical tests; The status of this disease in Northern Nigeria, another cocoyam producing region has not been reported. Molecular characterisation using polymerase chain reaction (PCR) based technique has proven to be a fast, sensitive and reliable method for determining genetic relationships among pathogenic organisms (10). However, genetic profile of *X. axonopodis* pv. *dieffenbachiae* on cocoyam using this technique has also not been reported anywhere in Nigeria.

There was the need to assess the status of this disease in Northern Nigeria where the crop occupies a very important position in the farming system. This research will provide valuable information that will form the basis for cocoyam bacterial leaf blight disease management decisions and breeding cocoyam cultivars with resistance to the pathogen. Molecular characterisation of the pathogen in the region will also provide information that could be used to map the pathogen in the region, a tool that could serve quarantine purposes.

## **2.0 MATERIALS AND METHODS**

### **2.1 Incidence and Severity of Bacterial Leaf Blight of Cocoyam**

Incidence and severity of bacterial leaf blight of cocoyam were determined between June and July November, 2015 in Kaduna and Kano States (Appendices 1 and 2). In each State, three Local Government Areas (LGAs) and three farming communities per LGA were visited. Three fields were surveyed in each community. In each field, cocoyam leaves were examined

for symptoms of bacterial leaf blight, thereafter, a 2 by 2 m<sup>2</sup> quadrat was set each at four corners and at the centre of the field. The total number of plants in the quadrat was recorded. Disease incidence was assessed by counting the number of plants showing bacterial leaf blight symptoms in each quadrat and the total number of plants in that quadrat. The disease incidence was calculated using the formula:

$$\text{Disease incidence} = \frac{\text{Number of plants infected}}{\text{Total number of plants examined}} \times 100$$

Disease Severity was recorded per plot using a disease scale of 0-5 developed by Opara *et al.* (9):

0 = no disease symptom visible on the plant;

1 = less than 10% of plant affected;

2 = 10-30 % of plant affected;

3 = 31-50% of plant affected;

4 = 51 - 70% of plant affected;

5 = above (>) 70% of plant affected;

Disease Severity Index (DSI) was evaluated using the following formula (11):

$$\text{DSI} = \frac{\text{Sum of individual disease scores}}{\text{Total number of plants scored} \times \text{maximum score}} \times 100$$

Leaf samples were collected from each field, wrapped in paper bags, labeled and taken to the Bacteriology Laboratory of the Department of Crop Protection, Ahmadu Bello University, Zaria for isolation. Global position system (GPS) parameters for each location were recorded using a Handheld GPS Navigator manufactured by Garmin, USA. Unstructured questionnaire was used to obtain some cropping information from the surveyed areas.

## **2.2 Characterisation of *Xanthomonas* spp**

### **2.2.1 Isolation of organism, hypersensitive reaction (HR) and pathogenicity tests**

Sampled leaves were surface-sterilised with 3.5 % w/v sodium hypochlorite and washed thoroughly with sterile water for 3 minutes before isolation. Small pieces (3 mm) of advancing lesions on leaves were excised from the boundary between diseased and the healthy tissues using sterile scalpel. Excised tissue was rinsed with sterile distilled water (SDW) for 1 minute before it was placed in a Petri dish containing 3.0 ml of SDW, shredded with sterile needles and allowed to stand for 10 minutes,. The suspension was then streaked on nutrient

agar (Difco) plates, placed upside down in an incubator set at 27 °C for 48-72 hours. Pure cultures were obtained after sub-culturing twice. The pure cultures were then stored in SDW in McCartney bottles at 4 °C.

Twenty-four-hour old cultures of the bacterial colony obtained were adjusted with spectrophotometer to an optical density of 0.3 at 600 nm of light, corresponding to a bacterial concentration of  $10^8$  cfu/ml. Inoculum concentration of  $10^8$  cfu/ml was used for HR test on tobacco leaves according to Wick (12). Cocoyam corms were planted into 30 cm diameter pots filled with heat sterilised soil at the rate of one corm per pot. The bacteria isolate was tested for pathogenicity on 15 day-old cocoyam plants (*Colocasia esculenta*).

### **2.2.2 Biochemical characterization**

Biochemical characterization was done by subjecting the bacterial isolates to the following tests (13).

#### ***Test for Gram reaction***

For Gram reaction, the alternative potassium hydroxide (KOH) test was done for all samples. Twenty-four hour old culture was added to 3% KOH solution on clean glass slide using flame-sterilized needle. The bacterial suspension was stirred against the KOH solution while occasionally lifting the loop for slimy and thickened slurry.

#### ***Test for catalase***

A drop of 3% hydrogen peroxide ( $H_2O_2$ ) was placed on glass slide, onto which a loopful of bacteria from a 24 hour old culture was suspended and observed for bubbling and production of gases.

#### ***Oxidase reaction test***

One ml 1 % aq. w/v solution of N'-tetramethyl-p-phenylene-diaminedihydrochloride solution was prepared and 100 µl of the solution were placed on a new piece of Whatman No. 1 filter paper with a clean Pasteur pipette. Part of a colony was removed with a sterile toothpick and smeared onto the moistened paper. This was observed for colour changes which should occur within 30 seconds for positive results.

#### ***Test for Pectolytic reaction***

Bacterial isolates were grown on sodium pectate gel medium which contained pectate substrate and incubated at 28 °C for three days. Production of degrading enzymes was

detected by observing depressions in the gel around the colony where the substrate has been degraded.

#### ***Test for amylolytic reaction***

Twenty-four hour old cultures of *Xanthomonas* spp. isolates were transferred to nutrient agar plates amended with 0.2% (w/v) soluble starch. Cultures that emerged were incubated at 30 °C for 72 hours. Plates were flooded with Lugol's iodine solution and observed for any change of colour which signifies starch hydrolysis.

#### ***Acid production***

Solutions of Dye's medium C and carbohydrate sources were prepared in tubes and inoculated with the isolates. This was incubated at 28 °C for 2-10 days and observed for colour changes. Yellow colour formation indicates acid production from carbohydrate. Substrates that were tested included glucose, fructose, mannose, lactose, galactose, sucrose, glycerol and arabinose.

#### **2.2.3 DNA extraction, Genomic finger printing and agarose gel electrophoresis**

Genetic profiling was conducted at the Department of Biochemistry, Entomology, Molecular Biology and Plant Pathology, Mississippi State University, USA. Bacteria cultures grown at 28°C on NBY medium (8 g liter<sup>-1</sup> nutrient broth, 2 g liter<sup>-1</sup> yeast extract, 2 g liter<sup>-1</sup> K<sub>2</sub>HPO<sub>4</sub>, 0.5 g liter<sup>-1</sup> KH<sub>2</sub>PO<sub>4</sub>, 15 g agar, 974 ml distilled water, 25 ml 20 % glucose, 1 ml 1 M MgSO<sub>4</sub>; pH 7.2) for 24 hours were cultivated in NBY liquid medium (NBY Medium above without Agar). 16S rDNA was extracted using Wizard Genomic DNA Purification Kit (14). Genomic DNA thus extracted was stored at -20 °C. PCR protocol was used to amplify the purified bacterial genomic DNA. 16S rRNA primers were used for the amplification, forward: 27F (5'-AGAGTTTGATCMTGGCTCAG-3') and reverse: 1492R (5'-GGYTACCTTGTTACGACTT-3') (15). The component of PCR master mix included buffer (10 µl), dNTP<sub>s</sub> (1.5 µl), MgCl<sub>2</sub> (5 µl), forward primer (1.5 µl), reverse primer (1.5 µl), Taq polymerase (0.5 µl) and sterile water (30 µl). Two microlitre of the purified DNA of each sample was pipetted into a separate PCR tube. Fifty microlitre of the master mix was added to each of the DNA samples, vortexed and centrifuged before placing in the thermocycler. The amplification program included initial denaturation at 94°C for 3 min, 32 cycles consisting of denaturation at 94°C for 20 s, annealing at 56°C for 30 s, and extension at 72°C for 1 min 10 s, and an extra extension step at 72°C for 5 min.

Amplification of the DNA was confirmed by gel electrophoresis. Agarose was prepared at the desired concentration (w/v) in 1xTris-Acetate-EDTA (TAE) buffer (1.0 % gel). 1.6 µl of loading buffer (15% Ficoll<sup>®</sup> 400, 0.03 % bromophenol blue, 0.03 % Xylene cyanol FF, 0.4 % orange G, 10 mM Tris-HCl (pH 7.5) ) was mixed with 8 µl of PCR product and loaded slowly into the wells. A known *Xanthomonas* strain (MSCTI) and sterile distilled water were also loaded to serve as positive and negative controls respectively. DNA molecular weight marker (1 kb) was also loaded. The order of sample loading in the gel was recorded. Ethidium bromide solution (0.5 µg/ml) was added to the buffer before running the gel.

Electrophoresis unit was then connected to the power pack and power supply turned on until the bromophenol blue dye reached the bottom of the gel (approximately 30 min at 100 V, for DNA to migrate 7 cm from the wells in a 1.0 % gel). The gel was removed and observed under UV Transilluminator. Photograph of the gel was taken using an orange filter fitted camera (Kumar 2009). The resulting amplicons were sequenced at the Eurofins Genomics Company, Eurofins MWG Operon LLC, Huntsville, Alaska, USA using Sanger sequencing. 16S rDNA gene sequences from the *Xanthomonas* spp. isolates were blasted against sequences of related xanthomonads from the NCBI online database ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) to determine genomic relatedness.

### 2.3 Statistical Analysis

Data from the survey were analyzed using descriptive statistics. The *Xanthomonas* spp. isolates were analysed for their evolutionary history using the maximum likelihood method. The analysis was done following the Tamura-Nei model (16). Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the maximum composite likelihood (MCL) approach, and then selecting the topology with the superior log likelihood value. Codon positions included were 1<sup>st</sup>+2<sup>nd</sup>+3<sup>rd</sup>+Noncoding. All positions containing gaps and missing data were eliminated. Evolutionary analyses were conducted in MEGA6 (17).

## 3.0 RESULTS

### 3.1 Incidence and Severity of Bacterial Leaf Blight

Mean bacterial blight incidence and severity observed in the Local Government Areas (LGAs) of the States are shown in Figures 1 and 2. In Kaduna State, the incidence in Kudan (93.7 %) was highest, followed by Sabon Gari (40.4 %) and the least disease incidence was recorded in

Giwa (16.6 %). For Kano State, Kiru LGA (30.0 %) had the highest disease incidence, followed by Bebeji (10.4 %) and there was no incidence of the disease in Rogo LGA. The disease severity in Kaduna State was highest in Kudan LGA (45.3 %), followed by Sabo Gari LGA (14.4 %), while Giwa LGA (6.3 %) had the least. In Kano State, Kiru LGA (12.7 %) had the highest disease severity, followed by Bebeji LGA (2.7 %). The mean bacterial blight incidence and severity for Kaduna and Kano States are shown in Figure 3, with Kaduna State having higher disease incidence and disease severity index than Kano State respectively.

The different symptoms observed in the locations were mostly spots and blight, consisting of small, water-soaked lesions, usually surrounded by prominent chlorotic halo (Plate I).

### **3.2 Characterisation of *Xanthomonas* spp**

#### **3.2.1 Tobacco Hypersensitive and Pathogenicity Tests**

The isolated bacteria formed yellow, convex, mucoid colonies on nutrient agar medium. Isolates corresponding to different locations were designated numerically with the prefix 'CE' to indicate cocoyam genus (*Colocasia esculenta*) origin. Four isolates namely CE2, CE3, CE4 and CE6 designating isolates from Tudun Sarki, Jaja, Yelwan Paki and Hunkuyi, respectively elicited hypersensitive reaction (HR) on tobacco and were found to be pathogenic on cocoyam. On tobacco, a clear HR was observed 48 hours after suspension of the inoculum was infiltrated into leaf tissues (Plate II). Control leaves did not show similar reaction. On cocoyam, symptoms appeared 2-5 days after inoculation on the leaves. Prominent chlorotic halos developed around the lesions which were similar to those observed in the field (Plate III).

#### **3.2.2 Biochemical characterization**

All the isolates were Gram negative, oxidase negative, catalase, amylolytic, pectolytic and producing acid from glucose, fructose, mannose, lactose, galactose, sucrose, arabinose and not from glycerol (Table 1).




Figure 1: Incidence and severity of bacterial leaf blight on cocoyam leaves in three Local Government Areas of Kaduna State, Nigeria during 2015 wet season.




Figure 2: Incidence and severity of bacterial blight on cocoyam leaves in three Local Government Areas of Kano State, Nigeria during 2015 wet season.






Figure 3: Incidence and severity of bacterial blight on cocoyam leaves in Kaduna and Kano States, Nigeria during 2015 wet season.



Plate I: Typical symptoms of bacterial leaf blight on cocoyam leaf in a field in Jaja,




Plate II: Necrotic portions on tobacco leaf inoculated with *Xanthomonas axonopodis* pv. *dieffenbachiae* showing hypersensitive reaction




Plate III: Leaf blight symptoms on cocoyam inoculated with *Xanthomonas axonopodis* pv. *dieffenbachiae* in the screen house

Table 1: Biochemical reaction of the isolates

| Biochemical tests     | Isolates |     |     |     |
|-----------------------|----------|-----|-----|-----|
|                       | CE2      | CE3 | CE4 | CE6 |
| Gram's reaction       | -        | -   | -   | -   |
| Oxidase activity      | -        | -   | -   | -   |
| Catalase Activity     | +        | +   | +   | +   |
| Amylolytic Activity   | +        | +   | +   | +   |
| Pectolytic Activity   | +        | +   | +   | +   |
| Acid production from: |          |     |     |     |
| Glucose               | +        | +   | +   | +   |
| Fructose              | +        | +   | +   | +   |
| Mannose               | +        | +   | +   | +   |
| Lactose               | +        | +   | +   | +   |
| Galactose             | +        | +   | +   | +   |
| Sucrose               | +        | +   | +   | +   |
| Arabinose             | +        | +   | +   | +   |
| Glycerol              | -        | -   | -   | -   |

+ and - indicate positive and negative reactions, respectively to the test indicated.

CE2 = Tudun Sarki isolate

CE3 = Jaja isolate

CE4 = Yelwan Paki isolate

CE6 = Hunkuyi isolate

### 3.2.3 Gel electrophoresis of amplified 16S rDNA

The 16S rDNA of four out of the six isolates were amplified using 16S rRNA primers (Plate IV) and the expected band size of 1465 bp was obtained.

### 3.2.3 Phylogenetic analysis

The sequences of the isolates were aligned and phylogenetic trees of sequenced isolates were constructed. The isolates obtained from Northern Nigeria form the same clade with three *Xanthomonas axonopodis* pv. *dieffenbachiae* strains KP247494, KM576803 and EU203153 on the dendrogram (Figure 4).




Plate IV: Electrograph of amplified 16S rDNA genes of *X. axonopodis* pv. *dieffenbachiae* isolates from Northern Nigeria.

M = molecular marker

Lanes: 1 = CE2 (Tudun Sarki isolate)

2 = CE3 (Jaja isolate)

3 = CE4 (Yelwan Paki isolate)

4 = CE6 (Hunkuyi isolate)

5 = MSCTI (positive control)

6 = sterile distilled water (negative control)




Figure 4: Phylogenetic tree showing relatedness of gene sequences of *Xanthomonas* isolates from Northern Nigeria to other *Xanthomonas* spp.

CE2 = Tudun Sarki isolate

CE3 = Jaja isolate

CE4 = Yelwan Paki isolate

CE6 = Hunkuyi isolate

## DISCUSSION

The bacterial leaf blight of cocoyam was recorded in the two States surveyed, with Kaduna State recording a significantly higher incidence and severity of the disease than Kano State. The high incidence and severity in Kaduna State could be as a result of higher rainfall and relative humidity of the areas surveyed in Kaduna State than in Kano State (Appendix 1). Most of the fields surveyed in Kaduna State, such as those in Kudan and Giwa LGAs were wetlands, a condition which favours the pathogen. *Xanthomonas axonopodis* pv. *dieffenbachiae* is favoured by warm and moist conditions. Hot, humid, and rainy weather conditions are ideal for disease development and the spread of *Xad* (18). The disease

incidence recorded in some Local Government Areas in Kaduna and Kano States was in line with the earlier report of Opara *et al.* (9), who reported over 65-68% incidence of BLB at Umudike, Abia State, South East Nigeria. A 74-100% incidence of BLB was also reported from a survey of commercial cocoyam fields in Florida, U. S. A. (19). However, the disease severity generally was low, averaged 2 or less, indicating that less than 30 % of the foliar area was damaged by *Xad*. For example, Pohronezny *et al.* (19) reported a less than 10 % disease severity of BLB in their survey of commercial cocoyam fields infected by *Xad* in Florida.

Bacterial organisms isolated from diseased leaf of cocoyam collected from Northern Nigeria fit biochemical descriptions of *Xanthomonas axonopodis*. Biochemical tests showed that the isolates are pectolytic, confirming Chase *et al.* (20) report which stated that strains from *Colocasia*, *Dieffenbachia* and *Philodendron* were highly pectolytic, while those obtained from *Xanthosoma* and *Synгонium* are non-pectolytic. Acids were not produced from glycerol by isolates obtained from various locations in Northern Nigeria which confirms the findings of Pohronezny *et al.* (19) and Opara *et al.* (9) who reported that strains from cocoyam did not produce acid from glycerol.

Gel electrophoresis plates showed that DNA of all isolates were amplified with 16S rRNA primers used. BLAST results for the isolates show 98-100 % genetic relatedness to *Xanthomonas axonopodis* pv. *dieffenbachiae* strains in the Genbank. On the phylogenetic trees, it is observed that all the strains form same clade with *XAD* reference strains KP247494, KM576803 and EU203153. This result re-affirms that isolates from Northern Nigeria are *Xanthomonas axonopodis* pv. *dieffenbachiae*. 16S rDNA profiling used, though reported by Vauterin *et al.* (21) to be inconsistent was able to distinguish the *XAD* as evidence from the PCR results and phylogenetic tree in this work.

## CONCLUSION

Bacterial leaf blight of cocoyam occurred in Kaduna and Kano States and the disease is more prevalent in Kaduna State. The nucleotide sequences of the isolates have been deposited in the Genbank (Accession numbers: KT758338.1, KT758339.1, KT758340.1 and KT758341.1) and can be accessed on <http://www.ncbi.nlm.nih.gov/nucleotide>. The sequences represent the first submission of *Xanthomonas axonopodis* pv. *dieffenbachiae* (causal organism of bacterial leaf blight of cocoyam) on any plant in West Africa and on cocoyam in Africa.

## **RECOMMENDATIONS**

Based on the findings of this study, the following recommendation is suggested:

- 1) An effective strategy which could help in the integrated management of the disease in the surveyed areas is recommended. This is because bacterial leaf blight is one of numerous diseases currently threatening cocoyam production in Northern Nigeria.

## **ACKNOWLEDGEMENTS**

We are very grateful to the United States Agency for International Development (USAID) and the University of California, Davis for sponsoring this research through the Norman E. Borlaug International Agricultural Science and Technology Fellows Leadership Enhancement in Agriculture Programme (Borlaug-LEAP).

## **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

## **AUTHORS' CONTRIBUTIONS**

SEL Alao designed the study; A. Abdullahi performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. O.O. Banwo managed the analyses of the study while O. Alabi, S. Lu. And R. Bandyopadhyay managed the literature searches. All authors read and approved the final manuscript.

## REFERENCES

1. Adelekan BA. An evaluation of the global potential of cocoyam (*Colocasia* and *Xanthosoma* species) as an energy crop. *British Journal of Applied Science and Technology*. 2012; 2(1): 1-15.
2. Baruwa OI, Oke JTO. Analysis of the technical efficiency of small-holder cocoyam farms in Ondo State, Nigeria. *Tropicultura*. 2012; 30(1): 36-40.
3. Odebunmi EO, Oluwaniyi OO, Sanda AM, Kolade BO. Nutritional composition of selected tubers and root crops used in Nigeria food preparations. *International Journal of Chemistry*. 2007; 17(1): 37-43.
4. FAOSTAT. Food and Agriculture Organization of the United Nations Statistics (FAOSTAT Data Results). 2014. [www.fao.org](http://www.fao.org).
5. Ugwuoke KI, Onyeke CC, Tsopmbeng NG. The efficacy of botanical protectants in the storage of cocoyam (*Colocasia esculenta* (L) Schott). *Journal of Tropical Agriculture, Food, Environment and Extension*. 2008; 7: 93 -98.
6. Bandyopadhyay R, Sharma K, Onyeka TJ, Aregbesola A, Lava-Kumar P. First report of taro (*Colocasia esculenta*) leaf blight caused by *Phytophthora colocasiae* in Nigeria. *American Phytopathological Society Journal*. 2011; 95(5): 918
7. Zarafi AB, Chindo PS, Shenge KC, Alao SEL. Investigations on cocoyam diseases in north western Nigeria. In: *Progress report of research projects*, Institute for Agricultural Research, Samaru, Zaria, Nigeria. 2012. Pp211-212.
8. Amodu US, Akpa AD. Determination of the relative susceptibility of roots and tubers to the soft rot bacteria (*Pectobacterium* spp.). *New Clues in Sciences*. 2012; 2: 97-103.
9. Opara E, Njoku CT, Isaiah C. Potency of some plant extracts and pesticides on bacterial leaf blight diseases of cocoyam (*Colocasia esculenta*) in Umudike, South Eastern Nigeria. *Greener Journal of Agricultural Sciences*. 2013; 3(5): 312-319.
10. Khoodoo MHR, Jaufeerally-Fakim Y. RAPD-PCR fingerprinting and Southern analysis of *Xanthomonas axonopodis* pv. *dieffenbachiae* strains isolated from different aroid hosts and locations. *Plant Diseases*. 2004; 88: 980-988.
11. Waller JM, Lenne JM, Waller SJ. *Plant Pathologist's Pocketbook*. 3rd edn. CABI Publishing, New York. 2012. pp. 27.
12. Wick R. Tobacco Hypersensitive; the first test to screen bacteria for pathogenicity. National Plant Diagnostic Network, USDA. 2010. Retrieved from [www.npdn.org/webfm\\_send/1230](http://www.npdn.org/webfm_send/1230), on 8/7/2014, 5 pm.



13. Goszczynska T, Serfontein JJ, Serfontein S. *Introduction to Practical Phytobacteriology: A Manual for Phytobacteriology*. SAFRINET, the Southern African (SADC) LOOP of BioNET-INTERNATIONALBacterial. 2000; P.13-18.
14. Promega. Technical Manual-Wizard Genomic DNA Purification Kit. Promega Corporation, 2800 Woods Hollow Road, Madison, WI 53711-5399, USA. 2005. Pp 16-17.
15. Lane DJ. 16S/23S rRNA sequencing. In: E. Stackebrandt and M. Goodfellow. *Nucleic acid techniques in bacterial systematics*. New York, N.Y.: John Wiley & Sons, Inc. 1991. pp. 115-176.
16. Tamura K, Nei M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*. 1993; 10: 512-526.
17. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution*. 2013; 30: 2725-2729.
18. Deng Z, Seijo ET, Peres AN. Characterisation of strains of *Xanthomonas axonopodis* pv. *dieffenbachiae* from bacterial blight of caladium and identification of sources of resistance for breeding improved cultivar. *HortScience*. 2010; 45 (2): 220-224.
19. Pohronezny K, Volin RB, Dankers W. Bacterial leaf spot of cocoyam (*Xanthosoma caracu*), incited by *Xanthomonas campestris* pv. *dieffenbachiae* in Florida. *Plant Disease*. 1985; 69: 170-173.
20. Chase AR, Stall RE, Hodge NC, Jones JB. Characterisation of *Xanthomonas campestris* strains from aroids using physiological, pathological, and fatty acid analyses. *Phytopathology*. 1992; 82: 754-759.
21. Vauterin L, Hoste B, Kersters K, Swings J. Reclassification of *Xanthomonas*. *International Journal of Systematic Bacteriology*. 1995; 45: 472-489.

## APPENDICES

Appendix 1: Cropping information obtained from cocoyam farm surveyed in Kaduna State

| State  | LGA        | Location    | Field | Farm size (ha) | Cropping pattern | Associated crop | Age of crop (months) | Source of corm | Topography | GPS                             |
|--------|------------|-------------|-------|----------------|------------------|-----------------|----------------------|----------------|------------|---------------------------------|
| Kaduna | Kudan      | Jaja        | 1     | 0.40           | Mixed cropping   | Okra            | 4                    | SEN            | lowland    | N 11°14.065; E 007°38.663; 660m |
|        |            |             | 2     | 0.35           | Sole cropping    |                 | 4                    | SEN            | lowland    | N 11°14.030; E 007°37.203; 680m |
|        |            |             | 3     | 0.40           | Sole cropping    |                 | 3                    | LS             | lowland    | N 11°14.075; E 007°36.257; 630m |
|        |            | Hunkuyi     | 1     | 0.20           | Mixed cropping   | Maize           | 4                    | SEN            | lowland    | N 11°16.786; E 007°39.430; 630m |
|        |            |             | 2     | 0.10           | Mixed cropping   | Maize           | 5                    | SEN            | lowland    | N 11°16.715; E 007°40.222; 595m |
|        |            |             | 3     | 0.32           | Sole cropping    |                 | 4                    | SEN            | upland     | N 11°16.516; E 007°39.325; 650m |
|        |            | Musawa      | 1     | 0.14           | Mixed cropping   | Sugarcane       | 5                    | LS             | lowland    | N 11°13.477; E 007°40.409; 658m |
|        |            |             | 2     | 0.30           | Sole cropping    |                 | 5                    | LS             | lowland    | N 11°13.426; E 007°41.209; 662m |
|        |            |             | 3     | 0.12           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°13.417; E 007°40.312; 640m |
|        | Sabon Gari | Tudun Sarki | 1     | 0.4            | Mixed cropping   | Maize           | 4                    | LS             | lowland    | N 11°07.236; E 007°54.307; 625m |
|        |            |             | 2     | 0.35           | Mixed cropping   | Maize           | 3                    | LS             | upland     | N 11°07.255; E 007°54.106; 630m |
|        |            |             | 3     | 0.12           | Sole cropping    |                 | 4                    | MM             | upland     | N 11°07.245; E 007°54.113; 659m |
|        |            | Bomo        | 1     | 0.09           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°09.012; E 007°41.503; 630m |
|        |            |             | 2     | 0.10           | Mixed cropping   | Maize           | 3                    | LS             | upland     | N 11°09.032; E 007°41.303; 630m |
|        |            |             | 3     | 0.20           | Mixed cropping   | Maize           | 3                    | LS             | upland     | N 11°09.119; E 007°41.256; 630m |
|        |            | Yan Sarki   | 1     | 0.12           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°07.263; E 007°35.459; 702m |
|        |            |             | 2     | 0.08           | Mixed cropping   | Okra            | 5                    | LS             | upland     | N 11°07.269; E 007°35.692; 702m |
|        |            |             | 3     | 0.13           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°07.450; E 007°35.205; 695m |
|        | Giwa       | Shika       | 1     | 0.12           | Mixed cropping   | Okra            | 5                    | LS             | lowland    | N 11°11.725; E 007°34.532; 670m |
|        |            |             | 2     | 0.03           | Sole cropping    |                 | 5                    | LS             | Lowland    | N 11°11.458; E 007°33.139; 680m |
|        |            |             | 3     | 0.10           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°11.525; E 007°34.502; 640m |
|        |            | Giwa        | 1     | 0.15           | Mixed cropping   | sugarcane       | 3                    | LS             | lowland    | N 11°15.239; E 007°27.514; 630m |
|        |            |             | 2     | 0.15           | Sole cropping    |                 | 2                    | LS             | lowland    | N 11°15.489; E 007°27.704; 660m |
|        |            |             | 3     | 0.08           | Sole cropping    |                 | 2                    | MM             | lowland    | N 11°15.200; E 007°27.586; 670m |
|        |            | Yakawada    | 1     | 0.35           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°15.997; E 007°23.788; 638m |
|        |            |             | 2     | 0.2            | Mixed cropping   | Maize           | 4                    | LS             | upland     | N 11°16.005; E 007°23.715; 589m |
|        |            |             | 3     | 0.28           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°16.035; E 007°23.658; 625m |

LGA = Local Government Area; LS = Locally sourced; SEN = South East Nigeria; MM = Makarfi Market

Appendix 2: Cropping information obtained from cocoyam farms surveyed Kano State

| State | LGA    | Location       | Field | Farm size (ha) | Cropping pattern | Associated crop | Age of crop (months) | Source of corm | Topography | GPS                              |
|-------|--------|----------------|-------|----------------|------------------|-----------------|----------------------|----------------|------------|----------------------------------|
| Kano  | Kiru   | Yelwan-paki    | 1     | 0.25           | Sole cropping    |                 | 5                    | LS             | upland     | N 11°30.016, E 008°09.627; 586 m |
|       |        |                | 2     | 0.10           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°30.008, E 008°09.661; 582 m |
|       |        |                | 3     | 0.07           | Sole cropping    |                 | 2                    | LS             | upland     | N 11°30.001, E 008°09.667; 581 m |
|       |        | Dankargo       | 1     | 0.36           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°30.587; E 008°10.418; 584 m |
|       |        |                | 2     | 0.25           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°30.570; E 008°10.410; 586 m |
|       |        |                | 3     | 0.32           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°30.400; E 008°10.545; 586 m |
|       |        | Dangora        | 1     | 0.18           | Mixed cropping   | maize           | 4                    | LS             | upland     | N 11°32.473; E 008°09.822; 575 m |
|       |        |                | 2     | 0.20           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°32.203; E 008°09.830; 572 m |
|       |        |                | 3     | 0.30           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°32.403; E 008°09.836; 573 m |
|       | Bebeji | Bebeji         | 1     | 0.08           | Sole cropping    |                 | 5                    | LS             | upland     | N 11°37.991; E 008°19.390; 495 m |
|       |        |                | 2     | 0.11           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°37.982; E 008°19.415; 405 m |
|       |        |                | 3     | 0.10           | Sole cropping    |                 | 5                    | LS             | upland     | N 11°38.001; E 008°19.340; 500 m |
|       |        | Gwarmai        | 1     | 0.28           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°32.054; E 008°15.365; 576 m |
|       |        |                | 2     | 0.16           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°32.039; E 008°15.382; 575 m |
|       |        |                | 3     | 0.25           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°32.045; E 008°15.376; 575 m |
|       |        | Ungwan-dankali | 1     | 0.18           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°30.217; E 008°15.983; 485 m |
|       |        |                | 2     | 0.14           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°30.220; E 008°15.996; 476 m |
|       |        |                | 3     | 0.26           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°30.317; E 008°15.973; 483 m |
|       | Rogo   | Zarewa         | 1     | 0.24           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°23.895; E 007°48.895; 704m  |
|       |        |                | 2     | 0.02           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°24.156; E 007°48.943; 704 m |
|       |        |                | 3     | 0.29           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°24.602; E 007°49.632; 696 m |
|       |        | Fulatan        | 1     | 0.30           | Mixed cropping   | sugarcane       | 4                    | LS             | lowland    | N 11°23.220; E 007°51.973; 676 m |
|       |        |                | 2     | 0.26           | Sole cropping    |                 | 3                    | LS             | lowland    | N 11°23.242; E 007°51.953; 675 m |
|       |        |                | 3     | 0.12           | Sole cropping    |                 | 4                    | LS             | upland     | N 11°23.515; E 007°51.490; 689 m |
|       |        | Babbarika      | 1     | 0.32           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°24.941; E 007°50.211; 693 m |
|       |        |                | 2     | 0.10           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°24.906; E 007°50.227; 695 m |
|       |        |                | 3     | 0.18           | Sole cropping    |                 | 3                    | LS             | upland     | N 11°25.052; E 007°50.015; 695 m |

LGA = Local Government Area; LS = Locally sourced; SEN = South East Nigeria; MM = Makarfi Market