

Effect of Artificial Detasseling and Defoliation on Maize Seed Production

ABSTRACT

The objective of this work was to evaluate the effect of artificial detasseling and defoliation on the production of Pioneer 30F90 simple hybrid corn (*Zea Mays* L.) seeds. The experiment was set up in a randomized complete block design, consisting of seven treatments (Control; detasseling; detasseling + defoliation of the top leaf; detasseling + defoliation of the two upper leaves; detasseling + defoliation of the three upper leaves; detasseling + defoliation of the upper four leaves; detasseling + defoliation of the upper five leaves), with 4 replicates. The plots were composed of 6 spaced rows of 0.90m by 6.00m in length. At the time of the issue of 50% of the tassel were carried out the treatments. For the evaluations the two central lines of each plot were collected, eliminating 1.00 m from each end, totaling a useful area of 7.2 m². At harvest, the crop presented a final stand of 5 plants m² (50000 ha⁻¹). According to the results obtained, it was concluded that the detasseling positively influenced the production components, due to the decrease of the auto-shading. The higher the level of defoliation, the lower the active photosynthetic area of the plant, and consequently the lower the productivity. The most affected production components in defoliation are grain numbers in the row, grain numbers in the ear and weight of 1000 grains. detasseling and defoliation do not interfere with the germination of corn seeds.

Keywords: *Zea mays* L.; light interception; tassel; grain yield

1. INTRODUCTION

Maize (*Zea mays* L.) is one of the most widespread agricultural products in the world. This cereal is an input for the elaboration of products directed to human food, animal feed, fuel production and industrial uses [1]. To meet this growing market demand, there is a large investment in research for hybrid breeding and seed production, with high potential for production. Generally, grain yield is the decisive factor in the choice of the hybrid, since it is directly related to the adaptation to the production environment [2].

Biotic and abiotic factors can cause stress in plants, resulting in reduced productivity. As a biotic factor, one can cite the attack of pests and diseases that cause great changes in the physiological processes of plants. However, the factors that are most related to the reduction of maize crop productivity are the abiotic factors related to the climate (soil water availability, air temperature, relative air humidity and solar irradiation) [3]. Among these factors, it is cited

29 light deficiency or lack of leaf area in the development periods, corresponding to the period
30 of grain filling [4]. There are two characteristics in the corn plant that diminish the efficiency
31 potential of the leaves. The most limiting is the habit of growth, which provides high-shading
32 of the lower leaves. The other is the presence of the tassel, which remains inactive shortly
33 after fertilization, but shade about 19% of the plant, depending on the cultivar [5].
34

35 Detasseling in corn is the most commonly used practice in the control of crosses to obtain
36 hybrids, which may favor or impair the plant, depending on the method used. For example,
37 the pure and simple removal of the tassel, which is a strong drain, may favor the plant, since
38 it reduces competition for photoassimilates; and the removal of the cartridge may result in
39 damage to the plant, because it usually occurs a loss of 4 to 5 upper leaves [6,7], which
40 makes the plant less efficient in grain production, because leaf extension above the ear is
41 the most physiologically active [8]. Moreover, the intensity and period of defoliation in the
42 crops are also important factors in grain yield, due to the relationship between leaf area and
43 canopy transpiration [9]. Corn grain filling occurs mainly from photoassimilates produced
44 post anthesis [10].
45

46 A number of authors [11,12,13] have reported an increase in grain yield when the tassel is
47 removed shortly before the maydis stigma emission. However, studies show that defoliation
48 of the corn plant [14, 15, 16, 8, 17] leads to significant losses in productivity. It is also verified
49 that the detasseling in different hybrids show different behaviors in relation to the production
50 components [18,19,17], thus demonstrating the need for research on other materials, such
51 as 30F90 corn not yet studied.
52

53 Considering the above, we can understand that the variation of the production of a crop
54 depends on the genotype of the material, which establishes the level of maximum potential
55 of production; of the production environment, which imposes limits on the development of
56 the expression of crop potential; and the physiological responses of plants to the challenges
57 posed by the environment [3].
58

59 The objective of this work was to evaluate the influence of the detasseling and the simulation
60 of several defoliation levels in the production of 30F90 maize seeds.
61

62 **2. MATERIAL AND METHODS**

63 The experiment was conducted in the field in the municipality of Quatro Pontes in the State
64 of Paraná, Brazil, in an area called rural allotment 91. The climate is of the Subtropical Moist
65 Mesothermal type - Cfa, of hot summers, with rains more concentrated in these months, and
66 without defined dry season. The average temperature in the coldest months is below 18 °C
67 and the average temperature in the warmer months above 22 °C, according to Köppen
68 classification [20]. The precipitation values for the period of the survey, which comprises the
69 spring / summer harvest had mean monthly minimum of 125 and maximum of 330 mm of
70 rainfall.
71

72
73
74 The Pioneer 30F90 simple hybrid corn was used in the experiment. The experimental design
75 was a randomized block design with 7 treatments and four replications, with the following
76 treatments: 1 (Control); 2 (detasseling); 3 (detasseling + defoliation of the top leaf); 4
77 (detasseling + defoliation of the two upper leaves); 5 (detasseling + defoliation of the three
78 upper leaves); 6 (detasseling + defoliation of the upper four leaves) and 7 (detasseling +
79 defoliation of the upper five leaves).
80

81 Seeding was carried out in September, in the no tillage system, having as a previous crop
82 the consortium of black oat and forage turnip, previously desiccated. Each experimental plot
83 consisted of 6 sowing lines of 6.00 m in length, spaced 0.90 m, where the two central lines
84 were harvested, discounting 1.00 m from both ends, totaling a useful area of 7.20 m² per
85 plot. In the context, 5 seeds per linear meter were sown, totaling a population of
86 approximately 55,000 ha⁻¹ seeds.

87
88 The soil of the experiment site is called Red Eutrophic Latosol [21], whose chemical
89 characteristics are found in Table 1.

90

91

92 **Table 1. Chemical characteristics of the soil in the period of installation**
93 **of the experiment, in Quatro Pontes - PR**

94

P	K ⁺	Ca ²⁺	Mg ²⁺	Al ³⁺	H+Al
mg dm ⁻³	Cmol dm ⁻³				
10.3	0.32	7.04	2.64	0.20	5.44

95

96

97 Fertilization was carried out according to the soil analysis, following the recommendation of
98 Oliveira [22] for the corn crop.

99

100 During the experimental period, the herbicide atrazine 250 g L⁻¹ (PRIMATOP) was applied
101 10 days after emergence (DAE), for the control of weeds at a dose of 4 L ha⁻¹, and 25 DAE
102 the insecticide Lufenuron (MATH) at the dose of 0.7 L ha⁻¹ for the control of the caterpillar
103 *Spodoptera frugiperda*.

104

105 The corn plants were submitted to the 60 DAE treatments, when 50% of the tassels were
106 exposed, where manual detasseling and defoliation were performed in the four central lines
107 of each plot, leaving the lateral lines intact so that they promoted the release of pollen and
108 consequently fecundation of the lateral lines, which will give rise to the corn seeds. Two days
109 later a pass through was made to certify the efficiency of the treatments.

110

111 The characteristics evaluated at 81 DAE were: Leaf area measured on-the-spot, which is a
112 non-destructive method consisting of the sum of the area of the rectangle plus the area of
113 the triangle of each leaf of the plants. To obtain the average leaf area of the plants, five
114 plants were used per plot; Foliar Area Index (LAI) obtained by multiplying the average leaf
115 area of one plant by five (5 plants m²).

116

117 The harvest of all useful area of the plot was done manually, and the ears were then
118 threshed, with the help of a manual threshing machine. The characteristics of the production
119 components evaluated after harvesting were as follows: Productivity (kg ha⁻¹) estimated with
120 balance aid, after correction of humidity to 13%; length of ear (cm), measured with the aid of
121 a caliper; number of grains in the ear, obtained by multiplying the count of the average
122 number of grains per row by the average number of rows; grains/row, obtained from the
123 sample of five ears within the useful portion of each treatment and repetition; weight of 1000
124 seeds, according to the Seed Analysis Rule [23]. In order to carry out this determination,
125 pure seeds were used, where at random eight replicates of 100 seeds each were taken per
126 experimental plot; stand calculated from the number of plants at the end of each treatment;
127 Soluble solids (^oBrix) measured using a portable refractometer on dry thatch. The collection
128 of the thatch was done after the appearance of the black layer (physiological maturation) two

129 between node per plant (between node above the ear and the other just below the ear), of
130 two plants per plot; germination test performed in BOD regulated at 30 °C diurnal and 20 °C
131 nocturnal, remaining 5 days until the first evaluation, and another 4 days until the second
132 evaluation [23]. 100 seeds of each of the four replications of the seven treatments were
133 used, totaling 400 seeds of each treatment.

134
135 The data were submitted to analysis of variance and the means were compared by the
136 Tukey test at 5% probability.

137
138

139 3. RESULTS AND DISCUSSION

140
141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

3.1. Leaf area and leaf area index (LAI)

Analyzing the results obtained for the leaf area and leaf area index (LAI), it can be seen that only the control and detasseling treatments do not differ between them (Table 2). However, in the other treatments there was an increasing reduction in both leaf area and LAI, directly related to the number of leaves removed. Simulation of leaf loss and the results found in this work with 30F90 maize for seed production reflect the reality of commercial crops. According to Dias [17], the manual or mechanized detasseling also causes the removal of leaves from the apex, next to the tassel, which can reduce the productivity and the quality of the seeds. This is because leaf area per plant is one of the main factors determining the strength of the source [24], and its reduction affects the photosynthetic activity and consequently the production of photosimilates by the plant [6], mainly because these upper leaves are considered photosynthetically more efficient [8, 25].

Table 2. Leaf area and leaf area index (LAI) in function of the artificial defoliation of Pioneer 30F90 maize plants in Quatro Pontes - PR

Treatments	Leaf area (m ²)	(%)	LAI	(%)
C	0.90 a	100.00	4.50 a	100.00
D	0.89 a	99.58	4.48 a	99.58
D+1DF	0.88 b	97.69	4.40 b	97.69
D+2DF	0.84 c	93.78	4.22 c	93.78
D+3DF	0.78 d	86.79	3.91 d	86.79
D+4DF	0.72 e	80.22	3.61 e	80.22
D+5DF	0.65 f	72.45	3.26 f	72.45

159

160

161

162

163

164

165

166

167

168

169

Averages followed by the same letters in the column do not differ from each other by the tukey test at 5% probability.

C = control; D= detasseling; D+1DF detasseling + defoliation of the top leaf; D+2DF = detasseling + defoliation of the two upper leaves; D+3DF detasseling + defoliation of the three upper leaves; D+4DF = detasseling + defoliation of the upper four leaves; D+5DF = detasseling + defoliation of the upper five leaves.

3.2. Productivity

170 The yield values did not show a significant difference between control treatments,
 171 detasseling, and detasseling + defoliation of a top leaf. However, when only the detasseling
 172 was performed, there was an increase in productivity of 4.97% in relation to the control
 173 (Table 3). Similarly, Hunter et al. [26] observed an average increase of 6.9% in grain
 174 production, when only the tassels were removed. The positive effects of the detasseling on
 175 grain yield occur mainly due to the loss of apical dominance and reduction of shading in the
 176 upper portions of the plants, as demonstrated by Duncan et al. [27]. Emygdio et al. [2]
 177 emphasize that productivity is a decisive factor in the choice of hybrids.
 178

179 On the other hand, according to the results found in this experiment with 30F90 maize, it is
 180 possible to notice the decrease in productivity as the levels of defoliation increase,
 181 demonstrating similar behavior to those obtained by other authors, such as Alvim et al. [8]
 182 and Vaz et al. [28]. The results obtained are in accordance with Souza and Barbosa [3], who
 183 cite as a variation of the production of a crop, the physiological responses of the plants to the
 184 challenges imposed by the environment. In the defoliation the plant is less efficient in the
 185 production of grains, because the leaf extension above the spike is the most physiologically
 186 active [8].
 187
 188

189 **Table 3 - Mean productivity values for the Pioneer 30F90 maize crop, in**
 190 **Quatro Pontes – PR**
 191

Treatments	Productivity (Kg ha ⁻¹)	(%)
D	10849 a	104.97
C	10335 ab	100.00
D+1DF	10302 ab	90.68
D+2DF	9800 bc	94.81
D+3DF	9511 bc	92.02
D+4DF	9041 cd	87.48
D+5DF	8602 d	83.23

192 **Averages followed by the same letters in the column do not differ from**
 193 **each other by the tukey test at 5% probability.**

194 **C = control; D= detasseling; D+1DF detasseling + defoliation of the top**
 195 **leaf; D+2DF = detasseling + defoliation of the two upper leaves; D+3DF**
 196 **detasseling + defoliation of the three upper leaves; D+4DF =**
 197 **detasseling + defoliation of the upper four leaves; D+5DF = detasseling**
 198 **+ defoliation of the upper five leaves.**
 199

200
 201 **3.3. Number of grains per row and total number of grains per tassel**
 202

203 The average results of grains per row and total number of grains per tassel are given in
 204 Table 4.
 205

206
 207 **Table 4. Mean values of grains per row and number of grains per ear of**
 208 **corn Pioneer 30F90 in Quatro Pontes - PR**

Treatments	Grains/Row	(%)	Grains/Tassel	(%)
D	41.69 a	103.93	667.04 a	103.93
D+1DF	40.48 ab	100.93	647.80 ab	100.93
C	40.11 ab	100.00	641.85 ab	100.00
D+2DF	39.15 bc	97.60	626.44 bc	97.60
D+3DF	37.69 c	93.95	603.04 c	93.95
D+4DF	34.12 d	85.07	546.00 d	85.07
D+5DF	33.29 d	83.00	532.76 d	83.00

210 **Averages followed by the same letters in the column do not differ from**
 211 **each other by the tukey test at 5% probability.**

212 **C = control; D= detasseling; D+1DF detasseling + defoliation of the top**
 213 **leaf; D+2DF = detasseling + defoliation of the two upper leaves; D+3DF**
 214 **detasseling + defoliation of the three upper leaves; D+4DF =**
 215 **detasseling + defoliation of the upper four leaves; D+5DF = detasseling**
 216 **+ defoliation of the upper five leaves.**

217
218

219 In the use of isolated detasseling, there is an increase in the number of grains per row and
 220 consequently the number of grains per tassel. The accumulated photoassimilates reserve
 221 during the vegetative phase in the stem and its translocation for the filling of grains in the
 222 treatment with defoliation allowed the grain to finish its formation [8], but the withdrawal of
 223 leaves reflects in losses, especially of grain per tassel, mass of 1000 seeds and
 224 consequently of productivity, which can be explained by the fact that of the total
 225 carbohydrates accumulated in the corn grains, 50% are produced by the leaves located in
 226 the upper third of the plant, 30% represents the contribution of the leaves of the middle third
 227 and the rest of the leaves distributed in the most basal part of the stem [29].

228
229

230 3.4. Ear length

231
232

232 The average results obtained for the ear length are given in Table 5. In the present work, it is
 233 evident that the length of the ear was affected directly by the reduction of the leaf area, in the
 234 same way that the number of grains in the tassel was reduced (Table 4), which significantly
 235 affects productivity, as previously noted (table 3). Pereira et al. [30] verified that the
 236 defoliation was the major factor influencing the length of the ear, corroborates with the direct
 237 relation between the foliar apparatus and the production/growth of the ear. Still, the authors
 238 conclude that in the components of maize production, defoliation causes large losses, and
 239 there is no means of compensation for maize.

240
241

242 **Table 5. Mean values of corn ear length Pioneer 30F90, in Quatro**
 243 **Pontes - PR**

244

Treatments	length of ear (cm)	(%)
D	18.45 a	104.24

D+1DF	17.74 ab	100.24
C	17.70 ab	100.00
D+2DF	16.60 bc	93.77
D+3DF	15.99 cd	90.33
D+4DF	15.22 de	85.99
D+5DF	14.34 e	85.99

245 **Averages followed by the same letters in the column do not differ from**
 246 **each other by the tukey test at 5% probability.**

247 **C = control; D= detasseling; D+1DF detasseling + defoliation of the top**
 248 **leaf; D+2DF = detasseling + defoliation of the two upper leaves; D+3DF**
 249 **detasseling + defoliation of the three upper leaves; D+4DF =**
 250 **detasseling + defoliation of the upper four leaves; D+5DF = detasseling**
 251 **+ defoliation of the upper five leaves.**

252

253

254 **3.5. 1000 seed mass**

255

256 The mass production component of 1000 seeds was affected by the leaf reduction, and
 257 there was no differentiation of the treatment detasseling, with the control, but of the other
 258 treatments (Table 6). Similar results of reduction in the mass of 1000 seeds with the
 259 increase of defoliation were also observed by Dias [17] and Vaz et al [28]. The decrease of
 260 the leaf area influences the photosynthetic rate, changes the plant metabolism, reducing the
 261 content of sucrose and reducing sugars, as well as the starch content in the seeds, and
 262 consequently affects the weight of 1000 seeds [4].

263

264

265 **Table 6. Mean values of the masses of 1000 seeds of maize Pioneer**
 266 **30F90, in Quatro Pontes - PR**

267

Treatments	1000 seed mass (g)	(%)
D	362.83 a	101.25
C	358.35 ab	100.00
D+1DF	355.44 bc	99.18
D+3DF	351.60 cd	98.11
D+4DF	351.44 cd	98.07
D+2DF	350.19 cd	97.72
D+5DF	346.90 d	96.81

268 **Averages followed by the same letters in the column do not differ from**
 269 **each other by the tukey test at 5% probability.**

270 **C = control; D= detasseling; D+1DF detasseling + defoliation of the top**
 271 **leaf; D+2DF = detasseling + defoliation of the two upper leaves; D+3DF**
 272 **detasseling + defoliation of the three upper leaves; D+4DF =**
 273 **detasseling + defoliation of the upper four leaves; D+5DF = detasseling**
 274 **+ defoliation of the upper five leaves.**

275

276

277 **3.6. Stand**

278
279
280
281
282
283
284
285
286
287
288
289

There was no difference between treatments for stands (Table 7). The evaluation was performed at the time of harvest, counting the plants to confirm that the results were not affected by the stand. Alvim et al. [8] also found that the final stand of corn plants was not significantly altered by defoliation and reported if they were made in the R₂ stage did not predispose the plants to tipping or stem breakage, probably due to the good resistance of the stem at this.

Table 7. Mean values of stand of the Pioneer 30F90 maize, in Quadro Pontes – PR

Treatments	Stand	(%)
D	36.25 a	100.69
D+1DF	36.25 a	100.69
C	36.00 a	100.00
D+2DF	35.75 a	99.30
D+5DF	35.75 a	99.30
D+4DF	35.50 a	97.93
D+3DF	35.00 a	97.55

290 **Averages followed by the same letters in the column do not differ from**
291 **each other by the tukey test at 5% probability.**

292 **C = control; D= detasseling; D+1DF detasseling + defoliation of the top**
293 **leaf; D+2DF = detasseling + defoliation of the two upper leaves; D+3DF**
294 **detasseling + defoliation of the three upper leaves; D+4DF =**
295 **detasseling + defoliation of the upper four leaves; D+5DF = detasseling**
296 **+ defoliation of the upper five leaves.**

297
298
299
300

3.7. Soluble Solids (°Brix)

301 The average results for soluble solids (°Brix) are shown in table 8. The removal of only the
302 tassel, presented a relative brix greater than the others, except for the treatment detasseling
303 + of the two upper leaves, to which it did not present statistical difference. This higher
304 content of °Brix is due to the decrease of the interception of light due to the removal of the
305 tassel and consequently the plant "needed" to remove smaller amounts of sugars from the
306 stem for the filling of the grains, agreed with the results found by Vasconcelos et al. [31],
307 where they reported that there was an increase in the residual amounts found in the stalks,
308 sabugos, and ears, taking into account detasseling practices used.

309
310
311
312
313

Table 8. Mean values of Soluble Solids (°Brix) of Pioneer 30F90 maize in Quatro Pontes - PR

Treatments	Soluble Solids (°Brix)	(%)
D	11.42 a	109.07

D+2DF	10.65 ab	101.67
C	10.47 b	100.00
D+1DF	10.47 b	100.00
D+3DF	10.25 b	97.85
D+5DF	10.12 b	96.66
D+4DF	9.97 b	95.25

314 **Averages followed by the same letters in the column do not differ from**
315 **each other by the tukey test at 5% probability.**

316 **C = control; D= detasseling; D+1DF detasseling + defoliation of the top**
317 **leaf; D+2DF = detasseling + defoliation of the two upper leaves; D+3DF**
318 **detasseling + defoliation of the three upper leaves; D+4DF =**
319 **detasseling + defoliation of the upper four leaves; D+5DF = detasseling**
320 **+ defoliation of the upper five leaves.**

321
322

323 **3.8. Germination test**

324

325 In the production of corn seeds, besides the productivity in the formation of new hybrids [2],
326 the physiological quality of the seeds (germination and vigor) is a very important factor. The
327 benefits of a high quality seed include rapid and uniform germination, with seedlings being
328 able to withstand environmental stresses and uniformity in emergence [32].

329

330 According to the results of the statistical analysis for the germination factor, it was observed
331 that there was no statistical difference between the treatments (Table 9). These results are
332 in agreement with some authors who, when studying the effect of defoliation on maize plants
333 on the physiological quality of the seeds, did not detect influence on germination or vigor [33,
334 34], which is more related to maturity at harvest than to defoliation [4].

335

336

337 **Table 9. Mean values of germination of Pioneer 30F90 maize in Quatro**
338 **Pontes - PR**

339

Treatments	Germination (%)
D+1DF	98.50 a
D+4DF	98.25 a
D	98.00 a
C	98.00 a
D+2DF	98.00 a
D+3DF	98.00 a
D+5DF	97.50 a

340 **Averages followed by the same letters in the column do not differ from**
341 **each other by the tukey test at 5% probability.**

342 **C = control; D= detasseling; D+1DF detasseling + defoliation of the top**
343 **leaf; D+2DF = detasseling + defoliation of the two upper leaves; D+3DF**
344 **detasseling + defoliation of the three upper leaves; D+4DF =**

345 **detasseling + defoliation of the upper four leaves; D+5DF = detasseling**
346 **+ defoliation of the upper five leaves.**

347
348

349 **4. CONCLUSION**

350

351 The removal of the tassel positively favors the crop with respect to the components of
352 production, due to the reduction of problems with auto-shading.

353

354 Defoliation during detasseling negatively influences crop productivity due to the fall in
355 number of grains in the row, number of grains in the ear and weight of 1000 grains, caused
356 by the decrease of the active photosynthetic area of the plant.

357

358 Detasseling alone or associated with defoliation does not interfere with the germination of
359 corn seeds.

360

361

362 **REFERENCES**

363

364 1. Garcia JC, Duarte JO. Corn production and consumption. In: Borém A, Rios SA, editors.
365 Biofortified corn. Visconde do Rio Branco: Suprema; 2011;23-44.

366

367 2. Emygdio BM, Ignaczak JC, Cargnelutti Filho A. Yield potential of single, three-way and
368 double cross commercial hybrids maize. *Revista Brasileira de Milho de Sorgo*. 2007;6:95-
369 103. DOI: <https://doi.org/10.18512/1980-6477/rbms.v6n1p95-103>

370

371 3. Souza GM, Barbosa AM. Corn stress factors are diverse and require constant monitoring.
372 *Agricultural View*. 2015;13(1):30-34. Available:

373

374 https://www.esalq.usp.br/visaoagricola/sites/default/files/VA_13_Fisiologiaartigo3.pdf.

375

376 4. Büll LT. Mineral nutrition of maize. In: Büll LT, CANTARELLA H, editors. *Corn crop:*
377 *Factors affecting productivity*. Piracicaba: Potafós; 1993;63-145.

378

379 5. Magalhães PC, Durães FOM, Carneiro NP, Paiva E. Physiology of corn. Sete Lagoas:
380 EMBRAPA-CNPMS (Circular Técnica, n. 22); 2002; 23. Available:
381 <http://docsagencia.cnptia.embrapa.br/milho/circul22.pdf>. Accessed: 25 November 2018.

382

383 6. Magalhães PC, Durães FOM, Oliveira AC, Gomes EEG. Effects of different detasseling
384 practices on maize yield. *Scientia Agricola*. 1999;56(1): 77-82. DOI:
385 <https://dx.doi.org/10.1590/S0103-90161999000100012>

386

387 7. Martin TN, Tomazella AL, Cícero SM, Dourado Neto D, Favarin JL, Vieira Júnior PA.
388 Important questions about production of corn seeds - first part. *Revista da FZVA*.
389 2007;14(1):119-138.

390

391 8. Alvim KRT, Britol CH, Brandão AM, Gomes LS, Lopes MTG. Quantification of leaf area
392 and defoliation effect in corn crop components. *Ciência Rural*. 2010;40(5):1017-1022.

393

394 9. Shimada S, Kokubun M, Shibata H, Matsui S. Effect of water supply and defoliation on
395 photosynthesis, transpiration and yield of soybean. *Japanese Journal of Crop Science*.
396 1992;61(2):264-270. DOI: <https://doi.org/10.1626/jcs.61.264>

397

398 10. Borrás L, Slafer GA, Otegui ME. Seed dry weight response to source-sink manipulations
399 in wheat, maize and soybean: a quantitative reappraisal. *Field Crop Research*. 2004;86:131-
400 146. DOI: <https://doi.org/10.1016/j.fcr.2003.08.002>

401

402 11. Hunter RB, Daynard TB, Hume DJ, Tanner JW, Curtis JD, Kannenberg LW. Effect of
403 tassel removal on grain yield of corn (*Zea mays* L.). *Crop Science*. 1969;9:405-406.

404

405 12. Mostert AJ, Marais JN. The effects of detasselling on the yield of irrigated maize. *Crop*
406 *Production/Gewasproduksie*. 1982;11:163-167.

407

408 13. Craig WF. Production of hybrid corn seed. In: Sprague GF, editors. *Corn and*
409 *improvement*. 3th ed. Madison: American Society of Agronomy; 1988;680-693.

410

411 14. Hanway JJ. Defoliation effects on different corn (*Zea mays* L.) hybrids as influenced by
412 plant population and stay of development. *Agronomy Journal*. 1969;61:534-538.

413

414 15. Egharevba PN, Harrocks RD, Zuber MS. Dry matter accumulation in maize response to
415 defoliation. *Agronomy Journal*. 1976;68:43-49.

416

417 16. Pereira FH. Influence of detasseling and defoliation on corn seed production and quality.
418 *Dissertação (mestrado em agronomia): Universidade Estadual Paulista, Faculdade de*
419 *Ciências Agrárias e Veterinárias*; 2007;38.

420

421 17. Dias CRP. Influence of detasseling and elimination of dominated plants on seed quality
422 and yield of corn. *Dissertação (Mestrado em Agronomia): Universidade de*
423 *Brasília/Faculdade de Agronomia e Medicina Veterinária*; 2015;32.

424

425 18. Almeida IPC; Silva PSL; Negreiros MZ; Barbosa, Z. Baby corn, green ear and grain yield
426 of corn cultivars. *Horticultura Brasileira*. 2005; 23:960-964.

427

428 19. Moreira JN, Silva PSL, Silva KMB, DombroskiJLD, Castro RS. Effect of detasseling on
429 baby corn, green ear and grain yield of two maize hybrids. *Horticultura Brasileira*.
430 2010;28(4):406-411.

431

432 20. Álvares CA, Stape, JL, Sentelhas PC, Gonçalves JLM, Sparovek, G. Köppen's climate
433 classification map for Brazil. *Meteorologische Zeitschrift*. 2014;22(6):711-728. DOI:
434 <https://doi.org/10.1127/0941-2948/2013/0507>

435

436 21. EMBRAPA. Brazilian system of soil classification. 5th ed. Rio de Janeiro: EMBRAPA;
437 2006;306.

438

439 22. Oliveira EL. Suggestion of fertilization and liming for crops of economic interest in the
440 state of Paraná. Londrina: IAPAR (Circular Técnica, n. 128); 2003;22-23. Available:
441 http://www.iapar.br/arquivos/File/zip_pdf/ct_128.pdf. Accessed: 18 November 2017.

442

443

444

445 23. BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. Rules for seed analysis.
446 Brasília: Mapa/ACS; 2009;399. Available: [http://www.agricultura.gov.br/assuntos/insumos-](http://www.agricultura.gov.br/assuntos/insumos-agropecuarios/arquivos-publicacoes-insumos/2946_regras_analise__sementes.pdf)
447 [agropecuarios/arquivos-publicacoes-insumos/2946_regras_analise__sementes.pdf](http://www.agricultura.gov.br/assuntos/insumos-agropecuarios/arquivos-publicacoes-insumos/2946_regras_analise__sementes.pdf).

448 Accessed: 30 March 2018.

449

- 450 24. Magalhães PC, Jones R. Assimilate enhancement on the carbohydrates and nitrogen
451 content in maize. *Pesquisa Agropecuária Brasileira*. 1990;25:1755-1761.
452
- 453 25. Viecelli CA, Fillwock JM, Suzin V. Efecto de la defoliación de las plantas en la
454 productividad de maíz. *Revista Brasileira de Tecnologia Aplicada nas Ciências Agrárias*.
455 2011;4(3):179-190.
456
- 457 26. Hunter RB, Mortimore CG, Kannenberg LW. Inbred maize performance following tassel
458 and leaf removal. *Agronomy Journal*. 1973;65:471-472.
459
- 460 27. Duncan WG, Williams WA, Loomis RS. Tassels and the productivity of maize. *Crop*
461 *Science*. 1967;7: 37-39.
462
- 463 28. Vaz PFT, Simonetti APMM, Montiel CB. The effect of defoliation of maize plants on
464 productive parameters. *Acta Iguazu*. 2016;5(2):94-101.
465
- 466 29. Fornasieri-Filho D. A. The corn crop. Jaboticabal: FUNEP; 1992;273.
467
- 468 30. Pereira MJR, Bonan ECB, Garcia A, Vasconcelos RL, Giacomo KS, Lima MF.
469 Morphoagronomic characteristics of maize in response to different levels of
470 defoliation. *Revista Ceres*. 2012;59(2):200-205. DOI: <https://dx.doi.org/10.1590/S0034-737X2012000200008>
471
472
- 473 31. Vasconcelos CA, Magalhães PC, Durães FOM, Fernandes FT. Detasseling practices on
474 tropical maize and effect on mineral nutrition and nutritional efficiency. *Pesquisa*
475 *Agropecuária Brasileira*. 1995;30:353-358.
476
- 477 32. Carvalho NM, Nakagawa J. *Seeds: science, technology and production*. 2.ed. Campinas:
478 Fundação Cargill. 2000;565.
479
- 480 33. Wilhelm WW, Johnson BE, Schepers JS. Yields, quality, and nitrogen use of inbred
481 corn with varying numbers of leaves removed during detasseling. *Crop Science*.
482 1995;35:209–212.
483
- 484 34. Komatuda AS, Santos CM, Santana DG, Souza MA, Brito CH. Effects detasseling
485 methodologies on yield and quality of hybrid maize seed. *Revista brasileira de milho e sorgo*.
486 2006;5(3):359-368.
487