Original Research Article PREVALENCE AND CORRELATES OF POSTPARTUM DEPRESSION IN OSOGBO,

3 NIGERIA

1 2

4 ABSTRACT

- 5 Background: Postpartum depression is the most prevalent postpartum mental health problem. It is
- 6 associated with subsequent low adherence to child healthcare. In severe depression, especially in the
- 7 presence of psychotic symptoms, there is a substantial risk of suicide and of infanticide.
- 8 AIM: This study aimed to assess the prevalence, pattern and correlates of postpartum depression among
- 9 women attending postnatal and infant welfare clinics at a teaching hospital in Nigeria.
- 10 **Study Design**: This was a cross-sectional study.
- Place and Duration of Study: This study was conducted at LTH, Osogbo Nigeria between September
 and November 2015.
- 13 Methodology: Data were obtained from 220 consenting postpartum women using the Edinburgh
- 14 Postnatal Depression Scale (EPDS) and a socio-demographic questionnaire. All respondents found to be
- 15 EPDS positive for depression as well as 10% of those negative for depression were further assessed with
- 16 the MINI International Neuropsychiatric Interview (MINI) (depression subscale). Data were analyzed using
- 17 the Statistical Package for Social Sciences (SPSS) version 21. The level of statistical significance was set
- 18 at a p-value of less than 0.05.
- 19 Results: Prevalence of postpartum depression was 9.5% using the MINI. Postpartum depression was

significantly associated with age (χ 2=4.767, p=0.035), suicidal ideation (χ 2=17.292, p < 0.01), support

- from respondent's partner during pregnancy (χ 2= 6.593, p= 0.010), support from partner's relatives (χ 2=
- 22 4.403, p=0.036) and number of children (χ 2=4.247, p=0.039). Unemployed mothers had significantly
- 23 higher EPDS scores than those who were employed (F=3.020, p< 0.05).</p>
- 24 **Conclusion:** The prevalence of postpartum depression is high. An increased media campaign about 25 postpartum depression and preventive measures is urgently needed. Screening for depression

- 26 throughout the perinatal period is important for early diagnosis and prompt intervention in order to
- 27 improve clinical outcome.
- 28 Key words: Postpartum depression, Osogbo, Prevalence, Correlates, Nigeria
- 29 INTRODUCTION
- 30 Postpartum depression (PPD) is the most prevalent postpartum mental health problem.(1) Postpartum
- 31 depression is a clinically significant depressive episode that begins in the postpartum period, lasts two
- 32 weeks or more and requires medical attention. (2,3) Depression in the postnatal period contributes to
- 33 several problems for the individual, family and society. (1) It has been associated with early breastfeeding
- 34 discontinuation, mother-child bonding impairment, a reduction of positive parenting behaviors(1) and the
- 35 presence of thoughts of harm.(2, 3)
- 36 Maternal depression is also associated with poor adherence to preventive child health, including
- 37 vaccinations.(4) Moreover, impaired interactions between mothers and children due to depression in
- 38 mothers have been associated with long-term impairment in children's cognitive and emotional
- 39 development.(5) In severe depression, especially in the presence of psychotic symptoms, there is a risk
- 40 of suicide and infanticide.(6)
- 41 The prevalence of depression in the postpartum period is estimated to range between 5-20 % (7-9) This
- 42 wide range might be due to cross-cultural variables, reporting style, differences in perception of mental
- 43 health and its stigma, differences in socio-economic environments and biological vulnerability.(8, 9) In
- 44 Nigeria, the prevalence of postpartum depression has been found to be comparable to that of the western
- 45 world (14.6%).(10)
- 46 Factors for development of postpartum depression are psychological distress and stressful life events
- 47 during the previous year,(11) poor relationship with partner,(8) absence of social support,(12, 13)
- 48 previous or current abuse, unemployment, having a low income or unplanned pregnancy.(14) Maternal
- 49 age, parity and education do not seem to be factors associated with postpartum depression.(15)
- 50 There is a need to detect and manage mothers with postpartum depression in order to reduce the 51 negative effects it could have on care of babies. Identification of associated factors is also important in

- 52 order to facilitate adequate preventive measures. However, there is dearth of studies in this part of the
- 53 country and hence the need for this study.
- 54

55 MATERIALS AND METHODS

- 56 Study Location
- 57 The study was conducted at the infant welfare and postnatal clinics of Ladoke Akintola University of
- 58 Technology (LAUTECH) Teaching hospital (LTH) Osogbo, Osun State.
- 59 Study Population
- 60 The study population comprised of women of age group 18-45 years who were in the postpartum period
- 61 attending postnatal and infant welfare clinics of the hospital.
- 62 Inclusion Criteria
- 63 **1.** Subjects aged 18 years to 45 years.
- 64 2. Women within 6 months postpartum.
- 65 **Exclusion Criteria**
- 66 **1.** Women without live birth.
- 67 2. Women with previous history of depression.
- 68 Study Design
- 69 This was a hospital based cross-sectional descriptive survey
- 70 Sample Size Estimation
- 71 The prevalence of postpartum depression in a study by Adewuya et al was 14.6%.(10) This was used to
- 72 calculate the minimum sample size for the study.
- 73 The minimum sample size for the study was calculated using

74	Sample size (n) = <u>Z²pq</u> (Sample size for population > 10,000)(16)
75	d ²
76	Where n = sample size
77	Z = standard normal deviation = 1.96 corresponding to 95% confidence interval
78	p = the estimated proportion of an attribute that is present in the population (i.e known
79	prevalence of the condition being studied) = 14.6% (0.146)
80	q=1.0 – P
81	d= degree of accuracy desired, set at 0.05
82	n= [1.96] ² x0.146x0.854/[0.05] ² =191.6 approximately 192
83	An attrition rate of 10% gives 192 x 10/100 = 19.2
84	<mark>192 +19= 211</mark>
85	However, because the study population is below 10,000, the true sample size (nf) is estimated from the
86	above, as follows:
87	n _f = <u>n</u> .
88	1 + (n) / (N)(16)
89	Where n_f = the desired sample size when population is less than 10,000.
90	n = the desired sample size when the population is more than 10,000
91	N = the estimate of the population size, with the value of 1000, which is the population of
92	postpartum women from the age of 18 to 45 years in 2014 at the postnatal clinic and infant welfare clinic
93	of LAUTECH Teaching Hospital.
94	
95	<u>n_f= <u>211</u> = 174</u>
96	1 + (211)/(1000)
97	
98	n _f = 174
99	The sample size was increased to 220, to increase the power of the study.

101 Sampling Method

102 Women attending postnatal and infant welfare clinics were consecutively approached and those who met the inclusion criteria and gave informed consent were recruited for the study until the predetermined 103 104 sample size was achieved. A removable identification sticker was left on all patients' card until the 105 completion of the study to avoid a repeat selection. A resident doctor in the psychiatry department who 106 speaks and writes in Yoruba and English was recruited as a research assistant in order to help administer 107 questionnaires to those who could not read in Yoruba or English. She was trained in the administration of 108 the questionnaires for over 6 hours in 3 divided sessions each lasting 2 hours on 3 consecutive days prior 109 to data collection. 110 The self-administered questionnaires were filled by all mothers who met the inclusion criteria. For those who were not able to read in Yoruba or English, the research assistant helped to administer the 111

112 questionnaire. She collected the questionnaires, scored the Edinburgh Postnatal Depression Scale 113 (EPDS) and those with a cut-off score of 10 or higher were interviewed with the Mini International

- 114 Neuropsychiatric Interview (M.I.N.I) depression module by the researcher.
- 115
- 116 Instruments

117 Data collection was done using the following instruments:

Socio-demographic Questionnaire: The socio-demographic information of respondents,
 included age, residence, marital status, number of husband's wives, position among husband's wives,
 family settings, family size, sex of index child, sex of previous children, level of education of both
 participant and partner, employment status of respondent and partner's monthly income.

2. Questions on Pregnancy-Related Factors: This aspect of the questionnaire enquired about support
 during pregnancy, mode of delivery, duration of delivery and number of weeks since delivery.

124 3.Edinburgh Postpartum Depression Scale (EPDS): The Edinburgh Postpartum Depression Scale has 125 been used as a screening tool for assessment of depression in a variety of clinical settings.(17) It is a 126 self-administered questionnaire which consists of 10 questions with four response categories scored from 127 0 to 3, whereby the greatest values represent depressed moods.(18) The range of scores for EPDS is from 0 to 30. Mothers who obtain an EPDS total score of 10 or greater are considered to have postpartum 128 129 depression.(19) Since its publication in the 1980s, this scale has been used in a growing number of studies across a variety of patient groups.(10, 17, 20) It has been validated in Nigeria.(19, 20)Sensitivity 130 131 is reported as 75% and specificity as 97%.(20) The reliability of the EPDS has been reported as 0.83 132 using Cronbach's alpha.(20)

133 4. Mini International Neuropsychiatric Interview (M.I.N.I)

134 **Depression module:**

The M.I.N.I is a short structured diagnostic interview developed jointly by psychiatrists and clinicians in the United States and Europe in 1990 for DSM-IV and ICD-10 psychiatric disorders. It has acceptably high validation and reliability scores when compared with other structured diagnostic interview schedule, but can be administered in a much shorter period of time. It can be used by clinicians, after a brief training session. Lay interviewers require more extensive training. Administration time is approximately 15 minutes.

The M.I.N.I is divided into modules identified by letters each corresponding to a diagnostic category. The Major Depressive Episode module has six sections A1 to A6. In this study questions only A1 to A3 were used since the study assesses major depressive episode (current) and women with previous history of depression have been excluded from the study.

145 Ethical Consideration

Approval to undertake the study was obtained from the Ethics and Research Committee of LAUTECH Teaching Hospital to ascertain that the methodology does not contravene international guidelines for research involving human subjects. Informed consent was obtained from all subjects after explaining the aim and objectives of the study to them. Ethical issues like non-disclosure to others, opportunity to

- 150 decline interview at any stage and non-exposure to risk were discussed with each respondent. The
- 151 participants bore no financial burden for the study.
- 152 Number of Authorization of the protocol by the committee is LTH/EC/2015/03/206
- 153 Data Analysis
- 154 All data collected were analyzed using the Statistical Package for Social Sciences (SPSS) software
- 155 (version 21). Results were presented using frequency distribution tables and relevant statistics such as
- 156 percentages, means and standard deviations. Cross tabulations were done to compare the outcome
- 157 variables for postpartum depression. Chi square statistic, student t-test, ANOVA and logistic regression
- 158 were used to evaluate the association between variables. Statistical significance was set at P< 0.05.

159 RESULT

160 Socio-Demographic Characteristics of the Respondents

161 Two hundred and twenty questionnaires were administered to the study group and all the questionnaires

162 were completed, giving a response rate of 100%.

163 Table 1: Socio-Demographic Characteristics of the Respondents (N = 220)

	Frequency (n=220)	Percentage
Age (years)		
≤ 20	2	0.9
20 -29	89	40.4
30 -39	124	56.4
≥40	5	2.3
Mean age 30.12 (± 4.76)		
Marital Status	0.5	
Cohabiting	25	11.4
Married	195	88.6
Marriage/ Cohabitation		
Monogamous	200	90.9
Polygamous	20	9.1
Employed		
Yes	167	75.9
No	53	24.1
level of Education		
No formal education	1	0.5
Primary	11	5.0
		0.0

Secondary	57	25.9
Tertiary	151	68.6
Tribe		
Yoruba	216	98.2
labo	3	1.3
Others Specified (Ishan)	1	0.5
Place of Residence		
Urban	214	97.3
Rural	6	2.7
Religion	1.10	00.0
Christianity	140	63.6
Islam	79	35.9
Traditional	1	0.5
Incomo pottorn		
Income pattern	117	52.2
Income>18000	117	
	103	40.0

The socio-demographic characteristics of the respondents are shown in Table 1. The mean age of the respondents was 30.12 (± 4.76) years. The respondents were mainly urban dwellers of Yoruba ethnic group from monogamous family settings. Women whose ages ranged between 30 and 39 years constituted more than half of the entire respondent population. Christians constituted about two-third of the respondents. The majority of the women had received education beyond the primary school level. Three-quarter of the respondents were employed. More than half of the respondents earned less than the current minimum wage of 18,000 Naira.

¹⁷² Prevalence of Postpartum Depression



175 Figure 1: Screening for Postpartum Depression using EPDS

176 The mean EPDS score was 4.31 (5.34), median was 2.00 while the range was 20.00. The mean value for

177 respondents with score greater or equal to 10 was 13.89 (2.77), median was 14.00 while the range was

- 178 10. Furthermore, the mean score for respondents with EPDS score less than 10 was 4.31 (5.34), median
- 179 was 2.00 while range was 9.00.
- 180 Screening for postpartum using EPDS is as highlighted in figure 1 above. Among the respondents, 39
- 181 (17.7%) were considered depressed (EPDS score \geq 10).
- 182
- 183
- 184
- 185
- 186 Prevalence of Postpartum Depression Using MINI



188 Figure 2: Prevalence of postpartum depression using MINI.

Figure 2 above depicts the prevalence of postpartum depression using MINI. Among the 57 respondents (39 EPDS positive and 18 EPDS negative) on whom MINI were administered, 21 respondents (9.5%) met criteria for the diagnosis of major depressive episode. Sensitivity and specificity were 100% and 90.95% respectively while positive predictive and negative predictive values were 53.84% and 100% respectively. For the calculation of true prevalence, each of the participants in phase 1 represents 1 while each of those who screened negative in phase 2 represents 10 participants (since only 10% of those who screened negative were randomly chosen).

196Table 2: Association of Postpartum Depression with Socio-demographic and Clinical197Characteristics of Respondents

Variables		Postpartum depression		χ2	df	<i>p</i> value	
		Yes n%	No n%				
Age (years)				. = . = #		0.005*	
<30 ≥30		4 (4.4) 17 (13.2)	87 (95.6) 112 (86.8)	4.767"	1	0.035*	
Marital Status		, , ,	· · ·				
Cohabiting	3	3 (12) 18 (9 2)	22 (88.0) 177 (90 8)	0.197#	1	0.715	
Educational		10 (3.2)	117 (30.0)				
Status				o = 4 =#			
Primary education	or	2 (16.7)	10 (83.3)	0.745"	1	0.321	

less Secondary education and more	19 (9.1)	189 (90.9)			
Employment Status Employed Unemployed	14 (8.4) 7 (13.2)	153 (91.6) 46 (86.8)	1.084	1	0.298
Monthly income (Naira) Less than 18000 18000 and more	13 (11.1) 8 (7.8)	104 (88.9) 95 (92.2)	0.709	1	0.400
Support from Partner Yes No	14 (7.4) 7 (21.9)	174 (92.6) 25 (78.1)	6.593	1	0.010*
Support from partner's relatives Yes No	7 (5.8) 14 (14.1)	114 (94.2) 85 (85.9)	4.403	1	0.036*
Number of Children One Child Two or more children	4 (4.5) 17 (12.9)	84 (95.5) 115 (87.1)	4.247*	1	0.039*
FamilyhistoryofmentalillnessYesNo	2 (40.0) 19 (8.8)	3 (60.0) 196 (91.2)	5.496*	1	0.073
Witness sexual abuse while growing up Yes No	2 (50.0) 19 (8.8)	2 (50.0) 197 (91.2)	7.722#	1	0.046*
Presence of Suicidal Ideation Yes No	5 (45.5) 16 (7.7)	6 (54.5) 193 (92.3)	17.292	1	0.0001*

198 *Significant *Fisher's test used

Table 2 shows the socio-demographic and clinical factors associated with postpartum depression among
respondents. There was a significant association between postpartum depression and age of the

respondents. Seventeen (13.2%) of those whose ages were 30 years and above had postpartum depression while only 4(4.4%) of those whose ages were 30 years and below had postpartum depression (χ 2=4.767, p=0.035).

There was also statistically significant association between postpartum depression and suicidal ideation among the respondents. Five (45.5%) respondents who had suicidal ideation had postpartum depression while 7.7% of those who had no suicidal ideation had postpartum depression (χ 2=17.292, p=0.0001).

There was a significant association between postpartum depression and support from partner during pregnancy. Twenty two percent of those without husband's support had postpartum depression compared to 7.4 % of those with support from husband (χ 2=6.593, p=0.010). Likewise, support from partner's relatives during pregnancy was also found to be significantly associated with postpartum depression. Fourteen (14.1%) of those that were not supported by partner's relatives during pregnancy had postpartum depression while only seven (5.8%) of those whose partner's relatives supported them had postpartum depression. This difference was also statistically significant (χ 2=4.403, p=0.036).

There was also a significant association between postpartum depression and number of children. Four (4.5%) respondents who had one child were depressed compared to 17 (12.9%) respondents who had two or more children (χ 2=4.247, p=0.039).

The association between postpartum depression and witnessing sexual abuse while growing up was also significant. Two (50%) respondents who witnessed sexual abuse while growing up had postpartum depression while only 8.8% of those who did not witness sexual abuse 0while growing up had depression $(\chi^2 = 7.722, p=0.046)$.

The mean score of EPDS across the occupational status of the respondents revealed a significant difference (not tabulated). The unemployed respondents were observed to have significantly higher EPDS scores than others implying that they had higher depressive symptoms (F=3.020, p< 0.05).

There were no statistically significant associations between postpartum depression and other variables such as marital status, educational status, monthly income, family history of mental illness and experience of sexual abuse while growing up.

Table 3: Postpartum depression and Socio-demographic Characteristics (Continuous variables) of respondents

Variable	Postpartum Depression		Т	df	Р
	Present (N=21)	Absent (N=199)			
	Mean (SD)	Mean (SD)			
Age of partner (years)	36.47 (3.501)	35.18 (5.651)	1.495	218	.145
Weeks since delivery	11.51 (8.003)	10.05 (6.251)	.987	218	.325

230

As shown in Table 3 above, the continuous variables of "age of participant's partner" and "weeks since delivery" were compared across those that had postpartum depression and those that did not have postpartum depression using student t-test. There was no significant difference between the mean age of participants' partners who had postpartum depression compared to mean age of partners of those who did not have postpartum depression (p=0.145). Although the mean postpartum period (weeks since delivery) in those with postpartum depression was slightly higher than those who did not have postpartum depression, the difference was not statistically significant (p=0.325).

238

Table 4: Association between postpartum depression and other variables in respondents using
 logistic regression

Variables	В	Odds ratio	P value	95% CI for EX	P (B)
Age (years < 30 (ref) ≥30	1 1.169	1 3.218	0.060	0.950	10.901
Number Children 1 (ref) ≥2	of 1 0.824	1 2.322	0.171	0.695	7.763
Mode delivery	of				

Vaginal(ref) CS	1 -0.807	1 0.446	0.327	0.089	2.243
Witnessed sexual abuse while growing up No (ref) Yes	1 1.839	1 6.290	0.176	0.439	90.233
Support from Husband's relatives					
No (ref)	1	1	0.450	0.470	4.040
Yes Support from	-0.733	0.481	0.152	0.176	1.310
Husband					
No (ref)	1	1			
Yes	-1.183	0.306	0.043	0.096	0.977
Hours of					
delivery					
0 to 12	1	1			
hours(ref)	0.004	4 005	0.543	0.044	40.004
>12 nours	0.691	1.995	0.547	0.211	18.891

ref reference point which is the variable to which others are being compared

242

Association between postpartum depression and other variables in respondents using logistic regression

are as shown in Table 4 above.

245 Variables were individually entered into a binary logistic regression model with postpartum depression as

the outcome variable and the significant predictor of postpartum depression is as depicted in Table 4

above.

248 **DISCUSSION**

In this study, the prevalence of postpartum depression was 9.5% squarely in the centre of the global range of prevalence of postpartum depression.(7-9) It is close to the 14.6% that was observed by Adewuya et al.(10) The rate is also comparable to studies from other parts of the world.(21, 22) An Australian study among new mothers found a prevalence rate of 7.5%,(21) likewise a community based study, where the rate was found to be 5.5%.(22) The rate was however lower than 22% reported among Jordanian women in primary care setting(23) and 27.2% found at 6 weeks postpartum in a survey of 206 women using EPDS.(24) These differences could be as a result of methodological differences
 considering the fact that only screening instrument was used in the studies.

257 There was a statistically significant association between maternal age and postpartum depression. 258 Women who had postpartum depression were more likely to be older. This is similar to the findings by 259 Kheirabadi et al who also found an association between postpartum depression and advanced maternal 260 age.(25) It is however different from a study by Abiodun et al who found association between younger 261 maternal age and postpartum depression.(26) It is also not in keeping with other studies in which no 262 association was found between maternal age and postpartum depression. (10, 27, 28) This finding further 263 emphasizes the importance of routine screening for postpartum depression especially in mothers who are 264 more than 30 years.

265 There was no statistically significant association between marital status and postpartum depression. This 266 is in keeping with other studies in which marital status does not seem to be related to depression.(8, 29) 267 However, it is unlike a study by Segre et al in which higher prevalence of postpartum depression was 268 found among unmarried postpartum women. (30) Marital status in itself may not be related to depression 269 but single mothers may receive less social support or have more deprived socio-economic situation. 270 Considering the fact that majority of the respondents were married, this may not reflect the true status of 271 the community, hence complementary community based studies may be needed to give the complete 272 picture among the women in the community.

- There was a statistically significant relationship between employment status and postpartum depression in this study. This is similar to what was obtained in similar studies.(14, 31) Most studies showed statistically significant associations between employment status and postpartum depression.(8, 14, 32) Unemployment predicts low socioeconomic status and has been found to be associated with the development of postpartum depression.(33) This may be attributed to the increased amount of stress placed on a mother due to financial means necessary for raising an infant.
- There was no statistically significant relationship between educational status and postpartum depression
 which is in keeping with a study by Owoeye et al but in contrast to findings by Ozdemir et al who found

that low educational level predicted postpartum depression.(14, 28) In this study, majority of the respondents (94.5%) had post-primary school education. The high literacy level may not give a true representation of the educational status of the community. Hence complementary community based studies may be needed to give the complete picture among the women in the community.

285

286 There was also a significant association between postpartum depression and number of children. 287 Respondents with two or more children reported higher levels of depression compared with respondents 288 who had one child. This finding agrees with some other studies, (22, 34) however a study by Ludemir et al 289 found no significant association between parity and postpartum depression. (29) A possible explanation 290 for this is the fact that in a country like Nigeria where resources are limited, an extra child might well contribute to an increased stress level of the mother. Furthermore, new mothers who need to care for 291 292 more young children at home may experience more stress which may predispose to postpartum 293 depression.

294 There was a statistically significant association between support from husband during pregnancy and 295 postpartum depression. Women who reported a lack of support from husband were more likely to develop 296 postpartum depression. This is in keeping with other studies where lack of husband's support was found 297 to be associated with postpartum depression. (35, 36) In a Norwegian study on depression and sleep in 298 postpartum period, one of the variables most strongly associated with depressive symptoms was being 299 discontent with the relationship with the partner.(9) Also, women who had support from husband's 300 relatives were less likely to have postpartum depression which is in keeping with a study by Mohammad 301 et al.(23) Absence of social support or social isolation has also been found to be risk factors for 302 developing postpartum depression. (12, 13) Poor partner relationship has previously been found to be 303 associated with depression both among Indian(37) and Norwegian postnatal women.(11) In a qualitative 304 study from India, women reported poor marital relationship as an important cause of their postnatal 305 depression.(38)

306 An alternative interpretation of the association is that a woman's depression may influence the marital 307 relationship negatively. Depressed women may rate marital difficulties more negatively than non-308 depressed women. However, a longitudinal study of immigrant women in Canada found lower scores on 309 the marital adjustment scale in pregnancy to be predictive of depressive symptoms two months after 310 delivery.(39) Given assertions that strong, trusting partner relationship may be vital for a woman's 311 psychological health during the postpartum period, (40) it makes sense that intimate partner violence 312 during this period may have a strong detrimental effect on women's mental health. (41, 42) Lack of social 313 support contribute to high rates of mental health problems after childbirth.(43) Findings from Coker et al 314 indicate that the risk of negative mental health outcomes declines significantly among abused women 315 who report receiving social support. (44) In Nigeria, Fatoye et al in a study on emotional distress and its 316 correlates among Nigerian women in late pregnancy also found that lower spousal support contributes to 317 mental health problems in women.(45)

318 There was no statistically significant association between family history of mental illness and postpartum

319 depression. This is not in agreement with other studies that have shown that risk factors for postpartum

320 depression can be familial and genetic.(9, 46) The respondents may have been unaware or unwilling to

321 Odisclose a history of psychiatric illness in the family.

322 There was also a statistically significant relationship between suicidal ideation and postpartum 323 depression. About half (45.5%) of those who had suicidal ideation had postpartum depression compared 324 to 7.7% of those without suicidal ideation. This is in keeping with findings from other studies in which 325 suicidal ideation has been found to be an integral part of postpartum depression.(47, 48)Findings from a 326 postpartum depression screening program of 10,000 women, indicated postpartum women who screened 327 positive for depression had high rates of suicidal ideation.(49) Suicidal ideation is a recognized symptom 328 of depression(50) and may represent an aspect of hopelessness which is also a manifestation of 329 depression.(50) Depression has been described as the single largest risk factor for suicidal ideation.(51)

330

331 Strengths of the study

- 332 100% questionnaires filled; screening and diagnostic instrument used; good training of research
- 333 assistant; tertiary hospital
- 334 **Limitations**
- 335 1. The study is subject to both recall and reporting bias because instruments used were based on self-
- 336 report.
- 337 2. The study population was drawn from one hospital, and may therefore not reflect the characteristics of
- 338 postnatal depression of the general population of Nigeria.
- 339
- 340

341 CONCLUSION

- 342 The prevalence of postpartum depression in this study shows the high burden of the disorder. Increased
- 343 media campaign about postpartum depression and preventive measures is urgently needed. Screening
- 344 for depression throughout the perinatal period is important for early diagnosis and prompt intervention to
- 345 improve the clinical outcome of women affected.
- 346

347 COMPETING INTEREST: NONE DECLARED

- 348 **CONSENT**: All participants gave written informed consent
- 349 **ETHICS**

Approval to undertake the study was obtained from the Ethics and Research Committee of LAUTECH Teaching Hospital to ascertain that the methodology does not contravene guidelines for research involving human subjects. Ethical issues like non-disclosure to others, opportunity to decline interview at any stage and non-exposure to risk were discussed with each respondent. The participants bore no financial burden for the study.

355 The respondents with depression were properly counseled on the need for help and were referred 356 appropriately to a psychiatric facility for expert management.

358 **REFERENCES**

Desmarais SL, Pritchard A, Lowder EM, Janssen PA. Intimate partner abuse before and during
 pregnancy as risk factors for postpartum mental health problems. BMC Pregnancy and Childbirth.
 2014;14:132.

Cadzow SP, Armstrong KL, Fraser JA. Stressed parents with infants:reassessing physical abuse
 risk factors. Child Abuse and Neglect. 1999;23(9):845-53.

364 3. Wisner KL, Peindl KS, Gigliotti T, Hanusa BH. Obsessions and compulsions in women with 365 postpartum depression. Journal of Clinical Psychiatry. 1999;60(3):176-80.

3664.Rahman A, Iqbal Z, Bunn J, Lovel H, Harrington R. Impact of maternal depression on infant367nutritional status and illness: a cohort study. Archives of General Psychiatry. 2004;61(9):946-52.

Letourneau N, Morris CY, Stewart M, Hughes J, Critchley KA, Secco sL. Social Support needs
identified by mothers affected by intimate partner violence. Journal of Interpersonal Violence.
2013;29(14):2873-93.

Oates M. Perinatal psychiatric disorders: a leading cause of maternal morbidity and mortality.
 British Medical Bulletin. 2003;67:219-29.

373 7. Miller L, LaRusso E. Preventing Postpartum Depression. Psychiatric Clinics of North America.
374 2011;34:53-65.

O'Hara MW, Swain AM. Rates and risk of postpartum depression- a meta-analysis. International
 Review of Psychiatry. 1996;8(1):37-54.

9. Signe KD. Depression and sleep in the postnatal period: University of Bergen; 2009.

Adewuya AO, Eegunranti AB, Lawal AM. Prevalence of postnatal depression in Western Nigerian
women: a controlled study. International Journal of Psychiatry in Clinical Practice. 2005;9(1):60-4.

11. Eberhard-Gran M, Eskild A, Tambs K, Samuelsen SO, Opjordsmoens S. Depression in postpartum
 and non-postpartum women: prevalence and risk factors. Acta Psychiatrica Scandinavica.
 2002;106(6):426-33.

Baker D, Taylor H. The relationship between condition-specific morbidity, social support and
 material deprivation in pregnancy and early motherhood. ALSPAC Survey Team.Avon Longitudinal study
 of pregnancy and childhood. Social Science and Medicine. 1997;45(9):1325-6.

Brugha TS, Sharp HM, Cooper SA, Weisender C, Britto D, Shinkwin R, et al. The Leicester 500
 Project.social support and the development of postnatal depressive symptoms, a prospective cohort
 survey. Psychological Medicine. 1998;28(1):63-79.

389 14. Owoeye AO, Aina OF, Morakinyo O. Risk factors of postpartum depression and EPDS scores in a
 390 group of Nigerian women. Tropical Doctor. 2006;36(2):100-3.

Robertson E, Grace S, Wallington T, Stewart DE. Antenatal risk factors for postpartum
 depression: a synthesis of recent literature. General Hospital Psychiatry. 2004;26(4):289-95.

393 16. Araoye MO. Research methodology with statistics for health and social sciences. Ilorin:394 Nathadex; 2004.

17. Cox JL, Murray D, Chapman G. A controlled study of the onset, duration and prevalence of
 postnatal depression. British Journal of Psychiatry. 1993;163:27-31.

18. Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Development of the 10-item
Edinburgh Postnatal Depression Scale. British Journal of Psychiatry. 1987;150:782-6.

39919.Ukwakwe R. Affective(depressive) morbidity in puerperal Nigerian women:validation of400Edinburgh Postnatal Depression Scale. Acta Psychiatrica Scandinavica 2003;107:251-9.

Adewuya AO, Ola B, Dada AO, Fatoso O. Validation of the Edinburgh Postnatal Depression Scale
as a screening tool for depression in late pregnancy among Nigerian women Journal of Psychosomatic
Obstetrics and Gynaecology. Journal of Psychosomatic Obstetrics and Gynaecology. 2006;27(4):267-72.

Eastwood JG, Jalaludin BB, Kemp LA, Phing NH, Barnett BE. Relationship of postnatal depressive
symptoms to infant temperament, maternal expectations, social support and other potential risk
factors: Findings from a large Australian cross-sectional study. BMC Pregnancy and Childbirth.
201;12:148.

40822.Nielsen FD, Videbech P, Hedegaard M, Dalby SJ, Secher NJ. Postpartum depression:409identification of women at risk. British Journal of Obstetrics and Gynaecology. 2000;107:1210-7.

410 23. Mohammad KI, Gamble J, Creedy DK. Prevalence and factors associated with the development 411 of antenatal and postnatal depression among Jordanian women. Midwifery. 2011;27:238-45.

412 24. Ebeigbe PN, Akhigbe KO. Incidence and associated factors of postpartum depression in a tertiary
413 hospital in Nigeria. Nigerian Postgraduate Medical Journal. 2008;15:15-28.

414 25. Kheirabadi GR, Maracy MR. Perinatal depression in cohort study on Iranian women. Journal of 415 Research in Medical Sciences. 2010;15:41-9.

416 26. Abiodun OA. Postnatal depression in primary care populations in Nigeria. General Hospital
417 Psychiatry. 2006;28:133-6.

418 27. Dauda S, Kazeem AA, Lamaran MD, Bappah MA. Postnatal depression and its associated factors
419 among Northeastern Nigerian women. Annals of Tropical Medicine and Public Health. 2016;9(3):184-90.

420 28. Ozdemir h, Ergin N, Selmoglu K, Bilgel N. Postnatal depression mood in Turkish women.
421 Psychological Health and Medicine. 2014;10:96-107.

Ludermir A, Lewis G, Valongueiro S, Araujo T, Araya R. Violence against women by their intimate
partner during pregnancy and postnatal depression: a prospective cohort study. Lancet.
2010;376(9744):903-10.

30. Segre LS, O'Hara MW, Arndt S, Stuart S. The prevalence of postpartum depression: The relative
significance of three social status indices. Social Psychiatry and Psychiatric Epidemiology.
2007;42(4):316-21.

428 31. Warner R, Apple L, Whitton A, Faragher B. Demographic and obstetric risk factors for postnatal 429 psychiatric morbidity. British Journal of Psychiatry. 1996;168:607-11.

430 32. Lee DT, Yip AS, Leung TY, Chung TK. Identifying women at risk of postnatal depression:
431 prospective longitudinal study. Hong Kong Medical Journal. 2000;6:349-54.

432 33. Ahmed HM, Alalaf SK, Al-Tawil NG. Screening for postpartum depression using Kurdish version
433 of Edinburgh Postnatal Depression Scale. Archives of Gynecology and Obstetrics. 2012;285:1249-55.

434 34. Mapayi B, Makanjuola ROA, Mosaku O, Adewuya A, Afolabi O, Aloba O, et al. Impact of intimate
435 partner violence on anxiety and depression amongst women in Ile-ife, Nigeria. Archives of Women's
436 Mental Health. 2013;16(1):11-8.

437 35. Ege E, Timur S, Zincir H, Geckil E, Sunar-Reeder B. Social support and symptoms of postpartum
438 depression among new mothers in Eastern Turkey. Journal of Obstetrics and Gynaecology Research.
439 2008;34:585-93.

440 36. Yagmur Y, Ulukoca N. Social support and psotnatal depression in low-socioeconomic level 441 postpartum women in Eastern Tuurkey. International Journal of Public Health. 2011;55:543-9.

442 37. Patel V, Rodrigues M, Desouza N. Gender, poverty, and postnatal depression: a study of mothers
443 in Goa, India. American Journal of Psychiatry. 2002;159(1):43-7.

44438.Pereira B, Andrew G, Pednekar s, Pai R, Pelto P, Patel V. The explanatory models of depression in445low income countries: Listening to women in India. Journal of Affective Disorders. 2006.

39. Zelkowitz P, Saucier JF, Wang T, Katofsky L, Valenzuela M, Westrich R. Stability and Change in
depressive symptoms from pregnancy to two months postpartum in childbearing immigrant women.
Archives of Women's Mental Health. 2008;11(1):1-11.

449 40. Dennis C, Ross L. Women's perception of partner support and conflict in the development of 450 postpartum depressive symptoms. Journal of Advanced Nursing. 2006;56(6):588-99.

451 41. Antoniou E, Vivilak V, Dagles M. Correlation of domestic violence during pregnancy with 452 postnatal depression: systematic review of bibliography. Health Science Journal. 2008;2(1):15-9.

453 42. Abiodun AD, Mapayi B, Abiodun AB, Mosanya JT, Adeomi AA. The relationship between intimate 454 partner violence and postpartum depression in Osogbo Nigeria. International Journal of Reproduction, 455 Contraception, Obstetrics and Gynaecology. 2018;7(10):3911-8.

456 43. Beck C. Predictors of postpartum depression. Nursing Research. 2001;50(5):275-85.

457 44. Coker A, Smith P, Thompson M, Mokeown R, Bethea L, Davis K. Social support protects against 458 the negative effects of partner violence on mental heatlth. Journal of Women's Health and Gender 459 Based Medicine. 2002;11(5):465-76.

460 45. Fatoye FO, Adeyemi AB, Oladimeji BY. Emotional distress and its correlates among Nigerian 461 women in late pregnancy. Journal of Obstetrics and Gynaecology. 2004;24(5):504-9.

46. Treloar SA, Martin NG, Bucholz KK, Madden PA, Heath AC. Genetic influences on postnatal
depressive symptoms: findings from an Australian twin sample. Psychological Medicine. 1999;29(3):64554.

465 47. Sit D, Luther J, Buysse D, Dills JL, Eng H, Okun M, et al. Suicidal ideation in depressed postpartum 466 women:; Associations with childhood trauma, sleep disturbance and anxiety. Journal of Psychiatric 467 Research. 2015;66:95-104.

468 48. Paris R, Bolton RE, Weinberg MK. Postpartum depression, suicidality and mother-infant 469 interactions. Archives of Women's Mental Health. 2009;12(5):309-21.

470 49. Wisner KL, Sit D, McShea MC, Rizzo DM, Zoretich RA, Hughes CL, et al. Onset timing, thoughts of 471 self-harm, and diagnoses in postpartum women with screen-positive depression findings. Journal of 472 American Medical Association. 2013;70(5):490-8.

473 50. WHO. The ICD-10 classification f mental and Behavioural Disorders: clinical descriptions and 474 diagnostic guidelines. 10th ed. Geneva, Switzerland: World Health Organization; 1992.

475 51. Bernal M, Haro JM, Bernert S, Brugha TS, deGraaf R, Bruffaerts R, et al. Risk factors for suicidality
476 in Europe: Results from the ESEMED study. Journal of Affective Disorders. 2007;101:27-34.

477

478

479

480