

Original Research Article

Quantification of Amaroswerin and Amarogentin in different parts of *Swertia chirayita* chromatographic analysis

ABSTRACT

Aims: In the present investigation, our aim is to quantify major secoiridoids (amaroswerin and amarogentin) in different parts of *Swertia chirayita* using HPLC to distinguish its parts having high content of amaroswerin and amarogentin, so that these parts must be included in plant material when plant is used for extraction of these phytoconstituents.

Study design: Statistical comparison was performed using OP-STAT software with CRD and was considered statistically significant.

Place and Duration of Study: The plant material including all studied parts was procured from field plants grown at Medicinal and Aromatic Plants Farm, Shilly (latitude-N 30° 54' 30" and longitude E 77° 07' 30", elevation 1550m) under Department of Forest Products, UHF, Solan (H.P.) India. Study was undertaken in the Departmental laboratory and in the period between September 2016 and December 2016.

Methodology: Waters binary HPLC unit with Waters HPLC pump 515, dual λ absorbance detector 2487 and Empower II software was used for quantification of phytoconstituents under study. Samples of different plant parts were extracted by using soxhlet method using methanol as solvent.

Results: In different parts of *Swertia chirayita*, amaroswerin content (%) was reported highest in flowers (0.741%), followed by leaves (0.386%), roots (0.188%) and lowest in stem part (0.226%) and amarogentin content (%) was reported as highest in flowers (0.617%), followed by leaves (0.447%), stem (0.426%) and lowest in roots (0.369%) of field grown plants.

Conclusion: To conclude, amaroswerin and amarogentin are present in all studied parts of *Swertia chirayita*. Amaroswerin and amarogentin content ranged from 0.160% to 0.741% and 0.369% to 0.617% respectively in different plant parts under study.

Keywords: *Swertia chirayita*, Amaroswerin, Amarogentin, quantification, HPLC.

1. INTRODUCTION

Swertia chirayita (Roxb. ex Fleming) H. Karst. belonging to the family Gentianaceae, listed in critically endangered category of IUCN list and is native of Himalayan region[1]. It is known differently in different languages as Chirayata (Hindi), Kairata, Bhunimba (Sanskrit), Chairavata (Urdu) etc. Its distribution is from Kashmir to Bhutan at an altitude of

1200-3000m and in the Khasi hills at 1200-1500m. It can also be grown in sub-temperate regions between 1500-2100m altitude [2,3,4,5]. It has been used in traditional medicine mainly for skin diseases and chronic fever and also for malaria, anaemia, bronchial asthma, liver disorders, hepatitis, gastritis, constipation, worms, epilepsy, ulcer, scanty urine, hypertension, certain type of mental disorders, secretion of bile, blood purification and diabetes [6, 7, 8, 9, 10]. *Chirayita* is high prized medicinal herb in India and is used either alone or as one of the constituents in some polyherbal formulations like Ayush-64, Melicon V etc. The plant has also been reported to possess hypoglycaemic activity[11], anti-inflammatory activity[12], hepatoprotective activity[13], wound healing activity[14], anti-carcinogenic activity[10], anti-malarial activity[15] as well as antibacterial activity[16]. Amarogentin is a well reported compound for antileishmanial, anticancerous, anti-diabetic and gastroprotective activity by different researchers[17,18,19,20]. Also amaroswerin reported to be have gastroprotective activity[19]. Due to its intense medicinal importance demand for this plant is increasing at the rate of 10 percent per year [21].

Keeping this in view, present investigation was carried out to quantify the major active constituents *i.e.* amaroswerin and amarogentin in different parts (leaves, flowers, stems and roots) of *Swertia chirayita* using HPLC in order to know its parts having high content of both amaroswerin and amarogentin, so that these plant parts must be included in plant material when used for extraction of studied phytoconstituents.

2. MATERIAL AND METHODS

2.1 Sample source: The plant material including all studied parts was procured from field plants grown at Medicinal and Aromatic Plants Farm, Shilly (30°54'32.40" N latitude and 77°09'04.29" E longitude, elevation 1550m) Department of Forest Products, UHF, Solan (H.P.) India. Identification was done in Herbarium Section of above said department with reference number 13386.

2.2 Extraction, Separation and Purification of Standard compounds *i.e.* Amarogentin and Amaroswerin): Field grown plants of *Swertia chirayita* were harvested at flowering stage, dried in the open under shade, coarsely powdered and further dried in the oven for 12 hours at 35-40°C. The oven dried material (500gm) was then further powdered and repeatedly extracted (5x3 hours) with petroleum ether (40-60°C) through reflux method on a boiling water bath. The petroleum ether extracted plant material was dried in the open (for complete evaporation of solvent) and then repeatedly extracted (5x4 hours) with methanol (3lts) through reflux method on a boiling water bath. All the filtrates were combined and then total volume was reduced to about 400ml by partially distilling off the solvent. The contents were then kept overnight in the refrigerator (4°C) when some brown coloured solid mass separated out which was filtered under vacuum and thoroughly washed with methanol. All the methanol washings were combined with the main filtrate and then solvent was completely distilled off under vacuum. The residue thus obtained was mixed with water (250ml) and contents were repeatedly extracted with chloroform (5 x 200ml) in a separating funnel to remove less polar compounds. Each time, after clear separation of solvent layers, lower chloroform layer was withdrawn and upper aqueous layer was further extracted with chloroform. The aqueous layer containing bitter compounds was then extracted with n-butanol (5 x 100ml) in a separating funnel. Each time after clear separation of the solvent layers, the upper n-butanol layer was taken and aqueous layer re-extracted with n-butanol. All the n-butanol layers were combined and solvent was distilled off under reduced pressure in a rotary vacuum evaporator.

The residue containing bitter compounds (obtained after distillation of n-butanol) was dissolved in minimum volume of methanol and then thoroughly mixed with small quantity of column chromatography grade Silica Gel (60-120 mesh) in a beaker. The contents were thoroughly mixed and then solvent was allowed to evaporate in the open and finally the adsorbed mixture was dried in a vacuum desiccator. The adsorbed mixture was then loaded on a clean and dry glass column (5 cm x 100 cm) packed with silica Gel (250gm, 60-120 mesh) and eluted with chloroform: methanol (98: 2 to 85:15). The eluted fractions of 50 ml each were collected and after distilling off the solvent from each fraction, were monitored on silica Gel G pre-coated TLC plates developed in solvent system chloroform: methanol: water :: 65:20:10. The spots were visualized by iodine and fast red- B salt solution where amarogentin and amaroswerin spots appeared as red spots with the latter reagent and as brown spots with iodine. None of the fractions afforded pure bitter compounds only. All the fractions containing bitter compounds (amarogentin and amaroswerin) were then combined and further chromatographed on a Silica Gel packed narrow glass column (3cm x 60cm) using chloroform: methanol (98:2 to 95:5) as

the solvent system. This time fractions of 25ml each were collected and monitored through TLC. The fractions containing the bitter compound having higher R_f (0.39) were combined, concentrated, dried under vacuum and dissolved in minimum quantity of methanol. Ethyl acetate was then added drop wise till slight turbidity appeared and the flask was left at room temperature and the bitter compound were crystallized after two days and then dried. Authenticity of the crystallized compound (amarogentin) was ascertained by running TLC and HPLC with a sample of reference amarogentin. The other fraction containing bitter compound having lower R_f (0.34) in above solvent system could not be crystallized. This fraction was again column chromatographed on small sized column (1cm x15 cm) using chloroform: methanol (98:2 to 95:5) solvent system for elution. The fractions containing pure amaroswerin (R_f = 0.34) were combined, concentrated and the amaroswerin was crystallized using methanol and ethyl acetate. The authenticity was ascertained by running TLC and HPLC with reference amaroswerin sample. The purity of isolated compounds was established by using HPLC.

2.3 Plant Material preparation: Parts of the plant separated, shade dried and was grinded mechanically to form the uniform particle size of the plant material, which was used for quantification of both amaroswerin and amarogentin in the samples under study.

2.4 Extraction of different plant parts samples: Oven dried and powdered plant material samples (2gm) of each plant part were repeatedly extracted in three replications with dichloromethane (100ml) in soxhlet apparatus for the removal of dichloromethane soluble, less polar and non bitter compounds. The samples were then dried in the oven and extracted with methanol (100ml) for 8 hours extraction duration. After extraction, the solvent from each extracted sample was completely distilled off and the residue was completely dried. Dry residue then dissolved in distilled water (25 ml) and then repeatedly partitioned with n-butanol (20ml). After clear separation of layers, upper n-butanol layer was collected and aqueous layer was again partitioned with n-butanol. All the n-butanol layers combined and solvent distilled off in a rotary vacuum evaporator and residue collected. The residue was dried to a constant weight after that the samples for HPLC analysis were prepared with mobile phase (methanol : water :: 45:55, v/v), filtered through a 0.25µm filter.

2.5 Instrumentation: The system used is of Waters binary HPLC unit with Waters HPLC pump 515, dual λ absorbance detector 2487 and program used for data analysis was Empower II software.

2.6 Chromatographic method: HPLC method used was developed by Sharma, 2017 [22]. UV detection was done at 235 nm, mobile phase (methanol : water :: 45:55, v/v), isocratic elution at a flow rate of 1ml/min. and injection volume was set to 20 µl. The total run time of standard and sample was 30 min. and 35 min. respectively with retention time of amaroswerin and amarogentin was 14.449 ± 0.102 min. and 18.610 ± 0.140 min. (Mean ± standard deviation of triplicate analysis) respectively.

2.7 Statistical analysis: Statistical comparison was performed OP-STAT software and was considered statistically significant.

3. RESULTS AND DISCUSSION

The results of the study aimed at the determination of the amaroswerin and amarogentin contents in different plant parts of *Swertia chirayita*. The results of studied contents in different parts of the plant found in present study are given in Table 1. The HPLC chromatograms of standard compounds and sample of different plant parts under study are given in Figures 1-5. The amaroswerin content (%) was highest in flowers (0.741%), followed by leaves (0.386%), roots (0.188%) and lowest in stem part (0.226%). The amarogentin content (%) in different parts of *Swertia chirayita* was reported as highest in flowers (0.617%), followed by leaves (0.447%), stem (0.426%) and lowest in roots (0.369%) of field grown plants of *Swertia chirayita*. Similar results have also been found for amaroswerin and amarogentin content in *Swertia japonica* [23] and *Swertia chirayita* [24]. Quantification of studied chemical compounds in *Swertia chirayita* fruit/seed is not done in this study so further studies can be done in this context.

Table 1: Quantification of Amaroswerin and Amarogentin in different parts of *Swertia chirayita*

Sr. no.	Plant Part	Amaroswerin content (%)	Amarogentin content (%)
1	Flowers	0.741 (0.860)	0.617 (0.786)
2	Leaves	0.386 (0.621)	0.447 (0.669)

3	Stem	0.160 (0.399)	0.426 (0.653)
4	Roots	0.188 (0.433)	0.369 (0.607)
CD _{0.05}		0.017	0.021
SE(m)		0.006	0.007
Values in the parentheses are transformed values using square root transformed values			




Fig.1: HPLC chromatogram of standard compounds amaroswerin (95 µg/ml) and amarogentin (95 µg/ml).




Fig. 2 : HPLC chromatogram of flowers sample of *Swertia chirayita*




Fig.3 : HPLC chromatogram of leaves sample of *Swertia chirayita*




Fig. 4 : HPLC chromatogram of stems sample of *Swertia chirayita*




Fig. 5 : HPLC chromatogram of roots sample of *Swertia chirayita*

4. CONCLUSION

To conclude, amaroswerin and amarogentin were present in all studied parts of *Swertia chirayita*. Amaroswerin and amarogentin content ranged from 0.160% to 0.741% and 0.369% to 0.617% respectively. From the present investigation, it can be suggested that for extraction purpose, harvesting should be done near to mature stage of plant, so that good amount of flowers and leaves are present in harvested plant material. It will aid in conservation of resources as less plant material gives more yield.

REFERENCES

- [1] Joshi P and Dhawan V. *Swertia chirayita* - an overview. *Current Science*. 2005; 89: 635-640.
- [2] Bentley R, Trimen H. (Republished 2002) *Medicinal Plants*. Vol. III, 1880. IBD, Dehradun, 183p.
- [3] Clarke CB. *The Flora of British India*. 1885. Vol.- IV, Hooker, J.D. (ed.) L.
- [4] Kirtikar KR, Basu BD. *Indian Medicinal Plants*. 1975. Vol.III, 2nd ed., M/S Bishen Singh Mahendra Pal Singh, Deharadun, 2391p.
- [5] Gaur RD. *Flora of the District Garhwal*, North West Himalaya (with ethnobotanical notes). 1999. Transmedia, Srinagar (Garhwal). p. 408.

- [6] Karan M, Vashisht K, Handa SS. Antihepatotoxic activity of *Swertia chirayita* on carbon tetrachloride induced hepatotoxicity in rats. *Phytotherapy Research*. 1999; 13:24-30.
- [7] Banerjee S, Sur TK, Mandal S, Das PC, Sikdar S. Assessment of the anti-inflammatory effect of the *Swertia chirayita* in acute and chronic experimental models in male albino rats. *Indian Journal of Pharmacology*. 2000; 32: 21-24.
- [8] Airi S, Bhatt A, Dhar U. Characterisation of *Swertia chirayita* (Roxb.ex Fleming) Karsten and *S. angustifolia* Ham. ex D Don. using SDS-PAGE. *Plant Genetic Resources Newsletter* (FAO Biodiversity). 2002; 147: 72-77.
- [9] Gao-Feng S, Run-Hua R, Yunh-Shang Y, Chun-Lei L, Ai-Mei, Xiang Li. 2004. Isolation and crystal structure of xanthenes from *Swertia chirayita*. *Chinese Journal of Structural Chemistry*. **23**: 1164-1168.
- [10] Saha P, Mandal S, Das A, Das PC, Das S. Evaluation of anticarcinogenic activity of *Swertia chirayita* Buch Ham, an Indian medicinal plant on DMBA induced mouse skin carcinogenesis model. *Phytotherapy Research*. 2004; 18: 373-378.
- [11] Chandrasekar B, Bajpai MB, Mukherjee SK. Hypoglycemic activity of *Swertia chirayita* (Rox ex. Flem) Karst. *Indian Journal of Experimental Biology*. 1990; 28: 616-618.
- [12] Chowdhary NI, Banyopadhyay SK, Banerjee SN, Dutta MK, Das PC. Preliminary studies on the anti-inflammatory effects of *Swertia chirayita* in albino rats. *Indian Journal of Pharmacology*. 1995; **27**: 37-39.
- [13] Chakravarty AK, Mukopadhyay S, Moitra SK, Das B. Syringaresinol, a hepatoprotective agent and other constituents from *Swertia chirayita*. *Indian Journal of Chemistry*. 1994; 338: 405-408.
- [14] Manjunath KP, Kulkarni GT, Patil KS. Preliminary investigation and wound healing activity of the root of *Swertia chirata* Buch Ham. *Indian Drugs*. 2006; **43**:535-537.
- [15] Valecha N, Devi U C and Joshi H. 2002. Comparative efficacy of ayush -64 vs chloroquine in vivax malaria. *Current Science*. **78**: 1120-1122.
- [16] Bhargava S and Garg R. 2007. Evaluation of antibacterial activity of aqueous extract of *Swertia chirayita* Buch Ham. root. *International Journal of Green Pharmacy*. **2**: 51-52
- [17] Ray S, Majumder H K, Chakravarty A K, Mukhopadhyay S, Gil R R and Cordell G A. 1996. Amarogentin, a naturally occurring secoiridoid glycoside and a newly recognized inhibitor of topoisomerase I from *Leishmania donovani*. *Journal of Natural Product*. **59**: 27-29.
- [18] Pal D, Sur S, Mandal S, Das A, Roy A, Das S *et al*. 2012. Prevention of liver carcinogenesis by amarogentin through modulation of G1/Scellcycle check point and induction of apoptosis. *Carcinogenesis*. **33**: 2424-2431.

- [19] Niiho Y, Yamazaki T, Nakajima Y, Yamamoto T, Ando H and Hirai Y. Gastro protective effects of bitter principles isolated from Gentian root and *Swertia* herb on experimentally induced gastric lesions in rats. *Journal of Natural Medicines*. 2006; 60: 82-88.
- [20] Phoboo S, Bhowmik P C, Jha P K and Shetty K. 2010. Anti-diabetic potential of crude extracts of medicinal plants used as substitutes for *Swertia chirayita* using *in vitro* assays. *Journal of Plant Sciences*. 7:48–55.
- [21] Anonymous (2008). <http://www.nmpb.nic.in> : Official website of National Medicinal Plant Board (NMPB), New Delhi.
- [22] Sharma S. 2017. Studies on the comparative evaluation of different extraction methods and standardization of HPLC conditions for estimation of active contents in some commercially important medicinal plants. PhD Thesis. Dr Y S Parmar University of Horticulture and Forestry, Nauni, Solan.
- [23] Miura H, Ida M, Kitamura Y and Sugii M. 1978. Studies on the tissue culture of *Swertia japonica* Makino (II). A comparison of constituents between callus cultures of various organs of original plants. *Shoyakugaku Zasshi*. 32 : 90-95.
- [24] Thakur M. 2008. Studies on variation in bitter contents with plant growth and post harvest conditions in *Swertia chirayita* (Roxb. ex Flem.) Karst. PhD Thesis. Dr Y S Parmar University of Horticulture and Forestry, Nauni, Solan.

UNDER PEER REVIEW