

# Impacts of Artisanal Crude oil Refining Activities on Soil Microorganisms

## Abstract

**Aim:** To evaluate the effect of illegal crude oil refining activities on soil microorganisms using standard microbiological methods.

**Study design:** This study employs laboratory experimental design, statistical analysis of the data and interpretation.

**Place and Duration of Study:** Soil samples were taken once a month for three months (May-July, 2018) from Ke in Degema Local Government Area of Rivers State, Nigeria, where illegal crude oil refining activities are ongoing.

**Methodology:** Using standard microbiological methods, total culturable heterotrophic bacterial counts, total fungal counts, HUB and fungal counts were analysed to evaluate the effect of the activities. Total hydrocarbon content of the soil samples was also analysed.

**Results:** The populations of the total heterotrophic bacterial, fungal and hydrocarbon utilizing bacterial (HUB) and fungal (HUF) counts of the contaminated soil were enumerated. The mean total heterotrophic bacterial counts in Station 1 around the pot ranged from  $2.5 \times 10^5$  to  $1.8 \times 10^6$  cfu/g, fungal counts ranged from  $2.1 \times 10^3$  to  $4.4 \times 10^4$  cfu/g, HUB(HUB) counts ranged from  $4.2 \times 10^4$  to  $6.4 \times 10^5$  cfu/g and hydrocarbon utilizing fungal (HUF) counts ranged from  $1.5 \times 10^3$  to  $4.0 \times 10^3$  cfu/g. The results of soil samples taken 20m away from the Pot location ranged from  $7.0 \times 10^5$  to  $8.2 \times 10^6$  cfu/g for total heterotrophic bacterial counts, fungal counts ranged from  $2.3 \times 10^3$  to  $1.5 \times 10^4$  cfu/g, HUB ranged from  $4.7 \times 10^4$  to  $5.7 \times 10^5$  cfu/g and HUF ranged from  $2.0 \times 10^3$  to  $3.5 \times 10^3$  cfu/g. Also, the results of total heterotrophic bacterial counts for Station 2 ranged from;  $4.3 \times 10^5$  to  $3.3 \times 10^6$  cfu/g, fungi  $2.0 \times 10^3$  to  $3.3 \times 10^4$  cfu/g, HUB ranged from  $3.8 \times 10^4$  to  $5.4 \times 10^4$  cfu/g and HUF  $1.6 \times 10^3$  to  $3.5 \times 10^3$  cfu/g, while 20m away from the Pot total heterotrophic bacteria ranged from  $1.3 \times 10^7$  to  $6.5 \times 10^7$  cfu/g, fungi  $5.8 \times 10^3$  to  $1.4 \times 10^5$  cfu/g, HUB  $5.4 \times 10^4$  to  $1.1 \times 10^5$  cfu/g and HUF  $3.1 \times 10^3$  to  $4.7 \times 10^4$  cfu/g. While the control samples taken from inside the community where no such activity is on, ranged from  $2.6 \times 10^7$  to  $7.9 \times 10^7$  cfu/g for total heterotrophic bacteria counts, total heterotrophic fungal counts ranged from  $2.8 \times 10^4$  to  $5.3 \times 10^4$  cfu/g, HUB  $2.0 \times 10^2$  to  $3.1 \times 10^2$  cfu/g and HUF  $2.0 \times 10^1$  to  $2.3 \times 10^1$  cfu/g. twelve bacterial genera were identified and eight fungal genera: *Bacillus*, *Alcaligenes*, *Flavobacterium*, *Acinetobacter*, *Pseudomonas*, *Micrococcus*, *Proteus*, *Serratia*, *Enterobacter*, *Streptococcus*, *Escherichia*, *Staphylococcus*, *Penicillium*, *Aspergillus*, *Fusarium*, *Mucor*, *Rhizopus*, *Geotrichum*, *Candida*, and *Cladosporium*. Total hydrocarbon content range from 106 to 281mg/kg across the locations. When compared with the control, it was observed that the microbial population and diversity were adversely affected. These variations observed in the microbial population are indicative of the effect of the illegal refinery on the soil microorganisms.

37 **Conclusion:** The results of this study indicates that the continuous contamination of the soil  
38 environment by the activities of illegal crude oil refining, lead to a decrease in microbial  
39 population and diversity. This may result in devastating ecological damage, adversely affecting  
40 the ecological balance which may affect food chain and in turn animals and humans.

41 **Keywords:** illegal crude oil refining, soil bacteria, fungi, population, diversity

## 42 **Introduction**

43 The discovery and large scale production of crude oil in the Niger Delta region have exposed this  
44 region to great crude oil pollution challenge. This region in the past years have experienced the  
45 devastating effect of oil spills into both the terrestrial and aquatic environments( Chikere and  
46 Ekwuabu, 2014).This results from oil refining operations, transport, equipment failure, accident,  
47 bunkering activities and also illegal crude oil refining activities (Douglas, 2018). Research has  
48 shown that, between 200,000 – 300,000 barrels of oil are lost daily due to oil thefts out of which  
49 about 75% is sold offshore while the remaining 25% are refined locally (Obenade and  
50 Amangabara, 2014; Douglas, 2018). The soil ecosystem is directly affected since, most of these  
51 activities take place here, resulting in the discharge of crude oil and its products at various levels  
52 of refining and waste products released. These components greatly impact on plants, animals and  
53 microorganisms that depend on the nutrients in the soil for their survival. It reduces plant growth,  
54 affects aeration by blocking soil pores, thereby creating anaerobic conditions (Njoku *et al.*,  
55 2016). When crude oil is refined, various hydrocarbon fractions are produced, which have eco-  
56 toxicological impacts on the environment when spilled. These impacts include; reduction in  
57 biodiversity, changes in soil physicochemical characteristics, groundwater contamination,  
58 adverse effect on microflora, bioaccumulation and biomagnifications in environmental receptors,  
59 alteration of the habitat and cancer in humans (Obire and Anyanwu, 2009; Kalantary *et al.*,  
60 2014). Toxicity of these products varies, which depends on the concentration, composition, the  
61 prevailing environmental conditions and the biological state of the organism when the pollution  
62 occurs (Obire and Anyanwu, 2009).

63 Microorganisms play key role as indicators of the Health of aquatic and terrestrial ecosystems.  
64 This is due to their availability, abundance, their rapid growth, and ease of testing, which have  
65 made them an important tool in pollution monitoring. Microorganisms are very sensitive to  
66 changes or fluctuations in their environment, which is why they are used as microbial indicators  
67 of pollution ( Parmar *et al.*, 2016). The increased input of crude oil and petroleum products into  
68 the environment have produced an enriched microbial community, which is able to survive in  
69 such contamination (Chikere *et al.*, 2009). Microorganisms have the ability to respond to low  
70 levels of pollutants and other biological and physicochemical changes in the environment  
71 (Parmar *et al.*, 2016). The microbial communities in the soil ecosystem are responsible for food  
72 chain/web, nutrient recycling and biodegradation.

73 Research has revealed that bacteria have the highest population in the soil, and they are most  
74 adapted to use hydrocarbon as a source of carbon and energy (Das and Chandran, 2011; Chikere et

75 al., 2011; Kostka et al., 2011). Whenever, crude oil and petroleum products are spilled into the soil  
76 ecosystem, the microbial community structure is altered and diversity reduces due to  
77 environmental stress or alteration which results in the production of dominant populations within  
78 the altered communities which can withstand such contamination with improved substrate  
79 utilization and physiological abilities( Atlas and Philip, 2005; Kumar and Khanna, 2010, Chikere  
80 et al., 2009). This research was carried out to evaluate the impact of the illegal crude oil refining  
81 activities on soil microorganisms. The Ke axis of the Degema Local Government Area of Rivers  
82 State, Nigeria houses several illegal crude oil refining sites and also a market for the refined  
83 products and other oil businesses.

84

## 85 **Materials and Methods**

### 86 **Study Area**

87 The study was conducted in two illegal crude oil refinery sites (designated as Station 1 and 2) in  
88 Ke, Degema Local Government Area, of Rivers State, Nigeria. The GPS Coordinates for Station  
89 1 is Location  $04^{\circ} 45' 33.6''$  N,  $007^{\circ} 00' 01.0''$  E, and Station 2 is  $04^{\circ} 45' 33.6''$  N,  $007^{\circ} 00' 01.0''$  E as shown in Fig 1.

91

92

93

94

95

96

97

98

99

100

101

102

103


104

105

106

107

108



109 Fig 1 : location of study site in Ke, Degema Local Government Area, Rivers State, Nigeria.

### 110 **3.2. Scope of Study**

111 This study was carried out between May and July, 2018. Soil samples were collected at about 0-  
112 15cm depth using a soil auger into sterile bags, from four different points around the Pots. Pot  
113 here refers to the fabricated aluminum tanks used in the distillation process. For Station 1 it is

114 designed as Pot 1 and soil samples bulked for homogeneity. Then, 20m away from the pot a  
115 second set of soil samples were also taken. Same was done for Station 2, soil samples were taken  
116 around the pot and 20m away from the pot (Pot 2). Control soil samples were taken inside the  
117 community, away from the illegal refining sites. These samples were labeled properly and  
118 immediately transported to the laboratory for analyses.

119

## 120 **Enumeration of Total Heterotrophic Bacteria**

121 To determine the total heterotrophic bacterial counts spread plate method was used on nutrient  
122 agar. One gram of soil was taken from each soil sample and homogenized in 9mls of  
123 physiological saline. An aliquot of 0.1ml of the dilutions of  $10^{-4}$  and  $10^{-5}$  were plated out on the  
124 surface of the agar and evenly spread using a sterile hockey stick (spreader/inoculator). Plates  
125 were incubated at  $30^{\circ}\text{C}$  for 24 hours. The colonies that developed on the plates were counted and  
126 mean calculated for duplicate plates, results expressed in colony forming unit per gram (CFU/g)  
127 (Douglas and Green, 2015).

## 128 **Enumeration of the Hydrocarbon Utilizing Bacterial Population**

129 HUBpopulations in the soil samples were enumerated using mineral salt agar. The vapour phase  
130 transfer method using Mineral salt medium composition of Mills *et al.*, 1978 was used as  
131 modified by Okpokwasili and Okorie (1988). Aliquot (0.1ml) of the  $10^{-4}$  to  $10^{-5}$  dilutions,  
132 previously obtained during the serial dilution of the soil samples, were inoculated in duplicates  
133 on appropriately labeled mineral salt agar plates which was freshly prepared and dried. The  
134 vapour phase transfer method in which a sterile Whatman No. 1 filter paper, placed on the lid of  
135 the Petri plate is saturated with 5ml Bonny light crude oil. Plates were inverted and incubated for  
136 7days at  $30^{\circ}\text{C}$ . Colonies were counted after incubation, average counts calculated for duplicate  
137 plates and expressed as colony forming unit (CFU/g).

## 138 **Enumeration of Total Heterotrophic Fungi**

139 Spread plate method was used on Sabouraud Dextrose agar (SDA). An aliquot, 0.1ml of  $10^{-3}$  and  
140  $10^{-4}$  dilutions were inoculated onto the freshly prepared SDA plates, in which 0.5% Ampicillin  
141 has been added. This was done to inhibit bacterial growth while allowing the growth of fungi  
142 (Cheesbrough, 2000). The inoculum was spread evenly using sterile hockey stick. Plates were  
143 inverted and incubated at  $28^{\circ}\text{C}$  for 5days. Colonies that developed on the plates were counted,  
144 average counts on duplicate plates calculated and recorded as cfu/g.

145

## 146 **Enumeration of the Hydrocarbon Utilizing Fungi**

147 The MSA as composed by Mills *et al.*, 1978 as modified by Okpokwasili and Okorie, (1988) to  
148 which 5% tetracycline was added to prevent bacterial growth was used. This medium was used  
149 for the isolation and enumeration of hydrocarbon utilizing fungi. From dilutions of  $10^{-3}$  and  $10^{-4}$ ,  
150 0.1ml aliquot was transferred on the freshly prepared plates; evenly spread using the hockey  
151 stick. The vapour phase transfer method in which a sterile Whatman No. 1 filter paper, placed on

152 the lid of the Petri plate is saturated with 5ml Bonny light crude oil. Plates were inverted and  
153 incubated for 7days at 30°C. Colonies that developed on the plates after incubation were counted,  
154 average counts calculated for duplicate plates and expressed as colony forming unit (CFU/g) of  
155 soil.

156

### 157 **Purification and characterization of Organisms**

158 Discreet colonies that developed on the Nutrient and Mineral Salt agar plates were subcultured  
159 by streaking on Nutrient agar until pure cultures were obtained. Colonies on SDA and MSA for  
160 fungi were also subcultured by streaking on SDA until pure cultures were obtained. Pure cultures  
161 of bacterial and fungal isolates were preserved in bijoux bottles containing nutrient and SDA  
162 slants respectively. The pure isolates were then refrigerated and were used for other analyses.  
163 The pure bacterial isolates were further investigated by carrying out routine microbiological  
164 analyses including; cultural and biochemical characteristics. The following test were done; Gram  
165 staining, cell motility, oxidase, indole and catalase production, citrate utilization, methyl Red-  
166 Voges Proskauer test, acid/gas production from sugar fermentation, as described by Bergey's  
167 Manual for Determinative Bacteriology(Holt *et al.*, 1994).

### 168 **Identification of Fungal Isolates**

169 The fungal isolates were identified basically by both macroscopic and microscopic examination.  
170 Macroscopic identification was done by observing the morphology of the pure cultures in the  
171 plates. The microscopy was done by removing a small portion and placing on clean grease free  
172 slide. Lactophenol blue was dropped on the slide and smeared, it was covered using a cover slip  
173 and viewed under x10 and x40 objective lens (Cheesebrough, 2000). The observed  
174 characteristics were recorded and compared with the identification key in Barnett and Hunter,  
175 (1972).

### 176 **Determination of Total Hydrocarbon Content (THC)**

177 Total Hydrocarbon Content (THC) analyses were carried out on all soil samples using  
178 spectrophotometric method. The total hydrocarbon content of the soil samples were determined  
179 by shaking 10g of a representative soil sample with 20ml xylene and the oil extracted determined  
180 by measuring the absorbance using a spectrophotometer at 420 nm using a spectronic 20. A  
181 standard curve of the absorbance of different concentrations of hydrocarbon concentration in the  
182 soil sample was measured in g/g after reference to a standard curve and multiplying by the  
183 appropriate multiplication factor (Nrior *et al.*, 2017).

184

### 185 **Results**

186 The effect of illegal crude oil refining activities on soil microorganisms was investigated. The  
187 mean total heterotrophic bacterial counts in Station 1 around the pot ranged from  $2.5 \times 10^5$  to  $1.8$   
188  $\times 10^6$  cfu/g, fungal counts ranged from  $2.1 \times 10^3$  to  $4.4 \times 10^4$  cfu/g, HUB counts ranged from  $4.2 \times$   
189  $10^4$  to  $6.4 \times 10^5$  cfu/g and hydrocarbon utilizing fungal (HUF) counts ranged from  $1.5 \times 10^3$  to  $4.0$   
190  $\times 10^3$  cfu/g. The results of soil samples taken 20m away from the Pot location ranged from  $7.0 \times$

191  $10^5$  to  $8.2 \times 10^6$ cfu/g for total heterotrophic bacterial counts, fungal counts ranged from  $2.3 \times 10^3$   
192 to  $1.5 \times 10^4$ cfu/g, HUB ranged from  $4.7 \times 10^4$  to  $5.7 \times 10^5$ cfu/g and HUF ranged from  $2.0 \times 10^3$   
193 to  $3.5 \times 10^3$ cfu/g. Also, the results of total heterotrophic bacterial counts for Station 2 ranged  
194 from;  $4.3 \times 10^5$  to  $3.3 \times 10^6$ cfu/g, fungi  $2.0 \times 10^3$  to  $3.3 \times 10^4$ cfu/g, HUB ranged from  $3.8 \times 10^4$   
195 to  $5.4 \times 10^4$ cfu/g and HUF  $1.6 \times 10^3$  to  $3.5 \times 10^3$ cfu/g, while 20m away from the Pot total  
196 heterotrophic bacteria ranged from  $1.3 \times 10^7$  to  $6.5 \times 10^7$ cfu/g, fungi  $5.8 \times 10^3$  to  $1.4 \times 10^4$ cfu/g,  
197 HUB  $5.4 \times 10^4$  to  $1.1 \times 10^5$ cfu/g and HUF  $3.1 \times 10^3$  to  $4.7 \times 10^4$ cfu/g. While the control samples  
198 taken from inside the community where no such activity is on, ranged from  $2.6 \times 10^7$  to  $7.9 \times$   
199  $10^7$ cfu/g for total heterotrophic bacteria counts, total heterotrophic fungal counts ranged from  $2.8$   
200  $\times 10^4$  to  $5.3 \times 10^4$ cfu/g, HUB  $2.0 \times 10^2$  to  $3.1 \times 10^2$ cfu/g and HUF  $2.0 \times 10^1$  to  $2.3 \times 10^1$ cfu/g.  
201 Mean values of counts are showed in Table 1.

202 Figures 1 and 2 reveal the distribution of the various group of organisms identified during the  
203 period. In this study, twelve bacterial genera identified from the control site include: *Bacillus*,  
204 *Alcaligenes*, *Flavobacterium*, *Acinetobacter*, *Pseudomonas*, *Micrococcus*, *Proteus*, *Serratia*,  
205 *Enterobacter*, *Streptococcus*, *Escherichia*, and *Staphylococcus*. The following eight fungal  
206 genera were identified from the control: *Penicillium*, *Aspergillus*, *Fusarium*, *Mucor*, *Rhizopus*,  
207 *Geotrichum*, *Candida*, and *Cladosporium*. The most predominant bacterial species in the  
208 uncontaminated soil sample were *Bacillus*, *Acinetobacter*, and *Pseudomonas* species. The  
209 hydrocarbon utilizing bacteria from the control site include: *Bacillus*, *Pseudomonas*, and *Serratia*  
210 species. In this study, the microbial diversity between the uncontaminated soil and the  
211 contaminated soil samples were recorded (Table 2). In Station 1, the bacteria isolated around the  
212 Pot were; *Flavobacterium* sp, *Micrococcus* sp, *Bacillus* sp, *Pseudomonas* sp and *Acinetobacter*  
213 sp; and *Penicillium* sp, *Mucor* sp, *Rhizopus* sp and *Aspergillus* sp were the fungi isolated around  
214 Pot 1. *Bacillus*, *Pseudomonas* and *Acinetobacter* sp, *Mucor*, *Rhizopus*, *Penicillium*, *Fusarium* and  
215 *Aspergillus* sp were isolated 20 meters away from Pot 1. Station 2 Pot 2, had the following  
216 bacterial genera: *Serratia*, *Micrococcus*, *Bacillus*, *Pseudomonas* and *Acinetobacter* whereas  
217 *Penicillium*, *Aspergillus*, *Rhizopus*, were the fungal genera. Except *Acinetobacter* sp and  
218 *Micrococcus* sp, bacterial genera isolated from Station 2, Pot 2 were also isolated 20 meters  
219 away from the Pot. Also *Penicillium*, *Mucor* and *Aspergillus* were the fungal genera isolated  
220 from Station 2 Pot 2, which was slightly different from *Mucor*, *Rhizopus*, *Penicillium* and  
221 *Aspergillus* sp isolated 20 meters away from the Station 2, Pot 2. The results of total hydrocarbon  
222 contents range from 106 – 281 mg/kg across the sampling locations. The highest value of 281  
223 mg/kg was observed in the month of May around Pot 1, the least value of 106mg/kg was  
224 observed for Pot 2 in the month of July. It was observed that the concentration decreased across  
225 the stations during the sampling period. This may be due to surface runoff as a result of the rains  
226 since that is the peak of the rainy season.

## 227 Discussion

228 The impact of illegal crude oil refining activities on soil microbes were determined by the  
229 enumeration of total heterotrophic bacterial, total heterotrophic fungal, HUB and fungal counts



230 presented in Table 1. This observation could be attributed to the presence of vegetation cover,  
231 high nutrient content (especially nitrogen and phosphorus) as a result of decomposition of  
232 organic materials to release nutrients and other environmental factors required for the survival of  
233 these microorganisms in the soil (Gougoulas et al., 2014). Counts observed in this study is similar  
234 to that obtained by previous researchers in contaminated soil Obire and Anyawu 2009. The  
235 continuous refining activities releases crude oil and petroleum products into the soil, resulting in  
236 pollution which could be inhibitory to certain group of organisms while it becomes an enriched  
237 microbial community for the other group capable of survival in such contaminations (Obire and  
238 Anyanwu, 2009; Douglas, 2018). From the results obtained, it was observed that the difference  
239 between the THBC and HUB was not significant which means that most of the organisms found  
240 in the contaminated locations are hydrocarbon utilizing microorganisms which are capable of  
241 using these contaminants as a source of carbon.

242 Soil has been reported as a suitable medium that aids the growth and survival of microorganisms,  
243 but the introduction of these contaminants retard the activities of these organisms, thus giving  
244 room to organisms that have the ability of metabolizing such products and limiting the growth of  
245 non-metabolizers of the products (Chikere and Ekwuabu, 2014). Lower microbial counts were  
246 observed in the samples around the pots from both stations. This observation apart from the  
247 contamination from the crude oil and petroleum products may be attributed to the heat used for  
248 the distillation process. This is in conformity with previous studies who have reported that  
249 temperatures that exceed 70-80°C are capable of killing many soil microbes and that non-spore  
250 forming fungi will be killed at 70°C (Pattison *et al.*, 2009).

251  
252 Results of this study also show that microbial diversity was also affected by the oil refining  
253 activity. Douglas (2018) has also reported that higher concentrations of the illegal refined crude  
254 oil deposit lead to a uniform reduction in species diversity and population of soil fungi over time.  
255 Thus, continuous dumping of *kpo-fire* residue into the terrestrial environment would impact  
256 negatively on the crucial role played by these groups of organisms in decomposition and  
257 interfere with other metabolic activities of the organisms in the environment.

258 The fungal isolates in this study have been reported by previous scholars to be capable of  
259 metabolizing or utilizing crude oil pollutants (Obire *et al.*, 2008, Douglas, 2018). Also, Obire and  
260 Anyanwu (2009), in a previous study of soil samples contaminated with crude oil had isolated  
261 fourteen fungi genera belonging to *Alternaria* sp., *Aspergillus* sp., *Cephalosporium* sp.;  
262 *Cladosporium* sp.; *Fusarium* sp., *Geotrichum* sp., *Mucor* sp.; *Penicillium* sp.; *Rhizopus* sp.  
263 *Trichoderma* sp., *Candida* sp., *Rhodotolura* sp., *Saccharomyces* sp. and *Torulopsis* sp from the  
264 control soil, with five hydrocarbon utilizing fungi identified out of the fourteen. But in this study,  
265 *Alternaria* sp, *Cephalosporium* sp.; *Geotrichum* sp.; *Rhodotolura* sp.; *Trichoderma* sp, and  
266 *Fusarium* sp were not identified. The hydrocarbon utilizing bacteria identified by this study has  
267 been shown to have the ability to utilize crude oil as carbon source (Chikere and Ekwuabu, 2014;  
268 Douglas and Green, 2015).

269

270

## 271 **Conclusion**

272 This research has shown that the illegal crude oil refining activities has increased the quantities  
273 of crude oil, petroleum products and residue (waste) into the soil environment, with the  
274 accompanying heat, used for the distillation process greatly affecting both microbial load and  
275 diversity in the soil environment. The refining activities could exert a negative impact on the  
276 population, diversity as well as the activities of soil microorganisms. Since, the microbial  
277 diversity is important for soil health, community structure and functions. Thus, the continuous  
278 exposure of the soil to the indiscriminate illegal refinery activities, will not only hamper the  
279 texture or structure of the soil but, would also lead to a decline in microbial populations which  
280 could pose a serious threat to the food chain, decomposition, nutrient recycling, bioremediation  
281 and the ecological balance.

282

## 283 **Disclaimer:**

284

285 This paper is based on preliminary dataset. Readers are requested to consider this paper as  
286 preliminary research article, as authors wanted to publish the initial data as early as possible.  
287 Authors are aware that detailed statistical analysis is required to get a scientifically established  
288 conclusion. Readers are requested to use the conclusion of this paper judiciously as statistical  
289 analysis is absent. Authors also recommend detailed statistical analysis for similar future studies.

290

## 291 **Conflict of Interests**

292 The author(s) have not declared any conflict of interests.

## 293 **References**

294 American Public Health Association (APHA) (2008) Standard Methods for the Examination of  
295 Water and Waste Water. 21st ed. Washington DC.

296

297 Atlas, R. M. and Philp, J (2005) Bioremediation: applied microbial solutions for real-world  
298 environmental cleanup. American Society for Microbiology(ASM) press, Washington, DC, pp  
299 78-105

300 Aparna C, Saitha P, Himabindu V, Alok B, Anjaneyulu Y (2010). Evaluation of ioremediation  
301 effectiveness on sediments Contaminated with industrial wastes. International Journal Of  
302 Environmental Sciences. Volume 1: (4).

303 Barnett, H. L. and Hunter, B. B. (1972). Illustrated Genera of Fungi Imperfecti. 3<sup>rd</sup> Ed., Burgess  
304 Publication Co.

305 Cheesebrough, M. (2000) District Laboratory Practice in Tropical Countries. Part 2, Cambridge  
306 University Press, London, UK. Pp 143 – 156.



- 307 Douglas, S. I. and Green, D. I. (2015) Microbial Communities Found in Diesel Contaminated  
308 Soil. *International Journal of Physical & Applied Sciences*. 2(4): 38 – 48.
- 309 Chikere, C. B. and Ekwuabu, C. B. (2014). Culture-dependent characterization of hydrocarbon  
310 utilizing bacteria in selected crude oil-impacted sites in Bodo, Ogoniland, Nigeria. *African*  
311 *Journal of Environmental Science & Biotechnology*. 8(6): 401-406.
- 312 Chikere, C. B., Okpokwasili, G. C. and Ichiakor, O.(2009) Characterization of hydrocarbon  
313 utilizing bacteria in Tropical marine Sediments. *African Journal of Biotechnology*. 8(11): 2541-  
314 2544
- 315 Douglas, S. I. (2018). Effect of illegally Refined Crude Oil (Kpo-Fire) Residue on Soil Fungi.  
316 *International Journal of current microbiology and Applied Sciences*. 7(12):3309-3316.
- 317 Holt JG, Krieg NR, Sneath PHA, Stanley JT, Williams ST (1994). *Bergey's Manual of*  
318 *Determinative Bacteriology*. 9th Ed. Williams and Wilkins Company, Baltimore, USA. pp. 71-  
319 561.
- 320 Kalantary, R. R., Mohseni-Banpi, A., Esrafile, A., Nasser, S., Ashmogh, F.F., Jorfi, S. and  
321 Jafari, M.(2014) Effectiveness of biostimulation through nutrient content on the bioremediation  
322 of phenanthrene contaminated soil. *Journal of Environmental Health Science and Engineering*.  
323 12(143): 1-9.
- 324 Kumar, M. and Khanna, S. (2010) Diversity of 16S rRNA and dioxygenase genes detected in  
325 coal tar-contaminated site undergoing active bioremediation. *Journal of Applied Microbiology*  
326 108: 1252 – 1262.
- 327 Mills, A. L., Breuil, C. and Colwell, R. R. (1978) Enumeration of Petroleum Degrading marine  
328 and estuarine Microorganisms by most probable number Method. *Canadian Journal of*  
329 *Microbiology*.12:234-248.
- 330 Njoku, J. D., Ebe, T. E. and Akachukwu, O. E. (2016). Analysis of Heavy Metal Contamination  
331 by Artisanal Refining Plants in The Niger Delta Region, Southern Nigeria. *British Journal of*  
332 *Environmental Sciences*. 4(3): 39-48.
- 333 Citation: Renner R. Nrior, Salome Ibietela Douglas and Martha Tamunoemi Ojongo, 2017.  
334 "Effect of wastewater on biodegradability of drilling fluid", *International Journal of*  
335 *Development Research*, 7, (12), 17872-17876.
- 336 Obenade, M, and Amangabara, G. T. (2014) Perspective: The Environmental Implications of Oil  
337 Theft and Artisanal Refining in the Niger Delta Region. *Asian Review of Environmental and*  
338 *Earth Sciences*, 1(2): 25 - 29
- 339 Obire, O. and Anyanwu, E. C. (2009) Impact of various Concentrations of Crude Oil on Fungal  
340 Populations of Soil. *International Journal of Science & Technology*. 6(2): 211-218.

341 Okpokwasili, G. C. and Okorie, B. B. (1988), Biodegradation Potentials of Microorganisms  
 342 Isolated From car engine lubricating oil. *Tribol International* 21(4): 215 -220.

343 Parmar, T. K., Rawtani, D., and Agrawal, Y. K. (2016) Bioindicators: the natural indicator of  
 344 environmental pollution. *Frontiers in Life Science*, 9(2): 110 -118.

345 Pattinson, G.S., K.A. Hammill, B.G. Sutton and P.A. McGee (2009) Simulated fire reduces the  
 346 density of arbuscular mycorrhizal fungi at the soil surface. *Mycological Research journal*, 103:  
 347 491-496.

348 United Nations Environmental Programme (UNEP) 2011. [www.unep.org](http://www.unep.org). Accessed 12 June,  
 349 2012.

350 Nilanjana Das and Preethy Chandran, "Microbial Degradation of Petroleum Hydrocarbon Contaminants:  
 351 An Overview," *Biotechnology Research International*, vol. 2011, Article ID 941810, 13 pages, 2011.

352 Chikere, C. B., Okpokwasili, G. C., & Chikere, B. O. (2011). Monitoring of microbial hydrocarbon  
 353 remediation in the soil. *3 Biotech*, 1(3), 117-138.

354 Kostka, J. E., Prakash, O., Overholt, W. A., Green, S., Freyer, G., Canion, A., ... & Huettel, M. (2011).  
 355 Hydrocarbon-degrading bacteria and the bacterial community response in Gulf of Mexico beach sands  
 356 impacted by the Deepwater Horizon oil spill. *Applied and environmental microbiology*, AEM-05402.


357 Gougoulas, C., Clark, J. M., & Shaw, L. J. (2014). The role of soil microbes in the global carbon cycle:  
 358 tracking the below-ground microbial processing of plant-derived carbon for manipulating carbon  
 359 dynamics in agricultural systems. *Journal of the science of food and agriculture*, 94(12), 2362-71.

360  
 361  
 362  
 363

364 Table 1: Mean Microbial Counts from the Sampling Site

Sample Location	THBC(cfu/g)	THFC(CFU/G)	HUB(CFU/G)	HUFC(CFU/G)
Station 1(pot)	$2.3 \times 10^5$	$2.6 \times 10^3$	$2.2 \times 10^4$	$2.3 \times 10^3$
20m away	$7.1 \times 10^6$	$1.9 \times 10^4$	$7.2 \times 10^4$	$3.0 \times 10^3$
Station 2(Pot)	$8.8 \times 10^5$	$5.0 \times 10^4$	$5.9 \times 10^4$	$1.0 \times 10^4$
20m away	$3.6 \times 10^6$	$3.8 \times 10^5$	$3.3 \times 10^4$	$3.3 \times 10^3$
Control	$7.8 \times 10^7$	$4.1 \times 10^5$	$2.7 \times 10^2$	$2.2 \times 10$

365



366

367

368

369

370

371


372

373

374

375

376



377

378 Table 2: Microbial Diversity From the Sampling Locations

THB(control)	THB	HUB	THF	HUF
<i>Bacillus</i> sp	<i>Micrococcus</i> sp	<i>Pseudomonas</i> sp	<i>Aspergillus niger</i>	<i>Penicillium</i> sp
<i>Klebsiella</i> sp	<i>Bacillus</i> sp	<i>Micrococcus</i> sp	<i>Aspergillus flavus</i>	<i>Apergillus</i> sp
<i>Pseudomonas</i> sp	<i>Enterobacter</i> sp	<i>Acinetobacter</i> sp	<i>Cladosporium</i> sp	<i>Fusarium</i> sp
<i>Serratia</i> sp	<i>Micrococcus</i> sp	<i>Bacillus</i> sp	<i>Penicillium</i> sp	<i>Rhizopus</i> sp
<i>Enterobacter</i> sp	<i>Acinetobacter</i> sp	<i>Proteus</i> sp	<i>Fusarium</i> sp	<i>Mucor</i> sp
<i>Micrococcus</i> sp	<i>Flavobacterium</i> sp	<i>Serratia</i> sp	<i>Rhizopus</i> sp	<i>Cladosporium</i> sp
<i>Flavobacterium</i> sp	<i>Serratia</i> sp	<i>Flavobacterium</i> sp	<i>Geotrichum</i> sp	
<i>Proteus</i> sp	<i>Alcaligenes</i> sp		<i>Mucor</i> sp	
<i>Acinetobacter</i> sp	<i>Proteus</i> sp			
<i>Escherichia coli</i>				
<i>Alcaligene</i> sp				
<i>Streptococcus</i> sp				

379 Key: THB total heterotrophic bacteria, HUB Hydrocarbon utilizing bacteria, hydrocarbon  
 380 utilizing fungi