

**Insect Pest Diversity and Damage Assessment In
Field Grown Okra (*Abelmoschus esculentus* (L.)
Moench) In The Coastal Savannah Agro-Ecological
Zone Of Ghana.**

ABSTRACT

Aims: The specific objectives of this study were: to identify the diversity of insect species associated with ten okra cultivars, and to assess the abundance of the insect species and the extent of leaf damage during vegetative, flowering and fruiting stages of ten okra cultivars under field conditions.

Study design: The experimental treatments were deployed in a Randomized Complete Block Design (RCBD), replicated four times.

Place and Duration of Study: The research was conducted at Nuclear Agriculture Research Center (NARC) farms and the laboratories of Radiation Entomology and Pest Management Center (REPMC) of Biotechnology and Nuclear Agriculture Research Institute (BNARI), between July 2017 and March 2018. The study area is located at Kwabenya, Accra on latitude 5° 40' N, longitude 0° 13' W with Ochrosol (Ferric Acrisol) soil type, derived from quartzite Schist.

Methodology: Plant materials used for the study consisted of five local and five exotic okra cultivars. The local cultivars were Asutem (AS), Togo (TG), Labadi dwarf (LD), Kwab (K1) and Adom (AD). These were obtained from the market (Asamankese and Dome) and okra farmers' fields. The exotic cultivars were Lucky 19F1 (LF1), F1 Kirene (F1K), F1 Sahari (F1S), Kirikou F1 (KF1) and Clemson Spineless (CS). These cultivars were obtained from a commercial seed shop, Technisem, Accra. Land preparation of the research site involved plowing and harrowing. The prepared land was lined and pegged into 40 plots using a Randomized Complete Block Design with four replications. Each replicate measured 35m x 7m and separated by 2m from each other with 10 subplots within a block. Each subplot measured 3m x 3m and spaced from one another by 1m. The total size of the experimental area was 646m². The okra seeds were manually sown to a depth of 2cm directly at a spacing of 0.50m x 0.60m. Four seeds per hill were sown and later thinned to one seedling per hill after emergence. Field management practices such as weed control and watering were done as and when required. Data on insects were collected from five okra plants randomly selected from the middle rows. Okra leaves were carefully examined by observing both the abaxial and adaxial surfaces. Insects found on the surfaces of the leaves were identified, counted manually and recorded as either major or minor based on their incidence pattern. Data was taken daily because the ten cultivars have different vegetative, flowering and fruiting dates. Insects were counted between the hours of 6.00 am and 8.00 am when they are inactive and cannot fly. In order to determine the extent of leaf

damage, the following described scoring scale was designed for this work. Leaf damage was determined by counting the total number of perforations created by the insects in all leaves found on the five randomly selected test plants. This was then divided by the total number of leaves on the five selected test plants to obtain the average number of perforations per leaf. Leaves were visually assessed and scored for severity of damage using a damage rating where; **1** very mild damage (1 to 15 perforations); **2** mild damage (16 to 30 perforations); **3** moderately severe damage (31 to 45 perforations); **4** very severe damage (46 to 60 perforations); **5** extremely severe damage (more than 60 perforations).

Results: A total of thirteen insect **pests** belonging to six orders (Coleoptera, Homoptera, Lepidoptera, Hymenoptera, Orthoptera and Mantodea), and thirteen families Chrysomelidae, Coccinellidae, Pyrgomorphidae, Meloidae, Noctuidae, Nolidae, Cicadellidae, Aleyrodidae, Aphididae, Pseudococcidae, Mantidae, Formicidae, and Acrididae) were found to be abundant **in** the field. Among these, the highest number of insect species belonged to Homoptera group **viz.**, Green Peach Aphid (*Myzus persicae*) **Okra** leafhopper (*Amrasca biguttula*), Whitefly (*Bemisia tabaci*), and striped **mealybug** (*Ferrisia virgata*) followed by Coleoptera (Flea beetle (*Podagrica* sp.) and **Ladybird** beetle (*Cheilomenes lunata*). On the vegetative stage of the **okra**, Flea beetle had the highest number on Lucky 19F1 (36.00±9.66 insects/plant). During the flowering stage, **plants** of L-19F1 had the highest mean number of Flea beetles (32.25±10.30 insects/plant). On the fruiting stage, **plants** of LD had the highest mean abundance of flea beetles (47.50±13.53 per plant).

Conclusion: A total of 1,439 insects were recorded at the fruiting stage which was significantly higher than the flowering (855) and vegetative stages (693). Mean Whitefly counts **were** relatively low at the vegetative, flowering and fruiting stages of the cultivars. However, Flea beetle (*Podagrica* sp.) and Green Peach aphids (*Myzus persicae*) mean numbers increased progressively throughout all the stages. In the present study, **the severity** of leaf damage was significantly higher at the fruiting stage compared with the flowering and vegetative stages. Plants of cultivars LD and AS were the most promising recording the least leaf damage (111.95) and (119.10) respectively.

13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

Keywords: Okra, Abundance, Severity, Numbers, Stages, Accessions

1. INTRODUCTION

Okra (*Abelmoschus esculentus* (L.) Moench) is an annual cultivated mainly for the fresh fruits that are consumed immature as a vegetable in a variety of ways. The crop also serves important medicinal and industrial purposes. Besides, the cultivation, processing and marketing of okra present opportunities for income generation among rural small-holder farmers. All over the growing regions of the tropics and subtropics, production of the crop faces serious challenges of insect pest infestation and viral diseases responsible for considerable yield loss. Consequently, in West Africa in particular, the average yield has been very low at an estimated 2.5 t/ha [1].

30 In Ghana, low yields in okra are attributed to several production constraints among
31 which low soil fertility and damage caused by insect pests are most critical [2].
32 Damage caused by insect pests has been reported as the major constraint in the
33 production of okra [3;4]. Tindal [5] reported several insect species infesting okra in
34 Ghana. These include Leaf roller, *Sylepta derogata* (F.), Cotton stainer, *Dysdercus*
35 *superstitiosus* (F), Cotton aphid, *Aphis gossypii* (Glov.) and flea beetle, *Podagrica*
36 *uniformis* (Jac.). Critchley [4] reported of 22 insect pests from 12 families in four
37 orders (Coleoptera, Hemiptera, Lepidoptera and Orthoptera) infesting okra in Brong-
38 Ahafo region of Ghana. Of these, the most important are the *Podagrica uniformis*
39 Jacoby and *P. sjostedti* Jacoby, followed by *Aphis gossypii* Glover, *Dysdercus*
40 *superstitiosus* (Fab) and *Sylepta derogata* (Fab.) and *Heliothis armigera* (Hub.). The
41 blister beetle, *Mylabris* spp., feeds on the flowers, reducing the number of fruits
42 formed, while both adults and nymphs of *A. gossypii* suck sap from young leaves
43 and buds, thus reducing the efficiency of the leaves [6; 7].
44

45 Two flea beetle species, *P. uniformis* and *P. sjostedti* are responsible for heavy leaf
46 damage of the crop [8]. Extensive leaf damage in the form of feeding holes on the
47 leaves results in a significant reduction of the photosynthetic ability of the plant. The
48 insect pests also feed on fruits, stems, and flowers culminating in poor crop
49 performance and low yields. The *Podagrica* species have also been implicated in the
50 transmission of okra mosaic virus [9;10]. The other insect pests of economic
51 importance in okra production, whiteflies (*Bemisia tabaci*) feed on plant sap and
52 cause the okra leaf curl and yellow vein mosaic diseases [9;11]. The species
53 diversity of insects and their pest status varies from region to region with the
54 variation in agro-climatic conditions. Asare-Bediako *et al.* [12] indicated that there is
55 always a phenomenon of the continual significant increase in insect populations
56 globally. In Ghana, for instance, rising insect pest populations have been attributed
57 to poor agronomic practices such as the use of untreated seeds for cultivation and
58 the continuous practice of mono-cropping by the majority of local farmers in order to
59 meet the increasing demand of the various staples in the country [12].
60

61 As a prerequisite for putting in place an effective integrated insect pest management
62 regime, it is necessary to properly identify which species of insects are major pests,
63 establish their diversity, abundance, and severity of damage they cause to plant
64 parts. It is, therefore, worrying that information on insect pests of okra at its various
65 growth stages in the coastal savannah agro-ecological zone of Ghana is lacking.
66 Thus a specific objective was set out to identify the diversity of insect species
67 associated with ten cultivars of okra, to assess the abundance of the insect species
68 during the vegetative, flowering and fruiting stages and the extent of leaf damage
69 under field conditions. Such information will help farmers to know which insect pests
70 to target, the best time to target control practices and the appropriate approach to
71 use. In addition, the findings of this work will serve as a useful guide in the
72 development of an effective pest management system for okra production,
73 particularly within the coastal savannah agro-ecological zone.
74

75 **2. MATERIAL AND METHODS**

76

77 **2.1 Soil and Rainfall Pattern of Study Area**

78 The research was conducted in Biotechnology and Nuclear Agriculture Research
79 Institute (BNARI) of the Ghana Atomic Energy Commission (GAEC) between July
80 2017 and March 2018. The study area is located at Kwabenya, Accra on latitude 5°
81 40' N, longitude 0° 13' W with Ochrosol (Ferric Acrisol) soil type, derived from
82 quartzite Schist [13]. The maximum and minimum average temperatures for the
83 period of study were 30.7°C and 26.0°C respectively with **an average** annual rainfall
84 of 220mm. The highest and lowest relative humidity is between 75 and 60% [14;15].
85 The experimental site is also well drained and has an elevation of 76 m above sea
86 level within the coastal savannah agro-ecological zone.
87

88 **2.2 Plant materials and field design**

89 Plant materials used for the study consisted of five local and five exotic okra
90 cultivars. The local cultivars were Asutem (AS), Togo (TG), Labadi dwarf (LD),
91 Kwabenya (K1) and Adom (AD). These were obtained from Markets (Asamankese
92 and Dome) and okra farmers' fields. The exotic cultivars were Lucky 19F1 (LF1), F1
93 Kirene (FIK), FI Sahari (FIS), Kirikou F1 (KFI) and Clemson Spineless (CS). These
94 cultivars were obtained from a commercial seed shop, Technisem, Accra. The land
95 was **plowed, harrowed and lined** and pegged into 40 plots using a Randomized
96 Complete Block Design with four replications. Each **replicates** measured 35m x 7m
97 and separated by 2m from each other with 10 subplots within a block. Each subplot
98 measured 3m x 3m and spaced from one another by 1m. **The total** size of the
99 experimental area was 646m². The okra seeds were manually sown to a depth of
100 2cm directly at a spacing of 0.50m x 0.60m. Four seeds per hill were sown and later
101 thinned to one seedling per hill after emergence. Field management practices such
102 as weed control and watering were done as and when required.
103

104 **2.3 Data collection**

105 **2.3.1 Insect abundance**

106 Insect abundance was estimated from five okra plants randomly selected from the
107 middle rows. Okra leaves were carefully examined by observing both the abaxial and
108 ad axial surfaces. Insects found on the surfaces of the leaves were identified,
109 counted and recorded as either major or minor based on their incidence pattern.
110 Data was taken daily between the hours of 6.00 am and 8.00 am.
111

112 **2.3.2 Determination of leaf damage**

113 In order to determine the extent of leaf damage, a scoring scale was designed for
114 this work. Leaf damage was determined by estimating the total number of
115 perforations created by the insects in all leaves found on each of the five randomly
116 selected test plants. This was then divided by the total number of leaves on the plant
117 to obtain the average number of perforations per leaf. The five-point scoring scale is

118 as follows: **1** very mild damage (1 to 15 perforations); **2** mild damage (16 to 30
119 perforations); **3** moderately severe damage (31 to 45 perforations); **4** very severe
120 damage (46 to 60 perforations); **5** extremely severe damage (more than 60
121 perforations) (Figure 1).

122

123

124

125

126

127

128

129

130

131



132 *Figure 1: Rating of the severity of leaf damage*

133

134

2.3.3 Data analysis

135

136

137

138

139

140

141

142

143

144

145

146

3. RESULTS AND DISCUSSION

147

3.1 Diversity of insect species associated with okra under field conditions.

148

149

Insects present on the ten cultivars of okra under open field conditions observed during the growth stages of the crop are shown in figure 2.




Fig. 2. Some insect species identified in the studied okra cultivars; A = Variegated grasshopper, B = Ants, C = Cotton stainer, D = Lady bird beetle, E = Flea beetle, F = Aphid.

Results of the study revealed that all ten cultivars of okra were susceptible to insect pest infestation (Table 1). A total of thirteen insect species belonging to six orders (Coleoptera, Lepidoptera, Homoptera, Hemiptera, Hymenoptera, Mantodea and Orthoptera), and twelve families namely Chrysomelidae, Coccinellidae, Pyrgomorphidae, Meloidae, Noctuidae, Nolidae, Cicadellidae, Aleyrodidae, Aphididae, Pseudococcidae, Mantidae and Formicidae) were recorded at the study site. The highest number of insect species belong to the order Homoptera and included Okra leafhopper (*Amrasca biguttula*), Whitefly (*Bemisia tabaci*), and Green Peach aphid (*Myzus persicae*). Those found in order Coleoptera were Flea beetle (*Podagrica* sp.), Blister beetle (*Mylabris pustulata*) and ladybird beetle (*Cheilomenes lunata*). The Striped mealy bug (*Ferrisia virgata*) and the Black carpenter ants (*Camponotus* sp were the only insects found in the order Hemiptera and Hymenoptera respectively.). Members of the order Orthoptera identified in this study included the Variegated grasshopper (*Zonocerus variegatus*) and Tobacco grasshopper (*Atractomorpha crenulata*). The order Mantodea had Praying mantis (*Mantis religiosa*) while Lepidoptera included Cotton Semilooper (*Anomis flava*) and Transverse Moth (*Xanthodes transversa*). The Flea beetle and Aphids were the most abundant insect pests. The insect species were grouped into the major and minor following [12] (Table 1).

UNDER PEER REVIEW

Table 1. Insect species associated with the ten okra cultivars under field conditions.

FAMILY	COMMON NAME	SCIENTIFIC NAME	PLANT PART INFESTED	RELATIVE ABUNDANCE	PEST STATUS
Chrysomelidae	Flea beetle	<i>Podagrica</i> sp.	Leaf	11.97	Major
Aleyrodidae	Whitefly	<i>Bemisia tabaci</i> (Gennadius)	Leaf	6.1	Minor
Aphididae	Green peach aphids	<i>Myzus persicae</i> (Sulzer)	Leaf, flower buds, flowers	66.46	Major
Formicidae	Black carpenter ants	<i>Camponotus</i> sp.	None	8.51	None
Lycosidae	Spider	<i>Hogna lenta</i> (Hentz)	None	0.21	None
Coccinellidae	Ladybird beetle	<i>Cheilomenes lunata</i> (Fabricius)	None	0.54	None
Cicadellidae	Okra leaf hopper	<i>Amrasca biguttula biguttula</i> (Ishida)	Leaf	5.12	Minor
Mantidae	Praying mantis	<i>Mantis religiosa</i> (Burmeister)	None	0.04	None
Pseudococcidae	Striped mealybugs	<i>Ferrisia virgata</i> (Cockerell)	Leaf, fruit	0.94	Minor
Pyrgomorphidae	Variegated grasshopper	<i>Zonocerus variegatus</i> (Linnaeus)	Leaf	0.02	Minor
	Tobacco grass hopper	<i>Atractomorpha crenulata</i> (Fabricius)	Leaf	0.04	Minor
Meloidae	Blister beetle	<i>Mylabris pustulata</i> (Thunberg)	Flower buds and flowers	0.04	Minor
Noctuidae	Cotton semilooper	<i>Anomis flava</i> (Fabricius)	Leaf	0.004	Minor
Nolidae	Transverse moth	<i>Xanthodes transversa</i> (Guenee)	Leaf	0.002	Minor

Although all the insects identified can attack the crop, the level of damage and their abundance varied among the cultivars. Based on the level of damage and their abundance, *Podagrica* sp. and *B. tabaci* were observed as major insect pests of the okra. These findings are consistent with reports of Obeng-Ofori and Sackey [8] and Asare bediako *et al.* [12]. It was observed that the low numbers of ladybird beetle (*C. lunata*) were responsible for the increase in Whitefly (*B. tabaci*) population. Ladybird beetle (*C. lunata*) have been used as natural enemies to control whitefly population in many plants [16;17;18;19;20]

3.2 Mean abundance of insect species at the vegetative stage.

The average incidence of flea beetle, whitefly, ants, aphids, spider, ladybird beetle, okra leafhopper, praying mantis, mealy bug and variegated grasshopper at the vegetative stage differed significantly ($P=0.05$) among all okra cultivars. Flea beetle had the highest number on Lucky 19F1 (36.00 ± 9.66 insects/plant) followed by K-F1, LD, AS, CS, AD, TG, F1-S, F1-K and K1 with abundance level of 32.75 ± 22.88 , 28.25 ± 8.10 , 23.00 ± 19.06 , 19.00 ± 16.63 , 18.75 ± 5.12 , 17.50 ± 10.66 , 16.50 ± 4.20 , 11.00 ± 4.97 and 9.00 ± 2.83 respectively. With respect to whitefly, F1-S had the highest number of insects (6.25 ± 2.18) per plant, while LD had the least (1.25 ± 0.50). Cultivar TG recorded the highest mean number of ants (43.25 ± 23.94) and praying mantis (0.70 ± 1.00) per plant, whereas, cultivar AS and F1-S had the minimum infestation of ants (3.25 ± 3.10) and praying mantis (0.24 ± 0.10). Similarly, CS had the highest mean number of spiders and okra leafhopper of 0.75 ± 0.50 and 0.75 ± 1.50 per plant respectively, whilst cultivar F1-S had the least infestation of spider (0.20 ± 0.11) and okra leafhopper (0.25 ± 0.10). Cultivar LD had the highest mean number of mealy bugs (1.75 ± 0.26) and grasshopper (1.75 ± 0.40) whereas cultivar L-19F1 and K-F1 had the least mean numbers of mealybugs (0.25 ± 0.10) and grasshopper (0.20 ± 0.10) per plant. Cultivars K-F1 and LD had the highest mean abundance of aphids and ladybird (27.00 ± 12.76 and 1.50 ± 1.00 respectively) while the least was recorded on TG (14.60 ± 9.20) and F1-K (0.18 ± 0.10) [Table2]

1 **Table 2. Mean abundance of insect species of the ten okra cultivars at the vegetative stage.**

Accessions	Flea beetle	Whitefly	Ants	Aphids	Ladybird beetle	Okra leafhopper	Praying mantis	Mealybug	Grass hopper
AD	18.75±5.12 ^{ab}	2.00±1.15 ^a	7.00±4.54 ^b	25.65±7.10 ^a	0.25±0.10 ^{ab}	1.00±0.41 ^a	0.25±0.10 ^a	1.00±0.41 ^{ab}	1.50±0.20 ^{ab}
AS	23.00±19.06 ^{ab}	5.75±3.94 ^a	3.25±3.10 ^b	25.05±17.53 ^a	0.20±0.10 ^b	0.57±0.50 ^a	0.50±0.20 ^a	1.50±0.33 ^{ab}	0.25±0.10 ^{ab}
CS	19.00±16.63 ^{ab}	1.75±0.21 ^a	20.75±17.42 ^{ab}	22.30±7.40 ^a	0.25±0.10 ^{ab}	0.75±1.50 ^a	0.50±0.10 ^a	1.25±0.50 ^{ab}	0.50±0.07 ^{ab}
F1-K	11.00±4.97 ^b	3.50±2.90 ^a	28.25±14.63 ^{ab}	23.08±16.3 ^a	0.18±0.10 ^b	1.25±1.00 ^a	0.25±0.10 ^a	0.75±0.50 ^{ab}	0.50±0.20 ^{ab}
F1-S	16.50±4.20 ^{ab}	6.25±2.18 ^a	12.50±10.10 ^b	19.25±12.6 ^a	1.00±0.20 ^{ab}	0.25±0.10 ^a	0.24±0.10 ^a	0.75±0.10 ^{ab}	0.50±0.37 ^{ab}
K1	9.00±2.83 ^b	3.75±1.92 ^a	17.50±16.86 ^{ab}	25.25±15.10 ^a	0.25±0.20 ^{ab}	0.50±0.20 ^a	0.50±0.20 ^a	2.25±1.50 ^a	1.00±0.81 ^{ab}
K- F1	32.75±22.88 ^a	3.75±2.97 ^a	19.50±14.80 ^{ab}	27.00±12.76 ^a	0.25±0.20 ^{ab}	1.00±0.20 ^a	0.51±0.10 ^a	1.50±1.11 ^{ab}	0.20±0.10 ^b
L D	28.25±8.10 ^{ab}	1.25±0.50 ^a	9.25±3.40 ^b	28.10±10.44 ^a	1.50±1.00 ^a	1.25±0.50 ^a	0.50±0.20 ^a	1.75±0.26 ^{ab}	1.75±0.40 ^a
L-19F1	36.00±9.66 ^a	3.50±0.32 ^a	12.00±11.52 ^b	19.25±9.50 ^a	0.75±1.00 ^{ab}	2.00±0.44 ^a	0.60±0.30 ^a	0.25±0.10 ^b	1.50±1.30 ^{ab}
TG	17.50±10.66 ^{ab}	2.50±1.38 ^a	43.25±23.94 ^a	14.60±9.20 ^a	0.50±1.00 ^{ab}	2.50±1.10 ^a	0.70±0.10 ^a	1.25±1.00 ^{ab}	0.75±0.50 ^{ab}

2 Note: Means followed by the same superscript in the same column are not significantly different at 5% probability level according to Duncan's multiple
3 range tests.

3.3 Abundance of insect species at the flowering stage.

The mean abundance of insect species recorded at the flowering stage on the okra cultivars was significantly different ($P=.05$) from each other. Plants of L-19F1 had the highest mean number of Flea beetles/plant (32.25 ± 10.30) while cultivar F1-K had the least number per plant (13.25 ± 14.86). In the case of whitefly, cultivar TG had the highest mean abundance per plant (26.75 ± 19.94) while AD had the least. Cultivar K-F1 recorded the highest mean number of ants (21.50 ± 15.20) and spider (1.25 ± 1.00) per plant whilst cultivar AD recorded the least infestation of ants (8.00 ± 6.50) and spider (0.12 ± 0.10). Mealy bug infestation was highest in cultivar AS (2.75 ± 2.22) and least in LD (0.25 ± 0.10). Cultivar LD had the highest mean number of grasshoppers per plant (3.75 ± 0.40) whilst the least mean number (0.20 ± 0.10) was recorded on TG. Cultivar K1, F1-S, L-19F1 and CS had the highest mean number of Aphids, Ladybird beetle, okra leafhopper and Praying mantis of 17.77 ± 5.30 , 1.25 ± 0.50 , 14.0 ± 5.71 and 2.50 ± 1.73 respectively [Table 3]

UNDER PEER REVIEW

Table 3. Mean abundance of insect species of the ten okra cultivars at the flowering stage.

Cultivar	Flea beetle	Whitefly	Ants	Aphids	Ladybird beetle	Okra leafhopper	Praying mantis	Mealybug	Grass hopper
AD	31.75±24.66 ^a	13.00±5.64 ^a	8.00±6.50 ^b	10.57±0.44 ^a	1.00±0.82 ^a	9.75±6.60 ^{ab}	1.50±1.30 ^{ab}	0.75±0.50 ^{bc}	0.75±0.50 ^{ab}
AS	23.25±22.64 ^a	18.75±13.12 ^a	17.50±4.73 ^b	11.13±5.88 ^a	0.75±0.96 ^a	9.50±5.68 ^{ab}	0.75±1.50 ^b	2.75±2.22 ^a	2.75±2.21 ^a
CS	20.50±12.76 ^a	14.00±4.40 ^a	12.75±2.87 ^b	11.20±5.92 ^a	0.75±1.50 ^a	7.25±5.90 ^{ab}	2.50±1.73 ^a	0.50±0.27 ^c	1.25±0.50 ^{ab}
F1-K	13.25±3.00 ^a	14.25±5.85 ^a	9.00±6.10 ^b	18.20±7.55 ^a	0.75±0.96 ^a	5.00±3.56 ^b	1.50±1.30 ^{ab}	0.50±0.10 ^c	1.25±0.50 ^{ab}
F1-S	20.50±14.86 ^a	22.25±4.03 ^a	20.75±12.40 ^b	16.75±7.30 ^a	1.25±1.50 ^a	9.50±7.14 ^{ab}	0.10±0.00 ^b	0.10±0.00 ^c	2.75±2.22 ^a
K1	13.75±11.50 ^a	18.75±0.60 ^a	6.00±1.50 ^b	17.77±5.30 ^a	1.00±1.41 ^a	8.25±5.12 ^{ab}	0.50±0.10 ^b	1.25±1.50 ^{abc}	1.25±0.50 ^{ab}
K- F1	18.50±12.50 ^a	18.25±3.40 ^a	21.50±9.20 ^b	13.95±9.06 ^a	0.50±0.18 ^a	10.5±3.11 ^{ab}	0.50±0.10 ^b	0.75±0.16 ^{bc}	0.75±0.50 ^{ab}
L D	28.00±14.07 ^a	21.25±14.60 ^a	18.75±13.20 ^b	13.80±6.87 ^a	0.75±0.50 ^a	12.5±7.93 ^{ab}	1.50±0.73 ^{ab}	0.25±0.10 ^c	3.75±0.40 ^b
L-19F1	32.25±10.30 ^a	22.50±14.64 ^a	20.00±10.04 ^b	12.53±2.63 ^a	0.50±1.00 ^a	14.0±5.71 ^a	0.50±0.10 ^b	2.50±0.08 ^{ab}	2.25±1.70 ^{ab}
TG	22.75±12.78 ^a	26.75±19.94 ^a	51.50±20.73 ^a	12.07±3.78 ^a	1.25±1.26 ^a	8.25±3.77 ^{ab}	1.00±0.82 ^{ab}	0.26±0.10 ^c	0.20±0.10 ^b

Note: Means followed by the same superscript in the same column are not significantly different at 5% probability level according to Duncan's multiple range tests.

3.4 Mean abundance of insect species at the fruiting stage.

Significant differences ($P=.05$) were obtained in the abundance of insects among the ten okra cultivars at the fruiting stage. Plants of LD had the highest mean abundance of flea beetles per plant (47.50 ± 13.53) whereas F1-K recorded the least (13.50 ± 3.00). With respect to whitefly infestation, plants of L-19F1 had the highest mean abundance of 28.50 ± 13.23 per plant with K1 recording the least (9.75 ± 7.41). Plants of cultivar TG recorded the highest infestation of aphids (77.50 ± 1.64) and ladybird (2.50 ± 3.11) whereas K1 and F1-K had the least (12.50 ± 10.34 and 0.25 ± 0.50 respectively). Similarly, cultivar K-F1 recorded the highest mean number of okra leafhopper (17.00 ± 8.05) whilst CS had the highest number of grasshoppers (2.25 ± 2.06). Cultivar K1 recorded the highest mean incidence of ants (30.50 ± 11.73) and praying mantis (3.25 ± 2.63) [Table 4].

Table 4. Mean abundance of insect of ten okra accessions at the fruiting stage during the study.

Accessions	Flea Beetle	Whitefly	Ants	Aphids	Ladybird beetle	Okra leafhopper	Praying mantis	Mealybug	Grass hopper
AD	29.75±7.80 ^{bc}	22.50±9.45 ^a	29.25±4.55 ^{ab}	75.00±2.42 ^{ab}	1.00±1.15 ^a	15.00±13.08a	1.50±1.73 ^a	0.75±1.50 ^{ab}	0.50±1.00 ^a
AS	22.00±7.44 ^{bc}	19.00±4.25 ^a	19.50±5.05 ^{ab}	76.50±8.63 ^{ab}	0.50±1.00 ^a	14.25±12.20a	1.00±0.81 ^a	1.25±0.96 ^{ab}	0.40±1.00 ^a
CS	27.50±4.80 ^{bc}	18.00±4.31 ^a	15.50±11.00 ^{ab}	77.50±4.70 ^{ab}	1.50±1.30 ^a	5.75±6.18a	0.75±0.50 ^a	1.00±1.15 ^{ab}	2.25±2.06 ^a
F1-K	22.00±16.10 ^{bc}	11.25±3.77 ^a	13.50±12.01 ^{ab}	33.75±5.10 ^b	0.25±0.50 ^a	3.50±0.68a	1.50±1.73 ^a	1.25±1.50 ^{ab}	0.75±0.50 ^a
F1-S	32.00±12.30 ^{abc}	15.25±2.75 ^a	9.00±2.94 ^b	53.75±8.50 ^{ab}	1.00±1.41 ^a	4.25±1.43a	2.00±2.82 ^a	2.75±2.21 ^a	0.50±0.58 ^a
K1	21.75±2.90 ^{bc}	9.75±7.41 ^a	30.50±11.73 ^a	12.50±0.34 ^a	0.50±1.00 ^a	12.25±2.12a	3.25±2.63 ^a	0.75±0.96 ^{ab}	0.50±1.00 ^a
K- F1	33.25±14.30 ^{ab}	14.00±3.98 ^a	17.00±12.02 ^{ab}	71.75±9.54 ^{ab}	0.50±1.00 ^a	17.00±8.05a	2.75±2.21 ^a	0.25±0.50 ^b	1.50±1.91 ^a
L D	47.50±13.53 ^a	21.00±9.20 ^a	16.75±8.40 ^{ab}	75.00±4.90 ^{ab}	1.00±2.00 ^a	10.75±9.32a	1.00±1.41 ^a	1.75±2.22 ^{ab}	1.25±1.50 ^a
L-19F1	46.00±3.20 ^a	28.50±3.23 ^a	16.00±0.60 ^{ab}	56.75±2.74 ^{ab}	1.75±2.06	13.75±3.09a	2.00±0.82 ^a	0.75±0.96 ^{ab}	1.50±1.30 ^a
TG	17.25±14.86 ^c	22.00±6.77 ^a	13.75±7.37 ^{ab}	77.50±1.64 ^{ab}	2.50±3.11 ^a	12.50±8.50a	2.00±1.15 ^a	2.00±1.15 ^{ab}	0.75±0.96 ^a

Note: Means followed by the same superscript in the same column are not significantly different at 5% probability level according to Duncan's multiple range tests.

Green peach aphid (*M. persicae*), flea beetle (*Podagrica* sp.), Okra leafhopper (*A. biguttula biguttula*), whitefly (*B. tabaci*), striped mealybug (*F. virgata*), Black carpenter ants (*Camponotus* sp.), spider (*H. lenta*), ladybird beetle (*C. lunata*), variegated grasshopper (*Z. variegatus*) and praying mantis (*M. religiosa*) were the common insects observed at vegetative, flowering and fruiting stages. The total number of insects differed from one stage to another. A total of 1,439 insects were recorded at the fruiting stage which was significantly higher than the flowering (855) and vegetative stages (693). The variation in the number of insect species observed at the different developmental stages could be due to environmental changes as suggested by Abro *et al.*, [21]. Mean whitefly count was relatively low at the vegetative, flowering and fruiting stages of the cultivars. However, Flea beetle (*Podagrica* sp.) and Green peach aphid numbers increased progressively throughout all the stages.

3.5 Severity of leaf damage by flea beetle (*Podagrica* sp.).

Mean severity scores of leaf damage by flea beetles (*Podagrica* sp.) during the vegetative, flowering and fruiting stages of the various cultivars are shown in Table 5. The results showed that the severity of damage during the above stages differed significantly among the ten okra cultivars. Cultivar F1-S had the highest severity score (4.56.50 perforations) whilst the least score (2; 16.40 perforations) was associated with CS. There were significant differences in the leaf damage by the flea beetle, and among the okra cultivars during the vegetative stage. At the flowering stage, the highest damage was observed in L-19F1 (5; 68.60 perforations) whilst the least was observed in LD (3; 33.10 perforations). With respect to the fruiting stage, plants from accession CS recorded the highest mean leaf damage (5; 79.70 perforations) followed by L-19F1 (5;78.10) with AS having the least leaf damage, (3;41.10). According to Echezona and Offordile [9], the feeding activity of flea beetle (*Podagrica* sp.) causes damage consisting of characteristic perforations of leaves resulting in uneven holes which decrease the photosynthetic surface area of the leaves, culminating in high yield loss of okra. In the present study, leaf damage was significantly higher at the fruiting stage compared with the flowering and vegetative stages. These results are consistent with those of Eguatu and Taylor [22] and Schippers [23] who reported increase in leaf damage caused by *Podagrica* sp. at the reproductive stages than the vegetative stage due to abundance of food sources such as pods, flowers and buds that attract a lot of the flea beetle to the okra plant. Plants of cultivars LD and AS were the most promising, recording the least leaf damage. These cultivars exhibit a good inherent potential to withstand insect attack and as such would be good materials for cultivation by farmers and for breeding.

Table 5. The severity of leaf damage by Flea beetle (*Podagrica* sp.) during three developmental stages of ten okra accessions.

Accession	Number of perforations per leaf		
	Vegetative* stage	Flowering* stage	Fruiting* stage
AD	(2) 17.10	(4) 51.40	(5) 64.70
AS	(3) 38.00	(3) 40.10	(3) 41.10
CS	(2) 16.70	(5) 68.60	(5) 79.70
F1-K	(2) 16.40	(4) 54.90	(5) 68.10
F1-S	(4) 56.50	(4) 50.55	(5) 64.45
K1	(2) 20.00	(4) 59.85	(4) 57.55
K- F1	(3) 33.10	(4) 49.95	(4) 54.15
L D	(2) 22.20	(3) 33.10	(4) 56.65
L-19F1	(3) 39.50	(5) 61.65	(5) 78.10
TG	(2) 29.40	(3) 44.25	(5) 72.70

**Bolded value in bracket indicates damage level on a five-point scale whereas corresponding value represents the number of leaf perforations. The scoring scale is as follows: 1 very mild damage (1 to 15 perforations); 2 mild damage (16 to 30 perforations); 3 moderately severe damage (31 to 45 perforations); 4 very severe damage (46 to 60 perforations); 5 extremely severe damage (more than 60 perforations).*

4. CONCLUSION

A total of thirteen insect types belonging to five orders (Coleoptera, Homoptera, Hymenoptera, Mantodea and Orthoptera) and thirteen families (Chrysomelidae, Coccinellidae, Pyrgomorphidae, Meloidae, Noctuidae, Nolidae, Cicadellidae, Aleyrodidae, Aphididae, Pseudococcidae, Mantidae, Formicidae, and Acrididae) were identified **in** the field. Out **of** the thirteen families recorded, two beneficial organisms Ladybird beetle and Spider belonging to the Coccinellidae and Lycosidae respectively were also found to be present **in** the field. A total of 1,439 insects were recorded at the fruiting stage which was significantly higher than the flowering (855) and vegetative stages (693). Whitefly (*B. tabaci*) count was relatively low at the vegetative, flowering and fruiting stages of the cultivars. However, *Podagrica* sp. numbers increased progressively throughout all the stages., Leaf damage was significantly higher at the fruiting stage compared to the flowering and vegetative stages. Plants of cultivars LD and AS were the most promising recording the least leaf damage.

1 **COMPETING INTERESTS**

2
3 Authors have declared that no competing interests exist.
4
5

6 **REFERENCES**

- 7
8
9 1. SRID-MOFA. Statistical Research and Information Directorate, Ministry of
10 Food and Agriculture, (Ghana). Production Figures. 2007; 56 -57.
11 2. Hayase D. Field evaluation of neem seed extract for the control of insect
12 pests of okra (*Abelmoschus esculentus* (L) Moench). Bsc. dissertation
13 submitted to the Crop Science Department, Faculty of Agriculture, University
14 of Ghana. 2001;17-41.
15 3. Sinnadurai S. Vegetable production in Ghana. The Ghana Farmer, Ministry of
16 Agriculture, Ghana. 1971;20.
17 4. Critchley BR. Pests of Vegetables. Their Identification and control in Ghana.
18 Natural Resource Institute, Greenwich. Department of International
19 Development. University of Grenwich. 1997;282.
20 5. Tindal HD. Fruits and vegetables in West Africa. FAO plant production and
21 protection series 1965; 4:55-56.
22 6. Norman JC. Okra: Tropical vegetable crops. Arthur H. Stock well Ltd.
23 1992;56-58.
24 7. Obeng-Ofori D. Insect Pest of Vegetables and Plantation Crops. Their biology
25 and control. Asempa Publishers Ltd. 1998;87.
26 8. Obeng-Ofori D, Sackey J. Field evaluation of non-synthetic insecticides for the
27 management of insect pests of okra *Abelmoschus Esculentus* (L.) Moench in
28 Ghana. Ethiopian J. Sci. 2003; 26:145-150.
29 9. Echezona BL, Offordile JI. Responses of flea beetles (*Podagrica* spp.) and
30 okra plants (*Abelmoschus esculentus* L. Moench) to differently coloured
31 polyethylene shades. Intl. J. Pest Manag't. 2001;57(2):161-168.
32 10. Alegbejo MD. Effect of sowing date on the incidence and severity of Okra
33 mosaic Tymovirus. J. Veg. Crop Prod. 2001; 8:9-14.
34 11. Ali M, Hossain MZ, Sarkern NC. Inheritance of Yellow Vein Mosaic Virus
35 (YVMV) tolerance in a cultivar of okra (*Abelmoschus esculentus* (L.) Moench).
36 *Euphytica*. 2000;111(3):205-209.
37 12. Asare-Bediako E, Van der Puije GC, Taah KJ, Abole EA, and Baidoo, A.
38 (2014). Prevalence of Okra Mosaic and Leaf Curl Diseases and *Podagrica* sp.
39 Damage of Okra (*Abelmoschus esculentus*) Plants. Interna. J. current rev.
40 academ. res. 2014;2 (6):260-271.
41 13. FAO/UNESCO. "Soil Map of the World". Revised Legend, World Resources
42 Report 60. FAO, Rome, Italy. 1994.
43 14. Akaho EKH, Maakuu BT, Anim-Sampong S, Emi-Reynolds, G., Boadu HO,
44 Osae EK, Akoto-Bamford S, Dodoo-Amoo, D.N.A. Intermediate safety
45 analysis report (GAEC-NNRI-RT-90). 2003.
46 15. Dickson KB, Benneh G. A new geography of Ghana. Longmans Group
47 Limited, London. 2004.
48 16. Legaspi JC, Correa JA, Carruthers RI, Legaspi Junior BC, Nordlund DA.
49 Effect of short-term releases of *Chrysoperla rufilabris* (Neuroptera:

- 50 Chrysopidae) against silverleaf whitefly (Homoptera: Aleyrodidae) in field
51 cages. J. Ento. Sci. 1996a; 31:102-111.
- 52 17. Legaspi JC, Nordlund DA, Legaspi Junior BC. Tri-trophic interactions and
53 predation rates in *Chrysoperla* spp. attacking the silverleaf whitefly.
54 Southwestern Entomologist. 1996b;21:33-42.
- 55 18. Liu TX, Stansly PA. Morphology of *Nephaspis oculatus* and *Delphastus*
56 *pusillus* (Coleoptera: Coccinellidae), predators of *Bemisia argentifolii*
57 (Homoptera: Aleyrodidae). Proceedings of the Entomological Society of
58 Washington. 1996a;98:292-300.
- 59 19. Liu TX, Stansly PA. Oviposition, development, and survivorship of *Encarsia*
60 *pergandiella* (Hymenoptera: Aphelinidae) in four instars of *Bemisia argentifolii*
61 (Homoptera: Aleyrodidae). An. Ento. I Soci. Amer. 89 :96-102.
- 62 20. Pfadt RE. Fundamentals of applied entomology. Macmillan Company, New
63 York. 1980;99:24-126.
- 64 21. Abro GH, Memon AJ, Syed TS, Shaikh AA. Infestation of *Earia* sp. on cotton
65 and okra grown as mono and mix crops. Pakistan J. of Biol. Sci, 7.
66 2004;(6):937-942.
- 67 22. Egwuatu RI, Taylor TA. (1976). Aspects of the spatial distribution of
68 *Acanthomia Tomentosicollis* stal (Heteroptera, Coreidae) in *Cajanus cajan*
69 (Pigeon pea). J. Econ. Ento. 1975;69(5):591-594.
- 70 23. Schipper RR. African Indigenous Vegetables. An Overview of the cultivated
71 species. Chathaam, UK: Natural Resources Institute/ACP-EO Technical
72 Centre for Agricultural and Rural Cooperation. UK. 2000.
- 73
74
75
76
77
78