

## Response of Soil Microorganisms to Oilfield Wastewater

### ABSTRACT

Environmental pollution resulting from oil exploitation and exploration has impacted negatively on the biodiversity of the affected areas. Therefore, this study investigated the response of soil microorganisms to oilfield wastewater. The oilfield wastewater and soil samples were collected from an onshore oil producing platform fortnightly for a period of three months and microbiological analyses were performed using standard methods. Soil around pond average counts for THB and HUB ( $\text{Log}_{10}$  CFU/g) were 7.32 to 7.35 and 4.16 to 4.22 respectively. Soil 80m away from pond average range for THB and HUB were 7.38 to 7.40 and 3.32 to 3.34 respectively. For TF and HUF, Soil around the pond mean recorded 4.65 to 4.85 and 4.12 to 4.16 ( $\text{Log}_{10}$  CFU/g) respectively. Mean monthly counts for soil 80m away TF and HUF ( $\text{Log}_{10}$  CFU/g) were 5.03 to 5.05 and 3.26 to 3.34 respectively. Soil around the pond recorded lower total heterotrophic bacteria count and total fungi count but higher hydrocarbon utilizing bacteria and hydrocarbon utilizing fungi than soil 80 m away.

There was significant difference ( $P < 0.05$ ) between the THB and HUB in the various samples. For the fungi count there was no significant difference ( $P > 0.05$ ) between TF and HUF in all the samples except in sample obtained from soil 80 m away from pond.

Bacteria species isolated from the study include: *Bacillus*, *Aeromonas*, *Micrococcus*, *Staphylococcus*, *Chryseomonas*, *Proteus*, *Pseudomonas*, *Klebsiella*. Apart from *Aeromonas* and *Chryseomonas* the rest of the isolate were identified also as hydrocarbon utilizing bacteria. While fungi species isolated includes: *Aspergillus fumigates*, *Aspergillus niger*, *Aspergillus flavus*, *Saccharomyces cerevisiae*, *Geotricum*, *Trichoderma*, *Fusarium* and *Penicillium*. Hydrocarbon utilizing fungi that occurred includes: *Aspergillus niger*, *Aspergillus flavus*, *Fusarium*, *Penicillium* and *Saccharomyces cerevisiae*. The response of these microorganisms in the oil polluted environment suggests that the isolated bacteria and fungi could utilize the oil as energy and carbon source and could be effective in the cleanup of the polluted sites.

**Key words:** Response, Oilfield wastewater, Pond wastewater, Soil, Bacteria, Fungi

### INTRODUCTION

Crude oil exploration and exploitation activities in Nigeria are major contributors of environmental pollution in the Niger Delta. Oilfield wastewater is the associated wastewater of crude oil production activities. Oilfield wastewater may contain hydraulic fracturing (HF) fluids, naturally occurring salts, radioactive materials, heavy metals, and other compounds from the

formation such as polycyclic aromatic hydrocarbons, alkenes, alkanes, and other volatile and semi-volatile organics (Warner *et al.*, 2012; Kassotis *et al.*, 2015).

Increased Petroleum activities, particularly in the Niger Delta has led to pollution stress on soil and surface water, due to the discharging of large quantities of oilfield wastewater without adequate treatment techniques (Obire and Amusan, 2003; Wemedo and Obire, 2012). Oilfield wastewater containing high organic and inorganic chemicals poses environmental problems. The main crucial environmental issues of the oilfield wastewater are total petroleum hydrocarbon, total solids (TS), and inorganic chemicals including heavy metals and polycyclic aromatic hydrocarbons (PAHs), biochemical and chemical oxygen demand (BOD and COD), and pathogens (Pichtel, 2016).

In Nigeria, Oil exploitation companies are known to discharge oilfield wastewater into Streams or ponds which are also a threat to the surrounding soil and groundwater (Obire and Wemedo, 1996). Accelerated soil quality change due to oilfield wastewater discharging with large quantities of nutrients and toxic substances into the environment has long become an issue problem in Niger Delta. It is estimated that over 90% of wastewater from operations of oil industries in Nigeria is still discharged to soil, rivers and streams without adequate treatment. This is largely due to the fact that most of the oil companies have no wastewater treatment plants or where they exist the facilities are inadequate (Human Rights Watch, 1998; Van Dessel, 1996). Soil contaminated by industrial effluents has affected adversely both soil health and crop productivity. Heavy metals are one of the major pollutants of interest in the environment because of its toxicity, persistence and bioaccumulation problems (Zouboules *et al.*, 2004). Excessive accumulation of micronutrients and other heavy metals like cadmium, lead, and nickel in the plants operates as stress factors causing physiological constraints leading to decrease vigour and plant growth (Zouboules *et al.*, 2004) and therefore crop yield (Jaja and Obire, 2015). The effects of petroleum activities on the environment in the Niger Delta are evident through the pollution of soil and water bodies and human habitat in the major cities. The oilfield wastewater contains toxic and hazardous substances that are detrimental to human health if they enter the food chain (Rajaram and Ashutost, 2008).

The objective of this present study therefore was to investigate the response of soil microorganisms to oilfield wastewater by enumerating microorganisms in the oilfield wastewater, wastewater in pond, soil around the pond and in soil 80m away from the pond. This investigation was conducted for a period of three months (January–March, 2018) regarded as part of the dry season in the Niger Delta.

## **MATERIALS AND METHODS**

### ***Collection of Oilfield wastewater and soil samples***

Oilfield wastewater was collected from Ogbogu Flow Station; an onshore oil production platform located in Ogba Egbema Ndoni local government Area (ONELGA) of Rivers State, Nigeria. The Oilfield wastewater samples were collected using 4 Litre capacity plastic bottles and stored in an ice packed cooler.

On the other hand, the soil samples were collected from around the pond and 80 meters away from the pond at a depth of 0-15cm with a sterile spatula into sterile polythene bags and

stored in an ice packed cooler. The collected and appropriately labeled oilfield wastewater and soil samples were immediately transported to the laboratory for analysis within 24 hours for processing and analyses. Samples were collected twice in a month (1<sup>st</sup> and 3<sup>rd</sup> week) for a period of three months (January, 2018 to March, 2018).

### ***Media Preparation***

Nutrient Agar was used for Total Heterotrophic bacterial count; Potato dextrose agar was used for total fungal count while Mineral salt agar medium prepared according to the modified minimal salts medium (MSM) composition of Mills *et al.* (1978) was used for the isolation of total hydrocarbon utilizing bacteria. Minimal salts medium (MSM) composition is – [MgSO<sub>4</sub>.7H<sub>2</sub>O (0.42g), KCl (0.29g), KH<sub>2</sub>PO<sub>4</sub> (0.83g), Na<sub>2</sub>HPO<sub>4</sub> (1.25g) NaNO<sub>3</sub> (0.42), agar (20g)] in 1Litre of distilled water. The mixture was thoroughly mixed and autoclaved at 15psi at 121°C for 15mins and was allowed to cool to 45°C. The medium was prepared by the addition of 1% (v/v) crude oil sterilized with 0.22µm pore size Millipore filter paper Moslein France (Obire, 1988) to sterile MSM, which has been cooled to 45°C under aseptic condition. The MSM and crude oil were then mixed thoroughly and aseptically dispensed into sterile Petri dishes to set.

### ***Microbiological Analysis of the Oilfield Wastewater and Soil Samples***

#### ***Determination of Total heterotrophic bacterial (THB) count of oilfield wastewater and soils***

The total heterotrophic bacterial (THB) count was determined using the nutrient agar and spread plate technique as described by Prescott *et al.* (2005). An aliquot (0.1ml) of each serially diluted sample using dilution factors of 10<sup>-5</sup> for Raw wastewater, 10<sup>-2</sup> for wastewater in the pond, and 10<sup>-4</sup> for all the soil samples was separately inoculated onto different sterile nutrient agar plates in triplicates. The plates were incubated at 37°C in an inverted position for 24 hours. After incubation, colonies that developed on the plates were counted and only counts of between 30 and 300 were recorded. The average values of replicate plates was calculated and expressed as colony forming unit - CFU/ml for oilfield wastewater and CFU/g for soil samples.

#### ***Determination of total fungi count of samples of oilfield wastewater and soils***

The total count of fungi in the samples was also determined by the spread plate technique. An aliquot (0.1ml) of serial dilution (10<sup>-2</sup>) of each of the various samples was plated onto separate Potato dextrose agar plates to which 0.1 ml of streptomycin solution was incorporated to suppress bacterial growth. The plates were incubated at 28°C for 5-7 days and the discrete colonies that developed were enumerated as the viable counts (CFU) of fungi in the oilfield wastewater and soil samples (Obire and Wemedo, 1996).

#### *Hydrocarbon utilizing bacterial count (HUB) of samples*

The population of the hydrocarbon utilizing bacterial of oilfield wastewater and soil samples was determined by inoculating 0.1ml aliquot of the serially diluted ( $10^{-1}$  and  $10^{-2}$ ) samples of oilfield wastewater and  $10^{-1}$  of soil samples onto mineral salt agar media using the spread plate technique described by Odokuma (2003). The Vapour Phase Transfer method will be adopted by the use of sterile filter paper discs that will be soaked in filter sterilized crude oil which served as the only carbon source in the mineral salt agar. The sterile crude oil-soaked filter papers were aseptically transferred to the inside cover of the inoculated Petri dishes and incubated for 5 days at room temperature. Colonies that develop were counted, average of duplicate colonies calculated colony forming units per ml of wastewater or per gram soil calculated.

#### *Hydrocarbon utilizing fungal count (HUF) of samples*

Total hydrocarbon utilizing fungal count of oilfield wastewater and soil samples was determined by inoculating 0.1ml of the serially diluted samples -1 on mineral salt agar. The mineral salt medium will be supplemented with streptomycin (0.1ml) to suppress bacterial growth (Obire and Wemedo, 1996). The Vapour Phase Transfer method was adopted by the use of sterile filter paper discs that were soaked in filter sterilized crude oil which served as the only carbon source in the mineral salt agar. The sterile crude oil-soaked filter papers were aseptically transferred to the inside cover of the inoculated Petri dishes and incubated for 5 days at room temperature. Colonies that develop were counted, average of duplicate colonies calculated colony forming units per ml of wastewater or per gram soil calculated.

#### *Characterization and identification of bacterial and fungal isolates from samples*

The cultural, morphological, microscopic characteristics of the isolates from the study were observed and recorded. The morphological and biochemical tests conducted using the isolates included Gram staining, motility, catalase, oxidase, citrate utilization, sugar fermentation, hydrogen sulphide production, indole production methyl red and Voges Proskauer test. Results of the morphological and biochemical characteristics of the isolates were compared with those of known Taxa using Bergey's manual of determinative bacteriology (1994).

For the presumptive identification of fungal isolates, pure fungal cultures were observed while still on plates and after wet mount in lacto-phenol on slides under the compound microscope. Observed characteristics such as vegetative hyphae and reproductive structures were recorded and compared with the established identification key of Barnett and Hunter (1972) and Malloch (1997).

#### *Statistical analysis*

Statistical analysis was also conducted using Duncan Multiple Range test and Analysis of variance to determine whether there is significant difference between the total counts and the hydrocarbon utilizers.

## RESULTS

The results of the total heterotrophic bacteria (THB) and hydrocarbon utilizing bacteria (HUB) ( $\log_{10}$ CFU/ml) of the oil field wastewaters were: January 8.26 and 3.15; February 8.30 and 4.11; March 8.26 and 4.09 respectively. For wastewaters in pond the THB and HUB were: January 5.44 and 5.32; February 5.43 and 5.31; March 5.39 and 5.30 respectively. Soil around pond mean monthly counts for THB and HUB ( $\log_{10}$ CFU/g) were: January 7.35 and 4.22; February 7.33 and 4.16; March 7.32 and 4.16 respectively. Soil 80m away from pond counts for THB and HUB were: January 7.38 and 3.32; February 7.40 and 3.34; March 7.40 and 7.32 respectively. Heterotrophic bacteria counts were higher in the oilfield wastewater in the various months of study compared to other samples. The hydrocarbon utilizing bacteria count was higher in the wastewater pond than other samples. Soil 80 m away from the pond recorded higher total heterotrophic bacteria count compared to the soil around the pond. While higher hydrocarbon utilizing bacteria count were recorded in soil around the pond.




Fig. 1: Counts of total heterotrophic bacterial (THB) and hydrocarbon utilizing bacteria (HUB) of wastewater and soil.

Mean monthly counts of the total fungi (TF) and the hydrocarbon utilizing fungi ( $\text{Log}_{10}$  CFU/ml) for the oilfield wastewaters were: January 4.26 and 3.45; February 4.23 and 3.38; March 4.28 and 3.43 respectively. The mean monthly counts of the TF and HUF for wastewater in pond were: January 4.58 and 3.76; February 4.49 and 3.63; March 4.56 and 3.63 respectively. Soil around the pond mean monthly counts for TF and HUF ( $\text{Log}_{10}$  CFU/g) were: January 4.85 and 4.16, February 4.74 and 4.12; March 4.65 and 4.12 respectively. Mean monthly counts for soil 80m away TF and HUF were: January 5.05 and 3.26; February 5.03 and 3.32; March 5.05 and 3.34 respectively. **The soil 80m away from the pond had higher total fungi count than the soil around pond, while soil around the pond had higher hydrocarbon utilizing fungi count.**




Fig. 2: Counts of total fungi (TF) and hydrocarbon utilizing fungi (HUF) of wastewater and soil.

Table 1: Analysis of Variable

Report		THB	HUB	THF	HUF
Microbiology/locations					
Oilfield wastewater	Mean	8.2733	3.7833	4.2567	3.4200
	Std. Deviation	0.02309	0.54857	0.02517	0.03606
Pond water	Mean	5.4200	5.3100	4.5433	3.6733
	Std. Deviation	0.02646	0.01000	0.04726	0.07506
Soil 80m away from pond	Mean	7.3933	3.3267	5.0433	3.3067
	Std. Deviation	0.01155	0.01155	0.01155	0.04163
Soil around pond	Mean	7.3333	4.1800	4.7467	4.1333
	Std. Deviation	0.01528	0.03464	0.10017	0.02309

The predominant bacteria and fungi that were isolated from the various samples are as shown in Table 2 and Table 3 respectively. The signs indicates presence (+) and absence (-) of indicated organisms in the row, for the various samples in January, February and March.

*Bacillus*, *Micrococcus*, *Pseudomonas sp.* occurred in all the samples in the various months of sampling. *Proteus sp.* was not isolated in the wastewater samples but only occurred in the soil samples in all the months of sampling. *Staphylococcus*, *Chryseomonas* and *Klebsiella sp.* were not isolated in the soil samples in the months of sampling but were isolated in the wastewater samples. *Aeromonas sp.* only occurred in the soil around pond in the month of February but was isolated in the wastewater samples in all the months.

*Aspergillus niger*, *Aspergillus fumigates* and *Penicillium sp.* were isolated in the various samples and in all the months. *Fusarium sp.* and *Saccharomyces cerevisiae* were not isolated in the soil samples. *Geotricum sp.* was isolated in the oilfield wastewater but occurred in all other samples. *Trichoderma sp.* was only isolated in the soil samples.

Table 2: Bacteria isolated from Oilfield Wastewater and Soil Samples

Types of bacteria	Oilfield wastewater	Wastewater pond	in Soil around pond	Soil 80m away from pond
<i>Bacillus sp</i>	+++	+++	+++	+++
<i>Aeromonas sp</i>	+++	+++	+-	---
<i>Micrococcus sp</i>	+++	+++	+++	+++
<i>Staphylococcus sp</i>	+++	--+	---	---
<i>Chryseomonas sp</i>	+++	+ - +	---	---
<i>Proteus sp</i>	+++	++ -	+++	+++
<i>Klebsiella sp</i>	+++	+++	---	---
<i>Pseudomonas sp</i>	+++	+++	+++	+++

Table 3: Fungi isolated from Oilfield Wastewater and Soil Samples

Types of fungi	Oilfield wastewater	Wastewater in pond	Soil around pond	Soil 80m away from pond
<i>A. niger</i>	+++	+++	+++	+++
<i>A. fumigatus</i>	+++	+++	+++	+++
<i>A. flavus</i>	++-	---	+++	+++
<i>Penicillium sp</i>	+++	+++	+++	+++
<i>Fusarium sp</i>	+++	+++	---	---
<i>Saccharomyces cerevisiae</i>	+++	+++	---	---
<i>Geotricum sp</i>	---	+++	++	+++
<i>Trichoderma sp</i>	---	---	+++	+++

## DISCUSSION

Microbial populations play a role in degradation of hydrocarbon contaminations, Atlas (1981) and Leahy and Colwell (1990) reported that the rate of petroleum hydrocarbon biodegradation in nature is determined by the populations of indigenous hydrocarbon degrading microorganisms. Leahy and Colwell (1990) concluded that hydrocarbon biodegradation depends on the composition of the microbial community and its adaptive response to the presence of hydrocarbons. Heterotrophic bacteria counts were higher in the oilfield wastewater in the various months of study compared to other samples. The occurrence of bacteria in oilfield wastewater despite the treatment with biocide during the treatment process could be attributed to organic and inorganic constituents that serve as nutrients for bacterial growth. This was also reported by Wemedo *et al.* (2012), and Marshall and Derinny (1988). The hydrocarbon utilizing bacteria count was higher in the wastewater pond than other samples. This could be as a result of continues discharge of petroleum products into the wastewater pond. It has been reported by several researchers that continuous discharges of crude oil into the ecosystem may result in selective increase or decrease in microbial population (Okpokwasili and Nnubia, 1995; Okpokwasili and Odokuma, 1996). There was significant difference ( $P < 0.05$ ) between the THB and HUB in the various samples. Soil around the pond recorded lower total heterotrophic bacteria count and higher hydrocarbon utilizing bacteria compared to the soil 80 m away from the pond. The lower total heterotrophic count could also be attributed to the toxicity of the oilfield wastewater on the soil around the pond resulting to the response of microorganisms that have the ability to withstand the toxins. Obire and Anyawu (2009) reported reduction in diversity of species with increase in concentration of added crude oil was an index for environmental stresses of oil hydrocarbons.

The soil 80m away from the pond had higher total fungi count than other samples. Higher HUF counts were observed in the soil around pond and could be attributed to the presence of residual crude oil in the polluted soil which boosts the carbon supply in the soil thereby favoring the growth of the hydrocarbon utilizing fungi which were adapted to the quantity of hydrocarbons in the



environment. It was also observed that total fungi counts were greater than HUF counts in all the samples in the various months. There was no significance difference ( $P>0.05$ ) between the HF and HUF in all samples except in soil 80m away from the pond. This indicates that samples were polluted with hydrocarbons. The result is in agreement with that reported by Chikere and Azubuikwe (2014).

Bacteria species isolated from the study include: *Bacillus*, *Aeromonas*, *Micrococcus*, *Staphylococcus*, *Chryseomonas*, *Proteus*, *Pseudomonas*, *Klebsiella*. Apart from *Aeromonas* and *Chryseomonas* the rest of the isolate were identified also as hydrocarbon utilizing bacteria. While fungi species isolated includes: *Aspergillus fumigates*, *Aspergillus niger*, *Aspergillus flavus*, *Saccharomyces cerevisiae*, *Geotricum*, *Trichoderma*, *Fusarium* and *Penicillium*. Hydrocarbon utilizing fungi that occurred includes: *Aspergillus niger*, *Aspergillus flavus*, *Fusarium*, *Penicillium* and *Saccharomyces cerevisiae*. The following organisms; *Staphylococcus sp*, *Fusarium*, *Chryseomonas sp*, *Klebsiella sp*, and *Saccharomyces cerevisiae* were not isolated in the soil samples while *Trichoderma sp* was only isolated in all soil samples. High numbers of certain oil-degrading microorganisms from an environment implies that those organisms are the active degraders of that oil and may have the ability to produce spores that may shield them from toxic effects of the hydrocarbon. The absence of certain species known to be soil microorganisms in favour of oil degraders could be detrimental to the ecosystem. Similar reports were published by Okerentugba and Ezeronye (2003), Onifade and Abubakar (2007), Boboye *et al.* (2010), Obire and Wemedo (1996) and Obire and Amusan, 2003.

## CONCLUSION AND RECOMMENDATIONS

The high population of hydrocarbon utilizers in the soil around pond suggests that the hydrocarbon utilizers were adapted to the quantity of hydrocarbons in the environment and thereby increased the number of hydrocarbon utilizers in the polluted area.

The study also revealed that most of the organisms isolated as total heterotrophic bacteria and total fungi were part of the utilizers. The response of these microorganisms in the oil polluted environment suggests that the isolated bacteria and fungi could utilize the oil as energy and carbon source and could be effective in the cleanup of the polluted sites.

## REFERENCES

- Atlas R.M. Microbial degradation of petroleum hydrocarbons: an environmental perspective. Microbiol. Rev. 1981;45:180–209.
- Barnett, H. L. and Hunter, B. B. (1972). *Illustrated Genera of Fungi Imperfecti*. 3<sup>rd</sup> Edition, Burgess Publishing Company, Minneapolis.
- Bergey, D.H., and John, G.H. (1994). *Bergey's manual of determinative bacteriology*. Baltimore Williams and Wilkins.

- Boboye, B., O.F. Olukunle and F.C. Adetuyi, 2010. Degradative activity of bacteria isolated from hydrocarbon-polluted site in Ilaje, Ondo State, Nigeria. *Afr. J. Microbiol. Res.*, 4: 2484-2491.
- Chikere, C.B. and Azubuike, C.C. (2014). Characterization of hydrocarbon utilizing fungi from hydrocarbon polluted sediments and water. *Nig. J. Biotech.* Vol. 27:49-54. ISSN:018917131.
- Human Rights Watch interview with Bruce Powell, zoologist and expert on Niger Delta ecology, London, June 20, 1998.
- Jaja, E. T and Obire, O. (2015). Impact of oilfield wastewater on soil quality and of soils planted with various plants. *Current studies in comparative education, science and technology*. 2(2): 179 – 196.
- Jaja, E. T and Obire, O. (2015). Phytotoxic effect of oilfield wastewater on the physiology and chlorophyll content of some crops. *Current studies in comparative education, science and technology*. 2(2): 198 - 212.
- Kassotis, C. D. Tillitt, D. E. Lin, C.-H. McElroy, J. A. and Nagel, S. C. (2015). “Endocrine-disrupting chemicals and oil and natural gas operations: potential environmental contamination and recommendations to assess complex environmental mixtures”. *Environmental Health Perspectives*.
- Leahy J.G., Colwell R.R. (1990). Microbial degradation of hydrocarbons in the environment. *Microbiol. Rev.* 54: 305–315.
- Malloch, D. (1997). *Moulds Isolation, Cultivation and Identification*. Department of Botany University of Toronto, Toronto USA.
- Marshall, T. R. and Derinny, J. S. (1988). The Microbial ecosystem in petroleum wasteland treatment. *Water Sci. Tech.* 20: 285- 291.
- Mills, A.L., C. Breuil and R.R. Colwell, 1978. Enumeration of petroleum-degrading marine and estuarine microorganisms by the most probable number method. *Can. J. Microbiol.*, 24: 552-557.
- Obire, O., (1988). Studies on the biodegradation potentials of some microorganisms isolated from water systems of two petroleum producing areas in Nigeria. *Nig. J. Bot.*, 1, 81-90 (10 pages).
- Obire, O. and Amusan, F.O. (2003). The Environmental impact of oilfield formation water on a freshwater stream in Nigeria. *J. Appl. Sci. Environ. Mgt.* 7(1): 61-66.
- Obire, O. and Anyanwu, E.C. (2009). Impact of various concentrations of crude oil on fungal populations of soil. *Int. J. Environ. Sci.* 6: 211-218.

- Obire, O and Wemedo, S. A (1996). The effect of Oil Field wastewater on the microbial population of soil in Nigeria. *Niger Delta Biologia*, 1: 77 – 85.
- Odokuma LO, Dickson AA (2003) Bioremediation of a crude oil-polluted tropical rain forest soil. *Glob J Environ Sci* 2:29–40.
- Okerentugba, P.O. and O.U. Ezeronye, 2003. Petroleum degrading potentials of single and mixed microbial cultures isolated from rivers and refinery effluent in Nigeria. *Afr. J. Biotechnol.*, 2: 288-292.
- Okpokwasili, G.C. and Nnubia, C. (1995). The effects of drilling fluids on marine bacteria from a Nigerian offshore oilfield. *Eviron.Manag.* 19: 923-929.
- Okpokwasili, G.C. and Odokuma, L.O. (1996). Effects of oil spill dispersant and drilling fluids on substrate specificity on marine bacteria. *Waste Manag.* 15:515-520.
- Onifade, A.K. and F.A. Abubakar, 2007. Characterization of hydrocarbon-degrading microorganisms isolated from crude oil contaminated soil and remediation of the soil by enhanced natural attenuation. *Res. J. Microbiol.*, 2: 149-155.
- Pichtel J. (2016). Oil and Gas Production Wastewater: Soil Contamination and Pollution Prevention. *Applied and Environmental Soil Science*. 2016: 1-24.
- Prescott, L.M., Harley, J.P. and Klien, D.A. (2005). *Microbiology*. 6<sup>th</sup> ed. McGraw Hill London. Pp. 135-140.
- Rajaram, T. and Ashutost, D. (2008). Water pollution by industrial effluents in India: discharge scenario and case for participatory ecosystem specific local regulation. *Environmental Journal*. 40 (1): 56-69.
- Ugochukwu, C.N.C and Ertel J. (2008). Negative impacts of oil exploration on biodiversity management in the Niger Delta area of Nigeria. *Impact Assessment and Project Appraisal*. 26(2): 139- 147.
- Van Dessel B. (1996). *The Environmental Situation in the Niger Delta, Nigeria*, section 5.5.1.
- Warner, N. R. Jackson, R. B. Darrah T. H. *et al.*, (2012). “Geochemical evidence for possible natural migration of Marcellus formation brine to shallow aquifers in Pennsylvania,” *Proceedings of the National Academy of Sciences of the United States of America*. 109(30): 11961–11966.
- Wemedo, S.A. and Obire, O. (2012). Acute toxicity test of oilfield wastewater on bacterial community of soil in Nigeria. *Research Journal of Environmental Toxicology*. 6:25-32.

Zouboulis, A.I., Loukidou, M.X and Matis, K.A (2004). **Bio-sorption** of toxic metals from aqueous solutions by bacteria strains isolated from metal polluted soils. Journal of Process Biochemistry 39: 90-95.

UNDER PEER REVIEW