

A note on corrections in approximation of the modified error function

Supriya Mandal, Debabrata Singh and M. M. Panja

Department of Mathematics, Visva-Bharati (A Central University),

Santiniketan-731235, West Bengal, India

Research Article

Received: 23 December 2018

Abstract

This article deals with the evaluation of some integrals involving error-, exponential- and algebraic functions with an objective to derive explicit expressions for the second and third order correction terms in the approximation of the modified error function, playing important role in the study of Stefan problem. The results obtained here appear to be new and resolve the lack of desired monotonicity property in the results presented by Ceretania et al.[1]. Results derived here seem to be useful for the researchers working with Stefan problems.

Keywords: Modified error function; Error function; Nonlinear ordinary differential equation; Approximation

1 Introduction

The understanding of phase-change processes are important for scientists and engineers due to their appearance in a wide variety of situations, both in natural and industrial processes. The mathematical modelling of such processes, melting and freezing in particular, as moving boundary problem and their analysis have been exercised since the nineteenth century(1-4).

Cho and Sunderland (5) in 1974 studied a phase-change process for a semi-infinite material (one-dimension) with a (linear) temperature dependent thermal conductivity and appears to be a good approximation for several materials (6). In such work, the solution to the (differential) equation involved in the model was obtained through an auxiliary function having resemblance to the error function $\text{erf}(x)$ (7). Consequently, such function was designated as modified error function (MEF). In spite of the applications in the studies of several phase-change processes (8-16), theoretical aspects e.g. existence and uniqueness, domain of definitions etc. of MEF was overlooked. In their studies (17,18) and references therein, Tarzia and his collaborators investigated the mathematical aspects of MEF as mentioned above.

In their study (18), the MEF has been represented as a series solution

$$\Phi_{\delta}(x) = \sum_{n=0}^{\infty} \phi_n(x) \delta^n, \quad x > 0. \quad (1.1)$$

*Corresponding author: E-mail: madanpanja2005@yahoo.co.in

of the nonlinear boundary value problem

$$[1 + \delta \Phi_\delta(x)] \Phi'_\delta(x) + 2x \Phi'_\delta(x) = 0, \quad (1.2)$$

$$\Phi_\delta(0) = 0, \quad \Phi_\delta(\infty) = 1. \quad (1.3)$$

An approximation $\Psi_{\delta,m}$ of the MEF, $\Phi_\delta(x)$ which is the m -th partial sum, is given by

$$\Psi_{\delta,m}(x) = \sum_{n=0}^m \phi_n(x) \delta^n, \quad x > 0, \quad m \in \mathbb{N}_0. \quad (1.4)$$

The authors in (18) obtained first and second order corrections $\phi_1(x)$ and $\phi_2(x)$ involved in the approximations $\Psi_{\delta,1}(x)$ and $\Psi_{\delta,2}(x)$ of the MEF for $\delta > -1$. But it is observed that $\Psi_{\delta,1}(x)$ appears to be better approximation than $\Psi_{\delta,2}(x)$, which is not desirable. The reason could not be addressed completely. Although the two corrections $\phi_0(x)$ and $\phi_1(x)$ was presented as explicit analytical function, the second order correction $\phi_2(x)$ could not be derived explicitly, rather it is obtained in terms of integrals involving products of exponential and error functions. The authors suggested that the numerical implementation of the integrals present in $\phi_2(x)$ might introduce non-negligible perturbations. During numerical experiment it is found that the order of magnitude of $\phi_2(x)$ was greater than that of $\phi_1(x)$, which raises uncertainty over the convergence of the series (1.1).

To avoid this undesirable property, the authors in this paper have derived explicit expression of $\phi_2(x)$ involving exponential and error functions. Furthermore, the explicit expression of the next order correction $\phi_3(x)$ has been obtained by the evaluation of some integrals involving error function and exponential function. In the course of this derivation some recurrence relation have been obtained, which are not available in the literature yet. With these expressions it is observed that the order of magnitude of the corrections decreases order by order, thus resolves the apparent problem of monotonicity of the successive correction terms that is necessary for the convergence of the series in (1.1). Hence, the inconsonance which arisen in (18) has been dispelled.

2 Approximate Solution

Use of expansion (1.4) in Eq.(1.2) with boundary condition (1.3) suggests that the leading order correction $\phi_0(x)$ is solution to the equation [1]

$$\phi_0''(x) + 2x \phi_0'(x) = 0 \quad (2.1)$$

with

$$\phi_0(0) = 0, \quad \phi_0(\infty) = 1. \quad (2.2)$$

The higher order corrections $\phi_n(x)$, $n \in \mathbb{N}$ are solutions to the equation

$$\phi_n''(x) + 2x \phi_n'(x) = A_{n-1}(x) \quad (2.3)$$

with

$$\phi_n(0) = 0, \quad \phi_n(\infty) = 0. \quad (2.4)$$

Here, $A_{n-1}(x)$ is given as

$$A_{n-1}(x) = - \sum_{k=1}^n \{ \phi_{k-1}'(x) \phi_{n-k}'(x) + \phi_{k-1}(x) \phi_{n-k}''(x) \}. \quad (2.5)$$

The value of the corrections $\phi_n(x)$ ($n \in \mathbb{N}$) in the expansion (1.1) can be calculated by using two fold integration given as

$$\phi_n(x) = \int_0^x e^{-t^2} \int_0^t e^{s^2} A_{n-1}(s) ds dt + c_{n,1} \operatorname{erf}(x). \quad (2.6)$$

Here, $c_{n,1}$ is the integration constant and $\text{erf}(x)$ is error function defined as (7)

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt \quad x > 0. \quad (2.7)$$

Now the solution of Eq.(2.1) with boundary condition (2.2) is given by

$$\phi_0(x) = \text{erf}(x). \quad (2.8)$$

With the use of $\phi_0(x)$, we can calculate $A_0(x)$ and $\phi_1(x)$ from (2.5) and (2.6) respectively as

$$A_0(x) = \frac{4 e^{-x^2}}{\pi} \left\{ \sqrt{\pi} x \text{erf}(x) - e^{-x^2} \right\} \quad (2.9)$$

$$\phi_1(x) = \frac{1}{2\pi} \left\{ 2 - 2 \sqrt{\pi} e^{-x^2} x \text{erf}(x) - \pi \text{erf}(x)^2 - 2 e^{-2x^2} \right\} + c_{1,1} \text{erf}(x). \quad (2.10)$$

Using the boundary condition (2.4) we have

$$c_{1,1} = \frac{(\pi-2)}{2\pi}. \quad (2.11)$$

Using this value of $c_{1,1}$ in (2.10) one gets

$$\phi_1(x) = \frac{1}{2\pi} \left[\left\{ \pi \text{erf}(x) + 2 \right\} \text{erfc}(x) - 2 \sqrt{\pi} x e^{-x^2} \text{erf}(x) - 2 e^{-2x^2} \right]. \quad (2.12)$$

Here $\text{erfc}(x) = 1 - \text{erf}(x)$ is the complimentary error function (7). To obtain the expression for $\phi_2(x)$ we present the following definition and theorems.

Theorem 1. *The integral involving exponential and error function can be represented as*

$$\int_0^x e^{-\lambda y^2} \text{erf}(y) dy = -2 \sqrt{\frac{\pi}{\lambda}} \left\{ T\left(\sqrt{2}\lambda x, \frac{1}{\sqrt{\lambda}}\right) - \frac{1}{2\pi} \tan^{-1}\left(\frac{1}{\sqrt{\lambda}}\right) \right\}. \quad (2.13)$$

Here $T(z, a)$ is Owen T-function defined as (7)

$$T(z, a) = \frac{1}{2\pi} \int_0^a \frac{\exp\left\{-\frac{z^2(1+t^2)}{2}\right\}}{(1+t^2)} dt. \quad (2.14)$$

Proof. Differentiating (2.14) w.r.t z and using the transformation $z t = \sqrt{2} \tau$, we get

$$\begin{aligned} \frac{dT(z, a)}{dz} &= -\frac{1}{2\pi} \int_0^a z \exp\left\{-\frac{z^2(1+t^2)}{2}\right\} dt \\ &= -\frac{1}{\sqrt{2}\pi} \int_0^{\frac{az}{\sqrt{2}}} \exp\left(-\frac{z^2}{2}\right) \exp(-\tau^2) d\tau \\ &= \frac{-1}{2\sqrt{2}\pi} \exp\left(-\frac{z^2}{2}\right) \text{erf}\left(\frac{az}{\sqrt{2}}\right). \end{aligned} \quad (2.15)$$

Further substitution of $az = \sqrt{2} y$ followed by integration with respect to y over $[0, x]$ provides

$$\int_0^x \exp\left(-\frac{y^2}{a^2}\right) \text{erf}(y) dy = -2 a \sqrt{\pi} \left\{ T\left(\frac{\sqrt{2}x}{a}, a\right) - \frac{1}{2\pi} \tan^{-1}(a) \right\}. \quad (2.16)$$

Choice of $a^2 = \frac{1}{\lambda}$ gives the result presented in the statement of the theorem. \square

Lemma 1.

$$T(z, 1) = \frac{1}{8} \left\{ 1 - \operatorname{erf} \left(\frac{z}{\sqrt{2}} \right)^2 \right\}.$$

Proof. We use the following property of Owen T-function (19)

$$T(z, 1) = \frac{1}{2} G(z) (1 - G(z)) \quad (2.17)$$

where $G(z) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^z e^{-\frac{t^2}{2}} dt$.

Now the substitution $\frac{t}{\sqrt{2}} = t'$ converts $G(z)$ into the following form.

$$\begin{aligned} G(z) &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\frac{z}{\sqrt{2}}} e^{-t'^2} dt' \\ &= \frac{1}{\sqrt{\pi}} \left[\int_{-\infty}^0 e^{-t'^2} dt' + \int_0^{\frac{z}{\sqrt{2}}} e^{-t'^2} dt' \right] \\ &= \frac{1}{2} \left[1 + \operatorname{erf} \left(\frac{z}{\sqrt{2}} \right) \right] \end{aligned}$$

Substituting the value of $G(z)$ in (2.17) one can obtain the relation stated in the lemma. \square

Lemma 2.

$$\int_0^x e^{-x^2} \operatorname{erf}(x) dx = \frac{\sqrt{\pi}}{4} \operatorname{erf}(x)^2.$$

Proof. For $\lambda = 1$ in Theorem 1 we have

$$\int_0^x e^{-x^2} \operatorname{erf}(x) dx = -2\sqrt{\pi} \left\{ T(\sqrt{2}x, 1) - \frac{1}{2\pi} \tan^{-1}(1) \right\}.$$

Using the result obtained in Lemma 1 the statement in this lemma can be proved. \square

Definition 1. We define the notation $I_{m,n,\lambda}(x)$ as follows

$$I_{m,n,\lambda}(x) = \int_0^x e^{-\lambda t^2} t^n \operatorname{erf}(t)^m dt. \quad (2.18)$$

Lemma 3. From Definition-1 it can be observed that

$$\int_0^x e^{-\lambda t^2} t^n \operatorname{erf}(bt)^m dt = \frac{1}{b^{n+1}} I_{m,n,\frac{\lambda}{b^2}}(bx) \quad (2.19)$$

where $b \in \mathbb{R}$ is constant.

Proof. Substitution of $bt = t'$ in the integral mentioned in the Lemma converts it into

$$\begin{aligned} &\frac{1}{b^{n+1}} \int_0^{bx} e^{-\frac{\lambda t'^2}{b^2}} t'^n \operatorname{erf}(t')^m dt' \\ &= \frac{1}{b^{n+1}} I_{m,n,\frac{\lambda}{b^2}}(bx). \end{aligned}$$

\square

Theorem 2. The integral $I_{m,n,\lambda}(x)$ given in (2.18) satisfies the recurrence relation

$$I_{m,n,\lambda}(x) = -\frac{1}{2\lambda} e^{-\lambda x^2} x^{n-1} \operatorname{erf}(x)^m + \frac{n-1}{2\lambda} I_{m,n-2,\lambda}(x) + \frac{m}{\lambda\sqrt{\pi}} I_{m-1,n-1,\lambda+1}(x) \quad (2.20)$$

where $\lambda > 0$, $m \geq 0$, $n \geq 1$. It may be observed that

$$I_{0,0,\lambda}(x) = \frac{\sqrt{\pi}}{2} \operatorname{erf}(x) \quad (2.21)$$

$$I_{0,1,\lambda}(x) = -\frac{1}{2\lambda} (e^{-\lambda x^2} - 1) \quad (2.22)$$

$$I_{1,0,\lambda}(x) = -2\sqrt{\frac{\pi}{\lambda}} \left\{ T\left(\sqrt{2\lambda}x, \frac{1}{\sqrt{\lambda}}\right) - \frac{1}{2\pi} \tan^{-1}\left(\frac{1}{\sqrt{\lambda}}\right) \right\} \quad (2.23)$$

$$I_{m,0,1}(x) = \frac{\sqrt{\pi}}{2(m+1)} \operatorname{erf}(x)^{(m+1)}. \quad (2.24)$$

Proof.

$$\begin{aligned} I_{m,n,\lambda}(x) &= \int_0^x e^{-\lambda t^2} t^n \operatorname{erf}(t)^m dt \\ &= -\frac{1}{2\lambda} \int_0^x \frac{d}{dt} (e^{-\lambda t^2}) t^{n-1} \operatorname{erf}(t)^m dt \\ &= -\frac{1}{2\lambda} \left[t^{n-1} \operatorname{erf}(t)^m \int \frac{d}{dt} e^{-\lambda t^2} dt \right]_0^x + \frac{1}{2\lambda} \left[\int_0^x \frac{d}{dt} (t^{n-1} \operatorname{erf}(t)^m) \int \frac{d}{dt} (e^{-\lambda t^2}) dt dt \right] \\ &= -\frac{1}{2\lambda} e^{-\lambda x^2} x^{n-1} \operatorname{erf}(x)^m + \frac{n-1}{2\lambda} \int_0^x e^{-\lambda t^2} t^{n-2} \operatorname{erf}(t)^m dt \\ &\quad + \frac{m}{\lambda\sqrt{\pi}} \int_0^x e^{-(\lambda+1)t^2} t^{n-1} \operatorname{erf}(t)^{m-1} dt \\ &= -\frac{1}{2\lambda} e^{-\lambda x^2} x^{n-1} \operatorname{erf}(x)^m + \frac{n-1}{2\lambda} I_{m,n-2,\lambda}(x) + \frac{m}{\lambda\sqrt{\pi}} I_{m-1,n-1,\lambda+1}(x). \end{aligned}$$

Results in (2.21), (2.22) are obtained by straightforward integration using the *Definition-1* while the result in (2.23) can be established by using *Theorem-1*.
Now,

$$\begin{aligned} I_{m,0,1}(x) &= \int_0^x \operatorname{erf}(t)^m e^{-t^2} dt \\ &= \frac{\sqrt{\pi}}{2(m+1)} \int_0^x \frac{d}{dt} (\operatorname{erf}(t)^{m+1}) dt \\ &= \frac{\sqrt{\pi}}{2(m+1)} \operatorname{erf}(x)^{m+1}. \end{aligned}$$

This completes the proof. □

Theorem 3. Using *Definition-1* we have the following recurrence relation for $\lambda = 0$

$$I_{m,n,0}(x) = \frac{x^{n+1}}{n+1} \operatorname{erf}(x)^m - \frac{2m}{\sqrt{\pi}(n+1)} I_{m-1,n+1,1}(x). \quad (2.25)$$

Proof. We use the formula for integration by parts to get

$$\begin{aligned} I_{m,n,0}(x) &= \int_0^x t^n \operatorname{erf}(t)^m dt \\ &= \frac{t^{n+1}}{n+1} \operatorname{erf}(t)^m \Big|_0^x - \frac{2m}{\sqrt{\pi}(n+1)} \int_0^x \operatorname{erf}(t)^{m-1} e^{-t^2} t^{n+1} dt \\ &= \frac{x^{n+1}}{n+1} \operatorname{erf}(x)^m - \frac{2m}{\sqrt{\pi}(n+1)} I_{m-1,n+1,1}(x). \end{aligned}$$

□

Theorem 4. The explicit expression for the second order correction $\phi_2(x)$ in the approximation of modified error function $\Phi_\delta(x)$ can be obtained as

$$\begin{aligned} \phi_2(x) &= \frac{1}{4\pi^2} \left[-2\pi \operatorname{erf}(x)^2 \left\{ \pi \operatorname{erfc}(x) - 2 \right\} - 4e^{-2x^2} \left\{ \pi(x^2 - 3) \operatorname{erf}(x) + \pi - 2 \right\} \right. \\ &\quad + \sqrt{\pi} e^{-x^2} x \left\{ \operatorname{erf}(x) \left\{ \pi(9 - 2x^2) \operatorname{erf}(x) - 4\pi + 8 \right\} - 4 \right\} \\ &\quad \left. + \left\{ \pi(3\sqrt{3} - 8) + 8 \right\} \operatorname{erf}(x) - 3\sqrt{3}\pi \operatorname{erf}(\sqrt{3}x) - 2\sqrt{\pi} e^{-3x^2} x + 4\pi - 8 \right]. \end{aligned} \quad (2.26)$$

Proof. From (2.5) one can get,

$$A_1(x) = -\left\{ 2\phi'_0(x)\phi'_1(x) + \phi_1(x)\phi''_0(x) + \phi_0(x)\phi''_1(x) \right\} \quad (2.27)$$

Explicit expressions for $\phi_0(x)$, $\phi_1(x)$ given in (2.8) and (2.12) have been used to obtained

$$\begin{aligned} A_1(x) &= \frac{4e^{-3x^2}}{\pi^2} \left[-e^{x^2} \left\{ 2\pi(x^2 - 2) \operatorname{erf}(x) + \pi - 2 \right\} \right. \\ &\quad \left. + \sqrt{\pi} e^{2x^2} x \left\{ \operatorname{erf}(x) \left\{ \pi(x^2 - 3) \operatorname{erf}(x) + \pi - 2 \right\} + 1 \right\} - 3\sqrt{\pi} x \right]. \end{aligned} \quad (2.28)$$

Using the integral representation of $\phi_n(x)$ in (2.6) for $n = 2$, one can obtain the explicit x -dependence of ϕ_2 with the help of *Lemma-2* and *Theorem-2* as

$$\begin{aligned} \phi_2(x) &= \frac{e^{-3x^2}}{4\pi^2} \left[-4e^{x^2} \left\{ \pi(x^2 - 3) \operatorname{erf}(x) + \pi - 2 \right\} - 2\sqrt{\pi} x \right. \\ &\quad - \sqrt{\pi} e^{2x^2} x \left\{ \pi(2x^2 - 9) \operatorname{erf}(x)^2 + 4(\pi - 2) \operatorname{erf}(x) + 4 \right\} \\ &\quad \left. + e^{3x^2} \left\{ -2\pi \operatorname{erf}(x)^2 (\pi \operatorname{erfc}(x) - 2) - 3\sqrt{3}\pi \operatorname{erf}(\sqrt{3}x) + 4\pi - 8 \right\} \right] \\ &\quad + c_{2,1} \operatorname{erf}(x). \end{aligned} \quad (2.29)$$

Use of the boundary condition (2.4) for $n = 2$, properties of error- and complementary error-functions provides the integration constant

$$c_{2,1} = \frac{(3\pi\sqrt{3} - 8\pi + 8)}{4\pi^2}. \quad (2.30)$$

Relations in (2.29) and (2.30) simultaneously recover the statement of *Theorem-4*. □

The result obtained here appears to be new.

To derive third order correction term $\phi_3(x)$, we recall (2.5) to obtain

$$A_2(x) = -\left\{ \phi'_1(x)^2 + 2\phi'_0(x)\phi'_2(x) + \phi''_0(x)\phi_2(x) + \phi_1(x)\phi''_1(x) + \phi_0(x)\phi''_2(x) \right\}. \quad (2.31)$$

Use of explicit expression for $\phi_i(x)$, $i = 0, 1, 2$ into the above expression yields a large expression in x which has been split into five parts as

$$A_2(x) = \sum_{k=1}^5 A_{2,k}(x) \quad (2.32)$$

where

$$A_{2,1}(x) = -\frac{e^{-2x^2}}{\pi^2} \left(8x^2 + 6\sqrt{3} + \pi - 28 + \frac{20}{\pi} \right) - \frac{18e^{-3x^2}}{\pi^{5/2}} (\pi - 2)x - \frac{2}{\pi^2} e^{-4x^2} (7x^2 + 4), \quad (2.33)$$

$$A_{2,2}(x) = \left[\frac{e^{-x^2}}{\pi^{5/2}} x \left\{ \pi (8x^2 + 6\sqrt{3} + \pi - 36) + 20 \right\} - \frac{12e^{-2x^2}}{\pi^2} (\pi - 2) (x^2 - 2) - \frac{2e^{-3x^2}}{\pi^{3/2}} x (13x^2 - 32) \right] \text{erf}(x), \quad (2.34)$$

$$A_{2,3}(x) = \left[\frac{6e^{-x^2}}{\pi^{3/2}} x (\pi - 2) (x^2 - 3) - \frac{e^{-2x^2}}{\pi} (10x^4 - 51x^2 + 48) \right] \text{erf}(x)^2, \quad (2.35)$$

$$A_{2,4}(x) = \frac{e^{-x^2}}{2\sqrt{\pi}} (4x^4 - 36x^2 + 59) \text{erf}(x)^3, \quad (2.36)$$

$$A_{2,5}(x) = -\frac{3e^{-x^2}}{\pi^{5/2}} x \left\{ \sqrt{3} \pi \text{erf}(\sqrt{3}x) - 2\pi + 4 \right\}. \quad (2.37)$$

Accordingly, we write $\phi_3(x)$ as

$$\phi_3(x) = \sum_{k=1}^5 \phi_{3,k}(x) + c_{3,1} \text{erf}(x) \quad (2.38)$$

where

$$\phi_{3,k}(x) = \int_0^x e^{-t^2} \int_0^t e^{s^2} A_{2,k}(s) ds dt. \quad (2.39)$$

Using *Definition-1* and the values of $A_{2,k}(x)$, $k = 1, 2, \dots, 5$ given in (2.33)-(2.37) we can express the integrals $\int_0^t e^{s^2} A_{2,k}(s) ds$, $k = 1, 2, \dots, 5$ as follows.

$$\begin{aligned} \int_0^t e^{s^2} A_{2,1}(s) ds &= -\frac{8}{\pi^2} I_{0,2,1}(t) - \frac{1}{\pi^2} \left(6\sqrt{3} + \pi - 28 + \frac{20}{\pi} \right) I_{0,0,1}(t) \\ &\quad - \frac{18}{\pi^{5/2}} (\pi - 2) I_{0,1,2}(t) - \frac{14}{\pi^2} I_{0,2,3}(t) - \frac{8}{\pi^2} I_{0,0,3}(t), \end{aligned} \quad (2.40)$$

$$\begin{aligned} \int_0^t e^{s^2} A_{2,2}(s) ds &= \frac{8}{\pi^{3/2}} I_{1,3,0}(t) + \left(\frac{1}{\pi^{3/2}} (6\sqrt{3} + \pi - 36) + \frac{20}{\pi^{5/2}} \right) I_{1,1,0}(t) - \frac{12}{\pi^2} (\pi - 2) I_{1,2,1}(t) \\ &\quad + \frac{24}{\pi^2} (\pi - 2) I_{1,0,1}(t) - \frac{26}{\pi^{3/2}} I_{1,3,2}(t) + \frac{64}{\pi^{3/2}} I_{1,1,2}(t), \end{aligned} \quad (2.41)$$

$$\begin{aligned} \int_0^t e^{s^2} A_{2,3}(s) ds &= \frac{6}{\pi^{3/2}} (\pi - 2) I_{2,3,0}(t) - \frac{18}{\pi^{3/2}} (\pi - 2) I_{2,1,0}(t) - \frac{10}{\pi} I_{2,4,1}(t) \\ &\quad + \frac{51}{\pi} I_{2,2,1}(t) - \frac{48}{\pi} I_{2,0,1}(t), \end{aligned} \quad (2.42)$$

$$\int_0^t e^{s^2} A_{2,4}(s) ds = \frac{2}{\sqrt{\pi}} I_{3,5,0}(t) - \frac{18}{\sqrt{\pi}} I_{3,3,0}(t) + \frac{59}{2\sqrt{\pi}} I_{3,1,0}(t), \quad (2.43)$$

$$\int_0^t e^{s^2} A_{2,5}(s) ds = \frac{6}{\pi^{\frac{5}{2}}}(\pi - 2) I_{0,1,0}(t) - \frac{\sqrt{3}}{\pi^{\frac{3}{2}}} I_{1,1,0}(\sqrt{3}t). \quad (2.44)$$

We now derive the explicit expressions of $I_{m,n,\lambda}(t)$ for various values of m, n, λ appearing in (2.40)-(2.44) by the use of *Lemma-2*, *Lemma-3*, *Theorem-2* and *Theorem-3*. Then using the obtained results and the definition of $\phi_{3,k}(x)$ given in (2.39) one can find the expressions of $\phi_{3,k}(x)$, $k = 1, 2, \dots, 5$ in terms of $I_{m,n,\lambda}(x)$ as

$$\begin{aligned} \phi_{3,1}(x) &= \frac{9}{\pi^{\frac{5}{2}}} \left(\frac{\pi}{2} - 1 \right) I_{0,0,3}(x) + \frac{7}{3\pi^2} I_{0,1,4}(x) + \frac{4}{\pi^2} I_{0,1,2}(x) - \frac{9}{\pi^{\frac{5}{2}}} \left(\frac{\pi}{2} - 1 \right) I_{0,0,1}(x) \\ &+ \left(\frac{12}{\pi^{3/2}} - \frac{10}{\pi^{5/2}} - \frac{3\sqrt{3}}{\pi^{3/2}} - \frac{1}{2\sqrt{\pi}} \right) I_{1,0,1}(x) - \frac{31}{18\pi^{\frac{3}{2}}} I_{1,0,\frac{1}{3}}(\sqrt{3}x), \end{aligned} \quad (2.45)$$

$$\begin{aligned} \phi_{3,2}(x) &= \frac{13}{6\pi^2} I_{0,1,4}(x) + \frac{1}{\pi} \left(\frac{3\sqrt{3}}{\pi} + \frac{1}{2} - \frac{15}{\pi} + \frac{10}{\pi^2} \right) I_{0,1,2}(x) + \frac{2}{\pi^2} I_{0,3,2}(x) + \frac{6}{\pi} \left(1 - \frac{2}{\pi} \right) I_{1,1,2}(x) \\ &+ \frac{3}{\pi^{3/2}} \left(1 - \frac{2}{\pi} \right) I_{0,0,3}(x) - \frac{51}{4\pi^{3/2}} I_{1,0,3}(x) + \frac{13}{2\pi^{3/2}} I_{1,2,3}(x) + \frac{3}{\pi^{3/2}} \left(\frac{2}{\pi} - 1 \right) I_{0,0,1}(x) \\ &+ \frac{1}{\sqrt{\pi}} \left(-\frac{3\sqrt{3}}{2\pi} + \frac{15}{2\pi} - \frac{5}{\pi^2} - \frac{1}{4} \right) I_{1,0,1}(x) + \frac{1}{\sqrt{\pi}} \left(\frac{10}{\pi^2} + \frac{3\sqrt{3}}{\pi} - \frac{18}{\pi} + \frac{1}{2} \right) I_{1,2,1}(x) \\ &+ \frac{2}{\pi^{3/2}} I_{1,4,1}(x) + \frac{9}{\pi^{3/2}} \left(\frac{\pi}{2} - 1 \right) I_{2,0,1}(x) + \frac{35}{9\pi^{3/2}} I_{1,0,\frac{1}{3}}(\sqrt{3}x), \end{aligned} \quad (2.46)$$

$$\begin{aligned} \phi_{3,3}(x) &= \frac{5}{3\pi^2} I_{0,1,4}(x) + \frac{12}{\pi^{5/2}} \left(1 - \frac{\pi}{2} \right) I_{0,0,3}(x) - \frac{3}{\pi^{5/2}} \left(1 - \frac{\pi}{2} \right) I_{0,2,3}(x) - \frac{31}{2\pi^{3/2}} I_{1,0,3}(x) \\ &+ \frac{5}{\pi^{3/2}} I_{1,2,3}(x) + \frac{27}{\pi^2} \left(1 - \frac{\pi}{2} \right) I_{1,1,2}(x) - \frac{6}{\pi^2} \left(1 - \frac{\pi}{2} \right) I_{1,3,2}(x) - \frac{18}{\pi} I_{2,1,2}(x) \\ &+ \frac{5}{\pi} I_{2,3,2}(x) - \frac{12}{\pi^{5/2}} \left(1 - \frac{\pi}{2} \right) I_{0,0,1}(x) - \frac{27}{4\pi^{3/2}} \left(1 - \frac{\pi}{2} \right) I_{2,0,1}(x) \\ &+ \frac{18}{\pi^{3/2}} \left(1 - \frac{\pi}{2} \right) I_{2,2,1}(x) - \frac{3}{\pi^{3/2}} \left(1 - \frac{\pi}{2} \right) I_{2,4,1}(x) - \frac{5}{\sqrt{\pi}} I_{3,0,1}(x) \\ &+ \frac{44}{9\pi^{3/2}} I_{1,0,\frac{1}{3}}(\sqrt{3}x), \end{aligned} \quad (2.47)$$

$$\begin{aligned} \phi_{3,4}(x) &= -\frac{19}{6\pi^2} I_{0,1,4}(x) + \frac{1}{3\pi^2} I_{0,3,4}(x) + \frac{91}{4\pi^{3/2}} I_{1,0,3}(x) - \frac{10}{\pi^{3/2}} I_{1,2,3}(x) \\ &+ \frac{1}{\pi^{3/2}} I_{1,4,3}(x) + \frac{111}{4\pi} I_{2,1,2}(x) - \frac{11}{\pi} I_{2,3,2}(x) + \frac{1}{\pi} I_{2,5,2}(x) \\ &- \frac{37}{8\sqrt{\pi}} I_{3,0,1}(x) + \frac{59}{4\sqrt{\pi}} I_{3,2,1}(x) - \frac{9}{2\sqrt{\pi}} I_{3,4,1}(x) + \frac{1}{3\sqrt{\pi}} I_{3,6,1}(x) \\ &- \frac{127}{18\pi^{3/2}} I_{1,0,\frac{1}{3}}(\sqrt{3}x), \end{aligned} \quad (2.48)$$

$$\begin{aligned} \phi_{3,5}(x) &= -\frac{3}{2\pi^2} I_{0,1,4}(x) - \frac{6}{\pi^{5/2}} \left(1 - \frac{\pi}{2} \right) I_{0,2,1}(x) + \frac{1}{4\pi^{3/2}} I_{1,0,\frac{1}{3}}(\sqrt{3}x) \\ &- \frac{1}{2\pi^{3/2}} I_{1,2,\frac{1}{3}}(\sqrt{3}x). \end{aligned} \quad (2.49)$$

We again derive the expressions for $I_{m,n,\lambda}(x)$ for different m, n, λ present in (2.45)-(2.49) by using *Lemma-2*, *Lemma-3*, *Theorem-2* and *Theorem-3* to obtain the explicit expressions for $\phi_{3,k}(x)$, $k =$

1, 2, ..., 5 given by

$$\begin{aligned}\phi_{3,1}(x) = & -\frac{1}{216\pi^2} \left[63 e^{-4x^2} + 216 e^{-2x^2} - 62\sqrt{3}\pi - 279 + 27 \left\{ 6\pi (\sqrt{3} - 4) + \pi^2 + 20 \right\} \times \right. \\ & \text{erf}(x)^2 + 324 \sqrt{3} \text{erf}(\sqrt{3}x) - 162 \sqrt{3} \pi \text{erf}(\sqrt{3}x) + 6 \left\{ 31\sqrt{3} \pi \text{erf}(\sqrt{3}x) \right. \\ & \left. \left. + 81(\pi - 2) \right\} \text{erf}(x) + 744\sqrt{3}\pi T\left(\sqrt{6}x, \frac{1}{\sqrt{3}}\right) \right], \quad (2.50)\end{aligned}$$

$$\begin{aligned}\phi_{3,2}(x) = & \frac{1}{216\pi^3} \left[54 \left\{ \pi (6\sqrt{3} + \pi - 28) + 20 + e^{-2x^2} \left\{ -\pi (4x^2 + 6\sqrt{3} + \pi - 28) - 20 \right\} \right. \right. \\ & - \sqrt{\pi} e^{x^2} x \left\{ \pi (4x^2 + 6\sqrt{3} + \pi - 30) + 20 \right\} \text{erf}(x) \left. \right\} + 54(\pi - 2)\pi \times \\ & \left\{ -6 (e^{-2x^2} + 1) \text{erf}(x) + 3\pi \text{erf}(x)^3 + 4\sqrt{3} \text{erf}(\sqrt{3}x) \right\} + \pi \left\{ 117(1 - e^{-4x^2}) \right. \\ & - 234\sqrt{\pi} e^{-3x^2} x \text{erf}(x) + 140\sqrt{3}\pi \left\{ 24 T\left(\sqrt{6}x, \frac{1}{\sqrt{3}}\right) \right. \\ & \left. \left. + 3 \text{erf}(x) \text{erf}(\sqrt{3}x) - 2 \right\} \right] \right], \quad (2.51)\end{aligned}$$

$$\begin{aligned}\phi_{3,3}(x) = & \frac{1}{72\pi^{5/2}} \left[27(\pi - 2) \left\{ \pi e^{-x^2} x (9 - 2x^2) \text{erf}(x)^2 - 4\sqrt{\pi} e^{-2x^2} (x^2 - 4) \text{erf}(x) \right. \right. \\ & + 8\sqrt{\pi} \text{erf}(x) - 23\sqrt{\frac{\pi}{3}} \text{erf}(\sqrt{3}x) - 2e^{-3x^2} x \left. \right\} + \sqrt{\pi} e^{-4x^2} \left\{ -120\sqrt{\pi} e^{x^2} x \text{erf}(x) \right. \\ & - 45 - 9\pi e^{2x^2} (10x^2 - 31) \text{erf}(x)^2 \left. \right\} + \sqrt{\pi} \left\{ 45 - 45\pi^2 \text{erf}(x)^4 \right. \\ & \left. \left. + 176\sqrt{3}\pi \left\{ 12 T\left(\sqrt{6}x, \frac{1}{\sqrt{3}}\right) + \text{erf}(x) \text{erf}(\sqrt{3}x) - 1 \right\} \right\} \right] \right], \quad (2.52)\end{aligned}$$

$$\begin{aligned}\phi_{3,4}(x) = & \frac{1}{216\pi^2} \left[e^{-4x^2} (333 - 36x^2) - 54\sqrt{\pi} e^{-3x^2} x (2x^2 - 19) \text{erf}(x) \right. \\ & - 27\pi e^{-2x^2} (4x^4 - 40x^2 + 91) \text{erf}(x)^2 - 9\pi^{3/2} e^{-x^2} x (4x^4 - 44x^2 + 111) \text{erf}(x)^3 \\ & \left. - \left\{ 254\pi\sqrt{3} \left\{ 48 T\left(\sqrt{6}x, \frac{1}{\sqrt{3}}\right) + 3\text{erf}(x) \text{erf}(\sqrt{3}x) - 4 \right\} + 333 \right\} \right] \right], \quad (2.53)\end{aligned}$$

$$\begin{aligned}\phi_{3,5}(x) = & \frac{1}{12\pi^{5/2}} \left[9\sqrt{\pi} e^{-4x^2} + 9e^{-x^2} x \left\{ \sqrt{3} \pi \text{erf}(\sqrt{3}x) - 2\pi + 4 \right\} + \sqrt{\pi} \left\{ \sqrt{3}\pi - 9 \right. \right. \\ & \left. \left. - 12\sqrt{3}\pi T\left(\sqrt{6}x, \frac{1}{\sqrt{3}}\right) - 3\text{erf}(x) \left\{ \sqrt{3} \pi \text{erf}(\sqrt{3}x) - 3\pi + 6 \right\} \right\} \right] \right]. \quad (2.54)\end{aligned}$$

Use of (2.50-2.54) in (2.38) and the boundary condition in (2.4) for $n = 3$, gives

$$c_{3,1} = \frac{1}{24\pi^3} \left\{ (19\sqrt{3} - 42)\pi^2 + (224 - 90\sqrt{3})\pi - 120 \right\}. \quad (2.55)$$

Substituting the derived expressions of $\phi_{3,k}(x)$, $k = 1, 2, \dots, 5$ and the value of $c_{3,1}$ in (2.38) we obtain the explicit expression for the third order correction $\phi_3(x)$ involving error-, complementary error- and

Owen T -functions as

$$\begin{aligned}
 \phi_3(x) = & -\frac{(x^2-5)e^{-4x^2}}{6\pi^2} - \frac{xe^{-3x^2}}{4\pi^{\frac{5}{2}}} \left\{ 3(\pi-2) + 2\pi(x^2-4) \operatorname{erf}(x) \right\} \\
 & - \frac{e^{-2x^2}}{4\pi^3} \left[20 + \pi^2 \left\{ 1 + 6(x^2-3) \operatorname{erf}(x) + (2x^4-15x^2+30) \operatorname{erf}(x)^2 \right\} \right. \\
 & \left. + 2\pi \left\{ 6 + 3\sqrt{3} - 4x^2 + 6(x^2-3) \operatorname{erfc}(x) \right\} \right] \\
 & - \frac{xe^{-x^2}}{24\pi^{\frac{5}{2}}} \left[6 \left\{ 20 + \pi(4x^2 + \pi + 6\sqrt{3} - 30) \right\} \operatorname{erf}(x) + 9\pi(\pi-2)(2x^2-9) \operatorname{erf}(x)^2 \right. \\
 & \left. + \pi^2(4x^4 - 44x^2 + 111) \operatorname{erf}(x)^3 - 18(4-2\pi + \sqrt{3}\pi \operatorname{erf}(\sqrt{3}x)) \right] \\
 & + \frac{1}{24\pi^3} \left[120 + \pi \left\{ 90\sqrt{3} - 104 + \pi(12\pi + 17\sqrt{3} + 6) \right\} + 6\sqrt{3}\pi^2 \operatorname{erf}(\sqrt{3}x) \right] \operatorname{erfc}(x) \\
 & - \frac{1}{8\pi^2} \left\{ 20 + 6(\sqrt{3}+2)\pi + 13\pi^2 \right\} \operatorname{erfc}(x)^2 + \frac{1}{4\pi} (7\pi+6) \operatorname{erfc}(x)^3 - \frac{15}{24} \operatorname{erfc}(x)^4 \\
 & + \frac{\sqrt{3}}{8\pi^2} \left\{ (11\pi-18) \operatorname{erfc}(\sqrt{3}x) - 128\pi T\left(\sqrt{6}x, \frac{1}{\sqrt{3}}\right) \right\}. \quad (2.56)
 \end{aligned}$$

This explicit expression for $\phi_3(x)$ seems to be new, not available in literature.

To exhibit the usefulness of the results derived here we have compared the correction terms $\phi_n(x)$ in (1.4) with the results obtained by Ceretani et al (18) in Fig.1. It is observed that the lack of monotonicity of successive corrections of MEF presented in (18) disappears.

Over and above, the residual error

$$ER[\Psi](x) \equiv \left[\{1 + \delta \Psi(x)\} \Psi'(x) \right]^{\delta} + 2x \Psi'(x)$$

for the approximate solutions $\Psi_{\delta,1}(x)$ derived here and $y_{i,p}(x; \delta)$, $i = 1, 2$ obtained recently by Bougoffa (19) have been presented in Fig. 2a and Fig. 2b for $\delta = 0.15$ and 0.1 , respectively. From a careful analysis of these figures it appears that the residual errors corresponding to the approximate solution $\Psi_{\delta,1}(x)$ derived here seem to be significantly less than the same corresponding to the approximate solutions presented by Bougoffa (19).

The monotonicity of the $ER[\Psi_{\delta,m}](x)$ corresponding to the approximate solution involving higher order corrections ($m = 2, 3$) has also been maintained as evident from the Fig.4.

3 Discussion

The main goal of this paper is the derivation of the explicit expressions for the second and third order corrections in the approximation of the MEF which satisfies Eq.(1.2). Results presented here have been derived through the evaluation of integrals involving error-, exponential- and algebraic functions. The plots of successive corrections $\phi_i(x)$, $0 \leq i \leq 3$ show that the order of magnitude is decreasing term by term with $|\frac{\phi_{n+1}}{\phi_n}| \leq \frac{1}{2}$. It indicates that the series in (1.1) seems to converge for $\delta \leq 2$. Most of the results obtained here appear to be new and resolve the lack of monotonicity of successive corrections in the approximation of MEF appearing in (18). The results derived here may be useful for the researchers working in the field of Stefan problems e.g., in the approximations of generalized error function introduced by Ceretani et al in their recent work (20).

The limitation of this approximation scheme is that the derivation of the explicit expressions for the next order corrections ($\phi_n(x)$, $n \geq 4$) involves intricate calculations due to the presence of integrals containing Owen T -functions which are not even manageable with the help of symbolic computations in a straightforward way. So, an alternative approximation scheme for the modified- and generalized-error function $\Phi_\delta(x)$ with higher order accuracy may be desirable.




Figure 1: $\phi_n(x)$ for $n = 0, 1, 2, 3$ involved in the approximate solution (1.4). Curves with dashed-lines representing results of Ceretani et al (18) while the rests (red) represent corrections derived here.

Acknowledgment

Authors are thankful to the Reviewers for their constructive suggestions for improvement in the presentation of this paper. This work is supported partly by DST funded INSPIRE fellowship (No-IF170200) (DS) and the UGC assisted SAP program (DRS Phase II F.510/4/DRS/2009 (SAP-I) through the Department of Mathematics, Visva-Bharati.

References

- [1] G. Lame, B.P.E. Clapeyron, Memoire sur la solidification par refroidissement dn globe liquide, Ann. Chem. Phys. 1831;47:250-256.
- [2] J. Stefan, Über einige probleme der theorie der warmeleitung, Zitzungsberichte Kaiserlichen Akad. Wiss. Cl. 1889;98:473-484.
- [3] J. Stefan, Über die diffusion von sauren und basen gegen einander, Zitzungsberichte Kaiserlichen Akad. Wiss. Cl. 1889;98:616-634.




Figure 2: $ER[\Psi](x)$ for $y_{1,p}(x; \delta)$ (dotted-line), $y_{2,p}(x; \delta)$ (dashed-line) and $\Psi_{\delta,1}(x)$ (red-line) corresponding to a) $\delta = .15$ and b) $\delta = .1$.

- [4] J. Stefan, Über die verdampfung und die auflösung als vorgänge diffusion, Sitzungsberichte Kaiserlichen Akad. Wiss. Cl. 1889;98:1418-1442.
- [5] S.H. Cho, J.E. Sunderland, Phase-change problems with temperature-dependent thermal conductivity, J. Heat Transf. 1974;96:214-217.
- [6] V. Alexiades, A.D. Solomon, Mathematical Modeling of Melting and Freezing Processes, Hemisphere Pub. Corp. Washington, 1993.
- [7] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, C.W. Clark (Eds.), NIST Handbook of Mathematical Functions, Cambridge University Press, NY, 2010.
- [8] A.C. Briozzo, M.F. Natale, D.A. Tarzia, Existence of an exact solution for a one-phase Stefan problem with nonlinear thermal coefficients from Tirskaa's method, Nonlinear Anal. 2007;67.
- [9] C. Wigner, Diffusion of lead chloride dissolved in solid silver chloride, J. Chem. Phys. 1950;18:1227-1230.
- [10] J.I. Frankel, B. Vick, An exact methodology for solving nonlinear diffusion equations based on integral transforms, Appl. Numer. Math. 1987;3:467-477.
- [11] D.L.R. Oliver, J.E. Sunderland, A phase-change problem with temperature-dependent thermal conductivity and specific heat, Int. J. Heat Mass Transfer, 1987;30:2657-2661.
- [12] M. Countryman, R. Kannan, Nonlinear boundary value problem on semi-infinite intervals, Comput. Appl. Math. Appl. 1994;3:59-75.
- [13] D.A. Tarzia, The determination of unknown thermal coefficients through phase-change process with temperature-dependent thermal conductivity, Int. Commun. Heat Mass Transf. 1998;25:139-147.
- [14] A.C. Briozzo, M.F. Natale, D.A. Tarzia, Existence of an exact solution for a one-phase Stefan problem with nonlinear thermal coefficients from Tirskaa's method, Nonlinear Anal. 2007;67:1989-1998.
- [15] N.N. Salva, D.A. Tarzia, A sensitivity analysis for the determination of unknown thermal coefficients through a phase-change process with temperature dependent thermal conductivity, Int. Commun. Heat Mass Transfer, 2011;38:418-424.
- [16] N.N. Salva, D.A. Tarzia, Simultaneous determination of unknown coefficients through a phase-change process with temperature-dependent thermal conductivity, JP J. Heat Mass Transfer, 2011;5:11-39.




Figure 3: $ER[\Psi_{\delta,m}](x)$ for a) $\Psi_{\delta,2}(x)$ and b) $\Psi_{\delta,3}(x)$. In both cases, curves with green and red lines represent residual error corresponding to a) $\delta = .15$ and b) $\delta = .1$, respectively.

- [17] A.N. Ceretania, N.N. Salva, D.A.Tarzia, Existence and uniqueness of the modified error function, Appl. Math. Lett. 2017;70:14-17.
- [18] A.N. Ceretania, N.N. Salva, D.A.Tarzia, Approximation of the modified error function, Appl. Math. Comput. 2018;337:607-617.
- [19] D.B. Owen, A table of normal integrals, Commun. Stat. Simul. Comput. 1980;9:389-419.
- [20] L. Bougoffa, A note on the existence and uniqueness solutions of the modified error function, Math Meth Appl Sci. 2018;doi:10.1002/mma.5095.
- [21] A.N. Ceretania, N.N. Salva, D.A.Tarzia, An exact solution to a Stefan problem with variable thermal conductivity and Robin boundary condition, Nonlin. Anal.: Real World Appl. 2018;40:243-259.