

Paper title: Haematologic and Gastric Histological Changes Associated with Administration of Ground Nutmeg (*Myristica fragrans*) Seed on Adult Male Albino Wistar Rats

Names of authors: *Christopher Chiedozie Mbadugha,¹ Gabriel Donatus Edem,¹ Uka Emmanuel² Isaac Etim Udoudo¹

Affiliation: ¹Department of Anatomy, Faculty of Basic Medical Sciences, University of Uyo, P.M.B. 1017, Uyo, Akwa Ibom State, Nigeria.

²Department of Biochemistry, Faculty of Basic Medical Sciences, University of Uyo, P.M.B. 1017, Uyo, Akwa Ibom State, Nigeria

Authors' contributions

This work was carried out in collaboration between all authors. Authors CMC, DEG, UE and IUE designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors CMC, DEG, UE and IUE, also managed the analyses of the study, and the literature searches. All authors read and approved the final manuscript.

Corresponding author's name: Christopher Chiedozie Mbadugha
E – mail: christophermbadugha@uniuyo.edu.ng

Haematologic and Gastric Histological Changes Associated with Administration of Ground Nutmeg (*Myristica fragrans*) Seed on Adult Male Albino Wistar Rats

ABSTRACT

Introduction: Substantially unproven scientifically are numerous assumed health benefits of nutmeg, moreover, a few of the scientifically substantiated research done were on the nutmeg oil and the mace. There is paucity of data based on scientific protocol on research carried out using numerable varying ranges of dosage of ground nutmeg seed.

Aims: The aim is to evaluate the effects of daily administration of ground nutmeg seed on some hematological parameters. Histological changes in the stomach were also investigated.

Study design: Adult male albino wistar rats totaling thirty were divided into groups: A,B,C,D,E and F; each group had five rats. Group A, the control was given water and 100 g of rat feed. Groups B, C, D, E and F, were given 1g, 5g, 10g, 15g and 20 g/kg body weight, of the ground nutmeg seed, respectively, for 26 days; each mixed with 100 g of feed. Measurement of weight was done once a week. With the aid of ethyl ether, blood samples were harvested via cardiac puncture The stomachs were dissected following sacrifice, fixed in 10 % buffered formalin and

stained with Haematoxylin and Eosin. Data were expressed in mean and standard error of mean. 95 % level of significance ($P = .05$) was used for statistical analysis.

Results and discussion: The control (group A) and the lowest dose-treated group (group B, given 1 g of ground nutmeg seed) had normal and intact histological structures of the stomach. The longitudinal gastric folds/ruggae, the surface mucous cells, mucous neck cells and parietal cells were not affected in Group C, given 5 g; but there was significant ($P = .05$) degeneration of simple columnar epithelium and longitudinal folds. The result also showed significantly ($P = .05$) degenerated simple columnar epithelium, longitudinal folds, coil pyloric glands and associated hypertrophy of the surface mucous cells and mucous neck cells in groups D, E and F, given, 10 g, 15 g, and 20 g, respectively; which were not dose dependent. There was significant ($P = .05$) decrease of white blood cells and dose-dependent increase in blood sugar levels in the treated groups.

Conclusion: Ground nutmeg seed when consumed in large doses alters the histology of the stomach, slightly lowers the body's defense against infections and predisposes to diabetes.

Key words: Blood, Nutmeg, Rat, stomach. Testes, Water.

INTRODUCTION

Nutmeg is a brownish colored hard seed from a tropical evergreen tree botanically known as *Myristica fragrans*. It has a warm, spicy sweet flavor, and was regarded as a valuable spices. It is in most cases ground into powder before use. Nutmeg is used in many dessert dishes, and also in savory recipes. Butter known as nutmeg butter is usually gotten from the nut by expression. By the process of steam distillation, essential oil obtained from ground nutmeg is useful in pharmaceutical industries. Nutmeg seed importance is innumerable; its uses cut across among others, oleochemical industries and in traditional medicine, were the oil is employed in managing disorders associated with the nervous and digestive systems. In fact, its various usefulness and improvement of health in general, dates back to the ancient times [1]. Nutmeg is reputed to play important roles in the removal of liver toxins, dissolving of kidney stones and relieve from infections. It also plays a crucial role by helping to increase blood circulation and appetite encouragement [2]. Its antioxidant properties is attributed to presence of phenolic compounds [3 - 7].

There is a demonstrated case of Immunoglobulin E reactivity, as well as reported case of allergy in nutmeg; limonene and eugenol, known chemical constituents of nutmeg are implicated as contact allergens [8 - 9]. Throughout their studies of this spice, researchers have found it to have a carminative effect. Carminatives are implicated in the relaxation of stomach muscles and are helpful in treating diarrhea, gas, nausea and indigestion [10].

Nutmeg is reputed to have a relatively narrow margin of safety; outcome of studies indicated that two to three teaspoons is the toxic dose of nutmeg when used as a spice [11]. Acute psychosis and anticholinergic – like episodes, with their related symptoms are known to be associated with nutmeg intake.[12 - 16].

STOMACH

The word Stomach has its origin from the ancient Greek word, *στόμαχος*, stomacho. It is part of the gastrointestinal tract. It is a hollow organ, muscular and dilated in structure. Its location is between the oesophagus and the small intestine, both in humans and many other animals. In humans, the stomach walls comprised of mucosa, submucosa, muscularis externa, and serosa [17 - 18].

The stomach in humans is divided into four regions, each with its cells and functions that vary [19]. The cardia is the inlet to the stomach. The fundus, a latin word, whose English equivalent is bottom, is located in the upper curved region. The body is the central area of the stomach. The pylorus, a Greek word, whose English equivalent is gatekeeper; is located at the lower region of the stomach, from which stomach contents empty into the duodenum [20 - 21].

Mucous secreting cells also referred to as goblet cells, line the luminal surface of the stomach. Gastric pits and gastric glands are implicated in secretion of mucus and bicarbonate. Mucous neck cells produce mucin. Parietal cells, also known as oxyntic cells, secrete gastric acid. Chief cells also called, peptic or zymogenic cells, secrete pepsinogen, that digests protein [22].




Figure 1. Gross anatomy of human stomach [23].

The rat stomach is located at the abdominal side of the diaphragm, and is connected to the esophagus at the gastroesophageal junction, and to the duodenum at the pyloric ring. The rat stomach is divided into the forestomach, also known as *pars proventricularis*, and glandular stomach, referred to as *corpus* or *pars glandularis*. The forestomach occupies about three fifths of the stomach area, while the glandular stomach is divided into the fundus and pylorus/antrum and communicates with the duodenum at the pyloric ring [24 - 25]. Heavily cornified squamous epithelium covers the forestomach. The mucosa of the glandular stomach is made up of rows of columnar cells that form numerous gastric pits which are based at the lamina muscularis mucosa and open into the lumen [26].

Substantially unproven scientifically are numerous assumed health benefits of nutmeg, just as seen or assumed in other herbs and spices. Moreover, a few of the scientifically substantiated research done were on the nutmeg oil and the mace. There is paucity of data based on scientific protocol on research carried out using numerable varying ranges of dosage of ground nutmeg seed; which is the major form in which nutmeg is consumed as a spice, hence, the need to carry out this research work titled *Haematological and Gastric Histological Changes Associated with Administration of Ground Nutmeg Seed on Adult Male Albino Wistar Rats*.

MATERIALS AND METHODS

Nutmeg seed was obtained from Itam market, Itu Local Government Area of Akwa Ibom State, Nigeria. It was identified by a plant taxonomist in the Department of Botany and Ecological Studies, Faculty of Science, University of Uyo, Uyo, Akwa Ibom State, Nigeria and a voucher specimen deposited at the herbarium of the same department. The Nutmeg seeds were ground into powder at the Department of Pharmacognosy and Natural Medicines, Faculty of Pharmacy, University of Uyo, Uyo, Akwa Ibom State, Nigeria. Out of the total gram of ground nutmeg seed obtained, a given gram was subjected to phytochemical analysis to determine its phytochemical properties. The method of Odebiyi and Sofowora [27] was employed in the analysis.



Figure 2. Nutmeg seed

Thirty adult male albino wister rats weighing between 148 g to 200 g were used for the study. The rats were purchased from the animal house of the Faculty of Basic Medical Sciences, University of Uyo, Nigeria, and were fed standard growers mash feed, produced by grand cereals limited, Nigeria. The rats were divided into six equal groups, namely A,B,C,D,E and F. Group A served as the control and were administered only distilled water and 100 g of rat feed; while groups B,C,D,E and F, the treatment groups, were administered varying doses (1 g, 5 g, 10 g, 15 g and 20 g/kg body weight, respectively) of the ground nutmeg seed mixed with 100 g of rat feed, respectively, for 26 days.

The animals were weighed on the 26th day of the experiment and anaesthetized using light ether. The stomach were harvested and fixed immediately in 10% buffered formalin for routine histological techniques, while the blood samples were collected by cardiac puncture in EDTA bottle and plain bottle for hematological studies. The gastric tissues were stained using Hematoxylin and Eosin; Drury and Wallington, method [28].

This research received a go-ahead-order to be carried out, from the appropriate Research and Ethics Committee. All procedures involving animals in this study conformed to the guiding principles for research involving animals as recommended by the Declaration of Helsinki and the Guiding Principles in the Care and Use of Animals.

STATISTICAL ANALYSIS

Data were analyzed using descriptive statistical tool, Primer, version 3.0, and were expressed as mean \pm standard error of the mean ($M \pm SEM$) and subjected to one way analysis of variance. Significant difference between means was assessed by Student - Newman-Keuls post hoc test. 95% level of significance ($P = .05$) was used for the statistical analysis..

RESULT

Following the conduction of phytochemical analysis on ground nutmeg seed, to determine the active ingredients present, the various observations and inferences were made: Alkaloids, Tannins and Saponnins were moderately present. Cardiac Glycosides noted via Lieberman, Salkowski and Keller Kiliani test, were slightly present, while Flavonoids are strongly present.

The weight of rats in the control group (group A) steadily increased gradually from the first week up to the fourth week. But in the experimental groups, the trend was not consistent, as shown in figure 2. In group B, there was observed decrease in weight on the fourth week, after steady and proportional increase from the first week to the third week; the same trend was observed in group E. In groups C and D, there were marked decrease in weight on the third week, followed by gradual rise in weight as seen in the fourth week. In group F, the decrease in weight started on the third week right into the fourth week; after gradual increase from the first to the second week.




Figure 3. Changes in body weight of rats in control and experimental groups.

In table 1, the outcome of full blood count is shown.

Table 1. Full Blood Count Resulting From the Administration of Ground Nutmeg (*Myristica fragrans*) Seed on Adult Male Albino Wistar Rats.

Group	Full Blood Count							Differentials			
	PCV (%)	Hb (g/dl)	WBC (l)	RBC (l)	MCV (fl)	MCH (pg)	MCHC (g/dl)	N (%)	L (%)	M (%)	E (%)
A	43	14	15.6	8.7	49	16	32	32	60	05	03

B	38	12	10.3	8.0	47	15	31	30	61	04	04
C	45	14.3	11.0	9.1	49	16	32	20	67	07	06
D	47	13.9	9.6	9.1	51	14	28	40	56	02	02
E	45	14.5	12.2	9.0	50	16	32	30	60	05	05
F	41	14.3	12.2	8.7	47	17	35	40	50	04	06

KEYS

- PCV - Packed Cell Volume
Hb - Haemoglobin
WBC - White Blood Cell
RBC - Red Blood Cell
MCV - Mean Corpus
MCH - Mean Corpuscular Haemoglobin
MCHC - Mean Corpuscular Haemoglobin Concentration
N - Neutrophils
L - Lymphocytes
M - Monocytes
E - Eosinophils

Gradual increase in blood sugar level was observed across the groups, which varied with the administered dose of nutmeg seed. This was shown in table 2.

Table 2. Blood Sugar Level Resulting From the Administration of Ground Nutmeg (*Myristica fragrans*) Seed on Adult Male Albino Wistar Rats.

Group	Blood sugar (mmol/l)
A	3.8±0.04
B	5.0±0.03
C	5.2±0.03
D	5.8±0.03
E	6.0±0.03
F	6.1±0.03

The result of our histopathological analysis suggests that nutmeg consumption above 10 g/kg bw per day is toxic to the testes. These are depicted succinctly in figures 4 to 15.




Figure 4. Contro group A. (X100).




Figure 5. Control group A. (X400).

Histologic section of the gastric mucosa (cardia, fundus and body regions) of rat administered distilled water. The longitudinal gastric folds/rugae, surface mucous cells SMC, mucous neck cells MNC and parietal cells PC were unaffected.




Figure.6. Experimental group B. (X100).




Figure 7. Experimental group B. (X400).

Histologic section of the gastric mucosa (cardia, fundus and body regions) of rat administered 1g of ground nutmeg seed mixed with 100g of rat feed. The longitudinal gastric folds/rugae, surface mucous cells SMC, mucous neck cells MNC and parietal cells PC were unaffected.




Figure 8. Experimental group C. (X100).




Figure 9. Experimental group C. (X400).

Histologic section of the gastric mucosa (Cardia, fundus and body regions) of rat given 5g of ground nutmeg seed mixed with 100g of feed. The tissue Section showed unaffected, longitudinal gastric folds/ruggae with intact surface mucous cells SMC, mucous neck cells MNC and parietal cells PC in some part and degenerating simple columnar epithelium DSCE and degenerating longitudinal folds DLGF in another area. Inference: 5g of ground nutmeg seed had moderate effect.




Figure 10. Experimental group D. (X100).




Figure 11. Experimental group D (X400).

Histologic section of the gastric mucosa (Pyloric region) of rat administered 10 g of ground nutmeg seed mixed with 100g of feed. Section revealed degenerating simple columnar epithelium DSCE, degenerating longitudinal folds DLGF associated with hypertrophy of the surface mucous cells HSMC and hypertrophy of mucous neck cells HMNC. There is also degeneration of the coil pyloric glands DCPG. Inference: severely affected by 10 g of nutmeg.




Figure 12. Experimental group E. (X100).




Figure 13. Experimental group E. (X400).

Histologic section of the gastric mucosa (fundus and body region) of rat given 15 g of ground nutmeg seed mixed with 100g of feed. Section revealed degenerating simple columnar epithelium DSCE, degenerating longitudinal gastric folds DLGF associated with hypertrophy of the surface mucous cells HSMC and hypertrophy of mucous neck cells HMNC. Inference: severely affected by 15 g of nutmeg.



Figure 14. Experimental group F. (X100).




Figure 15. Experimental group F. (X400).

Histologic section of the gastric mucosa (Pyloric region) of rat administered 20 g of ground nutmeg seed mixed with 100g of feed. Section revealed hypertrophy of simple columnar epithelium HSCE, degenerating longitudinal folds DLGF associated with hypertrophy of the surface mucous cells HSMC and hypertrophy of mucous neck cells HMNC. Inference: severely affected by 20 g of nutmeg.

Discussion

Understanding of pathologic processes associated with toxicity, drugs and chemical compounds safety profile are some of the insights normally made available by hematology. In addition,

information gotten from microscopic examinations, biochemical assay, urine analysis and clinical signs are complemented by hematology [29]. According to Johnson-Delaney [30], rat hematologic reference ranges are: RBC $6.76-9.75 \times 10^6/\text{mm}^3$, PCV 37.6-50.6%, WBC $6.6-12.6 \times 10^3/\text{mm}^3$, Hemoglobin 11.5-16.1 g/dL, Neutrophils $1.77-3.38 \times 10^3/\text{mm}^3$, Lymphocytes $4.78-9.12 \times 10^3/\text{mm}^3$, Eosinophils $0.03-0.08 \times 10^3/\text{mm}^3$, Monocytes $0.01-0.04 \times 10^3/\text{mm}^3$, Basophils $0.00-0.03 \times 10^3/\text{mm}^3$, Platelets $150-460 \times 10^3/\text{mL}$. Information emanating from rat's basic biological data, by [31], as reported in Pass and Freeth 1993, stated as follows: Red blood cell count $7-10 \times 10^6/\mu\text{l}$, Hemoglobin 11-19 gm/dl (dl = deciliter), Packed cell volume 40.5-54%, Leukocyte count Total $9(6-18) \times 10^3/\mu\text{l}$, Neutrophils 14-20%, Lymphocytes 69-86%, Monocytes 1-6%, Eosinophils 1-4%, Basophils Rare, Platelets $500-1,000 \times 10^3/\mu\text{l}$. A.D., Stammers [32], in his study of rats' blood count and temperature, conveyed the following information concerning RBC count: 7.4, 7.5, 8.0, 8.4, 10.0, all in 1 case; 8.6, and 9.0, in 2 cases; 8.8, 9.1, 9.6, 9.8, and 9.9, in 3 cases; 8.7, 9.3, and 9.5, in 4 cases; 9.2, and 9.4, in 6 cases; 8.9, in 7 cases. Subtle observation, perusal and comparison of the figures above with the result of our own hematology, as stated in table 1, indicates no provable contradictions and point to the fact that ours fall within the hematologic ranges of rat; thus, can serve as a premise to warrant the valid conclusion that ground nutmeg seed administered within the ranges 1 g to 20 g has no negative effect on full blood count.

As regards issues emanating from blood sugar level following consumption of ground nutmeg seed, our findings suggest dose dependent marked elevation in blood sugar level across the experimental groups; as was clearly shown in table 2. The study results did not corroborate the previous reports of activity of nutmeg against diabetes. In their experiments, Shyni *et al.*, [33] indicated that nutmeg can bring about an improvement in the body's response to insulin. There is also a reported case of pharmacists in Pune and Sagar, India, who studied nutmeg scientifically, and showed that extracts of the spice significantly decreased blood glucose levels; and improved the lipid profile in the blood [34].

The outcome of our histopathological analysis suggest that nutmeg consumption at higher doses above 10 g per day negatively affects the histology of the stomach. Our histological findings are consistent with that of Greger [35], which stated that Large dosage (about 30 grams a day) of nutmeg can be toxic.

There are no provable contradictions of our work with that of Adjene and Igbigbi [36], who concluded and implicated high dose of nutmeg in varying degrees of distortion in the lining and glandular epithelial cells of the stomach. Adjene and Igbigbi, were also of the view that "neoplastic changes, proliferation, hyperplasia and atrophic changes were observed in the stomach with a high dose of nutmeg. Therefore, it is likely that function of the stomach may be adversely affected by high dose of nutmeg". However, there are discrepancies in the method of preparation of the ground nutmeg seed, in the duration of administration and in the quantity of ground nutmeg seed administered in grams per kilogram body weight.

CONCLUSION

Our findings suggest that consumption of 20 g and below of ground nutmeg seed has no negative effect on full blood count and does not show any insulin – like activity. The outcome of our research also suggests that consumption of 10 g and above is toxic to the stomach.

CONSENT

Not applicable.

ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the department of Anatomy ethics committee".

REFERENCES

1. Nutmeg Benefits, What is Nutmeg Side Effects, Reviews and Facts. www.ProphecyFilm.com
2. Rayment W.J. Nutmeg Health Effects; Indepth Information on Nutmeg. 2007 – 2018
3. Dorman HJ, Figueiredo AC, Barroso JG, Deans SG. In vitro evaluation of antioxidant activity of essential oils and their components. *Flavour Fragr J* . 2000;15:12-16.
4. Valderrama JC. Distribution of flavonoids in the Myristicaceae. *Phytochemistry* . 2000; 55(6):505-511.
5. Shan B, Cai YZ, Sun M, Corke H. Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *J Agric Food Chem* . 2005; 53(20):7749-7759.
6. Su L, Yin JJ, Charles D, Zhou K, Moore J, Yu L. Total phenolic contents, chelating capacities, and radical-scavenging properties of black peppercorn, nutmeg, rosehip, cinnamon and oregano leaf. *Food Chem* . 2005.
7. Duan L, Tao HW, Hao XJ, Gu QQ, Zhu WM. Cytotoxic and antioxidative phenolic compounds from the traditional Chinese medicinal plant, *Myristica fragrans* . *Planta Med* . 2009;75(11):1241-1245.
8. Brancaccio RR, Alvarez MS. Contact allergy to food. *Dermatol Ther* 2004;17(4):302-313.
9. Schöll I, Jensen-Jarolim E. Allergenic potency of spices: hot, medium hot, or very hot. *Int Arch Allergy Immunol* . 2004;135(3):247-261.
10. Laura, M. Nutmeg benefits and side effects. *BodyNutrition*. 2018
11. Greger, M., Nutmeg Toxicity. Nutmeg Benefits, What is Nutmeg Side Effects, Reviews and Facts. 2013; Krankeboda Norregarden 1:31; 341 73 Lidhult, SWEDEN
12. Brenner, N., Frank, O.S. and Knight, E. Chronic nutmeg psychosis. *J R Soc Med*.. 1993;86(3):179-180.
13. Demetriades, A.K., Wallman, P.D., McGuinness, A. and Gavalas, M.C. Low cost, high risk: accidental nutmeg intoxication. *Emerg Med J*. 2005;22(3):223-225.
14. Forrester, M.B. Nutmeg intoxication in Texas, *Hum Exp Toxicol* . 2005;24(11):563-566.

15. Kelly, B.D., Gavin, B.E., Clarke, M., Lane, A. and Larkin, C. Nutmeg and psychosis. *Schizophr Res.* 2003;60(1):95-96.
16. Sangalli, B.C. and Chiang, W. Toxicology of nutmeg abuse. *J Toxicol Clin Toxicol* 2000;38(6):671-678.
17. The Stomach at The Anatomy Lesson by Wesley Norman (Georgetown University)
18. Sherwood, Lauralee. *Human physiology: from cells to systems*. Belmont, CA: Wadsworth Pub. Co. 1997; ISBN 978-0-314-09245-8. OCLC 35270048
19. Anatomy photo:37:06-0103 at the SUNY Downstate Medical Center – "Abdominal Cavity: The Stomach"
20. Brunicaudi, F. Charles; Andersen, Dana K.; et al., eds. (2010). *Schwartz's principles of surgery* (9th ed.). New York: McGraw-Hill, Medical Pub. Division.2010; ISBN 978-0071547703.
21. Lenglinger, J.; et al. "The cardia: esophageal or gastric? Critical reviewing the anatomy and histopathology of the esophagogastric junction". *Acta Chir Iugosl.* 2012;59 (3): 15–26. PMID 23654002.
22. Sampurna, R. *Gastrointestinal Pathology. An outline of the Anatomy and Normal Histology of the Stomach.* Kolkata – India. 2017; www.histopathology-india.net
23. <http://en.wikipedia.org....stomach>
24. Greene EC.(1968) *Anatomy of the rat.* Hafner, New York 1968; Harding RK, Morris GP Cell loss from normal and stressed gastric mucosae of the rat: an ultrastructural analysis. *Gastroenterology*, 1977;1968;72:857–863,Google Scholar
25. Robert A. Proposed terminology for the anatomy of the rat stomach. *Gastroenterology*, 1971; 60:344–345
26. Katsuyama T, Spicer SS. Histochemical differentiation of complex carbohydrates with variants of the concanavalin A-horseradish peroxidase method. *J Histochem Cytochem.* 1978; 26:233–250
27. Odebibi, O.O. and Sofowora, E.A. Phytochemical Screening of Nigerian Medicinal Plants II. *Lloydia*,1978; 41, 234 - 236.
28. Drury, R.A. and Wallington, E.A. *Carltons Histological Techniques*, 5th edn. New York, Oxford University Press,1980; p. 206.
29. Bailly Y., Duprat P. Normal Blood Cell Values, Rat. In: Jones T.C., Ward J.M., Mohr U., Hunt R.D. (eds) *Hemopoietic System. Monographs on Pathology of Laboratory Animals.* Springer, Berlin, Heidelberg. 1990; pp 27-38

30. Baker *et al.* 1979 and Weihle 1987. Rat's basic biological data, as reported in Pass and Freeth 1993; <http://www.ratbehavior.org>
31. Stammers AD, The blood count and body temperature in normal rats. *J Physiol.* 1926; 61(3):329–336. PMCID: PMC1514839 PMID: 16993796
32. Shyni, G.L., Kavitha, S., Sajin, K.F., Arya, A.D., Mangalam, S.N., and Raghu, K.G.. Licarin B from *Myristica fragrans* improves insulin sensitivity via PPAR γ and activation of GLUT4 in the IRS-1/PI3K/AKT pathway in 3T3-L1 adipocytes. *RSC Advances.* 2016; 6, no. 83: 79859-79870
33. <http://www.rpsgb.org.uk/>
34. Adjene, J.O. and Igbigbi, P.S. Effect of chronic consumption of nutmeg on the stomach of Adult Wistar Rats. *Foyin Journal of Health Sciences.* 2010; Vol. 2 Issue 2 p 62 - 65

UNDER PEER REVIEW