

# Estimation of Variability, Heritability and Genetic Advance of Exotic Rice (*Oryza sativa* L.) Breeding Lines for Short Growth Development

Md. Manik Sarker<sup>1\*</sup>, Lutful Hassan<sup>2</sup>, Md. Mamunur Rashid<sup>3</sup>, Syed Nazrul Islam<sup>4</sup>, Tilak Kumar Ghosh<sup>1</sup> and Sharmin Sultana<sup>1</sup>

<sup>1</sup>Department of Agricultural Extension, Ministry of Agriculture, Government of the People's Republic of Bangladesh

<sup>2</sup>Department of Genetics and Plant Breeding, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

<sup>3</sup>Farm Management Division, Bangladesh Rice Research Institute, Gazipur, Bangladesh

<sup>4</sup>Bangladesh Jute Research Institute, Manik Mia Avenue, Dhaka -1207, Bangladesh

## ABSTRACT

This work aims to study the genetic parameters for yield and yield contributing characters in thirty-two early maturing rice (*Oryza sativa* L.) lines. The morphological data were analyzed by different statistical plant breeding software's that were reliable and faster in analysis. The line CT 18173-1-9-1-3-6-M had the tallest; IR 82489-594-3-2-2 had the earlier flowering line but BP 10620F-BB4-13-BB8 the early maturing one. The highest effective tillers hill<sup>-1</sup> and yield plant<sup>-1</sup> was found in BP 10620F-BB4-13-BB8. Number of filled grains panicle<sup>-1</sup> and number of unfilled grains panicle<sup>-1</sup> exhibited high Genotypic Coefficient of Variation (GCV) and Phenotypic Coefficient of Variation (PCV). Small differences between GCV and PCV were recorded for all the characters studied which indicated less environmental influence. The characters *viz.*, 1000 seed weight and days to maturity exhibited high heritability and number of filled grains panicle<sup>-1</sup> measured high genetic advance. The thirty-two lines were assigned into three clusters, which contained a different number of lines. Among the lines, BP 10620F-BB4-12-BB8 possessed the highest selection score index contrary to IR 79201-49-1-1-1 which had the lowest selection index score. The overall evaluation for the lines revealed that the best line was BP 10620F-BB4-12-BB8 and it can be used for future breeding program or farmer's field cultivation.

**Keywords:** Genetic advance, Heritability, Clustering, Selection Index and Rice

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is the staple food for about 2.5 billion world's population which may escalate to 4.6 billion by the year 2050 (Macleay, 2002). Rice production have significantly increased by high yielding varieties (HYV), modern varieties and improved agricultural practices during the past century. More than 2,000 modern varieties have been commercially released in twelve countries of South and Southeast Asia over the past 40 years (Cantrell and Hettel, 2004). In Bangladesh at present cropping intensity is 183 % (BBS, 2012). It can be increased by further short duration rice nearly 300 %. A critical analysis of genetic variability is a prerequisite for initiating any crop improvement program and also for adopting appropriate selection techniques. The selection of lines plays a vital role in developing ideal combinations for breeding. The prediction of genetic advance is a precondition for crop improvement breeding programs especially when large populations are subjected to selection. The progress in breeding for yield and its contributing characters of any crop is polygenically controlled, environmentally influenced and determined by the magnitude and nature of their genetic variability (Wright, 1935; Fisher, 1981). Hence there is a need to study variability. Variability for different traits in the source population is a prerequisite for crop improvement since all attempts of breeding and selection would be futile unless major portion of variability is heritable. It is difficult to judge whether observed variability is highly heritable or not. Moreover, knowledge of heritability is essential for selection based improvement, as it indicates the extent of transmissibility of a character into future generations (Sabesan *et al.* 2009). Mruthunjaya and Mahadevappa (1995) reported that the success of a crop improvement program depends on the definition and assembly of the required genetic variations and selection for yield through high heritable traits. Making of selection indices and their analysis would give the most appropriate selection score of each of two or more characters to be used simultaneously for selection. This study aims to assess the genetic parameters for yield and yield contributing characters of early maturing rice (*Oryza sativa* L.) lines collected from International Rice research Institute (IRRI), Philippines.

54 **2. MATERIALS AND METHODS**

55 Thirty-one early maturing rice lines along with check variety Binadhan-7 were used for the experiments. The  
 56 seeds of the rice lines were collected from the International Network for Genetic Evaluation of Rice (*INGER*)  
 57 and International Rice Research Institute (table 1).

58 **Table1. Name and origin of studied rice lines**(Sarker et. al.,2014)

Sl. No	Lines Name	Origin	Sl. No	Lines Name	Origin
1	IR 79246-105-2-2-4	IRRI	17	BP 1018F-BB8-13-BB4	INDONESIA
2	IR 73718-26-1-2-5	IRRI	18	IR 79525-20-2-2-2	IRRI
3	BP 10620F-BB4-13-BB8	INDONESIA	19	IR 80285-34-3-3-2	IRRI
4	IR 79538-1-1-1-1	IRRI	20	CT 18173-1-9-1-3-6-M	CIAT
5	IR 76494-28-1-2-2	IRRI	21	BP 10620F-BB4-2-BB4	INDONESIA
6	YN 2883-12-2-1	MYANMAR	22	PSB RC 64	INDIA
7	AD 02207	INDIA	23	IR 08N261	IRRI
8	BP 10620F-BB4-8-BB8	INDONESIA	24	RATNAGIRI 2	INDIA
9	C1-4-11-7P-2P-1P	CIAT	25	MTU-1113	INDIA
10	IR 79201-49-1-1-1	IRRI	26	KARJAT 5	INDIA
11	BP 10620F-BB4-12-BB8	INDONESIA	27	KHAZAR	IRAN
12	IR 82489-594-3-2-2	IRRI	28	IR 59552-21-3-2-2	IRRI
13	CT 18509-10-6-1VI-2	CIAT	29	C 2-9-9-2P-1P-3P	CIAT
14	IR 74052-153-5-3-1-3	IRRI	30	IR 39809-26-3-3	IRRI
15	PSD RC 2	IRRI	31	CT 18148-11-1-1-1-1-M	CIAT
16	IR 08N293	IRRI	32	Binadhan-7	BINA

59 The experiment was laid out in a RCBD with three replications. The row to row and plant to plant distances  
 60 were 20 cm and 15 cm respectively. The following data plant height (cm), days to flowering, days to maturity,  
 61 total tillers and effective tiller/hill, filled and unfilled grain/panicle, 1000 seed weight (g), yield plant<sup>-1</sup> (g) were  
 62 collected from randomly selected 5 plants of each unit plot. The data was analyzed by using the plant breeding  
 63 statistical program (PLBSTAT, Version 2N, Utz 2007) for mean performance analysis and least significant  
 64 difference (LSD). The genotypic and phenotypic variances were calculated as per the formulae proposed by  
 65 Burton (1952). The genotypic (GCV) and phenotypic (PCV) coefficient of variation was calculated by the  
 66 formulae given by Burton (1952). Heritability was calculated by the formula given by Hansen *et al.* (1956). For  
 67 the heritability estimates, the genetic advance (GA) was estimated by the following formula given by Johnson *et*  
 68 *al.* (1955). Stat Graphics plus for Windows 3.0 (Statistical Graphics Crop. Rockville, USA) software was used  
 69 for clustering and ward's dendrogram construction. Selection indices were constructed using the method  
 70 developed by Smith (1936) based on the discriminate function of Fisher (1936).  
 71  
 72

73 **3. RESULTS AND DISCUSSION**

74 **3.1 Variations and performance of the lines**

75 Analysis of variance resulted in significant variations among the lines for all studied traits (table 2). These  
 76 results indicated that there are sufficient scope for further improvement of existing lines through breeding  
 77 program.

78 Minimum days to 50 % flowering were observed by the lines IR 82489-594-3-2-2 (64 days). The remaining lines  
 79 had between 65 to 86 days. Similar results have been reported by Jamal *et al.* (2009) in rice.

80 The line KARJAT 5 took the highest days to maturity (130 days). In contrast, the line BP 10620F-BB4-13-BB8  
 81 took the lowest days to maturity (104 days). This result was in full agreement with Dutta (2001) who observed  
 82 days to maturity varied between 105 to 130 days in rice lines.

83 Among the 32 lines, CT 18173-1-9-1-3-6-M had the highest plant height (80 cm) and the line MTU-1113 had the  
 84 shortest plant height (53 cm). The results of the present study is supported by the result of different researchers  
 85 e.g. (Sharma and Sharma, 1983; Talukder, 2004).

86 The minimum panicle length (18.3 cm) was recorded in BP 10620F-BB4-13-BB8 while maximum panicle length  
 87 (24 cm) was recorded in IR 08N293. Such variation also reported by Tahir *et al.* (2002) in rice. This type of  
 88 variation might be due to the genetic makeup of the exotic lines and genotypic environmental interaction.

89 A total number of tillers hill<sup>-1</sup> of 4 lines were higher in number compared to remaining 28 lines and the highest  
 90 effective tillers hill<sup>-1</sup> was found in BP 10620F-BB4-12-BB8 which contains 17.00 tillers hill<sup>-1</sup> and 13.42 effective  
 91 tillers hill<sup>-1</sup>. The lowest total number of effective tillers hill<sup>-1</sup> recorded YN 2883-12-2-1(5.40). This observation is  
 92 supported by Zahid *et al.* (2005).

93 The highest number of grains panicle<sup>-1</sup> was found in RATNAGIRI 2 (185). Whereas, the lowest grains panicle<sup>-1</sup>  
 94 was found in BP 10620F-BB4-13-BB8 (93). Prasad *et al.* (2001) and Tahir *et al.* (2002) reported highly  
 95 significant variation for the grains panicle<sup>-1</sup> for different genotypes.

96 The highest 1000 seed weight of 23.3 g was recorded for BP 10620F-BB4-13-BB8. In contrast, the minimum  
 97 1000 seed weight was found in the line KHAZAR (12.7 g). Tahir *et al.* (2002), and Bharali *et al.* (1994) reported  
 98 the variation of 1000 seed weight.

99 The highest yield plant<sup>-1</sup> (29.5 kg) was recorded for BP 10620F-BB4-12-BB8. In contrast, the minimum yield  
 100 plant<sup>-1</sup> was found in the line CT 18173-1-9-1-3-6-M (16.6 g). The same variability were reported by Zahid *et al.*  
 101 (2005) who studied twelve genotypes of coarse rice to check their yield performance in Kallar tract and reported  
 102 highly significant variation for different traits. Most of the cases results are very much similar to others due to  
 103 the genotypic novelty and adaptability to the present environment or other factors i.e. soil fertility, plant nutrients  
 104 translocation, transplantation season, time and weather condition also responsible.

105 **Table 2. Mean performance of 32 lines based on different morphological characters**

Lines name	DF	DM	PH	TT	ET	PL	FG	UG	SWT	Y/P
IR 79246-105-2-2-4	79.3	109.3	61.6	12.3	10.6	19.6	67.6	77.5	18.1	23.5
IR 73718-26-1-2-5	79.6	108.7	64.6	11.6	10.6	22.6	95.1	55.5	19.5	24.1
BP 10620F-BB4-13-BB8	65.3	104.0	55.0	9.6	9.0	18.3	93.8	24.8	23.3	21.1
IR 79538-1-1-1-1	74.0	106.0	67.3	10.0	8.6	20.3	88.6	61.3	20.8	20.9
IR 76494-28-1-2-2	70.3	105.7	74.6	10.3	9.6	22.3	138	56.0	17.8	23.1
YN 2883-12-2-1	74.6	104.7	72.6	5.3	4.3	22.3	115.5	36.0	17.8	18.3
AD 02207	75.0	114.0	64.3	13.6	10.3	20.3	108.7	46.6	19.8	21.5
BP 10620F-BB4-8-BB8	65.0	105.0	59.6	9.6	7.6	19.3	108.2	25.8	21.7	19.2
C 1-4- 11-7P-2P-1P	75.6	111.7	61.0	9.3	7.6	22.0	130.2	47.3	16.2	20.3
IR 79201-49-1-1-1	66.3	110.7	60.0	16.3	13.0	22.0	165.3	26.0	16.2	23.1
BP 10620F-BB4-12-BB8	70.6	115.0	60.3	17.0	13.4	21.3	109.3	22.6	22.1	25.6
IR 82489-594-3-2-2	64.0	105.7	70.6	11.7	10.0	22.0	169.2	66.1	17.3	24.0
CT 18509-10-6-1VI-2	78.3	129.0	70.6	10.0	9.00	22.0	140.3	68.0	16.4	21.2
IR 74052-153-5-3-1-3	74.0	111.3	71.3	10.6	7.00	22.6	77.3	42.5	18.2	20.0
PSD RC 2	75.6	112.7	61.3	10.7	8.60	21.0	99.3	24.8	16.6	20.3
IR 08N293	74.6	114.7	74.0	12.3	10.6	24.0	136	42.8	16.9	22.5
BP 1018F-BB8-13-BB4	74.3	115.3	68.0	13.6	12.6	21.6	113.5	73.1	17.6	24.2
IR 79525-20-2-2-2	74.0	113.7	72.6	14.0	11.6	21.6	165.5	50.3	21.6	20.7
IR 80285-34-3-3-2	74.3	114.3	64.3	14.0	11.0	21.0	119.5	36.5	17.2	21.7
CT 18173-1-9-1-3-6-M	64.3	113.7	80.0	8.30	7.0	22.0	118.7	43.1	15.4	16.6
BP 10620F-BB4-2-BB4	71.3	112.0	59.3	11.0	8.30	19.3	140.7	32.0	19.6	19.7
PSB RC 64	77.3	115.0	67.0	12.0	11.0	21.0	99.3	35.3	22.0	23.2
IR 08N261	75.6	123.3	70.6	14.3	12.0	20.3	181.3	25.8	17.8	24.6
RATNAGIRI 2	75.6	123.3	70.6	14.3	12.0	20.3	185.0	30.0	17.8	23.6
MTU-1113	73.6	129.7	53.0	11.6	10.0	20.3	112.7	68.3	19.3	23.7
KARJAT 5	80.0	130.3	59.0	12.3	11.3	21.0	107.8	49.5	14.3	24.1
KHAZAR	86.0	129.0	69.6	13.3	11.3	19.6	80.1	67.5	12.7	24.2
IR 59552-21-3-2-2	73.6	112.7	69.0	10.3	9.00	21.3	92.5	54.8	21.1	20.1
C2-9-9-2P-1P-3P	73.6	114.3	78.6	14.6	13.0	23.6	175.3	54.1	18.5	23.7
IR 39809-26-3-3	79.3	129.0	59.6	12.0	9.60	21.0	89.3	40.0	16.2	22.9
CT 18148-11-1-1-1-1-M	75.0	114.0	67.6	10.6	9.60	22.0	106.0	59.3	17.2	23.9
Binadhan-7	65.3	114.0	65.0	8.30	6.60	21.0	115.3	53.0	19.6	17.0
Mean	73.7	114.7	66.3	11.7	9.80	21.2	120.1	46.7	18.3	21.9
CV %	5.92	1.67	4.90	21.9	20.2	5.99	17.7	24.6	0.61	6.20

LSD 0.05	4.54	3.48	5.33	4.22	3.25	2.07	34.5	19.5	0.18	2.22
----------	------	------	------	------	------	------	------	------	------	------

DM=Days to 50 % flowering, DM=Days to maturity, PH=Plant height (cm), TT=Total number of tillers hill<sup>-1</sup>, ET=Effective number of tillers hill<sup>-1</sup>, PL=Panicle length (cm), FG=Filled grains panicle<sup>-1</sup>, UG=Unfilled grains panicle<sup>-1</sup>, SWT=1000 seed weight (g), Y/P=Yield plant<sup>-1</sup> (g)

### 3.2 Estimation of genetic parameters

#### 3.2.1 Variability parameters

Greater variability in the initial breeding material ensures better chances of producing desired forms of a crop plant. The phenotypic coefficient of variation (PCV) were higher than genotypic coefficient of variation (GCV) for all the characters studied indicating that they all interacted that less influenced by environment (table 3). The characters studied in the present investigation exhibited low, moderate and high PCV and GCV values. Among the characters, highest PCV and GCV values were recorded for a number of unfilled grains panicle<sup>-1</sup>, followed by grain yield plant<sup>-1</sup> (g) indicating that these characters could be used as selection for crop improvement and the lowest PCV and GCV values were recorded for days to maturity. Similar results were obtained by Kuldeep et al. (2004), Sabesan et al. (2009) and Jayasudha and Sharma (2010).

#### 3.2.2 Heritability

Heritability is classified as low (below 30 %), medium (30-60 %) and high (above 60 %). Heritability in broad sense indicates days of 50 % flowering, days to maturity, plant height, 1000 seed weight (g) and yield plant<sup>-1</sup> (g) were high heritable and that measured 75.6, 92.9, 78.7, 99.7 and 70.8 % respectively (table 3). High heritability values indicate that the less influenced by the environment in their expression. It indicates the scope of genetic improvement of these characters through selection. Similar results had been reported by idem. On the other hand, the total number of tillers hill<sup>-1</sup>, number of effective tillers hill<sup>-1</sup>, panicle length (cm), number of filled grains panicle<sup>-1</sup>, number of unfilled grains panicle<sup>-1</sup> were medium heritable (table 3).

#### 3.2.3 Genetic advance

Genetic advance is a useful indicator that can be expected as result of exercising selection of the population. Heritability in conjunction with genetic advance would give a more reliable index of selection value (Johnson et al. 1955). The genetic advance was highest for number of filled grains panicle<sup>-1</sup> and lowest for panicle length (cm). The genetic advance as percent of mean was highest in case of number of unfilled grains panicle<sup>-1</sup>, while lowest recorded by days to maturity (table 3). In this study high heritability along with high genetic advance was noticed for the traits, number of filled grains panicle<sup>-1</sup> and number of unfilled grains panicle<sup>-1</sup>. Similar results were also reported idem. Other characters (viz: days to 50 % flowering, plant height, number of tillers hill<sup>-1</sup>, effective tillers hill<sup>-1</sup>, panicle length, 1000 seed weight, yield plant<sup>-1</sup> showed high heritability along with moderate or low genetic advance which can be improved by intermitting superior lines of segregating population developed from combination breeding.

**Table 3. Estimation of genetic parameters of 32 rice lines based on different morphological traits related to yield**

Characters	Phenotypic variance	Genotypic variance	PCV (%)	GCV (%)	Heritability (%)	GA	GA (%)
DF	84.8	82.4	10.9	10.2	87.0	27.6	19.6
DM	94.3	92.6	8.76	7.7	78.3	3.60	14.1
PL	96.7	95.8	12.1	11.6	91.1	26.1	22.8
TT	97.8	97.1	9.00	8.20	82.5	23.7	15.4
ET	96.9	95.9	21.0	20.2	92.2	5.5	40.0
PL	98.0	96.4	21.9	20.9	91.5	5.4	41.3
FG	201.1	177	31.2	31.0	98.7	78.6	63.4
UG	112.8	109.5	51.6	51.4	99.1	28.0	105.5
SWT	145.3	131.3	19.6	19.2	95.7	9.40	38.7
Y/P	3.4	1.10	37.8	37.5	98.7	24.4	76.9

DM=Days to 50 % flowering, DM=Days to maturity, PH=Plant height (cm), TT=Total number of tillers hill<sup>-1</sup>, ET=Effective number of tillers hill<sup>-1</sup>, PL=Panicle length (cm), FG=Filled grains panicle<sup>-1</sup>, UG=Unfilled grains panicle<sup>-1</sup>, SWT=1000 seed weight (g), Y/P=Yield plant<sup>-1</sup> (g)

### 3.3 Nature and magnitude of genetic diversity


149 The knowledge on the nature and magnitude of genetic diversity governing the inheritance of quantitative  
 150 characters like yield and its components is essential for effecting genetic improvement. Using Euclidean  
 151 distance following Ward's method, the studied accessions were grouped into distinct clusters. It's were grouped  
 152 into three clusters (table 4). Cluster I and II had maximum number of lines and the cluster III contained only nine  
 153 lines.  
 154

155 **Table 4. Clustering pattern of 32 rice lines and the member present in each cluster**

Cluster number	Number of lines	Percent	Sl. no of lines
I	12	37.5	1, 2, 31, 17, 7, 19, 22, 13, 25, 26, 30, 27
II	11	34.3	3, 8, 21, 4, 28, 14, 9, 15, 6, 20, 32
III	9	28.1	5, 12, 16, 29, 18, 10, 11, 23, 24

156  
 157 **3.3.1 Ward's dendrogram**

158 Dendrogram grouped of the 32 lines of rice into three clusters (fig 1). Binadhan-7 as control line is grouped in  
 159 cluster II. with lowest (0.00) genetic distance, IR 79246-105-2-2-4, IR 73718-26-1-2-5, CT 18148-11-1-1-1-M,  
 160 BP 1018F-BB8-13-BB4, AD 02207, BP 1018F-BB8-13-BB4, IR 80285-34-3-3-2, PSB RC 64, CT 18509-10-6-  
 161 1VI-2, MTU-1113, KARJAT 5, KHAZAR, IR 39809-26-3-3 were grouped in cluster I. IR 76494-28-1-2-2, IR  
 162 82489-594-3-2-2, IR 08N293, C 2-9-9-2P-1P-3P, IR 79525-20-2-2-2, IR 79201-49-1-1-1, BP 10620F-BB4-12-  
 163 BB8, IR 08N261, RATNAGIRI 2 were grouped on cluster III.  
 164



165  
 166  
 167  
 168  
 169  
 170  
 171  
 172  
 173 Fig.1. Dendrogram based on summarized data on differentiation among 32 lines of rice according to Ward's  
 174 method

175  
 176 **3.3.2 Characterization of individual clusters**

177 The mean values of each cluster for eleven characters are presented in table 5. There was a wide range of  
 178 variation in the cluster mean values for all the characters. The mean values of all characters were categorized  
 179 into low (L), intermediate (I) and high (H) classes. Cluster II required minimum 70.8 days for 50 % flowering and  
 180 109.8 days for maturity. Cluster I required maximum 77.6 days for 50 % flowering, 119.8 days for maturity and  
 181 cluster III showed intermediate values (71.6-114). In case of plant height, cluster I showed low value (64.1) and  
 182 cluster III showed high value (70.2). Cluster II, showed intermediate values (65.6). With regard of total number  
 183 of tillers hill<sup>-1</sup>, cluster II showed low value (9.39) and cluster III showed high value (13.8). With regard of  
 184 effective number of tillers hill<sup>-1</sup>, cluster II exhibited low value (7.64) and cluster III showed high value (11.7). For  
 185 panicle length (cm) cluster III showed low value (21.9) and cluster II showed high value (20.8). Number of filled  
 186 grains panicle<sup>-1</sup>, cluster III showed high value (158.3) and cluster I showed low value (103.3). Number of unfilled  
 187 grains panicle<sup>-1</sup>, cluster I showed high value (56.4) and cluster II showed low value (40.5). With regards to 1000  
 188 seed weight, cluster II demonstrated high value (19.1) and cluster I showed low value (17.5). With regards to  
 189 yield plant<sup>-1</sup>, cluster III showed high value (23.4) and cluster II showed low value (19.4). The cluster III showed  
 190 best performance, genotypes of that cluster can be proposed for breeding program.  
 191

192 **Table 5. Cluster mean for 10 yield and yield related characters of 32 rice lines**

Characters	I	II	III
Days to 50%flowering	77.6 (H)	70.8 (L)	71.6 (I)
Days to maturity	119.8 (H)	109.8 (L)	114 (I)
Plant height (cm)	64.1 (L)	65.6 (I)	70.2 (H)
Total number of tillers hill <sup>-1</sup>	12.2 (I)	9.30 (L)	13.8 (H)
Effective number of tillers hill <sup>-1</sup>	10.6 (I)	7.60 (L)	11.7 (H)
Panicle length (cm)	21.0 (I)	20.8 (L)	21.9 (H)
Number of filled grains	103.3 (L)	107.3 (I)	158.3 (H)
Number of unfilled grains	56.4 (H)	40.5 (L)	41.5 (I)
1000 seed weight (g)	17.5 (I)	19.1 (H)	18.4 (L)
Yield plant <sup>-1</sup> (g)	23.2 (I)	19.4 (L)	23.4 (H)

H= High value, I= Intermediate value, L= Low value

### 3.4 Selection index

Selection index was constructed to identify suitable lines among 32 lines of rice in order to recommend for cultivation. Following simultaneous selection model was found by considering 10 characters. Here, among the 32 lines BP 10620F-BB4-12-BB8 possessed the highest selection score (7636) and rank as the best followed by IR 79246-105-2-2-4, IR 73718-26-1-2-5, BP-10620F-BB4-13-BB8, IR 79538-1-1-1-1, IR 79538-1-1-1-1, IR 76494-28-1-2-2, YN 2883-12-2-1, AD 02207, BP 10620F-BB4-8-BB8 and C1 -4- 11-7P-2P-1P with 6813, 5952, 5869 5828 5952, 5869, 4759, 4228, -167 and -231 respectively (table 6).

The expected genetic gain ( $\Delta G$ ) was 143 at 5 % selection intensity i.e. 4-5 highest scoring lines from these 32 rice lines might be recommended for cultivation for better yield.

**Table 6. Selection best rice 10 lines considering ten characters**

Sl. No	Lines Name	Selection Score	Rank
1	BP 10620F-BB4-12-BB8	7636	1
2	IR 79246-105-2-2-4	6813	2
3	IR 73718-26-1-2-5	4716	3
4	BP 10620F-BB4-13-BB8	5828	4
5	IR 79538-1-1-1-1	5952	5
6	IR 76494-28-1-2-2	5869	6
7	YN 2883-12-2-1	4759	7
8	AD 02207	4228	8
9	BP 10620F-BB4-8-BB8	-167	9
10	C1 -4- 11-7P-2P-1P	-231	10

## 4. CONCLUSION

The overall results showed that BP 10620F-BB4-13-BB8 was early maturing among the studied rice lines. BP 10620F-BB4-12-BB8 yielded a maximum total number of tillers hill<sup>-1</sup>, effective number of tillers hill<sup>-1</sup>, 1000 seed weight (g) and yield plant<sup>-1</sup>. However, among the tested lines, IR 79201-49-1-1-1 resulted in 206 higher panicle length on the basis of overall results BP 10620F-BB4-12-BB8 was exhibited better performances than that of other exotic lines, this line also possessed the highest selection score index and it was the leader of cluster III which is characterize by accessions with higher panicle length. The origin of this line was Indonesia by considering overall performance it can be recommended for farmer cultivation after appropriate variety release proceeding.

## ACKNOWLEDGEMENTS

We acknowledge the International Network for Genetic Evaluation of Rice (INGER), International Rice Research Institute (IRRI) for providing *Oryza sativa* L. seeds. The authors are thankful to Dr. Mirza Mofazzal Islam, Chief Scientific Officer and Head, Plant Breeding Division, Bangladesh Institute of Nuclear Agriculture, Mymensingh for providing all the assets required in the experiment. We are also grateful for the critical comments and suggestions of Professor Dr. Arif Hasan Khan Robin, Department of Genetics and Plant Breeding, Bangladesh Agricultural University and Professor Dr. Mirza Hasanuzzaman, Department of Agronomy, Sher-e-Bangla Agricultural University.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

225

## 226 **AUTHORS' CONTRIBUTIONS**

227 Author Md. Manik Sarker designed the study, performed the statistical analysis, wrote the protocol, and wrote  
228 the first draft of the manuscript. Author Lutful Hassan supervised the study and managed the literature  
229 searches. Author Md. Mamunur Rashid and Author Syed Nazrul Islam managed the experimental process and  
230 performed data collection. Author Tilak Kumar Ghosh Author Sharmin Sultana contributed in data analysis and  
231 participated in drawing inference of findings. All authors read and approved the final manuscript.  
232

## 233 **REFERENCES**

- 234 BBS (Bangladesh Bureau of Statistics), 2012. Monthly Statistical Bulletin of Bangladesh. Stat. Div., Minst.  
235 Planning, Bangladesh Bur. Stat, Govt. People's Repub. Bangladesh, Dhaka. p. 182.
- 236 Bharali B. and Chandra K., 1994. Effect of low light on dry matter production, harvest index and grain yield of  
237 rice (*Oryza sativa* L.) in wet season. *Neo-Botanica*. 2(1): 11-14.
- 238 Burton G.W. and Devane E.H., 1952. Estimating heritability in tall fescue (*Festuca arundinaceae*) from  
239 replicated clonal material. *Agronomy J.* 45: 478-481.
- 240 Cantrell R.P and Hettel G.P., 2004. New Challenges and Technological Opportunities for Rice-Based  
241 Production Systems for Food Security and Poverty Alleviation in Asia and the Pacific. Presented at the  
242 FAO Rice Conference, FAO, Rome, Italy, February 12-13.
- 243 Dutta R.K., 2001. Assessment of the advanced mutant lines/varieties of BINA with emphasis on  
244 physiological criteria. Report on ARMP Project No. 512, BARC, Farmgate, Dhaka-1215.
- 245 Fisher R.A., 1981. The correlation among relative on the supposition of Mendelian Inheritance. *Trans. Royal*  
246 *Soc. Edinberg*.
- 247 Fisher R.A., 1936. The use of multiple measurements in taxonomic problems, *Ann, Eugen*.7: 179-188.
- 248 Hansen G.H. Robinson H.F. and Comstock R.E., 1956. Bio-metrical studies of yield in segregating population of  
249 Korean bespedese. *Agronomy J.* 48: 267-282.
- 250 Jamal I., Khalil H., Bari A., Khan S. and Zada I., 2009. Genetic variation for yield and yield components in rice.  
251 *J. Agric. Biol. Sci.* 4(6): 60-64
- 252 Jayasudha S. and Sharma D., 2010. Genetic parameters of variability, correlation and path-coefficient for grain  
253 yield and physiological traits in rice (*Oryza sativa* L.) under shallow lowland situation. *Electronic J. Plant*  
254 *Breed.* 1(5): 33-38.
- 255 Johnson H.W., Robinson H.F. and Comstock R.E., 1955. Estimation of genetic and environmental variability in  
256 soybean. *Agronomy J.* 47: 314-318.
- 257 Kuldeep T., Bathshwar K., Ramesh B. and Tomer A., 2004. Genetic variability and correlations for some  
258 seedlings and mature plant traits in 70 genotypes of rice. *Res. Crops.* 5(1): 60-65.
- 259 Maclean J.L., 2002 (Eds). *Rice Almanac*. Los Baños: International Rice Research Institute, Bouake; Ivory  
260 Coast: West Africa Rice Development Association; Cali: International Center for Tropical Agriculture;  
261 Rome: FAO.
- 262 Mruthunjaya C., Wali and Mahadevappa M., 1995. Genetic variability, heritability and genetic advance for yield  
263 and its contributing characters in ratoon crops rice (*Oryza Sativa* L.). *Mysore J. agric. Sci.*, 29: 285-288.
- 264 Sabesan T., Suresh R. and Saravanan K., 2009. Genetic variability and correlation for yield and grain quality  
265 characters of rice grown in coastal saline low land of Tamilnadu. *Electronic J. Plant Breed.* 1: 56-59.
- 266 Sharma S.K. and Sharma B., 1983. Mutagenic sensitivity and mutability in rice. *Intl. Congerss. Genet.* 5: 330-  
267 333.
- 268 Smith HF., 1936. A discriminant function for plant selection. *Annals of Eugenics.* 7: 240-250.
- 269 Tahir M., Wadan D. and Zada A., 2002. Genetic Variability of Different Plant Yield Characters in Rice. *Sarhad J.*  
270 *Agric.*18 (2).
- 271 Talukder M.S., 2004. Effect of gamma rays on large seeded rice. M.S. Thesis, Department of Genetics  
272 and Plant Breeding, Bangladesh Agricultural University, Mymensingh. pp: 24-27
- 273 Utz and H.F., 2007. PLABSTAT (Version 2N), A computer program for the computation of variances and  
274 covariances. Institute of Plant Breeding, Seed Science, and Population Genetics, University of  
275 Hohenheim, Stuttgart, Germany.
- 276 Wright S., 1935. The analysis of variance and correlations between relative with respect to deviations from an  
277 optimum. *J. Genet.* 30: 243-256.

- 278 Zahid A.M., Akhtar M., Sabar M., Anwar M. and Ahmad M., 2005. Interrelation-ship among Yield and Economic  
279 Traits in Fine Grain Rice. Proceedings of the International Seminar on Rice Crop. October 2-3. Rice  
280 Research Institute, Kala Shah Kau, Pakistan. pp. 21-24.  
281
- 282 Sarker, M. M., Hassan, L., Islam, M. M., Rashid, M. M., & Seraj, S. (2014). Correlation and path coefficient  
283 analysis of some exotic early maturing rice (*Oryza sativa* L.) lines. *Journal of Bioscience and Agriculture*  
284 *Research*, 1(1), 01-07.  
285