

Original Research Article

***Ricinus communis* L.: Water Use Efficiency, Carbon Assimilation and Water Relations on Deficit Irrigation**

ABSTRACT

Aims: This study evaluated carbon assimilation, water relations, intrinsic and instantaneous water use efficiency, and water consumption of two cultivars of *Ricinus communis* L. cv. BRS 188 Paraguaçu and BRS Energia, subjected to regulated-deficit irrigation.

Study Design: The experiment was arranged in a completely randomized scheme in a factorial arrangement of 5 x 2, with five replicates.

Place and Duration of Study: The experiment was conducted in a greenhouse at the Universidade Estadual de Santa Cruz, Ilhéus, Brazil from December 2008 to February 2009.

Methodology: The growing plants were subjected to different water conditions by predefined quantities of water, so as to maintain the substrate under the following matric potential (ψ_m) during the experimental period: -1.6 kPa (near field capacity), -3.0 kPa, -7.3 kPa, -26.7 kPa, and -183.0 kPa.

Results: The cultivars differed significantly ($P = .05$) in predawn leaf water potential and relative water content, showing that the tissues of BRS Energia remained more hydrated compared to BRS 188 Paraguaçu. Under -183.0 kPa, the intrinsic water use efficiency and instantaneous water use efficiency were significantly higher in BRS Energia than in BRS 188 Paraguaçu, suggesting a conservative behavior of the cultivar BRS Energia. Non-stomatal limitations to photosynthesis were observed in BRS 188 Paraguaçu. Under greater water stress, BRS 188 Paraguaçu and BRS Energia plants had the leaf area reduced by 75.58% and 23.13%, respectively compared with the control. The water use efficiency of biomass was significantly higher in BRS Energia than in BRS 188 Paraguaçu.

Conclusion: The cultivar BRS Energia was more promising in relatively drier conditions compared to BRS 188 Paraguaçu. The carbon assimilation decreased in both castor bean cultivars only under severe water stress (-183.0 kPa), suggesting that the use of the deficit irrigation technique may be viable leading to lower water consumption and higher photosynthesis efficiency.

Keywords: castor bean; water stress; gas exchange; biomass; Euphorbiaceae.

1. INTRODUCTION

Castor bean (*Ricinus communis* L.), one of the 7000 species of the family Euphorbiaceae [1]. Castor bean is an important oil-seed crop grown throughout the world [2]. Production is concentrated on India, China, Brazil and Mozambique [3]. In Brazil, small- and medium-scale farmers have been producing castor oil for more than a century, especially in the state of Bahia [4,5]. Cultivation of castor bean is a good alternative to those farmers, because this crop has a low production cost, is drought-tolerance can be easily cultivated [6,7], and can grow any where including in infertile soil considered unsuitable for food production [8]. The species shows satisfactory fruit production even in the semi-arid region of northeastern Brazil where rainfall is sparse [9]. Thus, castor bean may be an alternative source of income for farmers in northeastern Brazil [9], especially for family farmers [10,8], allowing them to remain economically viable [11].

Given the global climate changes that are increasing water scarcity, irrigation and rational use of water have become important objects of study [12]. Strategies to reduce irrigation-water consumption and to improve water use efficiency (*WUE*) have become a priority for water conservation in agriculture [13]. In the cultivation of *Pyrus* L., deficit irrigation has reduced water consumption by about 5 to 18%, i.e., this irrigation method has enabled a water saving from 13-25% compared to full irrigation [14]. Regulated Deficit Irrigation (*RDI*) is among the water-saving strategies based on the adaptive and specific responses of plants to drought [15], where supplying less water than the plants requires is an important tool for reducing consumption of irrigation water [16,17]. Several cases of success using this technique have been reported, with gains in productivity [16] of many species such as *Olea europaea* L. [18], *Dianthus caryophyllus* L. [19],

58 *Capsicum annum* L. [20], *Citrus sinensis* [21], *Prunus armeniaca* [22], *Pistacia vera* L. [23], *Vitis*
59 *vinifera* L. [24] and *Citrus paradisi* Mac. [25]. Deficit irrigation (50% of evapotranspiration) in *Vitis*
60 *vinifera* L. cultivation was sufficient to ensure a high yield, **to water use efficiency** - *WUE*
61 (yield/water applied in irrigation) and good fruit quality [26]. *WUE* can be optimized by increasing
62 the productivity of a crop in line with the volume of water applied, or by reducing irrigation without
63 significantly reducing productivity [27].

64 Energy crops such as castor beans have attracted **attention to** producing biofuels such as
65 biodiesel, in developed as well as developing countries **contributing** to reduce dependency on
66 fossil fuel [8]. Studies on castor bean production systems in the climate conditions of Brazil are
67 especially relevant with regard to irrigation conditions, in order to augment the income of
68 producers [28].

69 The castor bean cultivar BRS Energia has an earlier cycle in relation to the other cultivars, with
70 120-150 days between the germination and maturation of recent racemes, and the first raceme
71 appears about 30 days after germination [29]. Thus, the precocity associated with easy cultivation
72 makes a cultivar BRS Energy with great productive **potential for** great social and economic
73 importance to the semi-arid region of northeastern Brazil. The BRS 188 Paraguaçu has agronomic
74 and technological characteristics superior to those of commercial cultivars [30]. Thus, the
75 comparative study of the physiological characteristics of each cultivar under water restriction
76 conditions can aid in selecting the best cultivar in response to the minimum water availability
77 needed for higher productivity and lower costs.

78 Growing of drought-tolerant cultivars will contribute to stable castor bean production, while the
79 screening of cultivars or breeding **lines of** drought stress responses can be a crucial part of
80 breeding programs [2]. In the present study, our main objective was to evaluate carbon
81 assimilation, water relations, intrinsic and instantaneous water use efficiency, and water
82 consumption of two castor bean cultivars, BRS 188 Paraguaçu and BRS Energia, subjected to
83 regulated deficit irrigation.

84

85 **2. MATERIAL AND METHODS**

86 **2.1 Plant material and growing conditions**

87

88 The experiment was conducted in a greenhouse at the Universidade Estadual de Santa Cruz,
89 Ilhéus, Bahia, Brazil (14°47'00" S, 039°02'00" W) from December 2008 to February 2009.
90 According to the Köppen climate classification, the local climate is the Af type humid tropical
91 climate, with mean annual temperatures ranging from 22 to 25°C [31]. During the experimental
92 period inside the greenhouse the air temperature ranged from 24 °C to 31 °C and relative humidity
93 (RH) from 65% to 98% (Hobo H8 Pro sensors, Onset Computer, Massachusetts, USA), and
94 cumulative photosynthetically active radiation (PAR) from 4.9 to 33 mol photons m⁻² day⁻¹ (S-LIA-
95 M003 quantum sensors coupled to a Hobo Micro Station Data Logger, Onset Computer,
96 Massachusetts, USA).

97 Two cultivars of *Ricinus communis* L. (BRS 188 Paraguaçu and BRS Energia) with different
98 growing cycle were used in the study. In BRS 188 Paraguaçu, the mean period between seedling
99 emergence and emission of the first raceme (inflorescence) is 54 days and the whole growing
100 cycle last for 250 days. The mean oil content of its seeds is 48%, and the mean yield are 1,500
101 kg/ha in a longer 250-day cycle under the rain-fed semi-arid conditions of northeastern Brazil [32].
102 BRS Energia is a shorter cycle cultivar with 120 to 150 days between the germination and
103 maturation of recent racemes, whereas the first raceme emerges earlier from about 30 days after
104 germination [29]. The oil content of seeds is 48% and fruit productivity is 1,500 kg/ha, on average,
105 under rain-fed semi-arid conditions [33].

106 The seeds were soaked for 2 h and then treated with the systemic fungicide Derosal®. The plants
107 were grown for 66 days in 21L pots filled with a mixture of sand and soil (3:1); textural analysis
108 frank-sandy. The substrate was prepared based on its chemical composition (Table 1). Pots
109 similar to those used in the experiment were assembled to estimate field capacity of substrate.
110 After correcting the pH with 1.55 g dm⁻³ dolomitic limestone (PRNT 90.87%) and adding 1.37 g dm⁻³
111 triple superphosphate and 0.60 g dm⁻³ of ready commercial formulation containing (N -16%; K₂O
112 - 16%; S - 7%; B - 0.2%; Cu - 0.2%; MgO - 1%; Zn and Mn - 0,5%.

113 Top-dressing chemical fertilization was based on 80 mg dm⁻³ urea and 10 mg dm⁻³ potassium
114 chloride. Each pot was filled with a known weight of soil which was irrigated to field capacity and
115 then sown five seeds per pot. When the plantlets were approximately 0.10 to 0.12 m tall, they were
116 thinned by leaving only one plant per pot. The plantlets isolated from thinning were used to collect

117 zero (initial biomass). Each pot was fertilized monthly with 50 mL of nitrogen (urea) and potassium
 118 (potassium chloride) solutions at concentrations of 56.8 kg/ha⁻¹ and 20 kg/ha⁻¹, respectively.
 119

Table 1. Chemical analysis of the substrate used in the experiment

pH	cmol _c /dm ³							mg/dm ³			
	Al	H+Al	Ca	Mg	Ca+Mg	P	K	Fe	Zn	Cu	Mn
4.47	0.67	4.9	0.19	0.08	0.27	0.4	8	103	1.17	0.5	1.7

120

121 Regulated-deficit irrigation (*RDI*) was started at 32 days after sowing (DAS) and the growing plants
 122 were then subjected to different water conditions by predefined quantities of water, so as to
 123 maintain the substrate under the following matric potential (Ψ_m) during the experimental period: -
 124 1.6 kPa (near field capacity), -3.0 kPa, -7.3 kPa, -26.7 kPa, and -183.0 kPa. The substrate Ψ for
 125 each treatment was estimated using an equation derived from the soil water-retention curve (Table
 126 2).

127 **Table 2. Mean percentages of water content of substrate (WCS) 20, 16, 12, 9 and 7% and**
 128 **their corresponding matric potential (Ψ_m)**

129

Treatments	WCS (%)	Ψ_m (-kPa)
20	19.7	1.6
16	15.6	3.0
12	12.1	7.3
9	9.1	26.7
7	6.7	183.0

130

131 Before each irrigation, all the pots were weighed and the difference between the current weight
 132 and that corresponding to each treatment corresponded to the weight of replacement water
 133 (evapotranspiration). Water consumption was considered as the water lost by the plants via
 134 transpiration, and the evaporation from the substrate in the pot.

135

136 2.2 Water relations

137 The pre-dawn leaf water potential (ψ_{PD}) was evaluated 18 days after the RDI application (DAAT),
138 using a Pressure Chamber Instrument Model 1000 (PMS Instrument Company, USA).
139 Pressurization was carried out slowly, and the **time of** the leaf collection and the measurement
140 was as short as possible [34]. The measurements were performed between 02:00 and 04:00 h,
141 when the mean air temperature **was** around 23.3°C and the relative humidity was 74%.
142

UNDER PEER REVIEW

143 **2.3 Leaf relative water content**

144 Leaf samples were first weighed (P1) and then placed to hydrate in pots filled with water, for 12 h
145 in the dark, this time was enough to reach the max turgor. After hydration, the leaves were
146 weighed again to obtain the turgid weight (P2) and were then placed in a forced-air oven at 75°C
147 for 72 h to obtain the biomass dry weight (P3). Relative water **content** was calculated using the
148 following formula: $RWC = [(P1-P3)/(P2-P3)] \times 100$ [35].

149

150 **2.4 Leaf gas exchange**

151 Leaf gas **exchanges were** evaluated 18 days after the application of treatments (DAAT), between
152 08:00 and 12:00 h, in the middle part of fully expanded physiologically mature leaves from five
153 randomly selected plants from each treatment. Net photosynthesis rate (A), intercellular CO_2
154 concentration (C_i), stomatal conductance to water vapor (g_s), and transpiration (E) per unit of leaf
155 area were measured using the Li-6400 Portable Photosynthesis System (LI-COR Biosciences Inc.,
156 Nebraska, USA) with integrated fluorescence camera (LI-6400-40 leaf chamber fluorometer, LI-
157 COR). Photosynthetically active radiation (PAR), atmospheric CO_2 concentration (C_a), and block
158 temperature were set at $1200 \text{ mol photons m}^{-2} \text{ s}^{-1}$, $400 \text{ } \mu\text{mol mol}^{-1}$ and 26°C , respectively, using
159 the equipment controls.

160

161 **2.5 Water use efficiency**

162 Three **forms into** expressing water use efficiency were used in the analysis and interpretation of
163 experimental data: instantaneous water use efficiency (A/E), intrinsic water use efficiency (A/g_s)
164 and water use efficiency of biomass (kg m^{-3}), calculated as the ratio of biomass produced to water
165 consumed (evapotranspiration). The calculations were performed with data collected at 8 DAAT
166 (1stharvest) and 34 DAAT (2nd harvest).

167

168 **2.6 Biomass determination**

169 Two destructive **measurements of** the beginning (8 DAAT) and the end (34 DAAT) of the
170 experimental period were performed. The harvests were treated independently, since the plants
171 collected 8 DAAT were different from those collected 34 DAAT. Leaf area was estimated, both
172 non-destructively and destructively, using allometric coefficients (width and length of a mature leaf)

173 previously generated for this purpose as described by [36], and a LI-COR 3100 (Biosciences Inc.,
174 Nebraska, USA) automatic leaf area meter. The dry mass of plant organs (root, stem and leaves)
175 was used to estimate the variables for growth, such as relative growth rate (*RGR*) according to
176 Hunt (1990). Each plant was placed in paper bags and oven-dried in a forced-air oven at 75°C
177 until constant weight.

178

179 **Statistical analysis**

180 The experiment was arranged in a completely randomized scheme in a factorial arrangement of 5
181 x 2, wherein the factors were: five water regimes and two cultivars of *R. communis*, with five
182 replicates. Differences between the cultivars were assessed using a t-test at 5% probability.

183

184 **3. RESULTS AND DISCUSSION**


185 **3.1 Leaf water relations**

186 The effects of deficit irrigation on ψ_{PD} and *RWC* differed between the two cultivars (Fig.1 A, B).
187 *RWC* was significantly higher in BRS Energia, with mean values of 89, 85 and 76% at -3.0; -7.3
188 and -183.0 kPa soil matric potential, respectively (Fig.1 A); whereas the corresponding values for
189 BRS 188 Paraguaçu were 83, 79 and 64% (Fig.1A). These data showed that although both
190 species consumed the same amount of water (Fig. 4 C, D), the short-cycle cultivar BRS Energia
191 was able to maintain more-hydrated tissues compared to the longer-cycle BRS 188 Paraguaçu,
192 especially at higher water deficits. One can therefore infer that BRS Energia is the more
193 promising cultivar in relatively dry locations due to its ability to maintain higher *RWC* and ψ_w .

194 The *RWC* is probably the most appropriate measure of plant water status in terms of the
195 physiological consequences of cellular water deficit. According to [37], the restriction of leaf water
196 status resulting from a reduction in *RWC* affects plant growth and development as observed in
197 BRS 188 Paraguaçu.

198 As observed for the *RWC*, the ψ_{PD} of BRS Energia was significantly higher than that of BRS 188
199 Paraguaçu, with values of -0.49 and -0.89 MPa \square_{PD} in the former in contrast to -0.6 and -1.4
200 MPa \square_{PD} in the latter at -7.3 and -183.0 kPa, respectively (Fig.1 B). The non-significant difference
201 between the cultivars for *RWC* and the significant difference between ψ_{PD} in -26.7 kPa (Fig.1 A,

202 B) may suggest some degree of osmotic adjustment, which enabled the plants to maintain turgor
 203 in a relatively low water potential.



204

205

206 **Fig. 1. (A) Relative Water Content (RWC) and (B) pre-dawn leaf water potential (Ψ_{PD}) in**
 207 **plants of *Ricinus communis* cv. BRS 188 Paraguaçu and cv. BRS Energia subjected to**
 208 **different water conditions: -1.6; -3.0; -7.3; -26.7 and -183.0 kPa after 18 days of treatment**
 209 **application (DAAT). Points are mean (n=5), error bars are the standard error of the mean,**
 210 **and letters indicate significant differences between cultivars with the same water level, by**
 211 **t-test ($P = .05$)**

212

213 Studies with different hybrids of *R. communis* showed that this species accumulates high
 214 contents of proline, total soluble sugars, amino acids and potassium after 33 days under water
 215 stress, and the sugars are the key players in osmotic adjustment in castor bean leaves [38].
 216 Similarly, *Jatropha curcas* plants possess an efficient adaptive mechanism to prevent severe
 217 drought stress by maintaining good leaf water status and effective osmotic adjustment [39,40].


218 In soil matric potential for -3.0 kPa, both cultivars had significantly similar Ψ_w but with different
 219 RWC values (Fig.1 A, B). This indicates that although the status of the water within the cells was
 220 the same, the leaf hydration status and physiological water were different.

221

222 3.2 Leaf gas exchange

223 The cultivars showed different behaviors for A/g_s , A/E and C_i/C_a when subjected to -183.0 kPa,
 224 with higher values for BRS Energia than for BRS 188 Paraguaçu (Fig.2 C, D and F). Both

225 cultivars had A , g_s and E constant at approximately $26 \mu\text{mol m}^{-2} \text{s}^{-1}$, $0.45 \text{ mol H}_2\text{O m}^{-2} \text{s}^{-1}$ and 3.8
226 $\text{mmol H}_2\text{O m}^{-2} \text{s}^{-1}$, respectively, after 18 days under matrix potential for the substrate above -26.7
227 kPa (Fig.2 A, B, D), showing that gas exchange was not affected when the matrix potential for the
228 substrate exceeded -26.7 kPa , regardless of the cultivar. The reduction in the photosynthesis rate
229 observed at -183.0 kPa (Fig.2 A), in turn, was closely associated with the closure of stomata
230 (Fig.2 B). The reduction in g_s increases resistance to CO_2 diffusion into the leaves, affecting the
231 accumulation of photoassimilates (Fig.4 A, B) [41]. If the plant loses water at a faster rate than its
232 capacity to absorb and transport it, then the leaf water potential decreases, causing the closure of
233 stomata and the reduction of photosynthesis (Fig. 1 B, Fig. 2 A, B) [42]. Similarly, in *J. curcas*, net
234 photosynthesis was significantly reduced only when soil water availability dropped below 30% of
235 field capacity. However, g_s proved to be quite sensitive to soil water availability, and the strict
236 stomatal regulation in this species was evident after 11 days of stress [39]. Compared to BRS
237 188 Paraguaçu, higher A/g_s was observed in plants of BRS Energia subjected to increased water
238 deficit (Fig. 2 C). This behavior is attributable to the rapid stomatic closure observed in BRS
239 Energia to minimize water loss and thus maintain leaf ψ_w (Fig. 2 B). The stomatal closure
240 contributed to optimize the efficiency of water use for the plants under stress [43], allowing them
241 to optimize CO_2 fixation versus water loss. Stomatal closure is considered a drought-avoidance
242 mechanism [44].



243

244

245 Fig. 2. (A) Net Photosynthesis rate (A); (B) stomatal conductance for water vapor (g_s); (C)
 246 intrinsic water use efficiency (A/g_s); (D) transpiration; (E) instantaneous water use
 247 efficiency (A/E) and (F) ratio (intercellular and atmospheric CO_2 concentrations) (C_i/C_a) of
 248 two castor bean cultivars cultivated in substrate with -1.6; -3.0; -7.3; -26.7 and -183.0 kPa
 249 of matric potential for 18 days after treatment application (DAAT). Points are mean ($n=5$),
 250 error bars are the standard error of the mean, and letters indicate significant differences (P
 251 = .05) by t -test between cultivars with the same water level.

252

253 This difference in behavior between the two cultivars was also observed in *Lotus corniculatus*

254 where the transpiration rate, RWC and g_s reflect specific physiological mechanisms in each

255 cultivar, and allow for metabolic acclimatization to drought conditions [45]. [46] obtained similar
256 results, and stated that the castor bean drought-resistance mechanism appears to be related to
257 an initial response and increased growth, as well as efficient stomatal control, minimizing water
258 loss from transpiration. Although the studies of *J. curcas* by [47] revealed that a reduction in water
259 availability (100, 75, 50 and 25% field capacity) resulted in decreased g_s and E in order to avoid
260 loss of water, however, the WUE was reduced.

261 The rapid closing of stomata and the lower E observed in the lower matrix potential for the
262 substrate for BRS Energia in relation to BRS 188 Paraguaçu (Fig. 2 B, D) resulted in increased
263 A/g_s and A/E (Fig. 2 C, E). This improved the hydration of leaf tissue (Fig. 1 A), suggesting a
264 conservative approach [48, 49, 50].


265 The C_i/C_a ratio of both cultivars was maintained at 0.65 in substrates above -26.7 kPa. Water
266 contents below -26.7 kPa led to a behavior contrary to that observed for A/g_s (Fig.2 C, F); thus,
267 the low value of C_i/C_a followed by an increase in the A/g_s of BRS Energia plants are due to low
268 g_s [39]. On the other hand, the higher CO_2 concentration of intercellular spaces (C_i) subjected to
269 low g_s observed in BRS 188 Paraguaçu indicates that this cultivar was more sensitive to the RDI
270 compared to BRS Energia (Fig.2 F). This behavior suggests the occurrence of non-stomatal
271 limitations of photosynthesis, such as low mesophyll conductance, reduced activity and
272 concentration of ribulose-1,5-bisphosphatecarboxylase-oxygenase (Rubisco), photoinhibition, and
273 reduced photochemical efficiency of PSII [51,52,53].

274

275 3.3 Growth and biomass accumulation

276 Because the experiment consisted of two cultivars with different cycles, short-cycle BRS Energia
277 (120-150 days) and long-cycle BRS 188 Paraguaçu (250 days), only the reproductive cycle of
278 BRS Energia was evaluated. According to literature, the BRS 188 Paraguaçu cultivar begins the
279 reproductive stage at 53 DAS [33]; however, in our study, no flowering was observed up to 66
280 DAS.

281 At 8 DAAT, due to the dry conditions, plant height was gradually reduced, especially in plants
282 subjected to -183.0 kPa, with reductions of 38.81 and 33.28% compared to the controls in BRS
283 Energia and BRS 188 Paraguaçu, respectively (Fig.3 A). At 34 DAAT, the reductions were even
284 more significant, 51.48% and 40.17%, respectively (Fig.3 B).



285

286

287

288

289

290

291

292

Fig. 3. Plant height (cm), leaf area (cm²), shoot biomass (g) and root biomass (g) of two castor bean cultivars grown in substrate with -1.6; -3.0; -7.3; -26.7 and -183.0 kPa at 8 DAAT (A, C, E and G) and 34 DAAT (B, D, F and H). Points are mean (n=5), error bars are the standard error of the mean.

This indicates that the plant height of cultivars is determined, among other factors, by the water supply [54], which inhibits cell elongation more than division, affecting various physiological and

293 biochemical processes such as photosynthesis, respiration, translocation, absorption of ions,
294 carbohydrates, nutrient metabolism, and growth factors [55].

295 Reductions in height were also observed by [56] in cultivars BRS 149 Nordestina and BRS 188
296 Paraguaçu, with reductions of 40.24, 24.89 and 13.83% in treatments with 40, 60 and 80%
297 available water compared to plants in soil maintained at field capacity.

298 After 8 DAAT there was a reduction in leaf area with increasing water stress, soon after the plants
299 were subjected to the treatments (Fig. 3 C).

300 Similarly, [57] reported a leaf-area reduction of more than 60% in BRS 188 Paraguaçu under
301 excess water stress and deficiency in only six days, and stated that in the juvenile stage until the
302 first 52 days after seedling emergence, this cultivar is very sensitive to water stress.

303 At 34 DAAT, under greater water stress, the plants showed a quite compromised leaf area, with
304 reductions of 75.58% and 23.13% compared with the control, for BRS 188 Paraguaçu and BRS
305 Energia, respectively (Fig. 3 D). According to [58], the reduction in leaf area, due to selective leaf
306 senescence combined with decreases in A and A/g_s (Fig. 2 A, C), allows plants to maintain an
307 "above-lethal" water potential. The same authors observed a similar behavior in *J. curcas* after 18
308 days of water stress. The reduction in leaf area and gas exchange during dry conditions reduces
309 not only water loss but also carbon assimilation, with consequent slower growth [59]. The smaller
310 reduction in leaf area observed in BRS Energia compared to BRS 188 Paraguaçu, especially at -
311 183.0 kPa, resulted from the ability of the former to produce leaves, although small, whereas BRS
312 188 Paraguaçu lost leaves. According to [45], the regrowth process generates small turgid leaves
313 that are physiologically acclimated to drought, showing obvious morphological changes resulting
314 from changes in growth and leaf development. At 34 DAAT, the longer period of drought had
315 significantly affected the shoot biomass of plants of both cultivars. At -183.0 kPa, cultivars BRS
316 Paraguaçu and BRS Energia showed reductions of 79.02 and 85.44% respectively, compared to
317 control plants (Fig. 3 F).


318 The root development was also strongly influenced by growing conditions. At 34 DAAT, the root
319 biomass at -183.0 kPa was lower than in the control, with reductions of 61.25 and 56.04% in BRS
320 Energia and BRS 188 Paraguaçu, respectively (Fig. 3 H). This indicates that both cultivars
321 showed no root growth in the most intense drought conditions, reducing the shoot:root ratio. [60]
322 noted that root growth is usually less affected by drought stress than shoot growth. A decrease in

323 the shoot:root ratio is a common observation under drought stress, which results either from an
324 increase in root growth or from a relatively larger decrease in shoot growth than in root growth, as
325 a result of pre-conditioning deficit-irrigation processes. Furthermore, as the matric potential of the
326 substrate decreased, the percentage of shaded roots in the BRS 188 Paraguaçu plants increased
327 possibly the result of suberization of the exodermis to protect the roots from adverse conditions
328 [60].

329 Within a short period of time (8 DAAT), the plants subjected to water-deficit treatments showed a
330 significant decrease in total biomass (*TB*) due to the reduction of the matric potential in the
331 substrate (Fig.4 A), indicating high sensitivity of growth to reduced water availability. When
332 subjected to severe water deficit (-183.0 kPa), total biomass decreased by 56% in both cultivars
333 compared to the control (Fig. 4 B). Leaves comprised most of the *TB* (Fig.3 D). This reduction in
334 growth of biomass observed in both species is attributable to a survival strategy.

335 The reductions in growth and biomass accumulation observed in the plants subjected to water
336 deficit, especially in BRS 188 Paraguaçu, are due to decreases in Ψ_w , which has been
337 associated with a reduction in the coefficient of cell division and in cell expansion [61], mainly
338 driven by leaf turgor pressure (Ψ_p). Similar behavior was observed in *J. curcas* after 18 days of
339 stress [58].

340 After 34 DAAT (Fig. 4 B), water deficits below -3.0 kPa reduced (*TB*) production, by 18.21, 25.47
341 and 75.97% in BRS Energia and 3.57, 35.10 and 80.95% in BRS 188 Paraguaçu at -7.3; -26.7
342 and -183.0 kPa in comparison with the control, respectively. With the reduction in water
343 availability, the water consumption (evapotranspiration) decreased linearly to values of 11.71,
344 7.41, 6.43, 4.14 and 0.53 L (BRS 188 Paraguaçu) and 11.35, 7.60, 5.69, 3.98 and 0.71 L (BRS
345 Energia), with mean daily consumption of 1.46, 0.93, 0.80, 0.52 and 0.07 L (BRS 188 Paraguaçu)
346 and 1.41, 0.95, 0.71, 0.50 and 0.09 L (BRS Energia) at -1.6, - 3.0, - 7.3, - 27.7 and -183.0 MPa,
347 respectively, over 8 DAAT (Fig. 4 C). Even so, there were no significant differences between the
348 cultivars. Similar results were observed for the same castor bean cultivars where the highest
349 water consumption (2534 mm) occurred with 100% available water over the 180 days of the crop
350 cycle [62].



351

352 Fig. 4. Total biomass (*TB*), cumulative water consumption (*WC*), relative growth rate in
 353 biomass (*RGR*) and water use efficiency (*WUE*) of two castor bean cultivars cultivated in
 354 substrate with -1.6; -3.0; -7.3; -26.7 and -183.0 kPa for 8 DAAT (A, C, E and G) and 34 DAAT
 355 (B, D, F and H). Points are mean (n=5), error bars are the standard error of the mean, and

356 **letters indicate significant differences ($P = .05$) by t -test between cultivars with the same**
357 **water level.**

358 During the entire experiment (34 DAAT), the final water consumption was 47.47, 39.53, 33.40,
359 22.41 and 6.33 L in BRS 188 Paraguaçu and 42.31, 39.22, 30.69, 22.94 and 7.72 L in BRS
360 Energia at -1.6, - 3.0, -7.3, -26.7 and -183.0 MPa of soil water, respectively, with a mean daily
361 consumption of 1.40, 1.16, 0.98, 0.66 and 0.19 L (BRS 188 Paraguaçu) and 1.24, 1.15, 0.90,
362 0.67 and 0.23 L (BRS Energia) (Fig. 4 D). Despite the different plant architectures of the two
363 cultivars, there were no differences in evapotranspiration.

364 BRS 188 Paraguaçu had a reduced RGR when subjected to -1.60 kPa water in the substrate at 8
365 DAAT (Fig. 4 E). Similar results were found by [63], who attributed the delay in development and
366 consequent limitation of the respiratory process of BRS 188 Paraguaçu to the 4.80% reduction in
367 growth of the root system at the highest soil water content, which was 100% field capacity.

368 Reductions in RGR were evident after 34 DAAT, in particular in BRS 188 Paraguaçu, where the
369 RGR was negative ($-8.58 \text{ mg g}^{-1} \text{ day}^{-1}$) (Fig. 4 F). Considering that the RGR corresponds to the
370 amount of new material produced in relation to the pre-existing material over time [64], the
371 cultivar BRS 188 Paraguaçu had stopped growth, which explains why the RGR was negative.
372 BRS Energia, in contrast, still showed positive values of RGR ($9.8 \text{ mg g}^{-1} \text{ day}^{-1}$) even under a
373 severe soil water deficit (Fig. 4 F). Those results suggest that the cultivar BRS 188 Paraguaçu is
374 less tolerant to water deficit compared to BRS Energia.

375 The lower water availability resulted in a decrease in A (Fig. 2 A) and consequently in the
376 production of carbohydrates, contributing to a reduction in biomass accumulation (Fig. 4 E, F) of
377 the plants. Similar results were found in *J. curcas*, in terms of CO_2 assimilation, stomatal
378 conductance, transpiration, growth, biomass and water use efficiency which progressively
379 reduced in response to decreasing soil moisture content [47].

380

381 **3.4 Water use efficiency (WUE)**

382 The WUE was evaluated taking into account the evapotranspiration of water (soil evaporation +
383 leaf transpiration) and dry biomass production. For both, pots containing only substrate were
384 covered with plastic to estimate evaporation, but the estimate was very low and was therefore
385 disregarded. Shading of the pot's surface by leaves further reduced evaporation, so that the

386 evaporation was higher than the transpiration. The *WUE* of BRS Energia increased linearly with
387 decreased matric potential in the substrate at 8 DAAT, reaching a *WUE* of up to 6 kg m^{-3} (Fig.4
388 G). This behavior can be attributed to increased branching and length of the roots. This can
389 minimize the depletion of water around the roots, thereby minimizing resistance to transport of
390 water to the root system [65].

391 The substrate with a matric potential of -1.6 kPa reduced the *WUE* of BRS 188 Paraguaçu at 8
392 DAAT (Fig.4G). Our results are not consonant with those obtained by [62], who in studies
393 involving BRS 188 Paraguaçu found increased *WUE* in the treatment with 100% available water
394 in relation to the lowest level (40%), with values of 2.78 and 0.28 kg m^{-3} , respectively. This
395 discrepancy can be attributed to the time when the analyses were performed: in the studies
396 conducted by [62] the cultivation time was 180 days, and the present study lasted 66 days.

397 At 34 DAAT, only for *WUE*, indicating that the cultivars have different behaviors as a function of
398 watering regimes (Fig.4 H). In contrast, the *WUE* of the BRS Energia plants was significantly
399 higher ($2.1, 2.4, 2.6$ and 1.1 kg m^{-3}) than that of the BRS 188 Paraguaçu plants ($1.6, 2.0, 1.9$ and
400 -0.4 kg m^{-3}) at $-1.6, -3.0, -26.7$ and -183 kPa , respectively (Fig.4H). In the same period, the *WUE* of
401 the plants was reduced in soil with the highest water deficit, regardless of the cultivar. The lower
402 efficiency recorded for BRS 188 Paraguaçu in relation to BRS Energia may possibly be attributed
403 to the decrease in *g_s* during water deficiency, which reduces the assimilation efficiency (0.05
404 $\mu\text{mol m}^{-2} \text{ s}^{-1}$) through photosynthesis, since BRS Energia showed higher values than BRS
405 188 Paraguaçu at -26.7 and -183.0 kPa . Similarly, it was found in *J. curcas* a reduction in *WUE*
406 under dry conditions most likely due to the negative effect of the higher potentials on the
407 production of plant biomass [66]. However, in this study, soil with matric potential greater than $-$
408 183.0 kPa allowed the plants to maintain *WUE*.

409

410 **4. CONCLUSION**

411 Among the variables studied here, the relative water content, predawn leaf water
412 potential, biomass, and relative growth rate were more sensitive to regulated water deficits. The
413 cultivar BRS Energia was more promising in relatively drier conditions compared to BRS 188
414 Paraguaçu, since it was able to maintain a larger leaf area and more-hydrated tissues,
415 maximizing the efficiency of water use. The carbon assimilation decreased in both castor bean

416 cultivars only under severe water stress (-183.0 kPa), suggesting that the use of the deficit
417 irrigation technique may be viable leading to lower water consumption and higher photosynthesis
418 efficiency.

419

420

421 **COMPETING INTERESTS**

422 The authors declare that there is no conflict of interests.

423

424

425 **REFERENCES**

- 426 1. Beltrão NEM, Silva LC. The multiple uses of castor oil (*Ricinus communis* L.) and the
427 importance of its cultivation in Brazil. *Fibers and Oils*. 1999; (31): 7. English.
- 428 2. Radhamani T, Ushakumari R, Amudha R, Anjani K. Response to water stress in castor
429 (*Ricinus communis* L.) genotypes under *in vitro* conditions. *J Cer Oil Seeds*. 2012; 3 (4): 56-58.
430 Available: <http://dx.doi/10.5897/JCO12.016>
- 431 3. Severino LS, Auld DL. Study on the effect of air temperature on seed development and
432 determination of the base temperature for seed growth in castor (*Ricinus communis* L.). *Aust J*
433 *Crop Sci*. 2014; 8 (2): 290-295.
- 434 4. Beltrão NEM, Cartaxo WV, Pereira SRP, Soares JJ, Silva ORRF. The sustainable
435 cultivation of castor bean in the Brazilian semi-arid region. *Campina Grande: EMBRAPA-CNPA 23*
436 *p (EMBRAPA-CNPA, Technical Circular, 84); 2005. English.*
- 437 5. Obiero C, Birech R, Maling'a J, Ngetich K, Freyer B. Performance of maize and beans
438 under castor-based intercropping system. *Am J Exp Agric*. 2014; 4 (1): 101-113.
- 439 6. Beltrão NEM, Oliveira MIP. Targeting of the residues and co-products from the biodiesel
440 production from castor beans and jatropha. *Bahia anal data*. 2009; 18 (4): 613-619. English.
- 441 7. Fogaça JJNL, Silva RA, Santos JL, Nunes RTC, Ferreira LL, Morais OM. Physiological
442 quality of castor bean seeds var. Carrapatinho depending on the position of the dealer. *Rev. Bras.*
443 *Agrarian Ciênc*. 2017; 40 (1): 87-93. Available: <http://dx.doi.org/10.19084/RCA16086>. English.

- 444 8. Nahar K. Castor bean (*Ricinus communis* L.) - the biofuel plant: morphological and
445 physiological parameters propagated from seeds in Bangladesh. Asian Busin. Review. 2013; 2 (2):
446 64-66. Available: <http://dx.doi.org/10.18034/abr.v2i2.306>.
- 447 9. Azevedo DMP, Nóbrega LB, Lima EF, Batista FAZ, Beltrão NEM. Cultural Management.
448 In: Azevedo DMP, Lima EF, editors. The agribusiness of castor bean in Brazil. Campina Grande:
449 Embrapa Algodão, Brasília: Embrapa Technological Information. 2001: 121-160. English.
- 450 10. Beltrão NEM, Vale LS, Marques LF, Cardoso GD, Souto JS. Castor bean and peanut
451 consortium: option for family agriculture. Rev. Verde Agroec Des Sust. 2010; 5 (4): 222-227.
452 Portuguese.
- 453 11. Lopes JS, Andrade TCQ, Santana GC. Biodiesel: opportunities and challenges. Rev
454 Bahia Agríc. 2007;8:24-27. Portuguese.
- 455 12. Chaves MM, Oliveira MM. Mechanisms underlying plant resilience to water deficits:
456 prospects for water-saving agriculture. J Exp Bot. 2004;55(1):2365-2384.
457 Available:<http://dx.doi.org/10.1093/jxb/erh269>.
- 458 13. Tabarzad A, Ghaemi AA. Deficit irrigation and sowing date as strategies to maximize
459 water use efficiency and crop water productivity in semi-arid region. Biol Forum Int J.
460 2015;7(1):30-42.
- 461 14. Cui N, Du T, Kang S, Li F, Zhang J, Wang M et al. Regulated deficit irrigation improved
462 fruit quality and water use efficiency of pear-jujube trees. Agric Water Manage. 2008;95:489-497.
463 Available:<http://dx.doi.org/10.1016/j.agwat.2007.11.007>.
- 464 15. Ma S-C, Duan A-W, Wang R, Guan Z-M, Yang S-J, Ma S-T et al. Root-sourced signal and
465 photosynthetic traits, dry matter accumulation and remobilization, and yield stability in winter
466 wheat as affected by regulated deficit irrigation. Agric Water Manage. 2015;148:123-129.
467 Available:<http://dx.doi.org/10.1016/j.agwat.2014.09.028>.
- 468 16. Fereres E, Soriano MA. Deficit irrigation for reducing agricultural water use. J Exp Bot.
469 2007;58:147-159. Available:<http://dx.doi.org/10.1093/jxb/erl165>.
- 470 17. Ruiz-Sanchez MC, Domingo R, Castel JR. Deficit irrigation in fruit trees and vines in
471 Spain. Span. J Agric Res. 2010;8:5-20. Available:<http://dx.doi.org/10.5424/sjar/201008S2-1343>.

- 472 18. Masmoudi CC, Ayachi MM, Gouia M, Laabidi F, Reguya SB, Amor AO et al. Water
473 relations of olive trees cultivated under deficit irrigation regimes. *Sci Hort.* 2010;25:573–578.
474 Available:<http://dx.doi/10.1016/j.scienta.2010.04.042>.
- 475 19. Álvarez S, Navarro A, Bañón S, Sánchez-Blanco MJ. Regulated deficit irrigation in potted
476 *Dianthus* plants: Effects of severe and moderate water stress on growth and physiological
477 responses. *Sci Hort.* 2009;122(4):579–585.
478 Available:<http://dx.doi.org/10.1016/j.scienta.2009.06.030>.
- 479 20. Shao G-C, Liu N, Zhang Z-Y, Yu S-E, Chen C-R. Growth, yield and water use efficiency
480 response of greenhouse-grown hot pepper under Time-Space deficit irrigation. *Sci Hort.*
481 2010;126:172–179. Available:<http://dx.doi/10.1016/j.scienta.2010.07.003>.
- 482 21. Tejero IG, Zuazo VHD, Bocanegra JAJ, Fernández JLM. Improved water use efficiency by
483 deficit-irrigation programmes: Implications for saving water in citrus orchards. *Sci Hort.*
484 2011;128:274-282. Available:<http://dx.doi/10.1016/j.scienta.2011.01.035>.
- 485 22. Pérez-Pastor A, Domingo R, Torrecillas SA, Ruiz-Sánchez MAC. Response of apricot
486 trees to deficit irrigation strategies. *Irrig Sci.* 2009;27:231–
487 242. Available:<http://dx.doi/10.1007/s00271-008-0136-x>.
- 488 23. Iniesta F, Testi L, Goldhamer DA, Fereres E. Quantifying reductions in consumptive
489 water use under regulated deficit irrigation in pistachio (*Pistacia vera* L.). *Agric Water Manage.*
490 2008;95:877–886. Available:<http://dx.doi.org/10.1016/j.agwat.2008.01.013>.
- 491 24. Acevedo-Opazo C, Ortega-Farias S, Fuentes S. Effects of grapevine (*Vitis vinifera* L.)
492 water status on water consumption, vegetative growth and grape quality: An irrigation scheduling
493 application to achieve regulated deficit irrigation. *Agric. Water Manage.* 2010;97(7):956-964.
494 Available:<http://dx.doi.org/10.1016/j.agwat.2010.01.025>.
- 495 25. Navarro JM, Botía P, Pérez-Pérez JG. Influence of deficit irrigation timing on the fruit
496 quality of grapefruit (*Citrus paradisi* Mac.). *Food Chem.* 2015;175:329-
497 336. Available:<http://dx.doi/10.1016/j.foodchem.2014.11.152>.
- 498 26. Santos TP, Lopes CM, Rodrigues ML, Souza CR, Silva JMR, Maroco JP, Pereira JS,
499 Chaves MM. Effects of deficit irrigation strategies on cluster microclimate for improving fruit
500 composition of Moscatel field-grown grapevines. *Sci Hort.* 2007;12:321–330.
501 Available:<http://dx.doi/10.1016/j.scienta.2007.01.006>.

- 502 27. Coelho EF, Coelho-Filho MA, Oliveira SL. Irrigated agriculture: efficiency of irrigation and
503 water use. *Bahia Agríc.* 2005; 7 (1): 57-60. English.
- 504 28. Passion FJR, Beltrão NEM, Azevedo CAV, Pimentel JVF, Carlos JA. Production and
505 yield components of castor bean BRS Energia in function of different levels of irrigation and
506 nitrogen organic fertilization. *Braz J Appl Technol Agric Sci.* 2013; 6 (3): 27-37. Available: <http://dx.doi/10.5935/PAeT.V6.N3.03>.
507
- 508 29. Brazilian Company of Research and Agriculture. National Center for Cotton Research
509 (Campina Grande, PB). Folder BRS: (Energy) Campina Grande. 2008a. English.
- 510 30. Embrapa - Brazilian Company of Research and Agropecuária. National Center for Cotton
511 Research (Campina Grande, PB). Folder BRS 188 Paraguaçu Campina Grande. 1999. English.
- 512 31. Faria Filho AF, Araújo QR. Zoning of the physical environment of the municipality of
513 Ilhéus, Bahia, Brazil, using the geoprocessing technique. Ilheus, CEPLAC / CEPEC. Technical
514 Bulletin. 2003; 187, 20p. English.
- 515 32. Freire EC, Lima EF, Andrade FP, Milani M, Nóbrega MBM. Genetic Improvement. In:
516 Azevedo DMP, Beltrão NEM editors. *The castor agribusiness in Brazil*. Embrapa Algodão
517 (Campina Grande, PB) 2 ed. Brasília, DF: Embrapa Technological Information. 2007: pp..
- 518 33. Embrapa Embrapa, Brazilian Research and Agricultural Company. National Center for
519 Cotton Research (Campina Grande, PB). Folder BRS: (Paraguaçu). Campina Grande. 2008b.
520 English.
- 521 34. Scholander PF, Hammel HT, Bradstreet ED, Hemmingsen EA. Sap pressure in vascular
522 plants. *Science.* 1965; 148 (3668): 339-346. Available: <http://dx.doi/10.1126/science.148.3668.339>.
523
- 524 35. Kramer PJ, Boyer JS. *Water relations of plant and soils*, Academic Press, New York,
525 1995.
- 526 36. Severino LS, Cardoso GD, Vale LS, Santos JW. Method for determining the leaf area of
527 the castor bean. *Research and Development Bulletin*, 55 EMBRAPA COTTON (Campina
528 Grande, PB). 2005, 20p. English.
- 529 37. Lima GS, Gheyi HR, Nobre RG, Soares LAA, Xavier DA, Santos Júnior JA. Water
530 relations and gas exchange in castor bean irrigated with saline water of distinct cationic nature.
531 *Afr J Agric Res.* 2015;10(13):1581-1594. Available: <http://dx.doi/10.5897/AJAR2015.9606>.

- 532 38. Babita M, Maheswari M, Rao LM, Shanker AK, Rao DG. Osmotic adjustment, drought
533 tolerance and yield in castor (*Ricinus communis* L.) hybrids. Environ Exp Bot. 2010;69(3):243-
534 249. Available:<http://dx.doi.org/10.1016/j.envexpbot.2010.05.006>.
- 535 39. Sapeta H, Costa JM, Lourenço T, Maroco J, Van der Linde P, Oliveira MM. Drought
536 stress response in *Jatropha curcas*: Growth and physiology. Environ Exp Bot. 2013; 85:76-84.
537 Available:<http://dx.doi/10.1016/j.envexpbot.2012.08.012>.
- 538 40. Silva EN, Ferreira-Silva SL, Viégas RA, Silveira JAG. The role of organic and inorganic
539 solutes in the osmotic adjustment of drought-stressed *Jatropha curcas* plants. Envir ExpBot.
540 2010;69:279-285. Available:<http://dx.doi/10.1016/j.envexpbot.2010.05.001>.
- 541 41. Gholz HL, Ewel KC, Teskey RO. Water and forest productivity. For Ecol Manage.
542 1990;30:1-18. Available:[http://dx.doi.org/10.1016/0378-1127\(90\)90122-R](http://dx.doi.org/10.1016/0378-1127(90)90122-R).
- 543 42. Marengo RA, Lopes NF. **Plant physiology: Photosynthesis, respiration, water relations
544 and mineral nutrition.** 1th ed. UFV, Viçosa; 2005.
- 545 43. Gullías J, Cifre J, Jonasson S, Medrano H, Flexas J. Seasonal and inter-annual variations
546 of gas exchange in thirteen woody species along a climatic gradient in the Mediterranean island
547 of Mallorca. Flora – Morphology, Distribution, Funct Ecol Plants. 2009;204:169-181.
548 Available:<http://dx.doi/10.1016/j.flora.2008.01.011>.
- 549 44. Rouhi V, Samson R, Lemeur R, Van Damme P. Photosynthetic gas exchange
550 characteristics in three different almond species during drought stress and subsequent recovery.
551 Environ Exp Bot. 2007;59:117-129. Available:<http://dx.doi/10.1016/j.envexpbot.2005.10.001>
- 552 45. Inostroza L, Acuña H, Tapia G. Relationships between phenotypic variation in osmotic
553 adjustment, wateruse efficiency, and drought tolerance of seven cultivars of *Lotus corniculatus* L.
554 Chil J Agric Res. 2015;75(1):3-12. Available:[http://dx.doi.org/10.4067/S0718-
555 58392015000100001](http://dx.doi.org/10.4067/S0718-58392015000100001).
- 556 46. Sausen TL, Rosa LMG. Growth and carbon assimilation limitations in *Ricinus communis*
557 (Euphorbiaceae) under soil water stress conditions. Acta Bot Bras. 2010;24(3):648-654.
558 Available:<http://dx.doi/10.1590/S0102-33062010000300008>.
- 559 47. Verma KK, Vatsal S, Gupta RK, Ranjan S, Verma CL, Singh M. Influence of water
560 application on photosynthesis, growth and biomass characteristics in *Jatropha curcas*. Curr Bot.
561 2012;3(4):26-30.

- 562 48. Kudoyarova GR, Kholodova VP, Veselov DS. Current state of the problem of water
563 relations in plants under water deficit. Russian Plant Physiol. 2013;60(2):155-165.
564 Available:<http://dx.doi/10.1134/S102144371302014>.
- 565 49. Lima Neto MC, Martins MO, Ferreira-Silva SL, Silveira JAG. *Jatropha curcas* and *Ricinus*
566 *communis* display contrasting photosynthetic mechanisms in response to environmental
567 conditions. Sci Agric. 2015;72(3):260-269. Available:<http://dx.doi/10.1590/0103-9016-2014-0325>.
- 568 50. Maes WH, Achten WMJ, Reubens B, Raes D, Samson R, Muys B. Plant-water
569 relationships and growth strategies of *Jatropha curcas* L. seedlings under different levels of
570 drought stress. J Arid Environ. 2009;73:877-
571 884. Available:<http://dx.doi/10.1016/j.jaridenv.2009.04.013>.
- 572 51. Flexas J, Medrano H. Drought-inhibition of photosynthesis in C₃ plants: stomatal and non-
573 stomatal limitations revisited. Ann Bot. 2002; 89(2):183–189.
574 Available:<http://dx.doi.org/10.1093/aob/mcf027>.
- 575 52. Grassi G, Magnani F. Stomatal, mesophyll conductance and biochemical limitations to
576 photosynthesis as affected by drought and leaf ontogeny in ash and oak trees. Plant Cell Environ.
577 2005;28:834–849. Available:<http://dx.doi.org/10.1111/j.1365-3040.2005.01333.x>.
- 578 53. Lawlor DW. Limitation to photosynthesis in water stressed leaves: stomata versus
579 metabolism and the role of ATP. Ann Bot. 2002;89:871–
580 885. Available:<http://dx.doi.org/10.1093/aob/mcf110>.
- 581 54. Sampaio Filho OM, Silva SA, Bahia HF, Silva MS, Carvalho DS. Descriptive analysis of
582 castor bean cultivars in two years of cultivation in the Bahian recôncavo. Rev Bras Educ Amb.
583 2011; 6: 28-34. English.
- 584 55. Jaleel CA, Gopi R, Sankar B, Gomathinayagam M, Panneerselvam R. Differential
585 responses in water use efficiency in two varieties of *Catharanthus roseus* under drought stress. C
586 R Biol. 2008; 331 (1): 42-47. Available: <http://dx.doi/10.1016/j.crv.2007.11.003>
- 587 56. Formiga LA, Guerra HOC, Lacerda RD, Silva JEB, Araujo MS. Effect of available soil
588 water on the development of two castor bean cultivars in the first and second cycles. Eng Agríc.
589 2014; 34 (6): 1128-1138. Available: <http://dx.doi.org/10.1590/S0100-69162014000600009>.
590 English.

- 591 57. Beltrão NEM, Souza JG, Santos JW. Water stress (deficiency and excess) and its effects
592 on the initial growth of the castor bean cultivar BRS 188 Paraguaçu. Rev Bras Oleag Fibrosas.
593 2003; 7 (2/3): 735-741. Portuguese.
- 594 58. Fini A, Bellasio C, Pollastri S, Tattini M, Ferrini F. Water relations, growth, and leaf gas
595 exchange as affected by water stress in *Jatropha curcas*. J Arid Environ. 2013;89:21-29.
596 Available:<http://dx.doi.org/10.1016/j.jaridenv.2012.10.009>.
- 597 59. Bañón S, Ochoa J, Franco JA, Alarcón JJ, Sánchez-Blanco MJ. Hardening of oleander
598 seedlings by deficit irrigation and low air humidity. Environ Exp Bot. 2006;56(1):36-43.
599 Available:<http://dx.doi.org/10.1016/j.envexpbot.2004.12.004>.
- 600 60. Franco JA. Root development under drought stress. Technol Know Transfer e-Bull.
601 2011;2(6):1-3.
- 602 61. Cattivelli L, Rizza F, Badeck F-W, Mazzucotelli E, Mastrangelo AM, Francia E et al.
603 Drought tolerance improvement in crop plants: An integrated view from breeding to genomics.
604 Field Crops Res. 2008;105(1/2):1-14. Available:<http://dx.doi.org/10.1016/j.fcr.2007.07.004>.
- 605 62. Barros Júnior G, Guerra HOC, Cavalcanti MLF, Lacerda RD. Water consumption and use
606 efficiency for two castor bean cultivars submitted to water stress. Rev Bras Eng Agríc Ambient.
607 2008; 12 (4): 350-355. Available: <http://dx.doi.org/10.1590/S1415-43662008000400003>. English.
- 608 63. Nunes EN, Nascimento DAM, Alves AG, Suassuna JF, Nascimento R. Growth of castor
609 bean cultivars (*Ricinus communis* L.) at different soil water levels. Sci Plena. 2013; 9 (10): 1-10.
610 English.
- 611 64. Hunt R. Basic growth analysis, Unwin Hyman, London; nineteen ninety.
- 612 65. Franco JA, Martínez-Sánchez JJ, Fernández JA, Bañón S. Selection and nursery
613 production of ornamental plants for landscaping and xerogardening in semi-arid environments. J
614 Hortic Sci Biotech. 2006; 81: 3-17. Available: <http://dx.doi.org/10.1080/14620316.2006.11512022>
- 615 66. Santana TA, Oliveira PS, Silva LD, Laviola BG, Almeida A-AF, Gomes FP. Water use
616 efficiency and consumption in different Brazilian genotypes of *Jatropha curcas* L. subjected to soil
617 water deficit. Biomass Bioenergy. 2015; 75: 119-125. Available: <http://dx.doi.org/10.1016/j.biombioe.2015.02.008>
- 618