

# **Original Research Article**

## **Vitamin C, Antioxidant and Polyphenol Activity of Some Selected Potato Varieties as Influenced by Vermicompost**

### **ABSTRACT**

The application of vermicompost may improve the quality of potato. The present study revealed that vermicompost had a significant effect on most of the quality contributing parameters studied under the experiment. Results demonstrated that quality parameters increased with increasing vermicompost level. Among the sixteen treatments combination, BARI Alu-25 (Asterix) with vermicompost at the rate of 6 t/ha showed the highest ascorbic acid (Vitamin C), antioxidant and polyphenol content. In the case of ambient storage condition; ascorbic acid and polyphenol decreased with an increasing storing period while antioxidant content increased with the increasing storing period up to 60 days after storage (DAS). BARI Alu-25 (Asterix) and BARI Alu-29 (Courage) may store under ambient storage condition up to 60 DAS without imparting any significant quality losses just prior to the sprouting of the tuber. It may be concluded that the potato growers of Bangladesh may apply vermicompost on their field at the rate of 6 t/ha for maintaining the good quality of potato.

**KEYWORDS:** Antioxidant, Potato, Polyphenol, Vermicompost, Vitamin C.

### **1. INTRODUCTION**

Potato (*Solanum tuberosum* L.) belonging to the Solanaceae family is grown in nearly 150 countries and is the world's single major tuberous crop with a vital role in the global food system and food security [1]. It is the 4<sup>th</sup> world's largest produced crop after maize, wheat and rice. The total world potato production was estimated 376.83 million metric tons in 2016 [2]. It is the most highly produced non-grain staple crop in the world, with one-third of total production harvested in densely-populated developing countries, like China and India [3]. In the world top ten potato producing countries, Bangladesh ranks 7<sup>th</sup> position [4]. Nutritive value of potato is relatively high, because of protein content and composition of high percentage of essential amino acids: lysine, leucine, phenylalanine, threonine and valine [5]. Potato is also characterized by high amounts of starch and lower content of sugars. It is an excellent source of vitamin C and other biologically active substances, such as polyphenols and flavonoids, which are commonly described as antioxidants [6, 7]. These substances have a beneficial influence on the human being, as they protect against cardiovascular disease and cancer, as well as reduce blood cholesterol level. Ascorbic acid (L-AA, vitamin C) as an additive is largely accepted in human diets because of its antioxidative potential [8]. Potato peel is a good source of natural antioxidants. Its extract provides protection against acute liver injury [9] and oxidative damage to erythrocytes [10]. Due to the increasing demand of consumers and foreign importers on this important crop, special attention should be given to increase its quality. Now-a-days gradual deficiencies in soil organic matter and reduced yield of crop and quality are alarming problem in Bangladesh. The cost of inorganic fertilizers is very high. On the other hand, the organic manure is easily available to the farmers and its cost is low compared to that of inorganic fertilizers. Vermicompost is a good source of different macro and micronutrients particularly NPKS. The increased microbial activity also improves the availability of soil phosphorous and nitrogen, which is very helpful for increasing potato quality [4]. So, using different amount of vermicompost materials may put contribution for improving quality of potato in Bangladesh condition.

## 2. MATERIALS AND METHODS

The field experiment was conducted at the Agronomy research field at Sher-e-Bangla Agricultural University, Dhaka-1207 during the period from November 1, 2016, to April 30, 2017, in Rabi season. The experimental area was situated at the 23°77'N latitude and 90°33'E longitude at an altitude of 8.6 meters above the sea level [4]. Topsoil was silty clay in texture, olive-grey with common fine to medium distinct dark yellowish-brown mottles. Soil pH was 5.6 and has organic carbon 0.45%. The experimental area was flat having available irrigation and drainage system and above flood levels. The selected plot was medium high land. The experiment was provoked in a split-plot design with 3 replications. The variety was assigned to the main plot and vermicompost to subplot. The distance between row to row was 50 cm and plant to plant distance was 25 cm. The distance between plot to the plot was 75 cm. The size of the unit plot was 2 m x 2.5 m. The experiment consisted of two factors, were as follows: Factor A: Potato Varieties (V-4); V<sub>1</sub>: BARI TPS-1, V<sub>2</sub>: BARI Alu-28 (Lady Rosetta), V<sub>3</sub>: BARI Alu-25 (Asterix), V<sub>4</sub>: BARI Alu-29 (Courage). Factor B: Vermicompost level (M-4); M<sub>1</sub>: 0 t/ha (Control), M<sub>2</sub>: 2 t/ha, M<sub>3</sub>: 4 t/ha, M<sub>4</sub>: 6 t/ha. All varieties of seed potato (certified seed) were collected from Tuber Crops Research Centre (TCRC), Bangladesh Agricultural Research Institute (BARI), Joydebpur, Gazipur and from BARI sub-station. Individual weight of seed potato was 60-70 g. Collected seed tubers were kept in room temperature to facilitate sprouting. Finally, sprouted potato tubers were used as planting material. Recommended doses of fertilizer were used as Urea 250 kg/ha, TSP 150 kg/ha, MoP 250 kg/ha, Gypsum 120 kg/ha, Zinc Sulphate 10 kg/ha and Boric Acid 10 kg/ha [11]. Different intercultural operations were done as per when needed. After haulm cutting the tubers were kept under the soil for 10 days for skin hardening. After harvesting the potatoes were collected and stored at ambient storage condition. Data were collected up to the sprouting of maximum potatoes.

### 2.1 Ascorbic acid determination

5gm of tuber sample was taken after meshing in a mortar by using a pestle. Then 10ml of HPO<sub>3</sub> was added in the sample and volume up to the mark. After centrifugation (2000rpm, 10min) it was filtrated and taken 5ml of solution for titration. Then the solution was titrated against indophenol solution and titration record was collected in a notebook. After that in x-axis used concentration of Ascorbic acid (mg) and in y-axis value of titration to find out the standard curve. Then calculated the ascorbic acid as mg/g FW (Fresh Weight).

### 2.2 Antioxidant determination

Preparation of the sample was the same as polyphenol determination. Total antioxidant capacity of fresh potatoes was quantified using the DPPH (2,2-diphenyl-1-picrylhydrazyl) method [12]. The absorbance reading was used to calculate spectrophotometer at 517nm wavelength. All the values of antioxidant were expressed as Trolox micrograms per gram fresh weight (Trolox µg/g FW) using a nonlinear regression algorithm from Trolox standard curve.

### 2.3 Polyphenol determination

The content of total phenolic compounds was determined spectrophotometrically according to the Folin-Ciocalteu method [13] with slight modification. The absorbance of the reaction solution was measured at 760 nm against a blank sample. The measurement was compared to a standard curve of gallic acid solutions and expressed as microgram of gallic acid equivalents per gram fresh weight (µg GAE/g FW).

### 2.4 Statistical Analysis

The data collected for different quality characters were statistically analyzed following the analysis of variance techniques by using MSTAT-C computer package program. The significant difference among the treatment means were differentiated by Least Significant Difference (LSD) at 5% level of probability [14].

### 3. RESULT AND DISCUSSION

#### 3.1 Ascorbic acid content

##### 3.1.1 Effect of variety

Significant variation was found among different varieties to ascorbic acid content of tuber at different storage time. The highest ascorbic acid content of tuber was exhibited by V<sub>3</sub> (73.75 mg, 68.04 mg, 57.97 mg, 51.24 mg) and lowest was exhibited by V<sub>1</sub> (45.62 mg, 39.37 mg, 40.23 mg, 36.37 mg); at 0, 20, 40 and 60 DAS, respectively (Table 1).

**Table 1. Effect of variety on ascorbic acid at different days after storage of potato tuber**

Variety	Ascorbic acid (mg/g FW)			
	0 DAS	20 DAS	40 DAS	60 DAS
V <sub>1</sub>	45.62 d	39.37 d	40.23 d	36.37 d
V <sub>2</sub>	57.61 c	53.82 c	44.11 c	39.40 c
V <sub>3</sub>	73.75 a	68.04 a	57.97 a	51.24 a
V <sub>4</sub>	68.94 b	62.52 b	53.69 b	47.88 b
CV (%)	6.20	4.48	5.30	4.96
LSD <sub>0.05</sub>	2.59	2.51	2.59	2.16
Level of significance	**	**	**	**

Numbers in columns followed by the same letter are not statistically different at P<sub>0.05</sub>.

\*\* = Significant at 1% level of probability,

V<sub>1</sub> – BARI TPS-1, V<sub>2</sub> – Lady Rosetta, V<sub>3</sub> – Asterix, V<sub>4</sub> – Courage

##### 3.1.2 Effect of vermicompost

Profound variation was found among different levels of vermicompost on ascorbic acid content of tuber at different storage time. The highest ascorbic acid content was contained by M<sub>4</sub> (79.80 mg, 74.95 mg, 58.40 mg, 52.52 mg) and lowest was contained by M<sub>1</sub> (46.63 mg, 40.20 mg, 37.44 mg, 31.34 mg); at 0, 20, 40 and 60 DAS, respectively (Table 2).

**Table 2. Effect of vermicompost on ascorbic acid at different days after storage of potato tuber**

Vermicompost level	Ascorbic acid (mg/g FW)			
	0 DAS	20 DAS	40 DAS	60 DAS
M <sub>1</sub>	46.63 d	40.20 d	37.44 d	31.34 d
M <sub>2</sub>	55.25 c	48.79 c	48.04 c	43.06 c
M <sub>3</sub>	64.23 b	59.80 b	52.10 b	47.97 b
M <sub>4</sub>	79.80 a	74.95 a	58.40 a	52.52 a
CV (%)	3.76	4.57	5.33	5.29
LSD <sub>0.05</sub>	1.95	2.15	2.20	1.95
Level of significance	**	**	**	**

Numbers in columns followed by the same letter are not statistically different at P<sub>0.05</sub>.

\*\* = Significant at 1% level of probability

M<sub>1</sub> – Control, M<sub>2</sub> – 2 t/ha, M<sub>3</sub> – 4 t/ha, M<sub>4</sub> – 6 t/ha

##### 3.1.3 Combined effect of variety and vermicompost

Significant variation was found among different variety and vermicompost levels on ascorbic acid content of tuber at different storage time. The maximum ascorbic acid content of tuber was contained by V<sub>3</sub>M<sub>4</sub> (90.89 mg, 86.78 mg) and minimum was contained by V<sub>1</sub>M<sub>1</sub> (36.75 mg, 30.54 mg); at 0 and 20 DAS, respectively. At 40 DAS maximum ascorbic acid content of tuber was contained by V<sub>3</sub>M<sub>4</sub> (66.60 mg) which was statistically similar to V<sub>3</sub>M<sub>3</sub> and minimum was contained by V<sub>1</sub>M<sub>1</sub> (28.54 mg). At 60 DAS maximum ascorbic acid content of tuber was contained by V<sub>3</sub>M<sub>4</sub> (59.09 mg) which was statistically similar to V<sub>3</sub>M<sub>3</sub> and minimum was contained by V<sub>1</sub>M<sub>1</sub> (25.94 mg) (Table 3).

Ascorbic Acid (AA) content significantly decreases during storage condition of potato tuber. The reduction was occurred for the oxidation of ascorbic acid into dehydro ascorbic acid and afterward to diketo-gluconic acid. Being a water-soluble vitamin and susceptible to oxidation, AA contents rapidly decrease with increasing rates of respiration and water loss in storage [15].

**Table 3. Combined effect of variety and vermicompost on ascorbic acid at different days after storage of potato tuber**

Combination	Ascorbic acid (mg/g FW)			
	0 DAS	20 DAS	40 DAS	60 DAS
V <sub>1</sub> M <sub>1</sub>	36.75 i	30.54 i	28.54 h	25.94 g
V <sub>1</sub> M <sub>2</sub>	39.50 i	30.58 i	41.67 g	35.54 f
V <sub>1</sub> M <sub>3</sub>	46.44 h	42.10 g	39.69 g	37.47 f
V <sub>1</sub> M <sub>4</sub>	59.78 ef	54.24 e	51.01 d	46.54 de
V <sub>2</sub> M <sub>1</sub>	38.90 i	36.90 h	32.21 h	28.94 g
V <sub>2</sub> M <sub>2</sub>	47.94 h	42.33 g	38.01 g	33.31 f
V <sub>2</sub> M <sub>3</sub>	61.81 e	58.40 e	48.99 de	45.20 e
V <sub>2</sub> M <sub>4</sub>	81.78 c	77.64 b	57.22 c	50.13 cd
V <sub>3</sub> M <sub>1</sub>	56.01 fg	48.78 f	46.33 ef	37.27 f
V <sub>3</sub> M <sub>2</sub>	70.10 d	64.72 d	56.39 c	52.40 bc
V <sub>3</sub> M <sub>3</sub>	77.99 c	71.86 c	62.54 ab	56.20 ab
V <sub>3</sub> M <sub>4</sub>	90.89 a	86.78 a	66.60 a	59.09 a
V <sub>4</sub> M <sub>1</sub>	54.86 g	44.57 fg	42.70 fg	33.22 f
V <sub>4</sub> M <sub>2</sub>	63.48 e	57.54 e	56.10 c	51.00 c
V <sub>4</sub> M <sub>3</sub>	70.69 d	66.83 d	57.19 c	53.00 bc
V <sub>4</sub> M <sub>4</sub>	86.74 b	81.13 b	58.76 bc	54.32 bc
CV (%)	3.76	4.57	5.33	5.29
LSD <sub>0.05</sub>	3.89	4.31	4.40	3.89
Level of significance	**	**	**	**

Numbers in columns followed by the same letter are not statistically different at P<sub>0.05</sub>.

\*\* = Significant at 1% level of probability

V<sub>1</sub> –BARI TPS-1, V<sub>2</sub> – Lady Rosetta, V<sub>3</sub> – Asterix, V<sub>4</sub> – Courage

M<sub>1</sub> – Control, M<sub>2</sub> – 2 t/ha, M<sub>3</sub> – 4 t/ha, M<sub>4</sub> – 6 t/ha

### 3.2 Antioxidant content

#### 3.2.1 Effect of variety

In respect of antioxidant content of tuber at 0 and 40 DAS were found statistically non-significant. At 20 and 60 DAS significant variation was found among different varieties on antioxidant content of tuber. At 20 DAS the highest (22.44 µg) was exhibited by V<sub>3</sub> which was statistically similar to V<sub>4</sub> and the lowest (21.73 µg) was exhibited by V<sub>1</sub>. At 60 DAS the highest (27.74 µg) antioxidant content of tuber was exhibited by V<sub>3</sub> which was statistically similar to V<sub>4</sub> and the lowest (26.23 µg) was exhibited by V<sub>1</sub> (Table 4).

**Table 4. Effect of variety to antioxidant at different days after storage of potato tuber**

Variety	Antioxidant (Trolox µg/g FW)			
	0 DAS	20 DAS	40 DAS	60 DAS
V <sub>1</sub>	20.49	21.73 c	22.15	26.23 c
V <sub>2</sub>	20.83	21.96 bc	22.28	26.81 b
V <sub>3</sub>	21.43	22.44 a	22.58	27.74 a
V <sub>4</sub>	21.09	22.17 ab	22.44	27.34 a
CV (%)	5.21	1.55	2.32	1.72
LSD <sub>0.05</sub>	-	0.341	-	0.465
Level of significance	NS	**	NS	**

Numbers in columns followed by the same letter are not statistically different at  $P_{0.05}$ .

\*\* = Significant at 1% level of probability, NS = Not significant

$V_1$  – BARI TPS-1,  $V_2$  – Lady Rosetta,  $V_3$  – Asterix,  $V_4$  – Courage

### 3.2.2 Effect of vermicompost

Profound variation was found among different levels of vermicompost on antioxidant content of tuber at different storage time. The highest (21.82  $\mu\text{g}$ , 22.72  $\mu\text{g}$ ) antioxidant of tuber was contained by  $M_4$  and the lowest (20.22  $\mu\text{g}$ , 21.48  $\mu\text{g}$ ) was contained by  $M_1$ , at 0 and 20 DAS, respectively. At 40 DAS the highest (22.94  $\mu\text{g}$ ) antioxidant of tuber was contained by  $M_4$  which was statistically similar to  $M_3$  and the lowest (21.65  $\mu\text{g}$ ) was contained by  $M_1$ . At 60 DAS the highest (27.95  $\mu\text{g}$ ) antioxidant of tuber was contained by  $M_4$  and the lowest (26.00  $\mu\text{g}$ ) was contained by  $M_1$  (Table 5).

**Table 5. Effect of vermicompost on antioxidant at different days after storage of potato tuber**

Vermicompost level	Antioxidant (Trolox $\mu\text{g}/\text{g}$ FW)			
	0 DAS	20 DAS	40 DAS	60 DAS
$M_1$	20.22 c	21.48 d	21.65 c	26.00 d
$M_2$	20.73 b	21.81 c	22.26 b	26.80 c
$M_3$	21.07 b	22.28 b	22.60 ab	27.38 b
$M_4$	21.82 a	22.72 a	22.94 a	27.95 a
CV (%)	2.14	1.61	1.90	1.39
LSD <sub>0.05</sub>	0.377	0.299	0.357	0.317
Level of significance	**	**	**	**

Numbers in columns followed by the same letter are not statistically different at  $P_{0.05}$ .

\*\* = Significant at 1% level of probability

$M_1$ – Control,  $M_2$  – 2 t/ha,  $M_3$  – 4 t/ha,  $M_4$  – 6 t/ha

### 3.2.3 Combined effect of variety and vermicompost

No significant variation was found among different variety and vermicompost levels on antioxidant content of tuber at 0, 20 and 40 DAS. At 60 DAS due to different variety and vermicompost levels was found statistically significant. The maximum (28.43  $\mu\text{g}$ ) antioxidant content of tuber contained by  $V_3M_4$  which was statistically similar to  $V_3M_3$  and  $V_4M_4$ . The minimum (25.27  $\mu\text{g}$ ) antioxidant was contained by  $V_1M_1$  (Table 6).

The activity of total antioxidant increased from harvesting to eight weeks of storage time and its activity varied with genotypic responses over storage period [16].

**Table 6. Combined effect of variety and vermicompost on antioxidant at different days after storage of potato tuber**

Combination	Antioxidant (Trolox $\mu\text{g}/\text{g}$ FW)			
	0 DAS	20 DAS	40 DAS	60 DAS
$V_1M_1$	19.45	21.20	21.23	25.27 e
$V_1M_2$	20.31	21.53	22.12	25.62 e
$V_1M_3$	20.67	21.66	22.52	26.35 d
$V_1M_4$	21.51	22.51	22.74	27.67 b
$V_2M_1$	20.33	21.30	21.58	25.54 e
$V_2M_2$	20.61	21.58	22.19	26.46 d
$V_2M_3$	20.78	22.43	22.56	27.55 bc
$V_2M_4$	21.60	22.52	22.81	27.69 b
$V_3M_1$	20.61	21.84	21.97	26.91 cd
$V_3M_2$	21.20	22.27	22.45	27.62 b
$V_3M_3$	21.49	22.57	22.69	28.01 ab
$V_3M_4$	22.43	23.09	23.20	28.43 a

V <sub>4</sub> M <sub>1</sub>	20.51	21.57	21.84	26.28 d
V <sub>4</sub> M <sub>2</sub>	20.81	21.88	22.26	27.48 bc
V <sub>4</sub> M <sub>3</sub>	21.32	22.48	22.64	27.60 b
V <sub>4</sub> M <sub>4</sub>	21.72	22.74	23.01	28.00 ab
CV (%)	2.14	1.61	1.90	1.39
LSD <sub>0.05</sub>	0.755	0.598	0.715	0.635
Level of significance	NS	NS	NS	*

Numbers in columns followed by the same letter are not statistically different at P<sub>0.05</sub>.

\*\* = Significant at 1% level of probability, NS = Not significant


V<sub>1</sub> –BARI TPS-1, V<sub>2</sub> – Lady Rosetta, V<sub>3</sub> – Asterix, V<sub>4</sub> – Courage

M<sub>1</sub> – Control, M<sub>2</sub> – 2 t/ha, M<sub>3</sub> – 4 t/ha, M<sub>4</sub> – 6 t/ha

### 3.3 Polyphenol content

#### 3.3.1 Effect of variety

Significant variation was found among different varieties to polyphenol content of tuber at different storage time. At 0 DAS the highest (0.6542 mg) polyphenol content of tuber was exhibited by V<sub>3</sub> and the lowest (0.3525 mg) was exhibited by V<sub>1</sub>. At 20 DAS the highest (0.4371 mg) polyphenol content of tuber was exhibited by V<sub>3</sub> which was statistically similar to V<sub>4</sub> and the lowest (0.2734 mg) was exhibited by V<sub>1</sub>. At 40 DAS the highest (0.1610 mg) polyphenol content of tuber was exhibited by V<sub>3</sub> and the lowest (0.1091 mg) was exhibited by V<sub>1</sub>. At 60 DAS the highest (0.1303 mg) polyphenol content of tuber was exhibited by V<sub>3</sub> which was statistically similar to V<sub>4</sub> and the lowest (0.08275 mg) was exhibited by V<sub>1</sub> (Figure 1).




**Figure 1. Effect of variety to polyphenol content of tuber at different days after storage**

(LSD values 0.031, 0.0757, 0.0070 and 0.0170 for 0 DAS, 20 DAS, 40 DAS and 60 DAS, respectively). V<sub>1</sub>–BARI TPS-1, V<sub>2</sub>– Lady Rosetta, V<sub>3</sub>– Asterix, V<sub>4</sub>– Courage

#### 3.3.2 Effect of vermicompost

Profound variation was found among different levels of vermicompost on polyphenol content of tuber at different storage time. At 0 DAS the highest (0.6667 mg) polyphenol of tuber was contained by M<sub>4</sub> and the lowest (0.2875 mg) was contained by M<sub>1</sub> (Figure 2). At 20 DAS the highest (0.4146 mg) polyphenol of tuber was contained by M<sub>4</sub> which was statistically similar to M<sub>1</sub> and the lowest (0.2457 mg) was contained by M<sub>2</sub>. At 40 DAS the highest (0.1662 mg) polyphenol of tuber was contained by M<sub>4</sub> and the lowest (0.1108 mg) was contained by M<sub>1</sub> which was statistically similar to M<sub>2</sub>. At 60 DAS the highest (0.1411 mg) polyphenol of tuber was contained by M<sub>4</sub> and the lowest (0.0778 mg) was contained by M<sub>1</sub> (Figure 2).





**Figure 2. Effect of vermicompost on polyphenol content of potato tuber at different days after storage**

(LSD values 0.026, 0.0449, 0.0092 and 0.0162 for 0 DAS, 20 DAS, 40 DAS and 60 DAS, respectively). M<sub>1</sub> – Control, M<sub>2</sub>– 2 t/ha, M<sub>3</sub>– 4 t/ha, M<sub>4</sub> – 6 t/ha

### 3.3.3 Combined effect of variety and vermicompost

Significant variation was found among different variety and vermicompost levels on polyphenol content of tuber at Different storage time. At 0 DAS the maximum (1.08 mg) polyphenol of tuber contained by V<sub>3</sub>M<sub>4</sub> and the minimum (0.180 mg) was contained by V<sub>1</sub>M<sub>1</sub>. At 20 DAS the maximum (0.571 mg) polyphenol of tuber contained by V<sub>4</sub>M<sub>1</sub> which was statistically similar to V<sub>3</sub>M<sub>3</sub> and V<sub>3</sub>M<sub>4</sub>. The minimum (0.127 mg) was contained by V<sub>1</sub>M<sub>1</sub>. At 40 DAS the maximum (0.2290 mg) polyphenol of tuber contained by V<sub>3</sub>M<sub>4</sub> and the minimum (0.09100 mg) was contained by V<sub>1</sub>M<sub>1</sub>. At 60 DAS due to different varieties and vermicompost levels was found statistically non-significant (Table 7).

The total amount of phenolics in potato varies significantly among different varieties [17]. Polyphenol oxidase (PPO) activity increases in potato due to the availability of substrate and its subsequent oxidation during storage. The prominent increase in the PPO activity in different treatments after the 1<sup>st</sup> month till the end of the storage period might be associated with an increased substrate (polyphenol) oxidation. A high amount of phenolics during storage is attributed to low PPO and high antioxidant activity in potatoes [18]. Studies revealed that the phenolics content in potato continued to increase during storage period until the onset of PPO activity [19]. Kulen *et al.*, (2013) demonstrated that storage time is also very important for phenolic compounds. They found that the total phenolic content (TPC) of potatoes was high at harvest, declined after two months of ambient storage.

**Table 7. Combined effect of variety and vermicompost on a polyphenol at different days after storage of potato tuber**

Combination	Polyphenol (GA mg/g FW)			
	0 DAS	20 DAS	40 DAS	60 DAS
V <sub>1</sub> M <sub>1</sub>	0.180 k	0.407 b-d	0.09100 h	0.06
V <sub>1</sub> M <sub>2</sub>	0.330 hi	0.127 h	0.1060 gh	0.09
V <sub>1</sub> M <sub>3</sub>	0.400 fg	0.240 g	0.1123 fg	0.09
V <sub>1</sub> M <sub>4</sub>	0.500 de	0.319 d-g	0.1270 d-f	0.09
V <sub>2</sub> M <sub>1</sub>	0.260 j	0.375 c-e	0.1053 gh	0.08
V <sub>2</sub> M <sub>2</sub>	0.370 gh	0.252 g	0.1060 f-h	0.10

V <sub>2</sub> M <sub>3</sub>	0.456 ef	0.296 e-g	0.1190 e-g	0.10
V <sub>2</sub> M <sub>4</sub>	0.526 cd	0.372 c-f	0.1490 bc	0.13
V <sub>3</sub> M <sub>1</sub>	0.420 fg	0.436 bc	0.1260 d-g	0.09
V <sub>3</sub> M <sub>2</sub>	0.496 de	0.329 d-g	0.1380 c-e	0.11
V <sub>3</sub> M <sub>3</sub>	0.620 b	0.482 ab	0.1510 bc	0.14
V <sub>3</sub> M <sub>4</sub>	1.08 a	0.501 ab	0.2290 a	0.18
V <sub>4</sub> M <sub>1</sub>	0.290 ij	0.571 a	0.1210 e-g	0.08
V <sub>4</sub> M <sub>2</sub>	0.430 f	0.274 fg	0.1220 e-g	0.11
V <sub>4</sub> M <sub>3</sub>	0.530 cd	0.381 c-e	0.1427 b-d	0.14
V <sub>4</sub> M <sub>4</sub>	0.560 c	0.466 bc	0.1597 b	0.16
CV (%)	7.51	14.53	8.32	17.46
LSD <sub>0.05</sub>	0.053	0.089	0.018	-
Level of significance	**	*	**	NS

Numbers in columns followed by the same letter are not statistically different at P<sub>0.05</sub>.

\* = Significant at 5% level of probability, \*\* = Significant at 1% level of probability,

NS = Not significant

V<sub>1</sub> – BARI TPS-1, V<sub>2</sub> – Lady Rosetta, V<sub>3</sub> – Asterix, V<sub>4</sub> – Courage

M<sub>1</sub> – Control, M<sub>2</sub> – 2 t/ha, M<sub>3</sub> – 4 t/ha, M<sub>4</sub> – 6 t/ha

#### 4. CONCLUSIONS

Tuber produced from V<sub>3</sub>M<sub>4</sub> contained maximum ascorbic acid (Vitamin C), antioxidant and polyphenol; whereas the V<sub>1</sub>M<sub>1</sub> was exhibited worst one. In conclusion, the potato growers of Bangladesh may apply vermicompost on their field at the rate of 6 tons per hectare to improve quality in the combination of BARI Alu-25 (Asterix) and BARI Alu-29 (Courage); can also store at ambient condition up to 60 days after storage (DAS) without degrading significant quality parameters.

#### REFERENCES

1. Singh M. Projection of potato export from India: a Markov chain approach. *Potato J.* 2010;37:18-55.
2. Mostofa M. An Introduction to Bioethanol and Its Prospects in Bangladesh: A Review. *J Energy Res Rev.* 2019;2(2):1-12.  
Available from: <https://doi.org/10.9734/JENRR/2019/46018>
3. CIP. Potato growth in accelerate. Accessed on January 12, 2017.  
Available: <http://www.cipotato.org/potato/facts>
4. Mostofa M, Roy TS, Chakraborty R, Modak S, Kundu PK, Zaman MS, Rahman M., Shamsuzzoha, M. Effect of Vermicompost and Tuber Size on Processing Quality of Potato during Ambient Storage Condition. *Int J Plant Soil Sci.* 2019;26(3):1-18.  
Available: <http://dx.doi.org/10.9734/ijpss/2018/46554>
5. Gumul D., Ziobro R., Noga M., Sabat, R. Characterization of five potato cultivars according to their nutritional and pro-health components. *Act Sci Polon Technol Alimt.* 2011;10(1):77-81.
6. Leszczynski W. Jakość ziemniak konsumpcyjnego (Quality of low-starch potato). *Zywn Nauka Technol Jakosc.* 2000;4:5-27.
7. Wroniak J. Walory żywieniowe ziemniaka jadalnego (Nutritional benefits of potato). *Ziemniak Polski.* 2006;16(2):17-20.
8. Astley SB. Dietary antioxidants-past, present and future? *Trends Food Sci Technol.* 2003;14(3):93-98.  
Available: [http://dx.doi.org/10.1016/s0924-2244\(02\)00281-9](http://dx.doi.org/10.1016/s0924-2244(02)00281-9)
9. Singh N, Kamath V, Narasimhamurthy K, Rajini PS. Protective effect of potato peel extract against carbon tetrachloride-induced liver injury in rats. *Environ Toxicol Pharmacol.* 2008;26(2):241–246.  
Available from: <http://dx.doi.org/10.1016/j.etap.2008.05.006>
10. Singh N, Rajini PS. Antioxidant-mediated protective effect of potato peel extract in erythrocytes against oxidative damage. *Chem Biol Interact.* 2008;173(2):97–104.



Available from: <http://dx.doi.org/10.1016/j.cbi.2008.03.008>

11. Mondal MRI., Islam MS., Jalil MAB., Rahman MM., Alam MS., Rahman MHH. Krishi Projukti Hatboi (Handbook of Agro-technology), 5th edition. Bangladesh Agricultural Research Institute, Gazipur-1701, Bangladesh. 2011:307.
12. Ferreira ICFR., Barros L., Soares ME., Bastos ML., Pereira JA. Antioxidant activity and phenolic contents of *Olea europaea* L. leaves sprayed with different copper formulations. *Food Chem.* 2007;103(1):188-195.  
Available: <https://doi.org/10.1016/j.foodchem.2006.08.006>
13. Singleton VL., Orthofer R., Lamuela RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methd. Enzymol.* 1999;299:152-178.
14. Gomez KA, Gomez AA. Statistical procedure for agricultural research. 2nd Edition. John Wiley and Sons, New York, USA. 1984;139-180.
15. Hagg M., Häkkinen U., Kumpulainen J., Ahvenainen R., Hurme, E. Effects of preparation procedures, packaging and storage on nutrient retention in peeled potatoes. *J Sci Food Agric.* 1998;77(4):519-526.  
Available: [https://doi.org/10.1002/\(sici\)1097-0010\(199808\)77:4%3C519::aid-jsfa75%3E3.3.co;2-3](https://doi.org/10.1002/(sici)1097-0010(199808)77:4%3C519::aid-jsfa75%3E3.3.co;2-3)
16. Kumar D, Singh BP, Kumar P. An overview of the factors affecting sugar content of potatoes. *Ann Appl Biol.* 2004;145(3):247-256.  
Available: <http://dx.doi.org/10.1111/j.1744-7348.2004.tb00380.x>
17. Brandl W., Herrmann K. Occurrence of chlorogenic acid in potatoes. *Zeitschrift für Lebensmittel-Untersuchung und Forschung.* 1984;179:192-194.
18. Lachman J, Hamouz K, Orsák M, Pivec V, Dvořák P. The influence of flesh colour and growing locality on polyphenolic content and antioxidant activity in potatoes. *Sci Hortic.* 2008;117(2):109-114.  
Available: <http://dx.doi.org/10.1016/j.scienta.2008.03.030>
19. Madiwale GP, Reddivari L, Holm DG, Vanamala J. Storage Elevates Phenolic Content and Antioxidant Activity but Suppresses Antiproliferative and Proapoptotic Properties of Colored-Flesh Potatoes against Human Colon Cancer Cell Lines. *J Agric Food Chem.* 2011;59(15):8155-8166.  
Available: <http://dx.doi.org/10.1021/jf201073g>
20. Kulen O, Stushnoff C, Holm DG. Effect of cold storage on total phenolics content, antioxidant activity and vitamin C level of selected potato clones. *J Sci Food Agric.* 2013;93(10):2437-2444.  
Available: <http://dx.doi.org/10.1002/jsfa.6053>