

# Protecting effect of vitamin E against chromosomal damage induced by an extremely low-frequency electromagnetic field in murine bone marrow erythrocytes

## Abstract

Living organisms including human are exposed to electromagnetic field from natural and industrial sources. Genotoxic effects of electromagnetic fields have been reported by several studies. Vitamin E is a lipid-soluble antioxidant that plays an important role in maintaining the integrity and function of cell membranes, nervous system, and reproductive system, among others, by functioning as a defense against oxidative damage. In this research the protective effects of vitamin E against the chromosomal damages induced by extremely low frequency electromagnetic field (ELEM) on bone marrow erythrocytes of adult male BALB/c mouse has been studied. Adult male BALB/c mice were intraperitoneally injected with vitamin E, with the dose of 200 mg/kg for 8 days and then exposed to ELEM with the intensity of 50Hz for 4 hours for 4 contiguous days. The mice were dissected and micronucleus assay was performed on the polychromatic erythrocytes (PCE) of their bone marrow. Results show that ELEM could increase the chromosomal damages judged by the higher frequency of MN compared to control. Results also revealed that the frequency of micronucleated polychromatic erythrocytes (MNPCE) was decreased significantly ( $p < 0.001$ ) in vitamin E treated mice compared to untreated. Data suggest the protecting effect of vitamin E in bone marrow cells of Balb/C mice against the ELEM.

**Keywords:** electromagnetic fields, vitamin E, erythrocyte, BALB/c, micronucleus.

## Introduction

Development and industrialization of societies has led to utilization of electricity more than ever. In recent years, due to the abundant use of devices generating electromagnetic waves, study of their biological effects on human and other living organisms' health has become more and more important (1). These waves are generated by different devices and appliances, such as

refrigerator, television, microwave, computer displays, printers, and halogen lamps and so on. Several studies have been performed on the analysis of the effect of waves at the molecular, cellular levels and also on the field of biophysics and cancer protection (2). Many studies have proved that electromagnetic waves affect living organisms function at cellular and molecular level (3, 4, 5 & 6). Because of its proved harmful effects on living organisms, attempts are being made to find out the solutions to reduce or prevent its unwanted biological effects (2, 3).

Also there are evidences showing the relationship between exposure to electromagnetic fields and gene expression irregularities and damage to DNA molecule (7). Damages to this molecule are important because of its role in cell proliferation and viability, and also mutations and cancer.

Electromagnetic waves had genotoxic effects leading to chromosomal instability and increase of structural damages to chromosomes on peripheral blood of human and different cells of other mammals (8, 9).

Electromagnetic waves are shown to be a potent oxidative agent to the cells and living organisms. It can cause oxidative damages to DNA by increasing the extent of lipid peroxidation and the iron level, while decreasing total antioxidant status, copper, and GSH values through overproduction of free radicals/reactive oxygen species (10). These oxidative changes may lead to oncogenic transformation of the exposed cells.

Antioxidants can prevent the damages caused by free radicals and protect the macromolecules of the cells through their scavenging capabilities. One of these antioxidants is vitamin E, this vitamin is lipid-soluble and have essential role in biological membranes, lipoproteins, nervous and reproductive systems. There are many reports about the function of this vitamin in preventing or improvement of the diseases, for example, its role in cardiovascular disease and ovarian cancer (11,12). This vitamin acts as the most important lipid-soluble antioxidant and prevents reproduction and dissemination of harmful free radicals in biological system (13,14). Destructive effect of free radicals occurs in the lack of vitamin E and leads to peripheral nerve damage (13). Also, various studies have proved the role of vitamin E against induced genetic damages by genotoxic substances such as Cisplatin, beryllium, acrylamide (15,16)

Since antioxidant effects of vitamin E in inhibition of induced damages by electromagnetic waves with low frequencies have not been studied so far, the study was performed to investigate the protective effect of vitamin E against electromagnetic field with frequency of 50 Hz and intensity of 50 Gauss on mouse bone marrow cells (in vivo) using micronucleus assay.

## Materials and Methods

### Animals:

Protecting effect of vitamin E against electromagnetic waves were studied on polychromatic erythrocytes of bone marrow of adult Balb/C male mice, with the weight of about 30-25 grams (from Mashhad Razi Serum Institute) as experimental model.

The mice were kept in natural conditions with about 60 – 70% humidity,  $23 \pm 1$  °C temperature and photoperiod of 12 h light, 12 h darkness in polycarbonate cages. Food and water were also supplied **Ad libitum**.

### Experimental procedure:

Forty mice were randomly divided into 4 groups, including:

Control: mice were kept in normal conditions

Test 1: mice injected (ip) with vitamin E with dose of 200 mg / kg **bw** for 8 consecutive days

Test 2: mice in this group were exposed to electromagnetic wave using the device made at Azad Islamic University, Mashhad branch, with intensity of 50 Gauss for 4 days, 4 hours daily

Experimental group: vitamin E was injected with dose of 200 mg / **kg bw** for 8 consecutive days and at **5<sup>th</sup> day of injection, they were** exposed to electromagnetic waves with intensity of 50 Gauss for 4 days, 4 hours daily.

### Slide preparation:

All mice were anaesthetized by chloroform and were dissected. The bone marrow of two legs was removed by injection of 0.5 ml of fetal bovine serum (GIBCO) and was collected in test tube. Tube containing the suspension was centrifuged (Kokusan, Japan) for 10 minutes at 900 rpm. The cell

palette resuspended in 0.5 ml of FBS. Four to five drops of cell suspension were placed on slide. Air-dried slides were fixed by absolute **Alcohol** for 5 minutes. Staining was performed according by Schmid using Migranvald and Gimsa (Merck, Germany)

### **Scoring:**

In a good prepared slide, polychromatic erythrocytes are clearly observed as pink; these cells have no nucleus. In these anucleated cells, micronucleus was visible as a small purple core in the cytoplasm. In each slide, the number of micronucleus was counted in 1000 polychromatic erythrocytes (x1000).

polychromatic erythrocytes lack a nucleus, which implies that the presence of micronuclei in their normally anucleated cytoplasm signifies encapsulated DNA fragments, tell-tale indication of chromosome damage

### **Statistical Analysis**

Data analysis was performed in SPSS and the statistical significance was determined using one- or two-way analysis of variance and Tukey's multiple comparison test. In all the analysis a 95% confidence level ( $P < 0.05$ ) was considered statistically significant. All the data were presented as mean  $\pm$  standard deviation (SD).

The ethics of animal handling and treatment were considered throughout the experiment.

## **Results**

The mean frequency of MnPCE in all groups is presented in image1. Injection of vitamin E significantly **decrease** the MnPCE frequency compared to control.

However, the result revealed that the frequency of MnPCE was significantly higher in electromagnetic waves treated mice compared to control ( $P < 0.001$ ). Treatment of the mice with vitamin E four days before exposure to electromagnetic waves did reduce the frequency of MnPCE compared to electromagnetic waves treated mice ( $P < 0.01$ ) and brought it to the control level.




Figure1: The mean frequency of MnPCE in all groups related Control group. ( \*\*P<0.01, \*\*\*P<0.001)

## Discussion

In this study the genotoxic effect of extremely low electromagnetic fields was investigated on mice using micronucleus assay. Production of micronucleus is an index of the chromosomal damages. This test is the most common genotoxic test that shows chromosomal damages, including the loss or breakage of chromosomes. Micronucleus formation is the result of chromosomal damages (17,18,19). These damages in form of chromosome loss or breaks will produce micronuclei during cell division.

This test was first performed by Schmid at 1973 in the mice' bone marrow erythrocytes (20,21). This assay was proved to be sensitive and quickly performed technique to study chromosomal damages induced by electromagnetic field(22).

Extremely low electromagnetic field can induce chromosomal abnormalities which could be detectable by micronucleus assay. Increase in the frequency of micronucleus in lymphocytes of the workers exposed to it is an indication for its clastogenic effect (23)

Data of the present study revealed the genotoxic effect of electromagnetic field in bone marrow polychromatic cells. Exposure of the pregnant rats to electromagnetic field led to tissue and DNA damages in testis of male newborn rats(24). Also mice exposed to electromagnetic fields represented tissue damages in their kidneys and testis caused by oxidative stress(25).

The result of this study is in accordance with the findings of several studies showing relation between exposure to electromagnetic field and induction of DNA damages, such as chromosomal instability, increase of chromosomal breaks, sister chromatid exchange and increase in the frequency of micronucleus in mice or rats(26,27). Rats exposed to long treatment with 900 MHz electromagnetic field exhibited high frequency of micronucleus in Polychromatic erythrocytes(22).

There are a variety of experiments about the cause of damage by electromagnetic waves. Since the energy of electromagnetic waves isn't high enough to cause direct damage and cracking, it seems that fields affect indirectly. Some researchers believe that electromagnetic and radio waves affect by increasing the ROS reactive oxygen. Simko has proposed that these waves increase the free radicals level (28)

The clastogenic property of magnetic field is due to its ability of inducing oxidative stress in cells and reducing the total antioxidant status and glutathione value in the cells(25). Hence antioxidant treatment of the exposed cells to electromagnetic fields might reduce the clastogenic damages in those cells. In this experiment we showed that treatment of the mice

with vitamin E before exposure to extremely low magnetic field can reduce the frequency of micronucleus in polychromatic erythrocytes of bone marrow.

It was observed the waves increased cracking in DNA when rats' brain cells were exposed on the microwave waves 2/45 GHz for 35 days (26). Mice of bulb C with age of 7 weeks were exposed to electric fields 3 Tesla for 24 hours and intensity of 4/7 Tesla for 24, 48 and 72 hours. The results showed that frequency of micronucleus is dependent to the radiation time and intensity of field (27).

Ruedger has also expressed that the effects of RF-EMF genotoxic can be through the production of free radicals and reaction with DNA improvement mechanism (28). Zmysolonyi has been reported the formation of free radicals under the effect of electromagnetic waves is the main reason of destructive effects of these waves (29). So we can conclude that free radicals are the reason of production micronucleus. Eldamerdash with the study of mobile phone waves has shown that waves cause changes in antioxidant activity of rats' blood and brain (30).

Awad also stated in his study that ROS has an important role in tissues exposed to waves emitted from mobile, and leads to damage to antioxidant system (31). Erguder has reported the oxidative stress is the reason of damage to the samples exposed to waves emitted from the computer (32). Claycombe and his colleagues have stated that oxygen free radicals as a Mutagen cause many cancers and create defects in the immune system (13).

Evaluation of the effect of electromagnetic waves and peritoneal injection of vitamin E showed that mean number of micronucleus is significantly decreased. Vitamins as antioxidant are necessary for health. The effect of vitamin E on antioxidant system of liver, kidneys and muscle of rabbit is positive (33).

Vitamin E is a fat-soluble vitamin and plays a role in lipoprotein metabolism and biological membranes and can induce or prevent of gene or protein expression (10). The results of this 173research match with the reports of Ouanes, Robichova, Singha and Gonzalez that all have mentioned anti genotoxicity role of vitamin E. Ouanes stated that vitamin E reduces the induced micronucleus by Zearalenone (34,35). Robichova has also reported that vitamin E decreases the

damaging effects of N-nitrosomorpholin in human hepatic cells (36). Singha has improved the effects of genotoxic atrazine on rats' blood cells and liver with the effect of vitamin E (37). Dicamba is a genotoxic substance (SCE) which causes DNA damage with the production of free radicals. Gonzalez has stated that vitamin E improves the adverse effects of this substance (38).

It has been observed that antioxidants reduce the percentage of chromosomal elimination in chromosomal patients. Dusinska reported that antioxidants decrease the chromosomal damages (39). Using 300 mg / kg bw Vitamin E for 21 consecutive days decreases the number of induced micronucleus by Acrylamide in rats' bone marrow (40). Micro elements have interventional role on gene's function due to their role in vital path and different reactions, and their lack impact on gene function (41). Consumption of vitamins leads to the stability of genome and reduction of Micronuclues (42). Erguder has also suggested the consumption of vitamins and antioxidants for reducing the effects of computer (32).

Claycombe also expressed that vitamin E as an antioxidant decreases chromosomal damages by preventing the formation of free radicals. In fact, antioxidants converted free radicals into harmless compounds that easily leave the body before they damage to any part of body. Even antioxidants may improve previous damages (9). Results of this research do not match with Hsieh's experiments that reported the destructive effects of vitamin E. The reason of contradictory results can be the used dosage. He observed that vitamin with dose 500 mg/kg has genotoxicity role, and antioxidants act as Peru-oxidation in high concentrations and create Autommune' damages (43). He has also stated that Vitamin E is a cancer promoter in high concentrations. The amount of damages by free radicals or under effect of environmental factors and presence of other antioxidants can influence the effectiveness of this vitamin (44).

The study performed by Demsia reported that long-term radiation for 30 days, 3 hours per day increases micronucleus in polychromatic erythrocytes of rat's bone marrow three times in comparison with control group (45). Ruiz-Gomez's tests have shown that electromagnetic waves could act as a co-inductor of DNA damage (46).

This result is contradictory with some experiences. The experience of Vijayalaxmi has suggested that electromagnetic field with 42 GHz for 30 minutes does not significantly increase the number



of micronucleus in polychromatic erythrocytes of mice with race of BALB/c (47). Comparison of the effect of radio waves 1800 MHz on the number of micronucleus in rat cells exposed to field for 5 days and 20 minutes daily with control group showed no significant difference between two groups (48).

It seems that different results of various studies depends on changes in study conditions, especially frequency change, intensity and radiation duration, type of animals and type of magnetic field (constant or intermittent).

The findings of this study showed that vitamin E will reduce genotoxic effects by electromagnetic field with low-frequency of 50 Hz in polychromatic erythrocytes of mice bone marrow. It seems that the use of food sources of vitamin E is very important and significant for the persons living in the environment of electromagnetic waves.

## **References**

1. Hossman KA, Herman DM. Effects of electromagnetic radiation of mobile phones on central nervous system. *Bioelectromagnetic*. 2003; 24(1):49-62.
2. Crumpton MJ, Collins AR. Are environmental electromagnetic fields genotoxic?. *DNA Repair*. 2004; 3(10):1385–1397.
3. Carbonari K, Goncalves L, Roth D, et al. Increased micronucleated cell frequency related to exposure to radiation emitted by computer cathode ray tube video display monitors. *Genet Mol Biol*. 2005; 28(3):469-474.
4. Moulder JE, Foster KR, Erdreich LS, et al. Mobile phones , Mobile phone base stations and cancer. *Int J Radiat Biol*. 2005; 81(3):189-203.
5. Mashevich M, Folkman D, Kesar A, et al. Exposure of human peripheral blood lymphocytes to electromagnetic fields associated with cellular phones leads to chromosomal instability. 2003; 24(2):82-90.

6. Speit G, Schutz P, Hoffmann H. Genotoxic effects of exposure to radiofrequency electromagnetic fields (RF-EMF) in cultured mammalian cells are not independently reproducible. *Mutation Research*. 2007; 626(1-2): 42–47.
7. Milena Villarini , Maria Vittoria Ambrosini , Massimo Moretti , Luca Dominici , Elena Taha , Danilo Piobbico , Cristiana Gambelunghe & Giuseppina Mariucci .Brain hsp70 expression and DNA damage in mice exposed to extremely low frequency magnetic fields: A dose-response study. *International Journal of Radiation Biology*. 2013; 89(7): 562–570
8. Luzhna L, Kathiria P, Kovalchuk O. Micronuclei in genotoxicity assessment: from genetics to epigenetics and beyond. *Front Genet*. 2013; 4:1-17.
9. Balamuralikrishnan B, Vellingiri B, Suresh K, Nattan S, Varsha P, Subramaniam M . Evaluation of Chromosomal Alteration in Electrical Workers Occupationally Exposed to Low Frequency of Electro Magnetic Field (EMFs) in Coimbatore Population, India. *Asian Pacific J Cancer Prev*. 2012;13(6), 2961-2966.
- 10-Burlaka A, Tsybulin O, Sidorik E, Lukin S, Polishuk V, Tsehmistrenko S, Yakymenko I. Overproduction of free radical species in embryonal cells exposed to low intensity radiofrequency radiation. *Exp Oncol*. 2013 :35(3):219-225.
12. Dietrich M, Jacques PF, Pencina M, et al. Vitamin E supplement use and the incidence of cardiovascular disease and all-cause mortality in the Framingham Heart Study: Does the underlying health status play a role?. *Atherosclerosis*. 2009; 205(2): 549–553.
13. Claycombe KJ, Mydany SN. Vitamin E and genome stability. *Mutat Res*. 2001; 475(1-2): 37–44.
14. Nakamura YK, Omaye ST. Vitamin E-modulated gene expression associated with ROS generation. *J Funct Foods*. 2009; 1(3):241-252.
15. Nersesyan A, Muradyan R. Sea- Buckthorn Juice protects mice against genotoxic action of Cisplatin. *Experimental Oncology*. 2004; 26(2): 153-165.

16. Fahmy MA, Hassan NH, Farghaly AA, et al. Studies on the genotoxic effect of beryllium chloride and the possible protective role of selenium/vitamins A, C and E. *Mutation Research*. 2008; 652(2):103–111.
17. Baharara J, Haddad F, Ashraf AR, et al. The Effect of extremely low frequency electromagnetic field (50Hz) on induction chromosomal damage on bone marrow erythrocytes of male Balb/C mouse . *Journal of Arak University of Medical Sciences* 2008;11 (2):17-26.
18. Haddad F, Moghimi A, Salmani A, et al. Analysing the Radioprotective effect of *Cotoneaster Nummularia* in Mouse Bone Marrow Cells using Micronucleus Assay. *Cell and Molecular Biology Research*. 2009; 1(2):77-83.
19. Verschaeve L. Genetic damage in subjects exposed to radiofrequency radiation. *Mutat Res*. 2009; 681(2-3):259-270.
20. Schmid W. The micronucleus test. *Mutat Res*. 1975; 31:9-16.
21. Fenech M. Chromosomal biomarkers of genomic instability relevant to cancer. *Drug Disco Today*. 2002; 7(22):1128-1137.
- 22 Zülal Atlı Şekeroğlu, Ayşegül Akar & Vedat Şekeroğlu. Evaluation of the cytogenotoxic damage in immature and mature rats exposed to 900 MHz radiofrequency electromagnetic fields. *International Journal of Radiation Biology*. 2013; 89(11):985-992
- 23- Akyıl D, Özkara A, Feyza Erdoğan S, Eren Y, Konuk M. Micronucleus assay in human lymphocytes after exposure to alloxidim sodium herbicide in vitro. *Cytotechnology*. 2015 ;67(6): 1059–1066.
- 24 Hancı H, Odacı E, Kaya H, Aliyazıcıoğlu Y, Turan İ, Demir S, Çolakoğlu. The effect of prenatal exposure to 900-megahertz electromagnetic field on the newborn rat testicles. *Reprod Toxicol*. 2013; 42:203-209.
- 25 Özorak A, Nazıroğlu M Çelik O, Yüksel M, Özçelik D, Okan M. Wi-Fi (2.45 GHz)- and Mobile Phone (900 and 1800 MHz)-Induced Risks on Oxidative Stress and Elements in Kidney and Testis of

- Rats During Pregnancy and the Development of Offspring. *Biological Trace Element Research*. 2013; 156(1-3):221-229.
26. Paulraj R, Behari J. Single strand DNA breaks in rat brain cells exposed to microwave radiation. *Mutat Res*. 2006; 596(1-2):76-80.
27. Suzuki Y, Ikahata M, Nakamura K, et al. Induction of micronuclei in mice exposed to static magnetic fields. *Mutagenesis*. 2001; 16(6):499-501.
28. Ruedger HW. Genotoxic effect of radiofrequency electromagnetic fields. *Pathophysiology*. 2009; 16(2-3):82-100.
29. Zmyslony M, Jahre JM. The role of free radicals in mechanisms of biological function exposed to weak constant and net magnetic fields. *Med Pr*. 1998; 49(2):177- 186.
30. El-Demerdash M, El Sheikh E, Gharib AF. Hazardous effects of electromagnetic radiation emitted by mobile phones on the brain and cochlea of Albino Rats: role of melatonin and vitamin C. *JASMR*. 2009; 4(1): 89-100.
31. Awad SM, Hassan NS. Health Risks of Electromagnetic Radiation from Mobile Phone on Brain of Rats. *Journal of Applied Sciences Research*. 2008; 4(12): 1994-2000.
32. Erguder IM, Durak I. Effects of computer use on human Salivary Oxidant/Antioxidant Status. *Journal of Biological Sciences*. 2006; 6(1): 14-17.
33. Lo Fiego DP, Santoro P, Macchioni P, et al. The effect of dietary supplementation of vitamins C and E on the  $\alpha$ -tocopherol content of muscles, liver and kidney, on the stability of lipids, and on certain meat quality parameters The effect of dietary supplementation of vitamins C and E on the  $\alpha$ -tocopherol content of muscles, liver and kidney, on the stability of lipids, and on certain meat quality parameters. *Meat Science*. 2004; 67(2):319–327.
34. Ouanesa Z, Ayed-Boussemaa I, Baatia T. Zearalenone induces chromosome aberrations in mouse bone marrow preventive effect of 17 $\beta$ -estradiol, progesterone and Vitamin E. *Mutat Res*. 2005; 565(2):139–149.
35. Ouanes Z, Abid S, Ayed I. Induction of micronuclei by Zearalenone in Vero monkey kidney cells and in bone marrow cells of mice: protective effect of Vitamin E. *Mutat Res*. 2003; 538(1-2): 63–70.

36. Robichova S, Slamenova D, Chalupa I. DNA lesions and cytogenetic changes induced by N-nitrosomorpholine in HepG2, V79 and VH10 cells: the protective effects of Vitamins A, C and E. *Mutat Res.* 2004; 560(2): 91–99.
37. Singh M, Kaur P, Sandhir R, et al. Protective effects of vitamin E against atrazine-induced genotoxicity in rats. *Mutat Res.* 2008; 654(2):145-159.
38. Gonzalez NV, Soloneski S, Larramendy ML. Dicamba –induced genotoxicity in Chinese hamster ovary (CHO) cells is prevented by vitamin E. *J Hazard Mater.* 2009; 163(1):337-343.
39. Dusinska M, Kazimírova A, Barancokova M, et al. Nutritional supplementation with antioxidants decreases chromosomal damage in humans. . 2013; 18(4):371-386.
40. Amany A, El- Tohamy AA, Boyomy AA. Anti –genotoxic effect of vitamin E administration in acrylamide rat bone marrow cells. *Arab J Biotech.* 2008; 11(1):29-38.
41. Fenech M. Nutrition and genome health. *Forum Nutr.* 2007; 60:49-65.
42. Fenech M, Baghurst P, Luderer W, et al. Low intake of calcium, folate, nicotinic acid, vitamin E, retinol, beta-carotene and high intake of pantothenic acid, biotin and riboflavin are significantly associated with increased genome instability--results from a dietary intake and micronucleus index survey in South Australia. *Carcinogenesis.* 2005; 26(5):991-999.
43. Hsieh C, Lin FB. Opposite effects of low and high dose supplementation of vitamin E on survival of MRL/lpr mice. *Nutrition.* 2005; 21(9):940–948.
44. Mitchel RE, Mccann RA. Skin tumor promotion by Vitamin E in mice: amplification by ionizing radiation and Vitamin C. *Cancer Detect Prev.* 2003; 27(2): 102–108.
45. Demisia G, Vlastos D, Matthopoulos DP. Effect of 910-MHz electromagnetic field on rat bone marrow. *Scientific World Journal.* 2004; 4 (2):48-54.
46. Ruiz- Gomez MJ, Martinez – Morillo M. Electromagnetic fields and the induction of DNA strand breaks. *Electromagn Biol Med.* 2009; 28 (2):201-214.
47. Vijayalaxm I, Logani MK, Bhanushali A, et al. Micronuclei in peripheral blood and bone marrow cells of mice exposed to 42 GHz electromagnetic millimeter waves. *Radiat Res.* 2004; 161 (3):341-345.

48. Gurbuz N, Sirav B, Yuvaci HU, et al. Is there any possible genotoxic effect in exfoliated bladder cells of rat under the exposure of 1800 MHz GSM-like modulated radio frequency radiation (RFR)?. *Electromagn Biol Med.* 2010; 29(3):98-104.