

# Trend analysis of various ocular surgeries performed at University of Port Harcourt Teaching Hospital (UPTH), Nigeria over a ten-year period

## Abstract

**Aim:** To identify the pattern of ocular surgeries performed at UPTH and determine any significant changes in trends of these surgeries over a ten-year period - 2005 to 2016.

**Methods:** It was a hospital-based retrospective study over a ten-year period. Data were extracted from the ocular surgery register and inputted into an excel-spread sheet. Data analyzed with United States Centers for Disease Control and Prevention (CDC) Epi-Info version 7 software to determine significant differences in trends of the various eye surgeries. A p-value  $< 0.05$  was considered statistically significant.

**Results:** A total of 1,039 major and 1,322 minor ocular surgeries were done in a ten-year period. Most of the major cases (198) were done in 2012 while the least number of the major cases (27) were done in 2016. Significant variations in trends within this period occurred with cataract extraction, corneal repair, evisceration/enucleation and trabeculectomy ( $p < 0.0001$  respectively). Most of the minor cases (271) were done in 2008 while the least number (83) was done in 2011 with significant trend occurring with pterygium excision ( $p < 0.001$ ) and conjunctival mass excision ( $p < 0.009$ ). The most common major ocular surgery was cataract extraction (744) while the most common minor surgery was pterygium excision (597).

**Conclusion:** Ocular surgical load in UPTH is comparable to other parts of the country. There was a significant variation in the trend of cataract surgery, corneal repair, evisceration/enucleation, trabeculectomy, pterygium excision and conjunctival mass excision. These may have been affected by incessant breaks in services as a result of both local and national strike actions by health workers. In addition, patients' ignorance or refusal to accept some of the procedures offered may have also contributed to the trend seen. Enlightenment and regular service delivery are keys to improving ocular surgical uptake.

**Keywords;** Trends analysis, Ocular Surgeries, Tertiary Hospital, Nigeria

## Introduction

Some of the commonest reasons for most ocular surgeries are either to restore or improve vision, for cosmetic reasons, or to reduce morbidity and mortality from


malignant lesions. Patients of all ages may present with ocular conditions that might require surgical intervention. In our environment, the most common ocular surgery is cataract – cataract is the commonest cause of visual impairment and blindness and affects more than 5% of the population in spite of the available remedies.<sup>1,2</sup> Other commonly performed ocular surgeries include corneo-scleral laceration repair<sup>3</sup> and evisceration/enucleation.<sup>4,5,6</sup> Trabeculectomy which is aimed at preventing blindness or reducing the rate of deterioration of vision in glaucoma usually presents with some challenges because patients with apparently good vision refuse surgeries while some surgeons prefer not to operate on those with end stage disease for fear of wipe out syndrome.<sup>8</sup>

Cases of lid lacerations are usually under reported because majority of them are minor and are either managed at home, chemists or in peripheral clinics and fail to reach the tertiary centres.<sup>7</sup> Other ocular procedures routinely done are chalazion incision and curettage, pterygium excision, surgeries for cancers,<sup>9,10,11</sup> foreign body removal, punctal dilation, epilation<sup>12</sup> and squint surgeries. These are aimed at restoring good cosmetic appearance. The study set out to report any changing trends in the pattern of eye surgeries done at our Facility – a Tertiary Health Institution.

## Methods

This study was a hospital-based retrospective study involving retrieval of records of all eye surgeries performed at University of Port Harcourt Teaching Hospital (UPTH) over a ten-year period (data for 2013 were not available because they were missing since they were not electronically stored). The various ocular surgeries performed in UPTH are routinely entered into an Ophthalmology register. Data were extracted from the ocular surgery register and inputted into an excel-spread sheet. Double check was employed to avoid errors in data entry. Data on major eye surgical procedures spanning eleven years were retrieved from the hospital records. Data analysis was performed using United States Centers for Disease Control and Prevention (CDC) Epi-Info version 7 software. Data were expressed using line graph and Chi square for trend performed to determine significant differences in trends of the various eye surgeries. A p-value < 0.05 was considered statistically significant.

## 67 Results



68 *No data available for 2013*

69 Fig. 1. Line graph shows trends in Major Ocular Surgeries in UPTH (2006 – 2016)

70 Cataract surgeries accounted for most cases done as shown by the line graph. It  
 71 accounted for about 55% to over 70% of surgeries done between 2006 and 2012;  
 72 then dropped to less than 55% between 2014 and 2016. The trend for other  
 73 common eye surgeries - corneal repair, scleral repair, trabeculectomy, eyelid repair,  
 74 enucleation/evisceration and exenteration were similar over the years. This trend  
 75 over the years shows statistical significance. ( $P < 0.05$ )

76

77

78

79

80

81

82

83 Table 1. Frequency Distribution of Major Ocular Surgeries in UPTH (2006-2016)

Year\*\*

Major Ocular Surgeries	2006	2007	2008	2009	2010	2011	2012	2014	2015	2016	Chi Square $\ddagger$	p-value
Cataract extraction + PCIOL	83 (70.3%)	60 (65.9%)	119 (75.3%)	83 (73.4%)	78 (72.2%)	78 (67.2%)	164 (82.8%)	29 (65.9%)	36 (54.5%)	14 (19.7%)	39.95	<0.0001 *
Corneal Repair	18 (15.2%)	14 (15.4%)	26 (16.4%)	14 (12.4%)	15 (13.9%)	18 (15.5%)	14 (7.1%)	5 (11.4%)	14 (21.2%)	6 (8.4%)	85.64	<0.0001 *
Sclera Repair	5 (4.2%)	7 (7.7%)	10 (6.4%)	7 (6.2%)	5 (4.6%)	1 (0.9%)	1 (0.5%)	0 (0.0%)	3 (4.5%)	1 (1.4%)	0.24	0.6234
Evisceration/ Enucleation	7 (5.9%)	7 (7.7%)	0 (0.0%)	7 (6.2%)	7 (6.5)	8 (6.9%)	13 (6.6%)	1 (2.3%)	6 (9.1%)	4 (5.6%)	14.56	<0.0001 *
Exenteration	0 (0.0%)	1 (1.1%)	2 (1.3%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (1.4%)	0.81	0.3675
Lid Repair	2 (1.7%)	2 (2.2%)	0 (0.0%)	1 (0.9%)	1 (0.9%)	8 (6.9%)	3 (1.5%)	0 (0.0%)	6 (9.1%)	1 (1.4%)	0.00	0.9754
Trabeculectomy	3 (2.5%)	0 (0.0%)	1 (0.6%)	1 (0.9%)	2 (1.8%)	3 (2.6%)	3 (1.5%)	9 (20.4%)	1 (1.5%)	0 (0.0%)	24.17	<0.0001 *
<b>Total</b>	<b>118</b> <b>(100%)</b>	<b>91</b> <b>(100%)</b>	<b>158</b> <b>(100%)</b>	<b>113</b> <b>(100%)</b>	<b>108</b> <b>(100%)</b>	<b>116</b> <b>(100%)</b>	<b>198</b> <b>(100%)</b>	<b>44</b> <b>(100%)</b>	<b>66</b> <b>(100%)</b>	<b>71</b> <b>(100%)</b>		

84  $\ddagger$ Chi square for trend      \*statistically significant      \*\*No data available for 2013

85 The highest number of eye surgeries (n=198; 82.8%) was done in 2012 while the  
86 least (n=71; 19.7%) was in 2016. The trends analyses all show statistical  
87 significance except for scleral repair, exenteration and lid repair. (p = 0.6234, p =  
88 0.3675, p = 0.9754 respectively) The peak periods for the procedures were 2008 for  
89 corneal and scleral repairs, 2011 for lid repair, 2012 for cataract surgeries, and  
90 2014 for evisceration and enucleation; and trabeculectomy. In 2014, the least  
91 number of corneal repairs (5) and evisceration/enucleation (1) were done; there  
92 were no scleral repairs, exenteration and lid repair surgeries.




Fig. 2. Line graph shows trends in Minor Ocular Surgeries in UPTH (2005 - 2012)

In Figure 2, pterygium excision was the most common minor ocular surgical procedure done over the years. This was followed by foreign body removal and chalazion incision and curettage. The year 2009 recorded the highest number of chalazion incision and curettage, while for pterygium it was 2010. The least common procedure done was punctal dilation. This change in trend over time was statistically significant ( $p < 0.05$ ).

109 Table. 2. Frequency Distribution of Minor Ocular Surgeries in UPTH (2005-2012)

Minor Ocular Surgeries	Year								Chi Square <sup>†</sup>	p-value
	2005	2006	2007	2008	2009	2010	2011	2012		
Pterygium Excision	45 (44.1%)	95 (39.7)	122 (42.9%)	102 (37.6%)	40 (40.4%)	86 (60.9%)	36 (43.4%)	71 (56.3%)	9.83	0.0017*
Chalazion I & C	30 (29.4%)	35 (14.6%)	48 (18.4%)	65 (23.9%)	15 (15.2%)	21 (14.9%)	20 (24.1%)	25 (19.8%)	0.09	0.7627
Epilation	7 (6.9%)	10 (4.2%)	17 (6.5%)	35 (12.9%)	6 (6.1%)	4 (2.8%)	2 (2.4%)	6 (4.8%)	1.31	0.2527
FB Removal	15 (14.7%)	77 (32.2%)	53 (20.3%)	47 (17.3%)	35 (35.4%)	27 (19.1%)	20 (24.1%)	20 (15.9%)	1.95	0.1628
Conjunctival Mass Excision	5 (4.9%)	22 (9.2%)	19 (7.3%)	18 (6.6%)	3 (3.0%)	2 (1.4%)	5 (6.0%)	4 (3.2%)	6.78	0.0092*
Punctal Dilation	0 (0.0%)	0 (0.0%)	2 (0.8%)	4 (1.5%)	0 (0.0%)	1 (0.7%)	0 (0.0%)	0 (0.0%)	0.03	0.8576
<b>Total</b>	<b>102</b> <b>(100%)</b>	<b>239</b> <b>(100%)</b>	<b>261</b> <b>(100%)</b>	<b>271</b> <b>(100%)</b>	<b>99</b> <b>(100%)</b>	<b>141</b> <b>(100%)</b>	<b>83</b> <b>(100%)</b>	<b>126</b> <b>(100%)</b>		

110 <sup>†</sup>Chi square for trend      \*Statistically significant

111 Most of the minor surgeries (771) were done between 2006 and 2008, with the  
 112 highest number (271) being in 2008. The least number of eye surgeries (n=83)  
 113 were done in 2011. There were more foreign body removals (77) and conjunctival  
 114 mass excision (22) in 2006 compared to other years; and more pterygium excision  
 115 (122) and chalazion incision and curettage in 2007 compared to 2008 with more  
 116 epilation (35) and punctal dilation (4). The least number of pterygium excisions  
 117 (36), chalazion I&C (20) and epilation (2) were recorded in 2011. The trends for  
 118 pterygium excision and conjunctival mass excision were statistically significant with  
 119 p< 0.0017 and 0.0092 respectively.

120

121

122

## 123 Discussion

124 A total of 1,039 major and 1,322 minor surgeries were done within this ten-year  
125 period. Most of the major cases were done in 2012. In 2016, only 27 major surgeries  
126 were done. Most of the minor cases were done in 2008 while the least number were  
127 done in 2011. These fluctuating trends may be due to the constant interruption of  
128 services as a result of both local and National industrial strike actions embarked  
129 upon by staff of the Federal Ministry of Health. Over seventy percent of the patients  
130 who had major surgeries had cataract extraction with significant differences in the  
131 numbers done each year ( $p < 0.001$ ).

132 In many developing countries, cataract surgical services are not always accessed  
133 and there are many reasons which include but not limited to lack of awareness that  
134 cataract is treatable and cost implications. In our setting with a number of cataract  
135 camps organized yearly, there is availability of cataract surgical services but the  
136 trend seen may not be unrelated to the almost constant disruption of clinical  
137 services for all kinds of reasons including those already stated. One hundred and  
138 forty four cases of corneal repairs were done this period. Most of the corneal repairs  
139 were done in 2008 – this is not surprising because that was at the height of the  
140 militancy that plagued the Niger Delta Region.

141 Few cases of scleral repair (40) were done in the period under review with majority  
142 occurring between 2007 and 2009. Ignorance could be responsible for this low  
143 number as some patients in our environment prefer to self-medicate or patronize  
144 quacks. Except for 2012 with 13 cases, all other years under review recorded less  
145 than 10 cases of evisceration and enucleation; only four cases of exenteration were  
146 recorded in this ten-year period. Some traditional beliefs discourage the removal of  
147 the eyes for any reason and this may be responsible for the small number recorded.  
148 This is in contrast to the studies in Enugu and Ile Ife.<sup>4,5,6</sup> Only 24 cases of eyelid  
149 repair were done in this period. Minor lid lacerations are either managed at home,  
150 chemists or in peripheral clinics and do not reach the Tertiary Eye Centres. This  
151 may explain the low volume compared to other studies.<sup>7</sup>

152 The total number of trabeculectomies was 23 representing 2.1% of the total major  
153 surgeries. There is a general low uptake of trabeculectomy surgical services in our  
154 centre but this improved significantly in 2014 (9 trabeculectomies) as a result of  
155 additional training in glaucoma for some of our Comprehensive Ophthalmologists.  
156 Most glaucoma patients shy away from trabeculectomy until very late in the disease  
157 process when the glaucoma is either very advanced or end-stage. The reason may be  
158 because patients with apparently good vision refuse surgeries while some surgeons  
159 prefer not to operate on those with end stage disease for fear of wipeout

phenomenon. Our result compares well with that of Kyari et al. who observed that less than half of patients offered glaucoma surgery underwent the procedure and all those on consecutive anti-glaucoma medical therapy refused surgery<sup>8</sup>

It was observed that pterygium excision was the commonest minor surgical procedure that was performed. This is similar to other studies in Nigeria.<sup>9,10,11</sup> In their study, Odugbo et al. noted that within a 7-year study period pterygium excision accounted for over 55% of conjunctival surgeries.<sup>4</sup> Pterygium is a common ocular surface disease in tropical countries including Nigeria. In our study, most pterygium excisions were performed using the bare-sclera technique with adjunct intraoperative application of 5-fluorouracil or mitomycin-C to the sclera bed or primary conjunctival autologous grafting similar to other studies in Nigeria.<sup>9-11</sup>

Conjunctival mass excision peaked in 2006 and progressively declined afterwards. This may be attributed to the increasing use of HAART in the management of HIV/AIDS patients leading to a decrease in the number of ocular surface squamous neoplasia. Studies elsewhere collaborates our finding<sup>10</sup>.

The uptake of chalazion incision and curettage surgical services had a similar undulating trend over the ten-year period; however, this was not statistically significant ( $p=0.7627$ ). Epilation constituted 0.66% of all minor ocular procedures performed over this period. More cases were done in Northern Nigeria compared to our Centre and this may be due to the high prevalence of Trachoma in Northern Nigeria.<sup>12</sup> Seventy-eight cases of conjunctival masses were excised over this period.

## Conclusion

The ocular surgical load in our Centre is comparable with other tertiary hospitals in Nigeria. There is a significant variation in the trend of cataract surgery, corneal repair, evisceration/enucleation, trabeculectomy, pterygium excision and conjunctival mass excision. These may have been affected by the several breaks in services and patient's ignorance or poor uptake of some of the surgical procedures. A lot need to be done in the areas of public enlightenment and provision of uninterrupted eye care service as a way of improving uptake of ocular surgeries.

We were limited by the small data size and non-availability of 2013 data.



194 **References:**

- 195 1. Visual impairment and blindness fact sheet N"282" August 2014.  
196 Archived from the original on 12/5/2015.
- 197 2. Priority eye diseases: Cataract. Prevention of blindness and visual  
198 impairment. World Health Organization. Archived from the original on  
199 2015-05-24.
- 200 3. Ortega I, Ryan AJ, Deshpandea, MacNeila S, Claeysensa F. Combined  
201 microfabrication and electrospinning to produce 3-D architectures for  
202 corneal repair Author links open overlay panel Acta Biomaterialia 2013;  
203 9(3): 5511-20
- 204 4. Timothy NH, Freilich DE, Linberg JV. Evisceration versus enucleation  
205 from the ocularist perspective. Ophthal Plast Reconstr Surg 2003; 19:  
206 417-20.
- 207 5. Eze BI, Maduka-Okafor FC, Okoye OI, Okoye O. Surgical indications for eye  
208 removal in Enugu, South Eastern Nigeria Nigerian Journal of  
209 Ophthalmology 2007; 15(2): 44-48
- 210 6. Awe OO, Adeoye AO, Onakpoya OH. Surgical eye removal in Ile-Ife,  
211 Nigeria: Nigerian Journal of Ophthalmology 2016; 24 (1): 31-34
- 212 7. Nitin Vichare. Management of Lid Lacerations Ocular Trauma DOS Times -  
213 Vol. 20, No. 8 February, 2015
- 214 8. Kyari F, Nolan W, Gilbert C. Ophthalmologists' practice patterns and  
215 challenges in achieving optimal management for glaucoma in Nigeria:  
216 results from a Nationwide Survey. BMJ Open. 2016; 6(10): e012230.  
217 Published 2016 Oct 11. doi:10.1136/bmjopen-2016-012230
- 218 9. Odugbo OP, Wade PD, Obikili GA, Alada JJ, Mpyet CD. Pattern and  
219 outcome of conjunctival surgeries in Jos University Teaching Hospital, Jos,  
220 Nigeria Nigerian Journal of Ophthalmology 2016; 24: 25-30
- 221 10. Akinsola FB, Mbadugha CA, Onakoya AO, Adefule Ositelu AO, Aribaba OT,  
222 et al. Pattern of conjunctival masses seen at Guinness Eye Centre LUTH  
223 Idi Araba. Nig Q J Hosp Med 2012; 22: 39-43.

- 224 11. Fadamiro CO, Ajite KO, Ajayi AI, Omotoye OJ. Pterygium excision with  
225 conjunctival autograft: Our technique. Niger J Ophthalmol 2012; 20: 38-  
226 41
- 227 12. Burr SE, Sillah A, Sanou AS, Wadagni AC, Hart J, Harding-Esch EM, et al.  
228 Cross-Sectional surveys of the prevalence of follicular trachoma and  
229 trichiasis in The Gambia: Has Elimination Been Reached? Published:  
230 September 19, 2016<https://doi.org/10.1371/journal.pntd.0004906>

231