

MARKER-ASSISTED SELECTION IN FISH: A REVIEW

ABSTRACT

The important economical traits like body growth, resistance to diseases, meat quality, etc. highly influence the profitability of food animals including fishes. The main target of every selective breeding programme is to produce improved traits offspring's for these traits. However, improvement of performance traits through traditional phenotype-based selection needs several generations to optimise these characters. Marker-Assisted Selection (MAS) is a type of indirect method of selection of better performing breeding individuals. MAS is beneficial when the traits are difficult, expensive to measure and has both low heritability and recessive traits. MAS facilitates the exploitation of existing genetic diversity in breeding populations and can be used to improve desirable traits in livestock. MAS depends on identifying the link between a genetic marker and Quantitative Traits Loci (QTL). The distance between marker and target traits determines the association of the marker with the QTL. After identifying the markers linked to QTL, they can be used in the selective breeding programme to select the brooders having better genetic potential for the targeted trait. Improvement of performance traits through MAS is fast and more accurate and allows us to understand the genetic mechanism affecting performance traits.

Keywords: Marker-Assisted Selection, Quantitative traits loci, genetic diversity, trait

1.0 INTRODUCTION

Marker-Assisted Selection (MAS) is a type of biotechnology that uses molecular genetic markers as a criterion for selecting desired traits (Ashraf, 2012). Marker Assisted Selection (MAS) is an indirect selection process where a trait of interest is selected not based on the trait itself but on a marker linked to it (Ribaut and Ragot 2007).

Formatted: Font color: Red

34 MAS is considered a “revolutionary” approach to traditional tree breeding as it allows
35 breeders to select individuals based on their genotypes, rather than being restricted to
36 phenotypic characteristics (Boopathi *et al.*, 2013).

37 Sax (1923) was the first to show how genetic factors influencing quantitative traits can be
38 identified using markers.

39 Recently MAS became a very popular method of indirect selection for production of
40 the genetically improved offspring's in aquaculture breeding programme offspring's. Most
41 of the performance traits including growth or disease resistance are controlled by multiple
42 genes and are therefore inherited as quantitative traits, analysis of their associated
43 quantitative trait loci (QTL) is an essential part of aquaculture genomics (Liu and Cordes,
44 2004). QTLs are largely unknown genes that affect performance traits (such as growth rate
45 and disease resistance) and these are important to breeders.

46 MAS in a breeding context involves scoring indirectly for the presence or absence of
47 a desired phenotype or phenotypic component based on the sequences or banding patterns of
48 molecular markers located in or near the genes controlling the phenotype. The sequence
49 polymorphism or banding pattern of the molecular marker is indicative of the presence or
50 absence of a specific gene or chromosomal segment that is known to carry a desired allele
51 (Brumlop and Finckh, 2011).

52 Marker-assisted selection method (MAS) or genome-wide marker-assisted selection
53 method (G-MAS) was not widely used in aquaculture, but nowadays its use is increasing due
54 to its ease of use and quicker than traditional phenotype-based selection. Now it becomes a
55 fertile field of research for the aquaculture researchers to discover novel genetic marker that
56 can be used to link with the QTLs in selective breeding programmes (Hauser *et al.*, 2011;
57 Dichmont *et al.*, 2012; Abdul-Muneer, 2014).

58 In order to manage individual species effectively, identification of different species
59 from a mixed catch becomes important. DNA markers are widely being accepted not only to
60 obtain information about gene flow and allele frequencies in aquaculture practices but also to
61 identify hybrids. The majority of the markers, which are used in inter- and intra-specific
62 disparity, include Random Amplified Polymorphic DNA (RAPD) for species and sub-species
63 identification done in tilapia (Bardakci and Skibinski, 1994), and iso-enzyme used in
64 intraspecific variations in Sparidae species (Alarcón and Alvarez, 1999). Similarly, Nijman *et*
65 *al.*, (2003) reported the use of mtDNA markers as an important tool in rapid detection of
66 hybridization between species and subspecies of livestock.

Formatted: Font color: Red

67 Markers tend not to have any biological effect, but rather can be thought of as notable and
68 constant points of reference within the genome (Guimaraes, *et al.*, 2007). Markers can be
69 found within the desired gene or, more commonly, linked to a gene determining a trait of
70 interest (Brumlop and Finckh, 2011; Guimaraes *et al.*, 2007). Unlike genetic engineering,
71 MAS does not alter the original DNA (Vogel and Van Aken, 2009); instead it uses genetic
72 marker to identify naturally-occurring genetic variations among individuals, with the intent of
73 selecting those with the best potential to meet desired criteria and objectives.

74 Marker Assisted Selection (MAS) provides several other benefits to breeders, in that
75 it can select for genes that demonstrate low heritability, have recessive alleles, and are
76 difficult, expensive, or time exhaustive to determine phenotypically (Boopathi, 2013a;
77 Brumlop and Finckh, 2011; Xu and Crouch, 2008). MAS also allows for gene pyramiding or
78 combining multiple genes within the same breeding line, while having fewer unintentional
79 losses and fewer selection cycles (Boopathi, 2013a; Xu and Crouch, 2008).

80 Furthermore, MAS may be viewed by the public with more support than genetic engineering
81 as breeders are not manually manipulating the genes, and thus all offspring inheritance occurs
82 naturally (Vogel and Van Aken, 2009). It is also believed that genetic markers may be
83 important in the assessment, conservation and use of diversity in germplasm and varieties
84 (Brumlop and Finckh, 2011).

85 Molecular marker maps have been constructed for a number of aquaculture species,
86 e.g. tilapia, *Clarias*, giant tiger prawn, kuruma prawn, Japanese flounder and Atlantic salmon,
87 although their density is generally low (Nichols *et al.*, 2003). As many preferred traits are not
88 observed until maturity, MAS eliminates this waiting period by allowing for the early
89 selection of desired genotypes at the seedling stage (Yanchuk *et al.*, 2002).

90 The desirable phenotypic variations in the performance traits of fishes are used to
91 increase the aquacultural yield, improve incomes of farmers and enhances food security
92 through selective breeding by choosing better-performed individuals. However, phenotype-
93 based selection needed considerable time to optimise the traits, so researchers are now
94 moving from phenotype based selection to genotype-based selection. The absence of a
95 molecular marker is the main limiting factor for the realization of genotype based selection
96 potentials in fishes. However, with the advent of DNA-based genetic markers in the late
97 1970s and now the ease of the marker discovery through the next generation sequencing
98 allowed researchers to identify large numbers of markers spreads throughout the genome of
99 any species of interest. The markers are used to detect linkage with the traits of interest, thus

100 allowing MAS finally to become a reality (Peterson *et al.*, 1990). This paper aims to provide
101 information regarding the technical aspect of MAS and the current application in fisheries
102 and Aquaculture in other to increase high quality production within a period of time.

103 **2.0 Marker Assisted Selection**

104 Incorporation of marker information into breeding programs in aiding identification
105 and selection of superior individuals has been widely studied (Bernardo, 1994; Han *et al.*,
106 1997; Xie and Xu, 1998; Romagosa *et al.*, 1999; Ayoub *et al.*, 2003; Jordan *et al.*, 2003).

107 Molecular markers in aquaculture and fisheries have been used for over 50 years
108 (Ryman and Utter, 1987; Liu and Cordes, 2004) and their use has steadily increased over the
109 last two decades (Park and Moran, 1994; Chauhan and Rajiv, 2010; Dichmont *et al.*, 2012;
110 Abdul-Muneer, 2014).

111 An important factor in MAS is the accuracy of estimating the genetic effects related to
112 the trait of interest. In contrast to genetic engineering (GE), MAS does not alter the original
113 DNA. Rather, it identifies whether the desired trait(s) are being expressed, so that individuals
114 with the best potential can be selected (Andersson, 2001).

115 Molecular marker analysis allows the identification of genome segments, so called
116 Quantitative Trait Loci (QTL), contributing to the genetic variance of a quantitative trait and
117 thus to select superior genotypes as these loci (Cannai *et al.*, 2003). Allelic variation in
118 genetic markers can be linked to the variation in traits of economic interest, and thus the
119 marker provides DNA level information on the inheritance of the traits.

120 The practical use of markers in selection can be roughly divided into three classes:

- 121 1) Removing genetic disorders,
- 122 2) Marker breeding value-selection, and
- 123 3) Genomic selection.

124 **2.4 MAS versus Phenotypic Selection**

125 Marker-Assisted Selection (MAS) will probably never replace Phenotypic Selection
126 (PS) entirely. There is no general pattern by which it can be predicted whether MAS or PS
127 will be more useful. Empirical comparisons of MAS and PS for increasing gain from
128 selection have been made in several studies. The outcomes of these studies are conflicting. In
129 some studiesstudies, MAS is reported to be more effective/efficient than PS (Yousef and
130 Juvik 2001; Abalo *et al.*, 2009) while other studies considered the two methods equal (Van
131 Berloo and Stam 1999; Willcox *et al.*, 2002; Hoeck *et al.*, 2003; Moreau *et al.*, 2004). In a
132 third group of studies PS proved to be more effective/efficient than MAS (Davies *et al.*,

Comment [JCM1]: And 2.1, 2.2 and
2.3??????????????

Formatted: Font color: Auto, Highlight

Formatted: Highlight

Formatted: Font color: Red

133 2006; Wilde *et al.*, 2007) and in other comparisons the effectiveness/efficiency of MAS and
134 PS varied within the same study, depending on the populations or on the trait selected for
135 (e.g. FlintGarcia *et al.*, 2003b; Robbins and Staub 2009).

136 2.5 Limitations of MAS

- 137 • Cost
- 138 • Requirement of technical skill
- 139 • Automated techniques for maximum benefit

140 2.6 Advantages of MAS

141 In addition to the cost and time savings described above, for a number of breeding
142 scenarios, MAS methods are likely to offer significant advantages compared with
143 conventional selection methods. These scenarios assume the availability of markers for
144 multiple traits and take into consideration the advantages of MAS under optimum situations
145 (Dreher *et al.*, 2002; Dudley, 1993).

- 146 1. Gene stacking for a single trait: MAS offers potential savings compared with
147 conventional selection when it allows breeders to identify the presence of multiple
148 genes/alleles related to a single trait, and the alleles do not exert individually
149 detectable effects on the expression of the trait.
- 150 2. Early detection: MAS offers potential savings compared with conventional selection
151 when it allows alleles for desirable traits to be detected early, well before the trait is
152 expressed and can be detected phenotypically. This benefit can be particularly
153 important in species that grow slowly.
- 154 3. Heritability of traits: Up to a point, gains from MAS increase with decreasing
155 heritability. However, due to the difficulties encountered in QTL detection, the gains
156 are likely to decline beyond a certain threshold heritability estimate.

157 2.7 Disadvantages of MAS

158 Perhaps the greatest disadvantage of MAS is the time and financial **investment**
159 **required**
160 **to develop** markers that are widely applicable for traits of agronomic importance.
161 Often a marker developed in one or a few related genotypes will not work for
162 other genotypes in a breeding scheme due to allelic effects. Furthermore, development of
163 markers, particularly for QTL, is complicated by epistatic interactions and the critical need
164 for good quality phenotypic data.

165 2.8 Quantitative Trait Loci

Formatted: Font color: Red

166 In fish, several QTL studies have been published; in salmonids (Jackson *et al.*, 1998;
167 Johansen 1999; Robinson *et al.*, 1999; Sakomoto *et al.*, 1999; Marfyniuk 2001, Ozaki *et al.*,
168 2001 Somorger 2001. Tao and Baiding 2003), in catfish (Liu *et al.*, 2003), in tilapia (Cnaani
169 *et al.*, 2003) and in silver barb (Hussain *et al.*, 2002).

170 Marker Assisted Selection (MAS) is followed by two steps, detection of molecular markers
171 associated with quantitative trait locus (QTL) and application of those markers.

172 The position of the chromosome that controls the economical important trait is termed as
173 QTL.

174 The concepts for detecting QTL were developed more than 90 years ago (Sax, 1923). In
175 aquaculture species, much effort has been applied for QTL mapping. QTLs are mapped by
176 linkage disequilibrium with molecular markers exhibiting Mendelian segregation.
177 Economically important traits are controlled by the single or group of gene.

178 The basic concept of QTL studies is to know the number and location of loci
179 associated with phenotypic traits (Mackay, 2001; Mauric io, 2001; Burt and Hocking, 2002;
180 Erickson *et al.*, 2004). Thus, candidate gene or molecular markers, resulted by QTL mapping,
181 could be used in MAS (Groenen *et al.*, 2000). QTL detection is an ongoing effort in
182 aquaculture species. More than 37 important traits have been located in about 20 aquaculture
183 species.

184 QTL mapping is the practical application of marker-assisted selection in aquaculture
185 (Al-Samarai, 2015). With rapid advancement of molecular technology, it is now possible to
186 use molecular marker information to map major quantitative trait loci (QTLs) on
187 chromosomes (Paterson *et al.*, 1988, 1991; Hilbert *et al.*, 1991; Jacob *et al.*, 1991; Stuber *et*
188 *al.*, 1992). Once QTL for a trait are identified, individuals can be selected for breeding on the
189 basis of marker alleles that segregate with favourable phenotypes (Lande and Thompson,
190 1990). This strategy, known as marker-assisted selection (MAS), is particularly useful for
191 traits that cannot be measured on selection candidates directly, notably disease resistance or
192 meat quality traits (Sonesson, 2007a).

193 2.8.1 QTL Detection in Fish

194 A number of genetic maps have been developed specifically to locate QTL in several
195 fish species. The first of such map was produced in Zebrafish insert scientific name (Postleth
196 wairt *et al.*, 1994; Shimoda *et al.*, 1999), which is a non-aquacultural species. Among
197 cultivable fish groups low-density maps have been developed for salmonids (Sakamoto *et al.*,
198 2000; Ghabi 2001) for catfish (Liu *et al.*, 2003; Poompson and Na-Nakorn 2004) for tilapia

Formatted: Font: Not Italic, Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

199 (Kocher *et al.*, 1998; Cnaani *et al.*, 2003), for Japanese flounder (Sanchez *et al.*, 2003), for
 200 red sea beam (Sakamoto *et al.*, 2003), for Oyster (Yu and Geso 2003), and for shrimp
 201 ([Http://shrimppmap.tag.csiro.au](http://shrimppmap.tag.csiro.au)).

2.8.3 QTL Mapping in Fish

Comment [JCM2]: And 2.8.2????????????????????
Formatted: Highlight

202 Although in fish several studies have confirmed the existence of significant genetic
 203 variation for quantitative traits at commercial importance (Kause *et al.*, 2003) and have
 204 recognized the potential of MAS for their genetic improvement (Flint and Mott, 2001). Thus
 205 far, very few QTL for production traits have been identified in fish (Sonesson, 2003). Much
 206 effort is devoted to QTL mapping for growth, feed conversion efficiencies, disease resistance,
 207 fecundity, and spawning time (Dunham *et al.*, 2001).

208 Several QTL studies have been published in rainbow trout for temperature tolerance
 209 (Jackson *et al.*, 1998). Danzmann *et al.*, 1998, [Pperry 2001](#)), spawning time (Sakamoto *et al.*,
 210 1999; fish back et al 2000, O' Malley 2001); growth (Martynick 2001), disease resistance
 211 (Ozaki *et al.*, 2001), and fitness traits (Somorjai 2001). Other notable QTL studies published
 212 [in aquaculturalaquaculture fish](#) species include: in tilapia for temperature and salinity
 213 tolerance (Streadman and Kocher 2002; Cnaan *et al.*, 2003), in catfish for feed conversion
 214 efficiency and bacterial [septicemiasepticaemia resistance](#) (Liu 2003), in guppy for growth
 215 (Nakajima and Taniguchi 2002), in at [fautic salmon](#) for infectious [anemiaanaemia resistance](#)
 216 (Moen *et al.*, 2003 [and in Arctictia charrCharr](#) for growth rates and fitness traits (Johansen
 217 1999, Somorjai 2001).

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Comment [JCM3]: Please check this word

Formatted: Highlight

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

218 In salmonids, QTL have been found related to body weight and size (Martyniuk *et al.*,
 219 2003; O'Malley *et al.*, 2003; Reid *et al.*, 2005), for colouration pattern (Streelman, Albertson
 220 and Kocher, 2003) and for one form of albinism (Nakamura *et al.*, 2001). Zimmerman *et al.*,
 221 (2005) found QTL for pyloric caeca number, a trait related to feed conversion efficiency.

222 **Table 1: QTL studies in selected aquaculture species**

Species	Traits	Reference
Arctic charr	Body weight and sexual maturation; Salinity tolerance	Küttner <i>et al.</i> , 2011
Asian seabass	Resistance against viral nervous necrosis disease	▲Wang <i>et al.</i> , 2006▲
	Growth-related trait	▲Xia <i>et al.</i> , 2014▲
	Omega-3 fatty acids	

Formatted: Font: 11 pt, Spanish (Mexico)

Formatted: Spanish (Mexico)

Formatted: Font: 11 pt, Spanish (Mexico)

Formatted: Spanish (Mexico)

Atlantic salmon	Growth traits and flesh colour	Baranski <i>et al.</i> , 2010;	Formatted: Font: 11 pt, Font color: Red
	Resistance against IPN	Tsai <i>et al.</i> , 2014;	Formatted: Font: 11 pt, Spanish (Mexico)
	Late sexual maturation	Moen <i>et al.</i> , 2009 ;	Formatted: Spanish (Mexico)
		Houston <i>et al.</i> , 2008 ; 2010	Formatted: Font: 11 pt, Font color: Red, Spanish (Mexico)
Catfish	Columnaris disease resistance	Gutierrez <i>et al.</i> , 2014	Formatted: Font color: Red, Spanish (Mexico)
	ESC disease resistance	Geng <i>et al.</i> , 2015	Formatted: Font: 11 pt, Font color: Red, Spanish (Mexico)
	Hypoxia tolerance	Wang <i>et al.</i> , 2013; Zhou <i>et al.</i> , 2017	Formatted: Font: 11 pt, Font color: Red
	Heat stress	Wang <i>et al.</i> , 2016;	Formatted: Font color: Red
	Head size	Wang <i>et al.</i> , 2016;	Formatted: Font: 11 pt, Spanish (Mexico)
		Jin <i>et al.</i> , 2016	Formatted: Spanish (Mexico)
		Geng <i>et al.</i> , 2016	Formatted: Font: 11 pt, Spanish (Mexico)
Common carp	Common carp	Zhang <i>et al.</i> , 2011	Formatted: Spanish (Mexico)
	Morphometric traits	Boulton <i>et al.</i> , 2011	Formatted: Spanish (Mexico)
	Swimming ability	Laghari <i>et al.</i> , 2014	Formatted: Spanish (Mexico)
Eastern oyster	Disease resistance	Yu and Guo, 2006	Formatted: Spanish (Mexico)
European seabass	Growth, body weight	Louro <i>et al.</i> , 2016;	Formatted: Font: 11 pt, Spanish (Mexico)
	Morphometric traits and stress	Massault <i>et al.</i> , 2010	Formatted: Spanish (Mexico)
	Response	Massault <i>et al.</i> , 2010	Formatted: Font: 11 pt, Spanish (Mexico)
Pacific white shrimp	Growth parameters	Andriantahina <i>et al.</i> , 2013	Formatted: Spanish (Mexico)
Giant tiger prawn	Disease resistance and sex determination	Robinson <i>et al.</i> , 2014	Formatted: Spanish (Mexico)
Japanese flounder	<i>Vibrio anguillarum</i> resistance	Wang <i>et al.</i> , 2014	Formatted: Font: 11 pt, Italic, Font color: Red
Pacific oyster	Growth	Guo <i>et al.</i> , 2012	Formatted: Font: 11 pt, Font color: Red
	Resistance against summer mortality	Sauvage <i>et al.</i> , 2010	Formatted: Font: 11 pt, Spanish (Mexico)
	Viability	Sauvage <i>et al.</i> , 2010	Formatted: Spanish (Mexico)
		Plough and Hedgecock, 2011	Formatted: Spanish (Mexico)
		Plough <i>et al.</i> , 2016	Formatted: Font: 11 pt, Spanish (Mexico)
Gilthead seabream	Skeletal deformities	Negrín-Báez <i>et al.</i> , 2015	Formatted: Spanish (Mexico)
	Sex determination and body growth	Negrín-Báez <i>et al.</i> , 2015	Formatted: Font: 11 pt, Spanish (Mexico)
		Loukovitis <i>et al.</i> , 2011	Formatted: Spanish (Mexico)
		Massault <i>et al.</i> , 2011	Formatted: Font: 11 pt, Spanish (Mexico)
		Massault <i>et al.</i> , 2011	Formatted: Spanish (Mexico)

Rainbow trout	Growth related traits	Kocmarek <i>et al.</i> , 2015; Wringe <i>et al.</i> , 2010; Leder <i>et al.</i> , 2006.
Tilapia	Growth traits Sex	Liu <i>et al.</i> , 2014; Wang <i>et al.</i> , 2015 Palaiokostas <i>et al.</i> , 2015

224

225 **2.8.4 QTL analysis**

226 **1. QTL for growth traits**

227 Growth is one of the most important economic traits of all aquaculture species. Up to
 228 2012, QTL analyses have been conducted in more than 20 aquatic species (Yue *et al.*, 2014),
 229 and growth was the most popular trait studied. Wang *et al.*, (2006) used 380
 230 F1 Asian seabass to identify five major QTLs and 27 potential QTLs. Of them, three major
 231 QTLs for body weight, length, and body length were located at a similar linkage group 2
 232 (LG2) position with the nearby Lca287 microsatellite and accounted for 28.8%, 58.9%, and
 233 59.7% of the phenotypic variations. The other two major QTLs for body weight were located at another
 234 LG2 position. These five major QTLs have been confirmed in two other Asian
 235 seabass populations (Wang *et al.*, 2008). Further QTL fine mapping of the Asian seabass
 236 growth trait identified three candidate candidates “growth genes” (CATHEPSIN D,
 237 KCTD15, and CSMD2) affecting body weight, body length, and total length (Wang *et al.*,
 238 2011). The function of the *cathepsin D* gene in humans involves cell proliferation and cell
 239 growth; therefore, *cathepsin D* may also be a major “growth gene” in Asian seabass.
 240 O’Malley *et al.*, (O’Maller *et al.*, 2008) identified QTLs for body weight in rainbow trout on
 241 10 different LGs.

242
 243 Wringe *et al.*, (2010) used additional backcrossed families and SSR markers to
 244 confirm the O’Malley *et al.*,’s results and found several major candidate growth genes (e.g.,
 245 *GH2* and *Pax7*). Reid *et al.*, 2004 identified a QTL for body weight in two LGs (AS8 and 11)
 246 of Atlantic salmon, and reported that it was homologous to the growth QTL in rainbow trout.
 247 Houston *et al.*, (2009) identified QTLs for body weight in Linkage group 1 (LG1) and LG5 of
 248 Atlantic salmon. Gutierrez *et al.*, (2012) further used a 6.5 K Single Nucleotide
 249 Polymorphisms (SNP) chip to identify QTLs in six LGs at the genomic level. Cnaani *et al.*,
 250 (2004) identified a QTL for tilapia growth on LG23, which is the linkage group with the
 251 genetic sex-determining region. Song *et al.* (2012) used 1487 SSRs to produce a high-density

Comment [JCM4]: This must be 2.8.4.1?????? Or 3.0 ??????????

Formatted: Highlight

Formatted: Font color: Red

Comment [JCM5]: Year?????????

Formatted: Highlight

252 genetic linkage map and successfully identified a QTL affecting body weight in LG14 of
253 Japanese flounder.

254 Some reports have used a candidate gene approach to identify growth-related genes
255 and molecular markers in fish. Tao and Boulding (2003) found polymorphisms in the growth
256 hormone gene (*GH*) that were significantly associated with growth rate of Arctic charr
257 (*Salvelinus alpinus*). Li *et al.*, (2009) reported an SNP in the insulin-like growth factor-
258 (IGF)1
259 gene 5' untranslated region (UTR) of largemouth bass (*Micropterus salmoides*). Sun *et al.*,
260 (2012) reported that two SNPs in exon 3 of the myostatin (*MSTN*) gene were significantly
261 related to body weight and Fulton's factor in common carp. Liu *et al.* (2012) also found that a
262 SNP in the *MSTN* 3' UTR was very significantly associated with total length, body length,
263 and body weight of bighead carp.

264 **2. QTL for feed conversion rate**

265 Food Conversion Ratio (FCR) is one of the most important economic traits in fish, as
266 fish with a better FCR increase profits.

267 Liu (2005) used Amplified Fragment Length polymorphisms (AFLP) markers to
268 construct a catfish genetic map and found a QTL associated with FCR. Zimmerman *et al.*,
269 (2005) revealed three QTLs for the number of pyloric caeca in three LGs of rainbow trout,
270 and this is an important index associated with FCR.

271 Food Conversion Ratio (FCR) studies have also been reported in common carp from
272 the Heilongjiang Fisheries Research Institute of the Chinese Academy of Fishery Sciences
273 (Wang, 2012).

274 **3. QTL for sex determination**

275 Sex phenotype and sex determination in fish have specific evolutionary status and
276 diversity. Males and females of some species have significant differences in growth rate or
277 commercial value; therefore, mono sex fish culture is a promising strategy. The sex-
278 determining (SD) loci and QTLs have been studied in a limited number of fish, such as tilapia
279 (Lee *et al.*, 2004) rainbow trout (Alfaqih *et al.*, 2009) and salmonids (Davidson *et al.*, 2009).
280 Previous studies have demonstrated that sex QTLs are located on LG1, 2, 3, 6, and 23 of
281 tilapia (Cnaani *et al.*, 2004; Lee *et al.*, 2004; Cnaani *et al.*, 2008) Eshel *et al.* (2011) reported
282 a major candidate sex QTL that is considered the sex determining region in tilapia. Fifty-one
283 genes in this region have been annotated, and 10 have been confirmed.

Formatted: Font color: Red

284 The anti-Müllerian hormone gene is the most differentially expressed gene in male
285 and female tilapia. Sun *et al.*, (2014) recently published several sex-specific markers, and one
286 is tightly linked with the sex-determining region discovered by Eshel *et al.*, (2011) The sex-
287 determining locus in rainbow trout is located on the LG of RT10, and this locus also
288 significantly affects thermo-resistance and body length. The sex-determining regions in Artic
289 charr (Moghadam *et al.*, 2007) brown trout (Gharbi *et al.*, 2006) and Atlantic salmon (Gilbey
290 *et al.*, 2004) are located on the LGs of AC4, BT28, and AS1, respectively.

291 Woram *et al.*, (2003) compared LGs of sex-determining loci in four salmonids and
292 found that although the nucleotide sequences flanking the sex-determining loci were well-
293 conserved, the SD LGs were diverse, suggesting that the regions underwent different
294 recombination events.

295 Loukovitis *et al.*, (2011) located growth and sex-determining QTLs in gilthead sea
296 bream and showed that these two traits have similar genetic control in LG21. Martínez *et al.*,
297 (2009) located a sex QTL on LG5 of turbot and proposed a ZZ/ZW sex-determining
298 mechanism. Viñas *et al.*, (2012) also found a major sex QTL on turbot LG5. These findings
299 suggest that the sex-determining genes may occur on turbot LG5. Song *et al.*, (2012) used
300 high-density genetic maps to locate seven sex QTLs on the half-smooth tongue sole LG1f,
301 LG14f, and LG1m.

302 Additional study by Chen *et al.*, (2014) provided insight into ZW sex chromosome
303 evolution and identified sex-determining genes, such as *dmrt1* and *neurl3*.

304 **2.8.5 Factors affecting QTL analyses**

305 The power of mapping QTL can be influenced by a number of factors, such as genetic
306 properties of QTL, experimental design, environmental effects, marker density and
307 informativeness, genotyping errors and precision of trait measurement. Details about how
308 these factors influence the power of QTL mapping can be found in some very good reviews
309 (Crosses 2001; Flint and Mott 2001; Doerge 2002).

310 **2.8.6 Methods of Detecting QTL**

311 Basically, three methods are frequently used for mapping QTL and estimating their
312 effects, namely Single-Marker Association Analysis (SMAA), Simple Interval Mapping
313 (SIM) and Composite Interval Mapping (CIM) (Crosses 2001; Flint and Mott 2001; Doerge
314 2002).

315 **2.9 Current Status of Applications of MAS in Fish**

316 Molecular marker maps have been constructed for a number of aquaculture species,
317 e.g. tilapia, catfish, giant tiger prawn, kuruma_ prawn, Japanese flounder and Atlantic salmon,
318 although their density is generally low. Density is high for the rainbow trout, where the map
319 published in 2003 has over 1 300 markers spread throughout the genome – the vast majority
320 are AFLPs but it also includes over 200 microsatellite markers (Nichols *et al.*, 2003). Some
321 QTLs of interest have been detected (e.g. for cold and salinity tolerance in tilapia and for
322 specific diseases in rainbow trout and salmon).

323 In a recent review of MAS in fish breeding schemes, Sonesson (2003) suggested that
324 MAS would be especially valuable for traits that are impossible to record on the candidates
325 for selection such as disease resistance, fillet quality, feed efficiency and sexual maturation,
326 and concluded that MAS is not used in fish breeding schemes today and that the lack of dense
327 molecular maps is the limiting factor. Marker Assisted Selection (MAS) has become a
328 valuable tool in selecting organisms for desirable traits. MAS is expected to increase genetic
329 gain compared to traditional breeding programs and reduce the cost of progeny testing by
330 early selection. The application of MAS in breeding programmes depends on the knowledge
331 of breeders about variable marker information.

332 **REFERENCES** please check all bibliography to make it all uniform with respect

333 Journal

- 334 Alarcon, J.A., Alvarez, M. C. (1999). Genetic identification Sparidae species by isozyme
335 markers. Applications to interspecific hybrids. *Aquaculture*, 173, 95-103.
- 336 Anderson, J. L., Mari, A. R., Braasch, I., Amores, A., Hohenlohe, P., Batzel, P. (2001).
337 Multiple sex-associated regions and a putative sex chromosome in zebrafish
338 revealed by RAPD mapping and population genomics. 3, 427-437.
- 339 Ashraf, M., Akram, N. A., Foolad, M. R. (Eds.). (2012). Marker-Assisted Selection in Plant
340 Breeding for Salinity Tolerance. 913, 305–333. doi:10.1007/978-1-61779-986-0.
- 341 Bardakci, F., Skibinski, D. O. F. (1994). Application of the RAPD technique in tilapia fish
342 species and subspecies identification. 73, 117–123. doi:10.1038/hdy.1994.110.
- 343 Bernardo, R. (1994). Prediction of maize single-cross performance using RFLPs and
344 Biotechnology in Agriculture and Food
345 (<http://tandfonline.com/doi/book/10.1081/EEBAF>).
- 346 Boopathi, N. M. (2013a). Marker-Assisted Selection. In Genetic Mapping and Marker
347 Assisted Selection: Basics, Practice and Benefits. 173–186. doi:10.1007/978-81-
348 322-0958-4

Formatted: English (United States)

Formatted: Font color: Red

Formatted: Spanish (Mexico)

349 Brumlop, S., Finckh, M. R. (2011). Applications and potentials of marker assisted selection (
350 MAS) in plant breeding. Pp. 178. http://www.bfn.de/0502_skripten.html
351 Chen, J., Wang, Y., Yue, Y., Xia, X., Du, Q., Chang, Z. (2014). A novel male-specific DNA
352 sequence in the common carp, *Cyprinus carpio*. *Mol Cell Probes*. 23, 235–9.
353 Cnaani, A., Zilberman, N., Tinman, S., Hulata, G., Ron, M. (2004). Genome-scan analysis for
354 quantitative trait loci in an F-2 tilapia hybrid. *Mol Genet Genomics*. 272(2):162-
355 172.
356 Danzmann, R. G., Gharbi, K. (1994). Gene mapping in fishes: a means to an end. *Genetica*.
357 111(1-3):3-23.
358 Davidson, W. S., Koop, B. F., Jones, S. J. M., Iturra, P., Vidal, R., Mas, A., Jonassen, I.,
359 Lien, S., Omholt, S. W. (2009). Sequencing the genome of the Atlantic salmon
360 (*Salmo salar*). *Genome Biol* 11:403
361 Davies, J., Berzonsky, W., LEACH, G. (2006): A Comparison of Marker-Assisted and
362 Phenotypic Selection for High Grain Protein Content in Spring Wheat. 152, 117-
363 134.
364 Dekkers, J. C. M., Hospital, F. (2002). The use of molecular genetics in the improvement of
365 agricultural populations. *Nature Revs. Genet*. 3: 22–32.
366 Doyle, J. J., Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of
367 fresh leaf tissue. *Phytochem. Bull*. 19: 11–15.
368 Dreher, K., Khairallah, M., Jean-Marcel, R., Monis, M. (2002). Money matters (I): costs of
369 field and laboratory procedures associated with conventional and marker assisted
370 maize breeding at CIMMYT. *Molecular Breeding I I*: 221-234.
371 Dudley, J. W. (1993). Molecular markers in plant improvement: manipulation of genes
372 affecting quantitative traits. *Crop Sci*. 33: 660–668.
373 Edward, M. D., Stuber, C. W., Wendel, J. F. (1987a). Molecular markers facilitated
374 investigation of quantitative trait loci in maize. I. Numbers, genomic distribution
375 and types of gene action. *Genetics* 115: 113-125
376 Erickson, D. L., Fenster, C. B., Stenoien, H. K., Price, D. (2004). Quantitative trait locus
377 analyses and the study of evolutionary process. *Molecular Ecology* 13: 2505-2522.
378 Eshel, O, Shirak, A., Weller, J. I., Slossman, T., Hulata, G., Cnaani, A. (2011). Fine mapping
379 of a locus on linkage group 23 for sex determination in Nile tilapia (*Oreochromis*
380 *niloticus*). *Anim Genet*. 42: 222–4.

Formatted: Spanish (Mexico)

381 Ferguson, M. (1994). The role of molecular genetic markers in the management of cultured
382 fishes. *Reviews in Fish Biology and Fisheries*, 4(3): 351–373.
383 <https://doi.org/10.1007/BF00042909>

384 Flint-Garcia, S.A., Darrah, L. L., McMullen, M. D., Hibbard, B. E. (2003). Phenotypic versus
385 genomic selection in an established commercial layer breeding program. *Genetics*
386 *Selection Evolution* 45: 29.

387 Gharbi, K., Gautier, A., Danzmann, R.G., Gharbi, S., Sakamoto, T., Hoyheim, B., Taggart, J.
388 B., Cairney, M., Powell, R., Krieg, F., Okamoto, N., Ferguson, M. M., Holm, L. E.,
389 Guyomard, R. (2006). A linkage map for brown trout (*Salmo trutta*): chromosome
390 homeologies and comparative genome organization with other salmonid fish.
391 *Genetics* 172:2405–2419

392 Gilbey, J., Verspoor, E., McLay, A., Houlihan, D. (2004). A microsatellite linkage map for
393 Atlantic salmon (*Salmo salar*). *Anim Genet* 35:98–105

394 Gimelfarb, A., Lande, R. (1994). Marker-assisted selection and marker QTL association in
395 breeding programs. *Genetics*. 132: 1199-1210.

396 Gjedrem, T. (2009). Genetic improvement of cold-water fish species. *Aquac Res.* 31:25–33.

397 Groen, A. F., Crooijmans, R. P. M. A., Van Kampen, A. J. A., Van der Beek, S, Van der
398 Poel, J. J., Groenen, M. A. M. (2000). Microsatellite polymorphism in commercial
399 broiler and layer lines. *Proc 5th World Congr Genet Appl Livestock Prod* 21:95-98.

400 Gross, M., Schneider, J., Moav, N., Alvarez, C., Myster, S., Liu, Z., Hallerman, E., Hackett,
401 P., Guise, K., Faras, A., Kapuscinski, A. (1995). Molecular analysis and growth
402 evaluation of transgenic northern pike. *Aquaculture* 103: 253-273.

403 Guo, X., Hershberger, W.K., Cooper, K., Chew, K.K., 2012. Artificial gynogenesis with
404 Hayes, B. J., Chamberlain, A. J., Mcpartlan, H., Macleod, I., Sethuraman, L., Goddard, M. E,
405 (2007). Accuracy of marker-assisted selection with single markers and marker
406 haplotypes in cattle. *Genetics Research* 89: 215-220.

407 Houston, R. D., Bishop, S. C., Hamilton, A., Guy, D. R., Tinch, A. E., Taggart, J. B.,
408 Derayat, A., McAndrew, B. J., Haley, C. S. (2009). Detection of QTL affecting
409 harvest traits in a commercial Atlantic salmon population. *Anim Genet.* 40(5):753-
410 755.

411 Houston, R. D., Haley, C.S., Hamilton, A., Guy, D. R., Tinch, A. E., Taggart, J. B. (2009).
412 Major quantitative trait loci affect resistance to infectious pancreatic necrosis in
413 Atlantic salmon (*Salmo salar*). *Genetics*. 178:1109–15.

Formatted: Spanish (Mexico)

414 Hulata, G. (2003). Detection of a chromosomal region with two quantitative trait loci,
415 affecting cold tolerance and fish size, in an F-2 tilapia hybrid. 223(1-4):117-128.

416 Jackson, T.R., Ferguson, M.M., Danzmann, R.G., Fishback, A.G., Ihssen, P.E., O'Connell,
417 M., Crease, T.J. (1998). Identification of two QTL influencing upper temperature
418 tolerance in three rainbow trout (*Oncorhynchus mykiss*) half-sib families. Heredity
419 80: 143– 151.

420 Jacob, H. J., Lindpainter, K., Lincoln, S. E., Kusumi, R. K., Mao, Y. P., Ganten, D., Dzau, V.
421 J., Lander, E. S., (1991). Genetic mapping of a gene causing hypersensitive rat.
422 Cell. 67:213-224.

423 Jansen, R.C. (1993). Interval mapping of multiple quantitative trait loci. Genetics 135, 205-
424 211.

425 Jonasson, J., Stefansson, S. E., Gudnason, A., Steinarsson, A. (1999). Genetic variation for
426 survival and shell length of cultured red abalone (*Haliotis rufescens*) in Iceland.
427 *Journal of Shellfish Research* 18: 621-625.

428 Kause, A., Paananen, T., Ritola, O., Koskinen, H. (2003). Direct and indirect selection of
429 visceral lipid weight, fillet weight, and fillet percentage in a rainbow trout breeding
430 program. *Journal of Anim Sci.* 85: 3218-3227.

431 Kocher, T. D., Lee, W.J., Sobolewska, H., Penman, D., McAndrew, B. (1998). A genetic
432 linkage map of a cichlid fish, the tilapia (*Oreochromis niloticus*). *Genetics* 148:
433 1225–1232.

434 Lander, E. S., Botstein, D. (1989). Mapping Mendelian factors underlying quantitative traits
435 using RFLP linkage maps. *Genetics* 121: 185-1 99

436 Lee, M., Sharopova, N., Beavis, W. D., Grant, D., Katt, M., Blair, D. and Hallauer, A.
437 (2004). Expanding the genetic map of maize with the intermated B73 Mo17 (IBM)
438 population. *Plant Molecular Biology* 48: 453–461.

439 Li, J., Borojevich, K. A., Koop, B. F., Davidson, W. S. (2009). Comparative genomics
440 identifies candidate genes for Infectious Salmon Anemia (ISA) resistance in
441 Atlantic Salmon (*Salmo salar*). *Mar Biotechnol.* 13: 232-241.

442 Liu, F., Sun, F., Xia, J. H., Li, J., Fu, G. H., Lin, G. (2014). A genome scan revealed
443 significant associations of growth traits with a major QTL and GHR2 in tilapia. *Sci*
444 *Rep.* 4:7256

Formatted: Spanish (Mexico)

445 Liu, P., Li, J., He, Y. Y., Kong, J., Wang, Q. (2004). Present situation and protective
446 measures of genetic resources in *Fenneropenaeus chinensis*. *Mar. Fish Res* 25: 80–
447 85.

448 Liu, Z., Karsi, A., Li, P., Cao, D., Dunham, R. (2003). An AFLP-based genetic linkage map
449 of channel catfish (*Ictalurus punctatus*) constructed by using an interspecific hybrid
450 resource family. *Genetics* 165: 687–694.

451 Liu, Z.J., Li, P., Argue, B., Dunham, R., (1998a). Inheritance of RAPD markers in channel
452 catfish (*Ictalurus punctatus*), blue catfish (*I. furcatus*) and their F1, F2 and
453 backcross hybrids. *Anim. Genet.* 29: 58–62.

454 Liu, Z.J., Nichols, A., Li, P., Dunham, R., (1998b). Inheritance and usefulness of AFLP
455 markers in channel catfish (*Ictalurus punctatus*), blue catfish (*I. furcatus*) and their
456 F1, F2 and backcross hybrids. *Mol. Gen. Genet.* 258, 260–268.

457 Loukovitis, D., Sarropoulou, E., Tsigenopoulos, C. S., Batargias, C., Magoulas, A.,
458 Apostolidis, A. P., Chatziplis, D., Kotoulas, G. (2011). Quantitative trait loci
459 involved in sex determination and body growth in the gilthead sea bream (*Sparus*
460 *aurata* L.) through targeted genome scan. *PLoS ONE*, 6:e16599.

461 Luo, W., Zeng, C., Deng, W., Robinson, N., Wang, W., Gao, Z. (1997). Genetic parameter
462 estimates for growth-related traits of blunt snout bream (*Megalobrama*
463 *amblycephala*) using microsatellite-based pedigree. *Aquac Res.* 45:1881–8.

464 Mackay, T. F. C. (2001). The genetic architecture of quantitative traits. *Annu Rev Genet.*
465 35:303-339.

466

467 Martinez, P., Bouza, C., Hermida, M., Fernandez, J., Toro, M. A., Vera, M., Pardo, B.,
468 Millan, A., Fernandez, C., Vilas, R., (2009). Identification of the Major Sex-
469 Determining Region of Turbot (*Scophthalmus maximus*). *Genetics* 183(4):1443-
470 1452.

471 Martyniuk, C. J., Perry, G. M. L., Mogahadam, H. K., Ferguson, M. M., Danzmann, R. G.
472 (2003). The genetic architecture of correlations among growth related traits and
473 male age at maturation in rainbow trout. *Journal of Fish Biol* 63:746–764

474 Moen, T., Agresti, J. J., Cnaani, A., Moses, H., Famula, T. R., Hulata, G., Gall, G. A. E.,
475 May, B. (2004b). A genome scan of a four-way tilapia cross supports the existence
476 of a quantitative trait locus for cold tolerance on linkage group 23. *Aquaculture* 35:
477 893–904.

Formatted: Spanish (Mexico)

478 Moen, T., Fjalestad, K.T., Munck, H. and Gomez-Raya, L. 2003. A multistage testing
479 strategy for detection of quantitative trait loci affecting disease resistance in Atlantic
480 salmon. *Genetics* 167: 851–858.

481 Moen, T., Hoyheim, B., Munck, H., Gomez-Raya, L. (2004a). A linkage map of Atlantic
482 salmon (*Salmo salar*) reveals an uncommonly large difference in recombination rate
483 between the sexes. *Anim. Genet.* 35: 81–92.

484 Moghadam, H., Poissant, J., Fotherby, H., Haidle, L., Ferguson, M., Danzmann, R. (2007).
485 Quantitative trait loci for body weight, condition factor and age at sexual maturation
486 in Arctic charr (*Salvelinus alpinus*): comparative analysis with rainbow trout
487 (*Oncorhynchus mykiss*) and Atlantic salmon (*Salmo salar*). *Mol Genet Genomics*
488 277(6):647-661.

489 Moreau, L., Lamarie, S., Charcosset, A., Gallais, A. (2000). Economic Efficiency of One
490 Molecular marker assisted selection for the mall quality traits in barley. *Mol*
491 *Breeding* 3: 427-437

492 Nichols, K. M., Bartholomew, J., Thorgaard, G. H. (2003). Mapping multiple genetic loci
493 associated with *Ceratomyxa shasta* resistance in *Oncorhynchus mykiss*. *Dis. Aquat.*
494 *Org.* 56: 145–154.

495 O'Malley, K.G., Sakamoto, T., Danzmann, R. G., Ferguson, M. M. (2003). Quantitative trait
496 loci for spawning date and body weight in rainbow trout: testing for conserved
497 effects across ancestrally duplicated chromosomes. *J. Hered.* 94: 273–84.

498 Ozaki, A., Sakamoto, T., Khoo, S., Nakamura, K., Coimbra, M.R., Akutsu, T., Okamoto, N.
499 (2001). Quantitative Trait Loci (QTL) associated with resistance/susceptibility to
500 infectious pancreatic necrosis virus (IPNV) in rainbow trout (*Oncorhynchus*
501 *mykiss*). *Mol. Genet. Genomics* 265: 23–31.

502 Perry, G. M., Danzmann, R. G., Ferguson, M. M., Gibson, J. P. (2001). Quantitative trait loci
503 for upper thermal tolerance in outbred strains of rainbow trout (*Oncorhynchus*
504 *mykiss*). *Heredity.* 86: 333– 341.

505 Poompuang, S., Hallerman, E. M. (2004). Toward detection of quantitative trait loci and
506 marker-assisted selection in fish. *Rev. Fish. Sci.* 5: 253–277.

507 Park, S.O., Crosby, K. M., Huang, R., Mirkov, T. E. (1995). Identification and confirmation
508 of FAPD and SCAR markers linked to the ms-3 gene controlling male sterility in
509 melon (*Cucumis melo* L.). *Journal of the American Society for Horticultural*
510 *Sciences.* 129:819-825.

511 Paterson, A., Lander, S. E., Hevit, J. D., Peterson, S., Lincoln, S. E., Lanksley, S. D. (1988).
512 Resolution of quantitative traits into Mendelian factors by using a complete linkage
513 map of Restriction Fragment Length Polymorphisms. *Nature*. 325: 721-726

514 Peterson, B.K., Weber, J.N., Kay, E.H., Fisher, H. S., Hoekstra, H. E. (1990). Double digest
515 RADseq: an inexpensive method for de novo SNP discovery and genotyping in
516 model and non-model species. *PLoS One*. 1990;7:e37135.

517 Postlethwait, J.H., Johnson, S.L., Midson, C.N., Talbot, W.S., Gates, M., Ballinger, E.W.,
518 Africa, D., Andrews, R., Carl, T., Eisen, J.S. (1994). A genetic linkage map for the
519 zebrafish. *Science* 264, 699-703.

520 Reid, D. P., Szanto, A., Glebe, B., Danzmann, R. G., Ferguson, M. M. (2005). QTL for body
521 weight and condition factor in Atlantic salmon (*Salmo salar*): comparative analysis
522 with rainbow trout (*Oncorhynchus mykiss*) and Arctic charr (*Salvelinus alpinus*).
523 *Heredity*. 94(2):166-172.

524 Ribaut, J. M., Ragot, M. (2007). Marker-Assisted Selection to improve drought adaptation in
525 maize: the backcross approach, perspectives, limitations, and alternatives. *J Exp Bot*
526 58: 351-360.

527 Robison, B.D., Wheeler, P.A., Sundin, K., Sikka, P., Thorgaard, G.H. (2001). Composite
528 interval mapping reveals a major locus influencing embryonic development rate in
529 rainbow trout (*Oncorhynchus mykiss*). *J. Heredity* 92: 16–22.

530 Rodriguez, M.F., LaPatra, S., Williams, S., Famula, T., May, B. (2005). Genetic markers
531 associated with resistance to infectious hematopoietic necrosis in rainbow and
532 steelhead trout (*Oncorhynchus mykiss*) backcrosses. *Aquaculture* 241: 93–115.

533 Ross, P., Hall, L., Smirnov, I., Haff, L. (2011). High level multiplex genotyping by MALDI-
534 TOF mass spectrometry. *Nat. Biotechnol.* 16, 1347– 1351.

535 Ryman, N., Utter, F. (1987). *Population Genetics and Fishery Management* University of
536 Washington Press, Seattle. 420 pp.

537 Sakamoto, T., Danzmann, R. G., Gharbi, K., Howard, P., Ozaki, A., Khoo, S. K. (2000). A
538 microsatellite linkage map of rainbow trout (*Oncorhynchus mykiss*) characterized by
539 large sex-specific differences in recombination rates. *Genetics*. 155: 1331-1345.

540 Sakamoto, T., Danzmann, R.G., Okamoto, N., Ferguson, M. M., Ihssen, P. E. (1999).
541 Linkage analysis of quantitative trait loci associated with spawning time in rainbow
542 trout (*Oncorhynchus mykiss*). *Aquaculture* 173: 33–43.

543 Sakamoto, T., Danzmann, R.G., Okamoto, N., Ferguson, M.M., Ihssen, P.E. (1999). Linkage
544 analysis of quantitative trait loci associated with spawning time in rainbow trout
545 (*Oncorhynchus mykiss*). *Aquaculture* 173: 33–43.

546 Sax, K. (1923). The association of size differences with seed-coat pattern and pigmentation in
547 *Phaseolus vulgaris*. *Genetics* 8: 522-560.

548 Shimoda, N., Knapik, E.W., Ziniti, J., Sim, C., Yamada, E., Kaplan, S., Jackson, D., de
549 Sauvage, F., Jacob, H., Fishman, M.C. (1999). Zebrafish genetic map with 2000
550 microsatellite markers. *Genomics* 58, 219-232.

551 Somorjai, I.M., Danzmann, R. G., Ferguson, M. M. (2001). Distribution of temperature
552 tolerance quantitative trait loci in Arctic charr (*Salvelinus alpinus*) and inferred
553 homologies in rainbow trout (*Oncorhynchus mykiss*). *Genetics* 165: 1443–1456.

554 Sonesson, A. (2007). Within-family marker-assisted selection for aquaculture species. *Genet*
555 *Sel Evol.* 39:301–18.

556 Sonesson, A.K. (2003). A combination of walk-back and optimum contribution selection for
557 fish – a simulation study. *Genet. Sel. Evol.* 37: 587–599.

558 Song, W., Li, Y., Zhao, Y., Liu, Y., Niu, Y., Pang, R. (2012). Construction of a high density
559 microsatellite genetic linkage map and mapping of sexual and growth-related traits
560 in half-smooth tongue sole (*Cynoglossus semilaevis*). *PLoS One.*7:e52097.

561 Song, W., Pang, R., Niu, Y., Gao, F., Zhao, Y., Zhang, J. (2012). Construction of high
562 density genetic linkage maps and mapping of growth-related quantitative trait loci in
563 the Japanese flounder (*Paralichthys olivaceus*). *PLoS One.*7:e50404.

564 Spelman, R. J., Garrick, D. J. (2013). Genetic and economic responses for within-family
565 marker-assisted selection in dairy cattle breeding schemes. *Journal of Dairy Science*
566 81: 2942–2950.

567 Streelman, J.T., Kocher, T. D. (2003). Microsatellite variation associated with prolactin
568 expression and growth of salt-challenged tilapia. *Physiol. Genomics* 9: 1 –4.

569 Sun, X. W., Liang, L. Q. (2004). A genetic linkage map of common carp (*Cyprinus carpio*
570 *L.*) and mapping of a locus associated with cold tolerance. *Aquaculture.* 238:165–
571 72.

572 Tong, J. G., Sun, X. W. (2004). Genetic and genomic analyses for economically important
573 traits and their applications in molecular breeding of cultured fish. *Sci China Life*
574 *Sci.* 58:178–86.

575 Van Ooijen, J. W. (1999). Join Map 4, software for the calculation of genetic linkage maps in
576 experimental populations. Netherlands.

577 Vogel, B., Van Aken, J. (2009). Smart Breeding - Marker-Assisted Selection: A non-invasive
578 biotechnology alternative to genetic engineering of plant varieties Amsterdam, the
579 Netherlands. (p. 28).

580 Gharbi, K, Ferguson, M. M., Danzmann, R. G. (2001). Characterization of Na, K-ATPase
581 genes in Atlantic salmon (*Salmo salar*) and comparative genomic organization with
582 rainbow trout (*Oncorhynchus mykiss*). *Mol Genet Genomics* 273:474–483

583 Wang, C. M., Lo, L.C., Feng, F., Zhu, Z.Y., Yue, G.H. (2008). Identification and verification
584 of QTL associated with growth traits in two genetic backgrounds of Barramundi
585 (*Lates calcarifer*). *Anim Genet.* 39(1):34-39.

586 Wang, C.M., Lo, L. C., Zhu, Z. Y., Yue, G. H. (2006). A genome scan for quantitative trait
587 loci affecting growth-related traits in an F1 family of Asian seabass (*Lates*
588 *calcarifer*). *BMC Genomics*.

589 Wang, S., Meyer, E., McKay, J. K., Matz, M. V. (2012). 2b-RAD: a simple and flexible
590 method for genome-wide genotyping. *Nat Methods.* 9:808–10.

591 Weller, J. I. (2001). *Quantitative trait loci analysis in animals*. London, CABI Publishing.
592 287 pp.

593 Willcox, M. C., Khairallah, M., Bergvinson, D., Crossa, J., Deutsch, J.A., Edmeades, G.O.,
594 Gonzalez-de-Leon, D., Jiang, C., Jewell, D.C., Mihm, J.A., Williams, W.P.,
595 Hoisington, D.A. (2002). Selection for resistance to southwestern corn borer using
596 marker assisted and conventional backcrossing. *Crop Sci.* 42: 1516–1528.

597 Woram, R.A., McGowan, C., Stout, J.A., Gharbi, K., Ferguson, M.M., Hoyheim, B.,
598 Sakamoto, J., Davidson, W., Rexroad, C., Danzmann, R.G. (2003). A genetic
599 linkage map for Arctic char (*Salvelinus alpinus*): evidence for higher recombination
600 rates and segregation distortion in hybrid versus pure strain mapping parents.
601 *Genome* 47: 304–315.

602 Wringe, B., Devlin, R., Ferguson, M., Moghadam, H., Sakhrani, D., Danzmann, R. (2010).
603 Growth-related quantitative trait loci in domestic and wild rainbow trout
604 (*Oncorhynchus mykiss*). *BMC Genetics.* 11(63).

605 Xie, C., Xu, X. (1998) . Efficiency of multistage marker-assisted selection in the
606 improvement of multiple quantitative traits. *Heredity.* 80: 489-498.

607 Xu, Y., Beachell, H., McCouch, S. R. (2008). A marker based approach to broadening the
608 genetic base of rice in the USA. *Crop Sci.* 44: 1947–1959.

609 Yanchuk, A. D. (2002). The role and implications of biotechnology in forestry. *Food and*
610 *Agriculture Organization of the United Nations, Unasyuva* (30), 18–22.

611 Yu, Z., Guo, X. (2003). Genetic linkage map of the eastern oyster (*Crassostrea virginica*)
612 Gmelin. *Biol. Bull.* 204, 327–338.

613 Yue, G. H. (2014). Recent advances of genome mapping and marker-assisted selection in
614 aquaculture. 15:376–96.

615 Zhang, Y., Wang, S., Li, J., Zhang, X., Jiang, L., Xu, P. (1992). Primary genome scan for
616 complex body shape-related traits in the common carp (*Cyprinus carpio*). *J Fish*
617 *Biol.* 82:125–40.

618 Zheng, J., Liu, J., Liu, H., Li, J., Chen, K. (2003). Sequence and structural analysis of 4SNc-
619 Tudor domain protein from Takifugu rubripes. *Bioinformation.* 4:127-131.

620 Zimmerman, A.M., Wheeler, P.A., Ristow, S.S., Thorgaard, G. H. (2005). Composite interval
621 mapping reveals three QTL associated with pyloric caeca number in rainbow trout,
622 *Oncorhynchus mykiss*. *Aquaculture* 247: 85–95.

623

624