

**MICROBIAL GIMICS: STRATEGIES OF SUCCESFUL PATHOGENICITY BY
*STAPHYLOCOCCUS AUREUS***

ABSTRACT

Staphylococcus aureus is a major human pathogen associated with a variety of clinical diseases. It is the leading cause of wound infections, skin infections, respiratory infections as well as device-related infections. This review comprehensively covers the virulence determinants of the organism and the different mechanisms of antibiotic resistance in the organism. Recently, *Staphylococcus aureus* has become a serious threat because of its ability to evolve which has led to challenges in the treatment of infections caused by the organism.

Keywords; antibiotic resistance, pathogenicity, infection.

INTRODUCTION

Staphylococcus aureus is a Gram-positive, non-motile, non-spore forming microorganism. It is present in the normal flora of the human nasopharynx and skin and makes up about 30% in a healthy human population [1]. It does not cause disease as a component of the normal flora but a break in the skin causes the bacterium to enter a wound and colonize it, thereby causing infections. However, *S. aureus* has the potential of being an opportunistic pathogen, producing a broad variety of diseases in humans, starting from a minor skin infection to a fatal form of pneumonia resulting in human mortality. *S. aureus* has a typical evolutionary nature which makes it a successful pathogen. It is associated with a variety of diseases **Examples** include; acute sepsis, respiratory infections, wound infections amongst others. It has also been implicated in different skin infections such as boils, impetigo, carbuncles, folliculitis etc. *S. aureus* is a major cause of bloodstream infections which occurs following a puncture on the mucosal membrane or on the surface of the skin following surgery, injury and the use of the catheter in hospital settings. Once inside the bloodstream, it has the capacity to infect numerous organs in the body and as well produces different pigments and molecules that help it to escape the host immunity and establish an infection such as protein A, staphyloxanthin etc. it, however, produces biofilms by producing different adhesins that enable it to adhere to host surfaces.

S. aureus is gradually evolving in animals (Livestock-associated Methicillin resistant *S. aureus*). This group of *S. aureus* heavily colonize pigs and calves in farms and because of this, the farmworkers and veterinarian are susceptible to infection by LA-MRSA [2]. It also encodes different virulence factors such as toxins, enzymes which are mediated by horizontal


37 gene transfer, and this, however, contributes to the emergence of antibiotic resistance to
38 multiple classes of antimicrobial drugs.

39 This review would be focused on the different mechanisms by which *S. aureus* acquires
40 resistance to antibiotics (horizontal gene transfer), some virulence determinants that are
41 mediated through this means and some antibiotic resistance in the organism.

42

43 HORIZONTAL GENE TRANSFER IN *S. AUREUS*.

44 This is a mechanism by which *S. aureus* can transfer DNA (mobile genetic elements) (MGE)
45 from one bacterial cell to another. This mechanism enhances the circulation of (MGE) which
46 encodes for virulence as well as antibiotic resistance. There are diverse ways by which the
47 genetic information can be transferred or acquired from other cells or the environment as
48 shown in Fig.1. [3]. They include through conjugation, generalized transduction, plasmids,
49 transposons, bacteriophages, genomic islands, staphylococcal cassette chromosome (SCC),
50 transformation etc.



51

52

53 **Fig.1:** The Acquisition of mobile genetic elements by *Staphylococcus aureus*: 1.
54 Incorporation of plasmids into a bacterial DNA. 2. Plasmid integrated into the chromosome
55 of a bacterium. 3. Plasmids as an independent circular DNA. 4. Transfer of a transposon

56 between plasmid and genomic DNA. 5. Transfer of transposons between plasmids. 6.
57 Transfer of transposons from a genomic DNA to a plasmid. [3].

58

59

60

61 **GENERALIZED TRANSDUCTION**

62 Transduction is the transfer of DNA from one cell to another through a bacteriophage (Fig.1)
63 During replication, the bacteriophage gets integrated into the chromosome and can be
64 transferred to its daughter cells (Fig.1). A prophage can be instigated by stress, resulting in
65 the cutting of the phage DNA, reproducibility of the prophage DNA, synthesis of novel
66 prophage proteins etc. The size of the prophage is typically 45kb and they are known to code
67 for virulence determinants like the Pantone-Valentine Leucocidin (PVL), chemotaxis
68 inhibitory protein amongst others. The phage particles can either kill the recipient host (lytic
69 pathway) or get integrated into the recipient's chromosome as a prophage (lysogenic
70 pathway). The lysogenic pathway is common in *S. aureus* where isolates carry between 1-4
71 different prophage types [1]. However, in generalized transduction, the new growing phage
72 particles package the bacterial chromosomal DNA instead of the phage DNA. It has been
73 shown that some bacteriophages do this while some others do not, but the mechanism is still
74 not known.

75 However, this could be a natural mechanism of conserving its host DNA as well as
76 transferring its genetic element to like or non-like bacterial cells. The phage particles that are
77 released during lysis binds to the *Staphylococcus aureus* recipient's receptor and introduces
78 its DNA into the cell [4,5]. Because the DNA is not a phage, it does not get integrated into
79 the chromosome like a lysogenic phage will do nor does it kill the recipient cell as the lytic
80 cell will do. However, some host DNA seems to be selectively packaged by the phage
81 leading to an elevated level of transfer.


82 **CONJUGATION AND PLASMIDS**

83 This is a mechanism whereby DNA is transferred from one cell to another through a pilus or
84 a pore [6]. In *S. aureus*, it is assumed that the pores are made between cells that are in close
85 contact with each other because the pili are not seen. A range of plasmids carrying resistance
86 genes is transferred during the process of conjugation. as shown in Fig 2. (Adopted by
87 evolution website). These conjugative plasmids are too large, and they carry an extensive
88 range of antibiotic resistance genes and virulence factors which they transfer from one
89 organism to another [7,8]. Most of the staphylococcal strains contain plasmids with 1-60kbp.
90 *S. aureus* plasmids are made up of three classes. Class I is made up of tiny multi copies of
91 plasmids per cell carrying resistance genes. The plasmids in this class do not have

92 transposons nor prophages. Class II plasmids are known to be larger in size and they appear
93 in lesser copy numbers. This class of plasmids includes the penicillinase, aminoglycosides
94 resistance plasmids. Class III plasmids consist of bigger plasmids which carry conjugative
95 transfer genes. The class III plasmids most often possess transposons including many copies
96 of insertion sequences. Before these plasmids get integrated into the host chromosome, they
97 are usually free DNA. They are known to code for some virulence factors such as exfoliative
98 toxin and bacteriocin [9]. They also encode resistance to various organic and inorganic ions
99 that are usually toxic to living cells and thermostable genes [10].

100

101



102

103

104 Fig 2: An illustration of a single plasmid (pLW1043) which carries resistance genes to be
105 conferred on different antibiotics.

106 **TRANSFORMATION**

107 This is a horizontal gene transfer mechanism that involves the uptake/intake of free DNA
108 from the environment by a competent bacterium. Some bacteria are readily competent such as
109 *Bacillus subtilis*, *Streptococcus pneumonia* whereas some are not readily competent such as
110 *E. coli*. For bacteria that are not readily competent, competence can be induced chemically
111 (addition of calcium ions) or through electroporation. However, previous studies show that *S.*
112 *aureus* has low transfer efficiency in taking up free DNA from the environment. This transfer
113 requires phage proteins (tail proteins) and is dependent on the presence of a lytic phage. The
114 phage proteins bind to the cell when DNA is present, and this facilitates the transfer of the
115 DNA into the cell. It has now been shown that *S. aureus* can engage in natural transformation
116 through a bacterial encoded protein [11]. In this case, its ability to take up DNA is being
117 controlled by sigma H factor which is needed for the maintenance of the lysogenic phage.
118 Interestingly, the *S. aureus* sigma H gene does not switch on competence until it is able to
119 duplicate itself and change its promoter region. This impulsive chromosomal arrangement

120 happens at low frequencies so that a tiny proportion of the population will finally express the
121 sigma factor. The expression of the sig H gene also requires specific nutritional requirements,
122 and this was known using a lysogenic bacterium which carries the sig H on a plasmid and
123 was able to take up the plasmid demonstrating that the process is phage independent.

124 **TRANSPOSONS**

125 *S. aureus* genome is also made up of transposons, insertion sequences, and transposon-like
126 elements. These mobile genetic elements contribute to the evolutionary nature of the
127 bacterium and can be found in the chromosome or in close contact with other mobile genetic
128 element either as single or multiple copies [3]. Insertion sequences are involved in carrying
129 genetic information that is needed for transposition. They don't encode for resistance but
130 oversee the recombining and upkeep of these resistance genes. Because of this, they are vital
131 in the development of *S. aureus* genome by promoting alterations in the bacterial gene
132 expression. Insertion sequences are also capable of inactivating numerous genes through
133 direct insertion or through a polar effect on close gene transcription [12]. Insertion elements
134 are mostly in a combination form e.g. Insertion sequence 256 and Insertion sequence 257 are
135 moderated by Transposons 4001 and 4003 forming a pair which mediates resistance to some
136 antibiotics like gentamicin and kanamycin. The insertion of Insertion 256 and 257 into *S.*
137 *aureus* chromosome function in the rearrangement of its genome. *Staphylococcus aureus*
138 transposons are little genetic elements which code genes that are resistant to a wide range of
139 antibiotics such as erythromycin, macrolide-lincosamide, spectinomycin, methicillin amongst
140 others. Various copies are found being integrated into plasmids or Staphylococcal cassette
141 chromosome [2].

142

143 **STAPHYLOCOCCAL CASSETTE CHROMOSOME (SCC)**

144 This is another mobile genetic element of *S. aureus* family. The SCC elements can insert into
145 the 'orfX' gene in the *staphylococcus* chromosome and are responsible for methicillin
146 resistance in *S. aureus*. Its integration requires a specific attachment site (attB_{sc}) in the orfX
147 region. They are classified into two groups; the mec-staphylococcal chromosome and the non
148 -mec staphylococcal chromosome.

149 **Mec- Staphylococcal cassette Chromosome**

150 All MRSA strains contain the SCC mec element. One of the genes it encodes is the mecA
151 gene. The 'mecA' genes confer resistance to all beta-lactam antibiotics most notably the
152 methicillin [13]. *S. aureus* can resist the methicillin antibiotic because of the production of a
153 modified penicillin-binding protein (PBP2a) which has a low affinity for beta-lactams
154 thereby rendering them clinically ineffective. There are various types of SCC mec ranging
155 from type I to type XI and they all encode resistance genes (Table 1). About six different
156 classes have been shown about their arrangement and associated genes [14,15].

157 **Non-mec staphylococcal cassette chromosome**

158 These are SCC elements that are not limited to encoding for only methicillin resistance. They
159 also contain virulence or survival determinants and have been identified in *S. aureus*. They
160 share some characteristics with the **major mec sec (WHY IN RED???)** such as the integration
161 into the staphylococcal chromosome, the presence of flanked repeated sequences. Regarding
162 the nomenclature of these elements, it was proposed to include a suffix that describes the
163 gene functions. Examples include SCCcap1 which is a type 1 capsule gene cluster, SCCfur
164 (which harbours the resistance for fusidic acid) and SCChg which carries an operon for
165 mercury resistance [15].

166

167

168 **Table 1: (WHERE IS THE TABLE TITLE???) (WHY THE SPACES BETWEEN THE**
169 **LINES ARE IN DOUBLE???)**

Scc mec types	Mec gene complex	Structure of the mec gene complex	Reference
---------------	------------------	--------------------------------------	-----------

I	Class B	IS1272 Δ mec RI- mec A IS431	[16]
II	Class A	mec I-mec RI-mec A-IS431	[16]
III	Class A	mec I-mecRI-mec A-IS431	[16]
IV	Class B	IS431-mec A- Δ mec RI- IS1272	[17]
V	Class C2	IS431-mecA- Δ mec RI-IS431	[18]
VI	Class B	ISI 272- Δ mec RI- mec A- IS431	[19]
VII	Class CI	mec- mecRI-mec A-IS431	[20]
VIII	Class A	IS431-mecA Δ mec RI- IS431	[20]
IX	Class C2	IS431-mecA-	[21]

[21]

X Class CI

XI Class E bla Z-mec A-mec RI-mecI [22]

170

171

172

173 **GENOMIC ISLANDS**

174 They are mobile genetic elements that are present among the core genes of a bacterium either
175 in the chromosome or in a plasmid and they are usually acquired by horizontal gene transfer.
176 [23,24]. Among the *S. aureus* strains that have been sequenced, three families or groups of
177 genomic islands are present [1,25, 26) known as the VSAa, VSAb, and VSAy. The VSAa

178 family carry a lipoprotein gene and a staphylococcal enterotoxin gene (SEI) [27]. The VSA_b
179 family encodes for bacteriocin, enterotoxins, hyaluronate lyase in addition to a serine
180 protease gene group [26, 28,29]. The VSA_y family comprises of genes coding Beta type
181 phenol soluble modulins (PSM) and a group of staphylococcal enterotoxin gene (SEI). [25].
182 These islands are usually flanked by 16-20 base pair direct repeats. These repeats are as a
183 result of the integration of the island into a specific site for it to exert its enzymatic function.
184 The genomic island's stability is enhanced by an upstream and downstream flanking of DNA
185 segments. However, most of the islands are not seen to be mobile since they have to
186 degenerate before they can be transferred.

187 **BACTERIOPHAGES**

188 Phages also play a key role in *S. aureus* adaptation and evolution, and they are transferred
189 through horizontal gene transfer. They are also involved in the induction, packaging, and
190 transfer of genomic islands. *S. aureus* phage is classified into three families known as
191 Siphoviridae, Myoviridae, and Podoviridae. The Podoviridae family contains the lytic and
192 chronic phages, and they harbour the smallest set of genomes compared to the genomes
193 present in the other families. The Myoviridae also contains the lytic and chronic phages but
194 the Siphoviridae family contains all the temperate phages and they are capable of living for a
195 very long time in the host. The virulent phages present in Myoviridae and Podoviridae are
196 used as a phage therapy in humans against *S. aureus* infections and for food preservation as
197 well. These phages also encode different virulence factors such as staphylokinase,
198 enterotoxins amongst others and these genes are located close to the attachment site in the
199 host chromosome [30]. In *S. aureus* pathogenicity islands (SaPI), helper phages are needed
200 for its mobilization and the helper phages that can perform this function include the temperate
201 phages which belong to the Siphoviridae family [31]. They help to increase the mobility of *S.*
202 *aureus* pathogenicity island to other staphylococci [32]. The SaPI are not mobile on their own
203 therefore they depend on a helper phage for its replication between different *S. aureus*
204 isolates [30,33]. For example, the Pantone Valentine leucocidin is transferred through a helper
205 phage from a PVL-positive to a PVL- negative *S. aureus* strain. It is also of importance to
206 note that only certain helper phages can increase the mobility of certain SaPI.

207

208 **EXPRESSION OF VIRULENCE OR SURVIVAL DETERMINANTS IN *S. AUREUS*.**

209 *S. aureus* produces a wide range of virulence factors which helps it to establish infections in
210 humans either by adhering to surfaces or tissues, by invading the immune system and by
211 causing lethal toxic effects to the host. As we have seen from above that some of these
212 virulence factors are encoded by the horizontal gene transfer mechanisms.

213 **PANTON VALENTINE LEUKOCIDIN (PVL)**

214 PVL is encoded by bacteriophages which enables them to be transferred from one organism
215 to another. It is classified as a cytotoxin, one of the beta forming toxins. It has been reported
216 to be present in community-associated methicillin-resistant *S. aureus* (CA-MRSA), a major
217 cause of necrotizing pneumonia. It lyses neutrophils, leading to the release of enzymes that
218 damages the surrounding tissues [34].

219 **ENTEROTOXINS**

220 *S. aureus* enterotoxins belong to a family of pyrogenic toxin superantigens (SAG). These
221 superantigens bind to the MHC Class II molecules in host animals, therefore, forming a
222 complex with the T cell receptor. The formation of the complex activates the T cell to
223 proliferate in a non-specific manner resulting in host immune suppression [34-35]. The
224 superantigen genes are the major cause of acute clinical syndromes such as toxic shock
225 syndrome, food poisoning etc. The superantigens have been classified into two groups:
226 classical and new enterotoxins (Argudin *et al.*, 2010, Hennekinne *et al.*, 2012, Wilson *et al.*,
227 2011). However, about 23 types of *S. aureus* enterotoxins have been reported and they are all
228 encoded on horizontal gene transfer mechanisms [10,36,37]. Moreover, they have also
229 contributed to the evolution of *S. aureus* as a pathogen. Some of these enterotoxins are
230 components of the enterotoxin gene cluster which is found on genomic islands.

231 **TOXIC SHOCK SYNDROME TOXIN (TSST)**

232 TSST is a superantigen that is produced by a small percentage of *Staphylococcus aureus*
233 isolates. Once these toxins are released into the bloodstream, they cause the over stimulation
234 of the immune system which subsequently leads to symptoms of toxic shock syndrome
235 (TSST). They are however known to live in the vagina of women that are infected which is
236 highly encouraged using a tampon [38]. They are also present in other sites of the body. It
237 has been reported that children, men, and non-menstruating women also have the potential of
238 developing TSST. TSST also has the capacity to stimulate the release of cytokines enhancing
239 the leakage of endothelial cells in low concentrations thereby producing a cytotoxic effect at
240 high concentrations. It also causes systemic infection by penetrating mucosal barriers even
241 though the infection is localized in the vagina or at any other location in the body.

242 **STAPHYLOKINASE**

243 This is another virulence factor of *S. aureus* which is encoded by lysogenic bacteriophage. It
244 is present in the DNA of some bacteriophage and can be transferred from one organism to
245 another. Staphylokinase interacts with plasminogen and α -defensins which enhances *S.*
246 *aureus* invasion into the host tissues. It has been shown that *S. aureus* that carries the
247 staphylokinase- plasminogen complex on their surface can lyse extracellular matrix by
248 activating the metalloproteinases present in the host. Staphylokinases also encourages
249 bacterial resistance in *S. aureus* especially to phagocytosis which is mediated by the
250 interaction of HNPs (Human neutrophil peptides), an important part of the innate immunity.

251 Most importantly, the production of staphylokinase enables *S. aureus* to persist longer on the
252 host skin and mucosa [38].

253 However, there are several other virulence factors produced by *S. aureus* which makes it a
254 versatile pathogen, having the ability to induce a wide range of infections. (Table 2).

255 **Table 2:** Other virulence factors of *S. aureus* [38].

256 <u>Virulence factors</u>	<u>Biological effects</u>
Structural components	
Capsule	Inhibits chemotaxis and phagocytosis; inhibits proliferation of mononuclear cells
Slime layer	Facilitates adherence to foreign bodies; inhibits phagocytosis
Teichoic acid	Binds to fibronectin.
Protein A	Inhibits antibody-mediated clearance by binding to IgG
Toxins	
Exfoliative toxins	Serine proteases that split the intercellular bridges in the stratum granulosum epidermis
Cytotoxins	Toxic for many cells including erythromycin, fibroblasts, leucocytes, macrophages and platelets.

Enzymes

Coagulase

Converts fibrinogen to fibrin

Hyaluronidase

Hydrolyses hyaluronic acids in connective tissues, promoting the spread of staphylococci in tissues

fibrinolysin

Dissolves fibrin clots

Lipases

Hydrolyses lipids

Nucleases

Hydrolyses DNA

257

258

259

260

261 **ANTIBIOTIC RESISTANCE IN *STAPHYLOCOCCUS AUREUS***

262

263 Several antimicrobial resistance genes are also carried on the mobile genetics' elements
264 discussed such as transposons and plasmids. The resistance genes confer resistance to a wide
265 range of antibiotics such as penicillin, macrolides, aminoglycosides, tetracyclines,
266 chloramphenicol, linezolid etc. The capacity of *S. aureus* to easily acquire these resistance
267 genes is one of the characteristics that make it successful in establishing infection, thereby
268 making the control of infection more difficult and complicated. *S. aureus* has been shown to
269 develop resistance to β lactam antibiotics such as penicillin, methicillin and glycopeptide
270 such as vancomycin amongst others.

271

272 **BETA-LACTAM RESISTANCE**

273 *S. aureus* resistance to beta-lactam antibiotics was first seen in penicillin which was mediated
274 by the production of penicillinase (a beta-lactamase) which hydrolyses the beta-lactam ring
275 present in penicillin. Thereby rendering it ineffective. However, methicillin was introduced to
276 subdue penicillin resistance, but it was not possible because *S. aureus* has a way of evolving
277 and adapting to new or nearly or classes of antibiotics which were used to treat it. Therefore,
278 Methicillin-resistant *S. aureus* (MRSA) strains evolved and this has been shown to be
279 mediated by the *mecA* gene. The *mecA* gene is present on the mobile genetic element which
280 is known as staphylococcal cassette chromosome (SCCmec) [39]. The methicillin resistance
281 is not acquired during infection as it has not been observed. However, studies have shown the
282 horizontal transfer of the staphylococcal cassette chromosome at the time of infection giving
283 rise to the emergence of methicillin-resistant *S. aureus* strains [40]. The MRSA strains
284 become resistant to beta-lactam antibiotics by producing a modified penicillin-binding
285 protein (PBP2a) which has a low affinity for beta-lactam antibiotics thereby rendering them
286 clinically ineffective. MRSA has been identified in hospitals; Hospital-associated MRSA
287 (HA-MRSA). Several clones accounted for most of the HA-MRSA include ST22, ST36,
288 ST239, and ST5. These clones successfully evolve and establish themselves mostly due to the
289 intensive use of antibiotics, mutations and poorly registered regimens [41]. MRSA has also
290 been identified in communities; Community associated MRSA (CA-MRSA). Previously,
291 CA-MRSA greatly affects immunocompromised individuals with predisposing factors and
292 those with health care exposure. However, in recent times, it affects healthy hosts particularly
293 children and middle-aged adults. This could be attributed to increased transmission of
294 infection, activation of more virulence genes and an increased pathogenicity during infection
295 [40]. Interestingly, MRSA has now been identified in animals: Livestock-associated MRSA
296 (LA-MRSA) as a cause of infection in humans. Infections due to LA-MRSA occur in persons
297 who have close access to farm animals such as pigs, poultry, dogs, cats etc. it affects mostly
298 the farmers and veterinarians. LA-MRSA was identified in a cow in 1972. In 2005, CC398
299 MRSA lineage was reported in pigs in Europe showing that the livestock was a good
300 reservoir for MRSA. The main reservoir for CC398 is in pigs but it has also been found in
301 veal calves, poultry, horses, dogs, cats and to an extent, in cows. There has been a general
302 agreement that CC398 is increasing worldwide although information on prevalence rate has
303 been difficult to obtain. Other complex MRSA lineages in livestock that have been found
304 include the CC9, CC1, CC5, CC97, CC121, CC130, and ST 425 [42]. It is of interest that a
305 human CA-MRSA type descended from bovine MSSA after bovine-host adaptation [43-44].
306 Risk factors for its transmission are not fully understood although one of the important risk
307 factors is the trade of pigs that are MRSA positive. However, some farmers have been found
308 positive even without buying new animals before the MRSA CC398 was detected. In these
309 exceptional cases, it could be that they become MRSA positive from MRSA-positive humans
310 like veterinarians. The use of antibiotics amongst farmers most notably the beta-lactams and
311 tetracyclines also induce selective pressure on the clones [45]. The most crucial risk factor for
312 LA-MRSA in humans is the close occupational access with animals which are MRSA
313 positive which depends on the contact time and intensity. In a study at Denmark in 2013,

314 most of the new cases, (about, 70%) that were reported had to do with direct contact with
315 pigs, (17%) were linked with members of the house who had close access to pigs while the
316 remaining 13% were those who had no contact with pigs but lived in places that had high
317 pig density indicating that transmission takes place probably from the people working at the
318 farms or through access with farm surroundings itself [46]. However, the comparative
319 contribution of transmission whether through the surroundings of the farm or through humans
320 hasn't been elucidated. Although, from the knowledge of *S. aureus* transmission in other
321 settings, human-human contact is predominant [47]. It has also been shown that MRSA has
322 been found on meat which raises the likelihood of MRSA being acquired through the food
323 chain. From the epidemiology of LA-MRSA, it clearly indicates that meat is not one of the
324 routes of transmission [46]. The increasing rate of LA-MRSA in pigs including humans who
325 have close access with pigs has resulted in an increase in cases in the communities, especially
326 in the immunocompromised persons. Therefore, it is possible that increasing numbers of
327 infections caused by LA-MRSA will be seen unless the epidemic is monitored. Furthermore,
328 if the human carriage of LA-MRSA clone is increased, then it would lead to a greater chance
329 of these clones undergoing adaptation which will enhance human-human transmissibility.
330 Measures to reduce the increasing reservoir in pigs is highly needed.

331

332 **GLYCOPEPTIDE RESISTANCE**

333 MRSA strains have also developed resistance to glycopeptide antibiotics such as
334 vancomycin. Vancomycin acts by binding to the D-ala D-ala residues of the peptidoglycan
335 thereby inhibiting cell wall synthesis. It is used in the treatment of infections caused by
336 MRSA such as osteomyelitis, endocarditis, bacteraemia [48]. Two mechanisms of
337 vancomycin resistance have evolved in *Staphylococcus aureus*. The first resistance to evolve
338 were *S. aureus* isolates which had decreased susceptibility to vancomycin known as
339 vancomycin- intermediate resistant *S. aureus* (VISA) strain. (with a MIC of 8µg/ml). These
340 strains have an excess binding site which can 'confine' the antibiotic [49]. They also show
341 characteristics of a decreased autolysis, attenuation of virulence and thickened cell wall [50].
342 The thickness of the VISA cell wall was first reported in a 4-month-old infant who had a
343 heart surgery; it showed that the VISA strain known as 'Mu50' which was isolated from the
344 discharge at the surgery site had a cell wall that was two times thick as the control strains
345 seen under the microscope. [49] demonstrated this and showed that the thickness of the cell
346 wall was a common characteristic of the VISA isolates. Due to the thickened cell wall,
347 present, it makes these strains more resistant because the antibiotic is being 'confined' by the
348 free D-ala residues in the cell wall [49]. Furthermore, VISA strains also show decreased
349 autolytic activity. It has been proved when cell assays were carried out in the VISA strain,
350 'Mu50' [51]. The reduced autolysis has been suggested to may have contributed to the
351 thickened cell wall thereby preventing the antibiotic from getting into its site of action.
352 However, the acquisition of resistance to antibiotics among VISA strains could be a
353 disadvantage towards its virulence [52]. Animal models have been used to ascertain the

354 extent of VISA pathogenesis; in an insect model, it was shown that the clinical VISA isolates
 355 had decreased virulence [53,50]. Also, in a rat model, the VISA isolate was shown to have a
 356 decreased virulence likewise in a mouse sepsis model, the VISA isolates had reduced
 357 infectivity and there was no capacity to cause liver abscesses. The VISA strains tend not to
 358 cause acute infections because of its attenuated virulence, however; this may be a ‘sneaky’
 359 strategy to evade host immune responses [54]. In addition, multiple mutations in different
 360 loci with VISA have also emerged and has been shown to contribute to its level of resistance
 361 to antibiotics. To identify these mutations, whole genome sequencing of the isolates has been
 362 carried out and it showed the presence of several mutations which were associated with
 363 resistance to other antibiotics such as β lactams, rifampicin including vancomycin [55].
 364 Recently, a second-high level vancomycin-resistant *S. aureus* (VRSA) emerged. The first
 365 case of VRSA was seen in a patient who was diabetic and had a co-infection of
 366 *Staphylococcus aureus* and *Enterococcus faecalis* [56]. Evidence has shown that resistance in
 367 MRSA strain was mediated by the acquisition of the Tn1546 transposon which encodes for
 368 vancomycin resistance factor (van A) in the *Enterococcus faecalis* strain. However, there
 369 hasn’t been a person-person spread, therefore, the importance of van-mediated resistance
 370 hasn’t been fully elucidated [57].

371 Mechanism of resistance of *S. aureus* to other antibiotics are also common and have been
 372 summarised in (Table 3). It is also important to note that resistance to new drugs like
 373 linezolid and daptomycin has been shown amongst MRSA in clinical settings.

374 **Table 3:** Mechanisms of *S. aureus* resistance to other antimicrobials [58, 3]

Antibiotic	Resistance genes	Mechanism of resistance	Location
Quinolones	par C, (a component of topoisomerase IV), gyrA,gyrB(a component of gyrase).	mutations in the QRDR region	Chromosome
Aminoglycosides	Modifying enzymes (acetyltransferase, phosphotransferase)	Acetylating or phosphorylating enzymes	Plasmids

Trimethoprim- Sulfamethoxazole	Sulfonamide: dihydropteroate synthase, dihydrofolate reductase	TMP;	Acetylating or phosphorylating enzymes overproduction of para amino benzoic acid decreased affinity for hydrofolate reductase.	Plasmids
Tetracyclines	Tetracyclines tetracycline, doxycycline and minocycline, TetM		Binding to the ribosome and removing the drug from its binding site.	Plasmids; Transposons
Erythromycin	msrA (efflux protein), erm (ribosomal methylase)		efflux pump and alteration of 23S rna Transposons	Plasmids
Linezolid	Cfr		methylation of the 23S rRNA that interferes with Ribosomal binding.	Plasmid
Daptomycin	mprF		increasing synthesis of total LPG	Chromosomal

translocation and
positive net
charges on the
cell membrane

375 The resistance of *Staphylococcus aureus* to beta lactam antibiotics as well as other antibiotics
376 such as Tetracyclines, Lincosamides and Gentamicin has led to the development of newer
377 drugs which are now exploited for the treatment of infections caused by the organism.
378 However, some of which are still undergoing clinical trials. Some of the promising molecules
379 such as Triclosan etc have been designed to target fatty acid biosynthesis, cell division
380 protein, the Clp P protease activator and the Lipid A moiety of lipid II.[59].

381

382 CONCLUSION

383 *S. aureus* is a successful pathogen due to its versatility and evolutionary nature and this has
384 contributed to its success in invading the human immune system thereby establishing an
385 infection. This has been seen from its ability to cause a wide range of mild infections and life-
386 threatening diseases in humans. There is a close relationship between the horizontal gene
387 transfer mechanisms and its virulence factors. These mechanisms not only encode for
388 resistance but also encodes for virulence determinants which are responsible for causing
389 infections in humans. This is important for our knowledge of how *Staphylococcus aureus* is
390 being shaped by selective pressures. This also allows us to understand the versatility of *S.*
391 *aureus* and discover ways by which its evolutionary nature can be genetically manipulated to
392 control infection and reduce its level of resistance to multiple antibiotics.

393

394 CONFLICT OF INTEREST

395 The authors declare no conflict of interest.

396

397 REFERENCES

398 1. Lindsay, J.A. and Holden, M.T., 2004. *Staphylococcus aureus*: superbug, super
399 genome? *Trends in microbiology*, 12(8), pp.378-385.

400

401 2. Lindsay, J.A., 2010. Genomic variation and evolution of *Staphylococcus*
402 *aureus*. *International Journal of Medical Microbiology*, 300(2), pp.98-103.

403

404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444

3. Malachowa, N. and DeLeo, F.R., 2010. Mobile genetic elements of *Staphylococcus aureus*. *Cellular and molecular life sciences*, 67(18), pp.3057-3071.
4. Xia, G., Corrigan, R.M., Winstel, V., Goerke, C., Gründling, A. and Peschel, A., 2011. Wall teichoic acid-dependent adsorption of staphylococcal siphovirus and myovirus. *Journal of bacteriology*, 193(15), pp.4006-4009.
5. Winstel, V., Liang, C., Sanchez-Carballo, P., Steglich, M., Munar, M., Bröker, B.M., Penadés, J.R., Nübel, U., Holst, O., Dandekar, T. and Peschel, A., 2013. Wall teichoic acid structure governs horizontal gene transfer between major bacterial pathogens. *Nature communications*, 4.
6. Grohmann, E., Muth, G. and Espinosa, M., 2003. Conjugative plasmid transfer in gram-positive bacteria. *Microbiology and molecular biology reviews*, 67(2), pp.277-301.
7. McCarthy, A.J., Witney, A.A. and Lindsay, J.A., 2012. *Staphylococcus aureus* temperate bacteriophage: carriage and horizontal gene transfer are lineage associated. *Front Cell Infect Microbiol*, 2(6).
8. Liu, M.A., Kwong, S.M., Jensen, S.O., Brzoska, A.J. and Firth, N., 2013. Biology of the staphylococcal conjugative multi resistance plasmid pSK41. *Plasmid*, 70(1), pp.42-51.
9. Bukowski, M., Władyka, B. and Dubin, G., 2010. Exfoliative toxins of *Staphylococcus aureus*. *Toxins*, 2(5), pp.1148-1165.
10. Argudín, M.Á., Mendoza, M.C. and Rodicio, M.R., 2010. Food poisoning and *Staphylococcus aureus* enterotoxins. *Toxins*, 2(7), pp.1751-1773.
11. Morikawa, K., Takemura, A.J., Inose, Y., Tsai, M., Ohta, T. and Msadek, T., 2012. Expression of a cryptic secondary sigma factor gene unveils natural competence for DNA transformation in *Staphylococcus aureus*. *PLoS Pathog*, 8(11), p.e1003003.
12. Needham, C., Noble, W.C. and Dyke, K.G.H., 1995. The Staphylococcal Insertion Sequence IS257Is Active. *Plasmid*, 34(3), pp.198-205.

445

446

447

448

13. Chambers, H.F. and DeLeo, F.R., 2009. Waves of resistance: *Staphylococcus aureus* in the antibiotic era. *Nature Reviews Microbiology*, 7(9), pp.629-641.

449

450

451

452

453

454

455

456

457

458

459

15. De Lencastre, H., Oliveira, D., and Tomasz, A., 2007. Antibiotic resistant *Staphylococcus aureus*: a paradigm of adaptive power. *Current opinion in microbiology*, 10(5), pp.428-435.

460

461

462

463

464

465

466

467

16. Ito, T., Katayama, Y., Asada, K., Mori, N., Tsutsumimoto, K., Tiensasitorn, C. and Hiramatsu, K., 2001. Structural comparison of three types of staphylococcal cassette chromosome mec integrated into the chromosome in methicillin-resistant *Staphylococcus aureus*. *Antimicrobial agents and chemotherapy*, 45(5), pp.1323-1336.

468

469

470

471

472

473

17. Kwon, N.H., Park, K.T., San Moon, J., Jung, W.K., Kim, S.H., Kim, J.M., Hong, S.K., Koo, H.C., Joo, Y.S. and Park, Y.H., 2005. Staphylococcal cassette chromosome mec (SCCmec) characterization and molecular analysis for methicillin-resistant *Staphylococcus aureus* and novel SCCmec subtype IVg isolated from bovine milk in Korea. *Journal of Antimicrobial Chemotherapy*, 56(4), pp.624-632.

474

475

476

477

478

18. Ito, T., Okuma, K., Ma, X.X., Yuzawa, H. and Hiramatsu, K., 2003. Insights on antibiotic resistance of *Staphylococcus aureus* from its whole genome: genomic island SCC. *Drug Resistance Updates*, 6(1), pp.41-52.

479

480

481

482

19. Oliveira, D.C., Milheiriço, C. and de Lencastre, H., 2006. Redefining a structural variant of staphylococcal cassette chromosome mec, SCCmec type VI. *Antimicrobial agents and chemotherapy*, 50(10), pp.3457-3459.

483

484

20. Zhang, K., McClure, J.A., Elsayed, S. and Conly, J.M., 2009. Novel staphylococcal cassette chromosome mec type, tentatively designated type VIII, harboring class A

- 485 mec and type 4 ccr gene complexes in a Canadian epidemic strain of methicillin-
486 resistant *Staphylococcus aureus*. *Antimicrobial agents and chemotherapy*, 53(2),
487 pp.531-540.
- 488
489
- 490 21. Li, S., Skov, R.L., Han, X., Larsen, A.R., Larsen, J., Sørum, M., Wulf, M., Voss, A.,
491 Hiramatsu, K. and Ito, T., 2011. Novel types of staphylococcal cassette chromosome
492 mec elements identified in clonal complex 398 methicillin-resistant *Staphylococcus*
493 *aureus* strains. *Antimicrobial agents and chemotherapy*, 55(6), pp.3046-3050.
- 494
- 495 22. García-Álvarez, L., Holden, M.T., Lindsay, H., Webb, C.R., Brown, D.F., Curran,
496 M.D., Walpole, E., Brooks, K., Pickard, D.J., Teale, C. and Parkhill, J., 2011.
497 Methicillin-resistant *Staphylococcus aureus* with a novel mecA homologue in human
498 and bovine populations in the UK and Denmark: a descriptive study. *The Lancet*
499 *infectious diseases*, 11(8), pp.595-603.
- 500
501
- 502 23. Hentschel, U., and Hacker, J., 2001. Pathogenicity islands: the tip of the
503 iceberg. *Microbes and infection*, 3(7), pp.545-548.
- 504
- 505 24. Dobrindt, U., Hochhut, B., Hentschel, U. and Hacker, J., 2004. Genomic islands in
506 pathogenic and environmental microorganisms. *Nature Reviews Microbiology*, 2(5),
507 pp.414-424.
- 508
509
- 510 25. Gill, S.R., Fouts, D.E., Archer, G.L., Mongodin, E.F., DeBoy, R.T., Ravel, J.,
511 Paulsen, I.T., Kolonay, J.F., Brinkac, L., Beanan, M. and Dodson, R.J., 2005. Insights
512 on the evolution of virulence and resistance from the complete genome analysis of an
513 early methicillin-resistant *Staphylococcus aureus* strain and a biofilm-producing
514 methicillin-resistant *Staphylococcus epidermidis* strain. *Journal of*
515 *bacteriology*, 187(7), pp.2426-2438.
- 516
- 517 26. Baba, T., Bae, T., Schneewind, O., Takeuchi, F. and Hiramatsu, K., 2008. Genome
518 sequence of *Staphylococcus aureus* strain Newman and comparative analysis of
519 staphylococcal genomes: polymorphism and evolution of two major pathogenicity
520 islands. *Journal of bacteriology*, 190(1), pp.300-310.
- 521
522
- 523 27. Lina, G., Bohach, G.A., Nair, S.P., Hiramatsu, K., Jouvin-Marche, E. and Mariuzza,
524 R., 2004. Standard nomenclature for the superantigens expressed by
525 *Staphylococcus*. *Journal of Infectious Diseases*, 189(12), pp.2334-2336.

526

527

28. Holden, M.T., Hsu, L.Y., Kurt, K., Weinert, L.A., Mather, A.E., Harris, S.R., Strommenger, B., Layer, F., Witte, W., de Lencastre, H. and Skov, R., 2013. A genomic portrait of the emergence, evolution, and global spread of a methicillin-resistant *Staphylococcus aureus* pandemic. *Genome research*, 23(4), pp.653-664.

531

532

29. Tsuru, T. and Kobayashi, I., 2008. Multiple genome comparisons within a bacterial species reveals a unit of evolution spanning two adjacent genes in a tandem paralog cluster. *Molecular biology and evolution*, 25(11), pp.2457-2473.

535

536

537

30. Novick, R.P., 2003. Mobile genetic elements and bacterial toxinoses: the superantigen-encoding pathogenicity islands of *Staphylococcus aureus*. *Plasmid*, 49(2), pp.93-105.

540

541

31. Deghorain, M. and Van Melderen, L., 2012. The Staphylococci phages family: an overview. *Viruses*, 4(12), pp.3316-3335.

543

544

545

32. Mir-Sanchis, I., Martínez-Rubio, R., Martí, M., Chen, J., Lasa, Í., Novick, R.P., Tormo-Más, M.Á. and Penadés, J.R., 2012. Control of *Staphylococcus aureus* pathogenicity island excision. *Molecular microbiology*, 85(5), pp.833-845.

548

549

33. Ram, G., Chen, J., Kumar, K., Ross, H.F., Ubeda, C., Damle, P.K., Lane, K.D., Penadés, J.R., Christie, G.E. and Novick, R.P., 2012. Staphylococcal pathogenicity island interference with helper phage reproduction is a paradigm of molecular parasitism. *Proceedings of the National Academy of Sciences*, 109(40), pp.16300-16305.

554

555

556

34. Pinchuk, I.V., Beswick, E.J. and Reyes, V.E., 2010. Staphylococcal enterotoxins. *Toxins*, 2(8), pp.2177-2197.

558

559

35. Ortega, E., Abriouel, H., Lucas, R. and Gálvez, A., 2010. Multiple roles of *Staphylococcus aureus* enterotoxins: pathogenicity, superantigenic activity, and correlation to antibiotic resistance. *Toxins*, 2(8), pp.2117-2131.

562

563

564

36. Tormo-Más, M.Á., Mir, I., Shrestha, A., Tallent, S.M., Campoy, S., Lasa, Í., Barbé, J., Novick, R.P., Christie, G.E. and Penadés, J.R., 2010. Moonlighting bacteriophage

565

- 566 proteins derepress staphylococcal pathogenicity islands. *Nature*, 465(7299), pp.779-
567 782.
- 568
- 569 37. Schelin, J., Wallin-Carlquist, N., Thorup Cohn, M., Lindqvist, R. and Barker, G.C.,
570 2011. The formation of *Staphylococcus aureus* enterotoxin in food environments and
571 advances in risk assessment. *Virulence*, 2(6), pp.580-592.
- 572
- 573
- 574 38. Todar, Kenneth. (2012). "Bacterial Protein Toxins". Todar's Online Textbook of
575 Bacteriology. Madison, Wisconsin.
- 576
- 577 39. Strommenger, B., Bartels, M.D., Kurt, K., Layer, F., Rohde, S.M., Boye, K., Westh,
578 H., Witte, W., De Lencastre, H. and Nübel, U., 2013. Evolution of methicillin-
579 resistant *Staphylococcus aureus* towards increasing resistance. *Journal of*
580 *Antimicrobial Chemotherapy*, p. dkt413.
- 581
- 582
- 583 40. Stryjewski, M.E., and Corey, G.R., 2014. Methicillin-resistant *Staphylococcus*
584 *aureus*: an evolving pathogen. *Clinical infectious diseases*, 58(suppl 1), pp. S10-S19.
- 585
- 586 41. Liebowitz, L.D., and Blunt, M.C., 2008. Modification in prescribing practices for
587 third-generation cephalosporins and ciprofloxacin is associated with a reduction in
588 methicillin-resistant *Staphylococcus aureus* bacteraemia rate. *Journal of hospital*
589 *infection*, 69(4), pp.328-336.
- 590
- 591
- 592 42. Fitzgerald, J.R., 2014. Evolution of *Staphylococcus aureus* during human colonization
593 and infection. *Infection, genetics, and evolution*, 21, pp.542-547.
- 594
- 595 43. Price, L.B., Stegger, M., Hasman, H., Aziz, M., Larsen, J., Andersen, P.S., Pearson,
596 T., Waters, A.E., Foster, J.T., Schupp, J. and Gillece, J., 2012. *Staphylococcus aureus*
597 CC398: host adaptation and emergence of methicillin resistance in
598 livestock. *MBio*, 3(1), pp.e00305-11.
- 599
- 600
- 601 44. Spoor, L.E., McAdam, P.R., Weinert, L.A., Rambaut, A., Hasman, H., Aarestrup,
602 F.M., Kearns, A.M., Larsen, A.R., Skov, R.L. and Fitzgerald, J.R., 2013. Livestock
603 origin for a human pandemic clone of community-associated methicillin-resistant
604 *Staphylococcus aureus*. *MBio*, 4(4), pp. e00356-13.
- 605

- 606 45. Moodley, A., Nielsen, S.S. and Guardabassi, L., 2011. Effects of tetracycline and zinc
607 on the selection of methicillin-resistant *Staphylococcus aureus* (MRSA) sequence
608 type 398 in pigs. *Veterinary microbiology*, 152(3), pp.420-423.
609
610
- 611 46. Larsen, J., Petersen, A., Sørum, M., Stegger, M., van Alphen, L., Valentiner-Branth,
612 P., Knudsen, L.K., Larsen, L.S., Feingold, B., Price, L.B. and Andersen, P.S., 2015.
613 Methicillin-resistant *Staphylococcus aureus* CC398 is an increasing cause of disease in
614 people with no livestock contact in Denmark, 1999 to 2011. *Euro surveillance:
615 bulletin Europeen sur les maladies transmissibles= European communicable disease
616 bulletin*, 20(37).
617
- 618 47. Lekkerkerk, W.S.N., Van Wamel, W.J.B., Snijders, S.V., Willems, R.J., van
619 Duijkeren, E., Broens, E.M., Wagenaar, J.A., Lindsay, J.A. and Vos, M.C., 2015.
620 What is the origin of livestock-associated methicillin-resistant *Staphylococcus aureus*
621 clonal complex 398 isolates from humans without livestock contact? An
622 epidemiological and genetic analysis. *Journal of clinical microbiology*, 53(6),
623 pp.1836-1841.
624
- 625 48. Rubinstein, E. and Keynan, Y., 2014. Vancomycin revisited—60 years later. *Frontiers
626 in public health*, 2, p.217.
627
628
- 629 49. Cui, L., Iwamoto, A., Lian, J.Q., Neoh, H.M., Maruyama, T., Horikawa, Y. and
630 Hiramatsu, K., 2006. Novel mechanism of antibiotic resistance originating in
631 vancomycin-intermediate *Staphylococcus aureus*. *Antimicrobial agents and
632 chemotherapy*, 50(2), pp.428-438.
633
- 634 50. Howden, B.P., McEvoy, C.R., Allen, D.L., Chua, K., Gao, W., Harrison, P.F., Bell, J.,
635 Coombs, G., Bennett-Wood, V., Porter, J.L. and Robins-Browne, R., 2011. Evolution
636 of multidrug resistance during *Staphylococcus aureus* infection involves mutation of
637 the essential two component regulator WalKR. *PLoS Pathog*, 7(11), p. e1002359.
638
639
- 640 51. Utaida, S., Pfeltz, R.F., Jayaswal, R.K. and Wilkinson, B.J., 2006. Autolytic
641 properties of glycopeptide-intermediate *Staphylococcus aureus* Mu50. *Antimicrobial
642 agents and chemotherapy*, 50(4), pp.1541-1545.
643
- 644 52. Shang, W., Hu, Q., Yuan, W., Cheng, H., Yang, J., Hu, Z., Yuan, J., Zhang, X., Peng,
645 H., Yang, Y. and Hu, X., 2016. Comparative fitness and determinants for the

646 characteristic drug resistance of ST239-MRSA-III-t030 and ST239-MRSA-III-t037
647 strains isolated in China. *Microbial Drug Resistance*, 22(3), pp.185-192.

648

649

650 53. Peleg, A.Y., Monga, D., Pillai, S., Mylonakis, E., Moellering, R.C. and Eliopoulos,
651 G.M., 2009. Reduced susceptibility to vancomycin influences pathogenicity in
652 *Staphylococcus aureus* infection. *Journal of Infectious Diseases*, 199(4), pp.532-536.

653

654 54. Gardete, S., Kim, C., Hartmann, B.M., Mwangi, M., Roux, C.M., Dunman, P.M.,
655 Chambers, H.F. and Tomasz, A., 2012. Genetic pathway in acquisition and loss of
656 vancomycin resistance in a methicillin resistant *Staphylococcus aureus* (MRSA)
657 strain of clonal type USA300. *PLoS Pathog*, 8(2), p.e1002505.

658

659

660 55. Mwangi, M.M., Wu, S.W., Zhou, Y., Sieradzki, K., de Lencastre, H., Richardson, P.,
661 Bruce, D., Rubin, E., Myers, E., Siggia, E.D. and Tomasz, A., 2007. Tracking the in
662 vivo evolution of multidrug resistance in *Staphylococcus aureus* by whole-genome
663 sequencing. *Proceedings of the National Academy of Sciences*, 104(22), pp.9451-
664 9456.

665

666 56. Chang, S., Sievert, D.M., Hageman, J.C., Boulton, M.L., Tenover, F.C., Downes,
667 F.P., Shah, S., Rudrik, J.T., Pupp, G.R., Brown, W.J. and Cardo, D., Vancomycin-
668 Resistant *Staphylococcus aureus* Investigative Team. 2003. Infection with
669 vancomycin-resistant *Staphylococcus aureus* containing the vanA resistance gene. *N.*
670 *Engl. J. Med*, 348, pp.1342-1347.

671

672

673 57. Sievert, D.M., Rudrik, J.T., Patel, J.B., McDonald, L.C., Wilkins, M.J. and Hageman,
674 J.C., 2008. Vancomycin-resistant *Staphylococcus aureus* in the United States, 2002–
675 2006. *Clinical Infectious Diseases*, 46(5), pp.668-674.

676

677 58. Lowy, F.D., 2003. Antimicrobial resistance: the example of *Staphylococcus*
678 *aureus*. *The Journal of clinical investigation*, 111(9), pp.1265-1273.

679

680 59. Timothy, J., Foster, 2017. Antibiotic resistance in *Staphylococcus aureus*. Current
681 status and future prospects. *FEMS Microbiology Reviews*.

682

683

684

685

686

UNDER PEER REVIEW