

# **Prevalence of Asymptomatic Bacteriuria among Pregnant Women Attending Antenatal Care at the Korle-Bu Teaching Hospital Accra, Ghana: A cross-sectional study**

## **ABSTRACT**

**Background:** The significance of asymptomatic bacteriuria (ASB) relates to its potential to cause acute pyelonephritis, preterm labor and preterm rupture of the membranes. Additionally, it has been associated with clinical conditions such as anemia, preterm birth, low birth weight and perinatal mortality.

**Aim:** This study therefore determined the prevalence of ASB among pregnant women attending antenatal clinic.

**Materials and Methods:** A cross-sectional study using convenient sampling method was used to recruit 200 pregnant women attending antenatal care at the Korle-Bu Teaching Hospital (KBTH) from January – April, 2014. Bacteriological, urine chemistry and routine urine analyses as well as antibiotic susceptibility profile of bacterial isolates from mid-stream urine samples of pregnant women were carried out using appropriate standard methods. Variables were reported in mean, standard deviation, percentages and bar graph. Chi square test was used to establish statistical difference and association between variables where p-value <0.05 was considered statistically significant.

**Results:** Out of the total participants, 23 had asymptomatic bacteriuria but no prevalence of ASB was found among Muslims. The association between marital status and prevalence of ASB was significant (Pearson  $\chi^2 = 4.88$ , p-value = 0.027). Five bacterial isolates were obtained with *Escherichia coli* (43.6%) being the most prevalent organism. There were more negative reactions than positive reactions with regards to nitrite and leukocyte esterase determination. Both gram positive and negative isolates recorded high susceptibility to Gentamicin and Norfloxacin. However, Ciprofloxacin showed a high rate of resistance to only gram negative isolates whiles Chloramphenicol, Ampicillin and Amoxillin showed high resistance to gram positive isolates.

**Conclusions:** The overall prevalence of asymptomatic bacteriuria was 11.5% [95%CI: 7.4% - 16.8%] among the referral pregnant women attending KBTH with *Escherichia coli* being the most prevalent organism whiles the most susceptible antibiotics were Gentamicin and Norfloxacin.

**Keywords:** *Urinary tract infections, antibiotic susceptibility, asymptomatic bacteriuria (ASB), bacterial profile.*

16

## 17 1. INTRODUCTION

18

19 Urinary tract infections (UTI's) are diseases that affect any part of the urinary system being it  
20 the kidney, ureters, bladder or the urethra (Kriplain, Bukshee , & Rafan, 1993). Most UTI's  
21 affect the lower urinary tract (bladder and urethra) before they complicate to the upper tract  
22 level (Mezue, Ofong, Nmezi, & Ugochukwu-Obi, 2006). Urinary Tract Infections are a  
23 common problem encountered in pregnancy due to the morphological and physiological  
24 changes that take place in the genitourinary tract (Sharma, 1990). It is of two types,  
25 symptomatic and asymptomatic (Akinloye, Ogbolu, Akinloye, & Alli, 2006). Asymptomatic  
26 bacteriuria refers to a condition in which urine culture reveals a significant growth of  
27 pathogens greater than  $10^5$  bacteria per milliliter in the absence of urinary symptoms  
28 (Lavanya & Jogalakshmi, 2008).

29 Urinary Tract Infections affect all age group, but women particularly pregnant women are  
30 more susceptible than men due to their pregnancy condition and anatomically short urethra  
31 which is easily contaminated by fecal flora (Gupta, Sahm, Mayfield, & Stamm, 2001). A  
32 bacterium during pregnancy has a greater propensity of progressing to pyelonephritis due to  
33 the following; reduction of glycogen secretion to the intracellular cavity of the vagina  
34 (Patterson & Andriole, 1997), relaxation of muscles of the ureters (Aboderin, Ako-Nai,  
35 Zailani, Ajayi, & Adedosu, 2004) which causes backward flow of urine and the cortisol  
36 production from progesterone secretion (Maclean, 1997).

37 The significance of ASB relates to its potential to cause acute pyelonephritis, preterm labour  
38 and preterm rupture of the membranes, other clinical condition includes; anemia, preterm  
39 birth, low birth weight and perinatal mortality (J. Akerele, Abhulimen, & Okonofua, 2001;  
40 Patterson & Andriole, 1997). A study carried out on ASB showed a prevalence between 2-  
41 7% (Okpere, 2003) but may be as high as 10% in high risk population (Kacmaz, Cakir,  
42 Aksoy, & Biri 2006). Furthermore, prevalence of ASB conditions may be as twice as  
43 common in pregnant women with sickle cell trait and three times as common in pregnant  
44 women with diabetic mellitus status (Kiningham, 1993). Studies done so far show that the  
45 prevalence rate of ASB in developing countries is higher than that in the developed states  
46 (Hooton , Winter , Tiu , & Stamm 1995).

47 A hospital based study carried out in some developed states including Canada showed a  
48 prevalence rate of 4-7% as compared to that of developing countries like Nigeria (i.e. Benin  
49 City and Sagamu) and Ethiopia. In Ghana, a study conducted in Komfo Anokye Teaching  
50 Hospital (KATH) and University of Cape Coast (UCC) hospital showed prevalence rates of  
51 7.3% and 56.5% respectively (P. Akerele, Abhuliren, & Okonofua, 2001.; A. Alemu *et al.*,  
52 2013; A. Boye *et al.*, 2012; Turpin, Minkah, Danso, & Frimpong, 2007). Another report also  
53 showed that ASB occurs between 5-9% for both non-pregnant and pregnant women and if  
54 left untreated, can progress to symptomatic UTI's including acute cystitis and pyelonephritis;  
55 this progress to symptomatic UTI's is likely to be more in pregnant women than non-  
56 pregnant women. However, other conditions that increase the incidence include low socio-  
57 economic status, grand multi-parity and advance in age. A recommendation was made by  
58 the Infectious Diseases Society of America that there is a need to screen and treat all  
59 positive cultures during pregnancy. The study therefore determined the prevalence of ASB  
60 (Asymptomatic bacteriuria) in pregnant women attending antenatal clinic at the Korle-Bu  
61 Teaching Hospital. The research work also went ahead to ascertain the antibiotic  
62 susceptibility patterns of these significant bacteriuria.

63

## 64 2. MATERIAL AND METHODS

65

### 66 2.1 Study design / Study site

67 A cross-sectional study design was carried out to recruit pregnant women from January to  
68 April 2014 at the Obstetrics and Gynecology Department of Korle-Bu Teaching Hospital  
69 (KBTH). Korle-Bu Teaching Hospital is a premier health care facility in Ghana, it's one of the  
70 tertiary hospitals in the country and is located in the Accra metropolis in the Greater Accra  
71 region of Ghana.

### 72 2.2 Study participants/ Eligibility criteria

73 A total of 200 pregnant women attending antenatal clinic at KBTH within the study period  
74 were recruited using convenient sampling method. Pregnant women within the ages of 15-  
75 45years were included in this study, however, participants with signs and symptoms of  
76 Urinary tract infections (UTI's), kidney diseases, diabetes, urological surgery, carcinoma and  
77 spinal cord injuries were excluded.

### 78 2.3 Data Collection and Processing

#### 79 2.3.1 Urine Sample Collection

80 An interview was conducted to obtain information on age, marital status, level of education,  
81 area of residence, parity, gestational age, gravidity, body mass index, blood pressure,  
82 hemoglobin level, religion and many other socio-demographic characteristics.

83 Study participants after completing the interview were given labelled sterile universal  
84 containers by the interviewer and counselled on how to collect midstream urine.  
85 Approximately 15mls of urine were collected from each participant. The urine samples were  
86 transported immediately on ice to the Central laboratory, Department of Microbiology, KBTH  
87 for processing within four hours of collection to ensure that the pathogenic organisms  
88 present in the urine were isolated and also to avoid the possible proliferation of the  
89 pathogenic organisms.

#### 90 2.3.2 Urine Sample Processing

91 The samples were subjected to routine microscopy and culture. A dip stick test strip was first  
92 inserted into the urine samples to check for the presence of nitrites and leukocytes. The  
93 urine samples were further centrifuged and their sediments were observed under a  
94 microscope for the presence of pus cells, red blood cells, casts and epithelial cells. The  
95 Urine samples were also cultured on Cysteine Lactose electrolyte deficiency (CLED) agar  
96 media using the streak plate method by use of a calibrated inoculation loop of size  
97 0.001CFU/L, the plates were then incubated at 37°C and read within 18-24 hours for  
98 significant bacteriuria. Significant bacteriuria was defined as a single colony count of  $\geq 10^5$   
99 organisms. The organisms were isolated and identification of pathogens was done using  
100 standard biochemical and sugar fermentation tests. Antibiotic susceptibility tests were also  
101 conducted on them by plating on Muller Hinton agar employing the Kirby Bauer disc diffusion  
102 method.

103

104

## 2.4 Statistical Analysis

Data collected was entered to Microsoft Excel and further analyzed using SPSS version 22.0. Most variables were reported in frequencies and percentages while hemoglobin and bacteria isolates were reported in mean and bar graph respectively. Chi square and Fisher's exact test were also used to establish statistical difference and association between variables where  $p$ -value  $< 0.05$  was considered as statistically significant.

## 3. RESULTS AND DISCUSSION

**Table 1** depicts demographical data of 200 pregnant women enrolled in the study. Out of this, majority of participants (30.5%) were found to be in both 26-30 and 31-35 year groups and (90.5%) of participants being Christians. The proportions of married participants were also higher (88.0%). Additionally, (44.0%) and (49.0%) of participants had secondary education and in their second trimester respectively. The mean hemoglobin level was  $11.45 \pm 0.08$  while greater portions (42.0%) and (41.5%) of these pregnant women were overweight and obese correspondingly.

The overall prevalence of ASB was 23 (11.5%). Among the age ranges, 41-45 age group had the highest prevalence (28.6%) of ASB while the least (3.4%) was found among 36-40 age group. Prevalence of ASB was also found higher, (12.7%) and (25.0%) among pregnant women who were not married and Christians respectively. However, no prevalence of ASB was found among Muslims. Also, ASB prevalence was highest (18.8%) and (14.3%) among women who had no formal education and in their first trimester duration of pregnancy correspondingly. Pregnant women who were overweight had the highest (14.3%) prevalence of ASB. (**Table 1**)

The association between marital status and prevalence of ASB was significant (Pearson  $\chi^2 = 4.88$ ,  $p$ -value = 0.027). Also, the relationship between haemoglobin level and prevalence of ASB was significant ( $p = 0.018$ ). All the other demographic factors were not significant and had no association with prevalence of ASB. (**Table 1**)

133

**Table 1: Prevalence of ASB and demographic characteristics of study population**

Characteristic	Non-Bacteriuria N (%)	Bacteriuria N (%)	Total N (%)	$\chi^2$	$P$ -value
<b>Age (years)</b>					
15-20	11 (78.6)	3 (21.4)	14 (7.0)	6.513	0.259
21-25	25 (89.3)	3 (10.7)	28 (14.0)		
26-30	52 (85.2)	9 (14.8)	61 (30.5)		
31-35	56 (91.8)	5 (8.2)	61 (30.5)		
36-40	28 (96.6)	1 (3.4)	29 (14.5)		
41-45	5 (71.4)	2 (28.6)	7 (3.5)	4.884	0.027 <sup>*a</sup>
<b>Marital Status</b>					
Married	159 (90.3)	17 (9.7)	176 (88.0)		
Single	18 (75.0)	6 (25.0)	24 (12.0)	2.434	0.487
<b>Education</b>					
None	13 (81.2)	3 (18.8)	16 (8.0)		
Basic	55 (87.3)	8 (12.7)	63 (31.5)		

Secondary	81 (92.0)	7 (8.0)	88 (44.0)		
Tertiary	28 (84.8)	5 (15.2)	33 (16.5)		
<b>Trimester</b>				1.015	0.602
1 <sup>st</sup>	6 (85.7)	1 (14.3)	7 (3.5)		
2 <sup>nd</sup>	89 (90.8)	9 (9.2)	98 (49.0)		
3 <sup>rd</sup>	82 (86.3)	13 (13.7)	95 (47.5)		
<b>Religion</b>				2.728	0.099
Christian	158 (87.3)	23 (12.7)	181 (90.5)		
Muslim	19 (100.0)	0 (0.0)	19 (9.5)		
<b>BMI</b>				1.997	0.850
Underweight	1 (100.0)	0 (0.0)	1 (0.5)		
Normal	28 (87.5)	4 (12.5)	32 (16.0)		
Overweight	72 (85.7)	12 (14.3)	84 (42.0)		
Obese	76 (91.6)	7 (8.4)	83 (41.5)		
<b>ASB Status</b>	177 (88.5)	23 (11.5)	200 (100.0)	-	-
<b>Hb g/dl</b>	11.45±0.08	10.76±1.35	10.83±1.32		0.018 <sup>*a</sup>

135 **N: Number; \* Significant at  $p < 0.05$ ; ASB: Asymptomatic Bacteriuria; all p-values**  
136 **calculated from fisher's exact test except those denoted by a.**  
137  
138


139 **Table 2** shows the dip stick examination for urine chemistry which revealed majority of the  
140 participants testing negative for both nitrite and leukocyte esterase test. However, 24(12.0%)  
141 and 38(19.0%) tested positive for nitrite and leukocytes esterase respectively.  
142

143 **Table 2: Urinary parameters among pregnant women**

Urine Chemistry Parameters	N (%)
<b>Leukocytes</b>	
Positive	38 (19.0)
Negative	162 (81.0)
<b>Nitrite</b>	
Positive	24 (12.0)
Negative	176 (88.0)

144 **N: Number**  
145  
146  
147

148 **Figure 1** represents frequency of the bacteria isolates, the predominant isolate was *E coli*  
149 (43.6%), followed by *CoNS*, *Citrobacter spp*, *Enterococcus spp* and *Klebsiella spp* as  
150 (26.1%), (21.7%), (4.3%) and (4.3%) respectively.



**Figure 1:** Frequency of bacteriuria isolate among pregnant women

With respect to gram negative isolates, the antimicrobial susceptibility pattern showed Gentamicin and Norfloxacin to be the most effective antibiotics whereas Ciprofloxacin was found to be the most resistant antibiotic. On the other hand, for overall gram positive isolates, Gentamicin and Norfloxacin showed the most sensitivity rates while the most resistant drugs were Chloramphenicol, Ampicillin and Amoxillin. (Table 3)

163 **Table 3: Antimicrobial susceptibility pattern of bacteria isolates from urine culture of**  
164 **pregnant women**

ISOLATES	N		ANTIBIOTICS							
			GEN	NOR	AMO	OFLO	TET	CRP	AMP	COT
Gram Negative										
CoNS	6	S	6(100.0)	6(100.0)	1(16.7)	5(83.3)	1(16.7)	0(0.00)	2(33.3)	5(83.3)
		R	0(0.00)	0(0.00)	5(83.3)	1(16.7)	5(83.3)	6(100.0)	4(66.7)	1(16.7)
Gram Positive										
Citrobacter	5	S	5(100.0)	5(100.0)	0(0.0)	3(60.0)	2(40.0)	0(0.0)	0(0.0)	4(80.0)
		R	0(0.00)	0(0.00)	5(100.0)	2(40.0)	3(60.0)	5(100.0)	5(100.0)	1(20.0)
E.coli	10	S	10(100.0)	10(100.0)	0(0.0)	3(30.0)	3(30.0)	0(0.0)	0(0.0)	8(80.0)
		R	0(0.0)	0(0.0)	10(0.0)	7(70.0)	7(70.0)	10(100.0)	10(100.0)	2(20.0)
Enterococcus	1	S	1(100.0)	1(100.0)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	0(0.00)	1(100.0)
		R	0(0.0)	0(0.00)	1(100.0)	1(100.0)	1(100.0)	1(100.0)	1(100.0)	0(0.00)
Klebsiella spp.	1	S	1(100.0)	1(100.0)	0(0.0)	1(100.0)	0(0.00)	0(0.00)	0(0.00)	1(100.0)
		R	0(0.0)	0(0.0)	1(100.0)	0(0.00)	1(100.0)	1(100.0)	1(100.0)	0(0.00)
TOTAL	17	S	17(100.0)	17(100.0)	0(0.0)	7(41.2)	5(29.4)	0(0.00)	0(0.00)	14(82.4)
		R	0(0.0)	0(0.0)	17(100.0)	10(58.8)	12(70.6)	17(100.0)	17(100.0)	3(17.6)

165 **N-Number: S-Sensitivity: R-Resistance: CoNS- Coagulase Negative Staphylococcus:**  
166 **E. coli- Escherichia coli: GEN-Gentamicin: NOR- Norfloxacin: AMO-Amoxicillin: OFLO-**  
167 **Ofloxacin: CIP-Ciprofloxacin: TET-Tetracycline: CRP- Chloramphenicol: AMK-**  
168 **Amikacin: AMP-Ampicillin: COT: Cotrimoxazole and CEF-Ceftriaxone**

173 Prevalence of ASB revealed from the study among pregnant women was 11.5% [95%CI:  
174 7.4% - 16.8%] which was coherent with a similar study conducted in Dhaka by Jubaida and  
175 colleagues whose study found a prevalence of 10.2% (Jubaida *et al.*, 2013). However, other  
176 studies revealed contrast results; a recent similar study conducted in Iran revealed a  
177 prevalence of 6.5% for significant ASB which is relatively lower compared to the prevalence  
178 from this study (Etminan-Bakhsh, Tadi, & Darabi, 2017). This relatively low prevalence might  
179 be due to screening and treatment being carried out irrespective of symptoms by the  
180 subjects which is not usually practiced in our setting. Again, a study by Labi and colleagues  
181 predicted a far lower prevalence of 5.5% per their study (Labi, Yawson, Ganyaglo, &  
182 Newman, 2015). Other studies showed relatively higher prevalence rates compared to that  
183 from this study (J. Akerele *et al.*, 2001; A. Boye *et al.*, 2012; Imade, Izekor, Eghafona,  
184 Enabulele, & Ophori, 2010.; Mokube, Atashili, Halle-Ekane, Ikomey, & Ndumbe 2013) who  
185 reported 29.5%, 86.6%, 56.5%, 23.5%, and 45.3% respectively. Inconsistency in the

prevalence rates from these various studies may be due to differences in the socio-demographic characteristics of the study population.

Out of the overall prevalence, *Escherichia coli* emerged the predominant bacteria isolated with a prevalence of 43.6%. This result is comparable with other studies which also revealed *E. coli* as the most prevalent isolate (Kerure, Surpur, Sagarad, & Hegadi, 2013; Senthinath *et al.*, 2013; Sheiner, Mazor-Drey, & Levy, 2009). On the contrary, the result is dissimilar to a study conducted in Nigeria where *Klebsiella species* was the most prevalent organism isolated (Izuchukwu, Oranu, Bassey, & Orazulike, 2017). Another study found *S. aureus* to be the predominant bacterium isolated (Olamijulo, Adewale, & Olaleye, 2016). There was a significant association between marital status and asymptomatic bacteriuria ( $P=0.027$ ), with the unmarried group having a relatively higher ASB compared to the married group, a finding different from a study by Olamijulo *et al.* (2016). The high amount of bacterial isolates (BI) among the unmarried group could be associated with sex activities before and after women conceived.

The laboratory analysis for Antibiotic Susceptibility Test (AST) showed that all five (5) isolates were most susceptible to Gentamicin and Norfloxacin. However, among the twelve (12) different drugs used for the study, the most resistant drugs were Ciprofloxacin, Chloramphenicol, Ampicillin and Amoxillin; Ciprofloxacin showed high resistance among the gram positive organisms while the rest predicted relatively low resistances among gram positive organisms. However, comparatively the highly susceptible antibiotics (Gentamicin, Norfloxacin, Ofloxacin, Cotrimoxazole and Ceftriaxone) were also found to be highly effective from studies by Okalla, Shu, Etoga, Mengue, and Adiogo (2018) in Cameroon, Alemu, Dagnew, Alem, and Gizachew (2013) in Ethiopia, Onanuga, Omeje, and Eboh (2018) in Bayelsa state Nigeria, A. Boye *et al.* (2012) and Turpin *et al.* (2007) in Ghana, and Mokube, Atashili, Halle-Ekane, Ikomey, and Ndumbe (2013) in Buea State.

On the contrary, Sujatha and Nawani (2014) and Alex Boye *et al.* (2012) found Tetracycline, Ampicillin, and Amikacin as susceptible to both gram positive and negative organisms. Ferrieres, Hancock, and Klemm (2007) stated that clear alternatives as a first line drug against UTI before culture and sensitivity were Gentamicin and Norfloxacin but Gentamicin was relatively safer in pregnancy and effective against most urinary tract infections (Kenechukwu, Chinekwu, Davidson, & Golibe 2005; Miglioli, Silini, Carzeri, Grabocka, & Allerberger 1999) but may cause nephrotoxicity and negative fetal conditions when used adversely and close to term during pregnancy (Noel, Tufon, Waindim, Akwo, & Bong, 2013). The high resistance among Penicillin related antibiotics might be associated with high rate of its abuse unlike the aminoglycosides and the fluoroquinolones whose low resistance may be due to their less availability and the wide range of antibiotic therapy.

The number of leukocyte esterase positive reactions of the urine dip stick was greater than nitrite reactions which is mostly used in most hospitals, clinics and some health centers for bacteria identification during routine urine examination. Detection of presence of bacteria by nitrite reaction using the urine dipstick was highly specific but lacked sensitivity resulting in consequence of having a number of false negatives. A method comparative study by Habiba Ibrahim and Yunusa, 2015 revealed similar results supporting the false negativity of the urine dipstick in detecting bacteria in urine. However, other studies showed contrasting results to that from this study (Okusanya, Aigere, Eigbefoh, Okome, & Gigi, 2014; Shelton, Boggess, Kirvan, Sedor, & Herbert, 2001). Clearly this shows that the dip stick cannot identify non-nitrate producing bacteria as well as urine that stays in the bladder for a short period before it is voided, thereby making it present with shortfalls as a routine test kit.



The results of this study support the suggestion that urine culture should be the method of choice for screening among pregnant women. Therefore, ASB identification among pregnant women should be treated as per antimicrobial sensitivity pattern of the isolate to prevent overlook of some bacterial growth thereby leading to maternal and perinatal morbidity.

Another observation was the presence of 23 Bacteria isolates among the Christian participants compared to none among the Muslims. This might be attributed to effective rinsing of the vulva following every episode of urination among Muslims, a practice in the religion. Furthermore, high bacteriuria was found among the 41-50 age group. This general revelation of ASB isolation being more in the older women compared to young females might be due to the action of hormones leading to increased deposition of glycogen within the smooth muscles. This then attracts lactobacillus activity which is glycogenphilic and thus colonizing the vaginal epithelium, making it acidic and hence preventing invasion of the region by other pathogens. Therefore, decreasing glycogen deposition, and reduction in the lactobacillus colonies as part of ageing process enhance bacterial adherence and invasion by pathogens, making them more susceptible to UTI (Eschenbach *et al.*, 1992). However, a similar study found high bacteria isolates among 21-25 age group which was likely to be an active sex group (A. Boye *et al.*, 2012). The anatomical relationship of the female urethra to the vagina makes it liable to trauma during sexual intercourse and this can increase bacteriuria. (A. Boye *et al.*, 2012).

Prevalence of ASB was found to be high among pregnant women in their 1<sup>st</sup> and 3<sup>rd</sup> trimesters from this study. A previous study carried out by Olusanya and Olutiola (1984) also confirms the high prevalence among the third trimester group while another study also found a high ASB isolates among the 2<sup>nd</sup> trimester group (A. Boye *et al.*, 2012). Finally, hemoglobin was found to be significantly associated with bacteriuria ( $P=0.018$ ). This clinical condition might be due to the production of polysaccharides, endotoxins and particularly hemolysis by most of the bacteria which causes tissue such red blood cells damage.

#### 4. CONCLUSION AND RECOMMENDATIONS

Overall prevalence of asymptomatic bacteriuria was 11.5% [95%CI: 7.4% - 16.8%] among the referral pregnant women attending KBTH with *E. coli* 43.6% being the predominant isolate followed by CoNS 26.7% which were all susceptible to Gentamicin and Norfloxacin. The study showed clearly that urine dip stick method is not a “gold” process for bacteria identification since it cannot isolate most of the gram positive bacteria.

The study recommends that screening and treatment of asymptomatic bacteriuria in pregnancy should be an integral part of obstetric care and should be included in all antenatal guidelines. Also, all urine samples from pregnant women should be cultured for ASB isolation and hence an appropriate drug selection via AST process. Furthermore, urine culture should serve as the gold standard as against the urine dipstick method and be practiced among all facilities to help reduce incidence of misdiagnosis. Lastly, from this study we recommend and advocate that all Penicillin related antibiotics should not be used for the treatment of ASB.

#### LIMITATION

Some participants might have not been able to adequately follow the mid-stream urine collection.

291 **COMPETING INTERESTS**

292

293 The authors hereby declare that there are no competing interests regarding the publication  
294 of this article.

295

296

297 **CONSENT (WHERE EVER APPLICABLE)**

298

299 Written informed consent was obtained from participants before the study took place and  
300 these documents are preserved by the authors.

301

302 **ETHICAL APPROVAL (WHERE EVER APPLICABLE)**

303

304 Ethical clearance was sought from the Institutional Review Board of the University of Cape  
305 Coast (IRB/UCC) and administrators of the Hospital before the study was commenced. In  
306 addition, all pregnant women recruited into the study gave their informed consent after  
307 thorough explanation of the rationale of the study.

308

309 **REFERENCES**

310

311

312

Aboderin, A. O., Ako-Nai, A. K., Zailani, S. B., Ajayi, A., & Adedosu, A. N. (2004). A Study of Asymptomatic  
Bacteriuria in Pregnancy in Ile-Ife, Southwestern Nigeria. . *Afr J Exper. Microbiol*, 5, 252-259.

314

Akerele, J., Abhulimen, P., & Okonofua, F. (2001). Prevalence of asymptomatic bacteriuria among pregnant  
women in Benin City, Nigeria. *J Obstet Gynaecol*, 21(2), 141-144.

316

Akerele, P., Abhulimen, F., & Okonofua, J. (2001.). Prevalence of asymptomatic bacteriuria among pregnant  
women in Benin city. Nigeria . *J. Obstruct. Gynaecol*, 21(2), 141-144.

318

Akinloye, O., Ogbolu, D. O., Akinloye, O. M., & Alli, O. A. T. (2006). Asymptomatic bacteriuria of pregnancy in  
Ibadan, Nigeria: a re-assessment. . *Br J Biomed Sci.*, 63.

320

Alemu, A., Dagnaw, M., Alem, M., & Gizachew, M. (2013). Uropathogenic bacterial isolates and their  
antimicrobial susceptibility patterns among HIV/AIDS patients attending Gondar University Specialized  
Hospital, Gondar, Northwest Ethiopia. *Journal of Microbiology Research and Reviews*, 1(4), 42-51.

323

Alemu, A., Moges, F., Shiferaw, Y., Tafess, K., Kassu, A., Anagaw, B., & Agegn, A. (2013). Prevalence and  
antimicrobial susceptibility of asymptomatic significant bacteriuria among new antenatal enrollees in  
Southwest, Nigeria. *International Research Journals of Microbiology*, 4(8), 197-203.

326

Boye, A., Siakwa, P. M., Boampong, J. N., Koffuor, G. A., Ephraim, R. K. D., Amoateng, P., . . . Penu, D.  
(2012). Asymptomatic urinary tract infections in pregnant women attending antenatal clinic in Cape  
Coast, Ghana. *E3 J Med Res*, 1(6), 74-83.

329

Boye, A., Siakwa, P. M., Boampong, J. N., Koffuor, G. A., Ephraim, R. K. D., Amoateng, P., . . . Penu, D.  
(2012). Asymptomatic urinary tract infections in pregnant women attending antenatal clinic in Cape  
Coast, Ghana. *E3 Journal of Medical Research*, 1(6), 74-83.

332

Eschenbach et. al. (1992). Asymptomatic Bacteriuria among pregnant women. *Clinical Microb. J*, 27, 251-256.

333

Etminan-Bakhsh, M., Tadi, S., & Darabi, R. (2017). Asymptomatic bacteriuria in pregnant women attending  
Boo-Ali Hospital Tehran Iran: Urine analysis vs. urine culture. *Electronic physician*, 9(11), 5760.

335

Ferrieres, L., Hancock, V., & Klemm, P. (2007). Biofilm exclusion of uropathogenic bacteria by selected  
asymptomatic bacteriuria Escherichia coli strains. *Microbiology*, 153(6), 1711-1719.

337

Gupta, K., Sahm, D. F., Mayfield, D., & Stamm, W. E. (2001). Antimicrobial resistance among uropathogens  
that cause community-acquired urinary tract infections in women: a nationwide analysis. *Clin Infect  
Dis.*, 33, 89-94.

338

339

- Hooton , T. M., Winter , C., Tiu , F., & Stamm , W. E. (1995). Randomized comparative trial and cost analysis of 3-day antimicrobial regimens for treatment of acute cystitis in women. *JAMA*, 273, 41-45.
- Imade, P. E., Izekeor, P. E., Eghafona, N. O., Enabulele, O. I., & Ophori, E. (2010,). Asymptomatic bacteriuria among pregnant women. *North American Journal of Medical Sciences*, 2, 263.
- Izuchukwu, K. E., Oranu, E. O., Bassey, G., & Orazulike, N. C. (2017). Maternofetal outcome of asymptomatic bacteriuria among pregnant women in a Nigerian Teaching Hospital. *Pan African Medical Journal*, 27(1).
- Jubaida, N., Kawsar, N. M., Elora, N., Rahimgir, M., Shapla, N. R., & Muid, S. A. (2013). Prevalence of asymptomatic bacteriuria in pregnant women. *Journal of Armed Forces Medical College, Bangladesh*, 9(2), 64-69.
- Kacmaz, B., Cakir, O., Aksoy, A., & Biri , A. (2006). Evaluation of rapid urine screening test to detect asymptomatic bacteriuria in pregnancy. 2006, 59(4), 261-263. (Put the name of the journal)
- Kenechukwu , M., Chinekwu , O., Davidson , N., & Golibe , U. (2005). Antibiotic sensitivity patterns in urinary tract infections at the tertiary Hospital. *Medikka(Journal of the University of Nigeria Medical Students)*.
- Kerure, S. B., Surpur, R., Sagarad, S. S., & Hegadi, S. (2013). Asymptomatic bacteriuria among pregnant women. *Int J Reprod Contracept Obstet Gynecol*, 2(2), 213-216.
- Kiningham, R. B. (1993). Asymptomatic bacteraemia in pregnancy. *Am Fam Physician*, 47(5), 1232-1238.
- Kriplain, A., Bukshee , K., & Rafan, A. (1993). Asymptomatic bacteriuria in pregnant women,India *Obstet Gynaecol India*, 43, 489-491.
- Labi, A., Yawson, A., Ganyaglo, G., & Newman, M. (2015). Prevalence and associated risk factors of asymptomatic bacteriuria in ante-natal clients in a large teaching hospital in Ghana. *Ghana medical journal*, 49(3), 154-158.
- Lavanya, S. V., & Jogalakshmi, D. (2008). Asymptomatic bacteriuria in antenatal women. *Indian J. Med. Microbiol.*, 20(2), 105-106.
- Macleane, A. B. (1997). Urinary tract infection in pregnancy,. *Br J Urol.*, 7(80), 10-30.
- Mezue, K., Ofong, C., Nmezi, D., & Ugochukwu-Obi, G. (2006). Antibiotic Sensitivity patterns in urinary tract infections at a Tertiary Hospital. *Journal of the University Of Nigeria Medical students*, 12(2).
- Miglioli , P. A., Silini , R., Carzeri , O., Grabocka , E., & Allerberger , F. (1999). Antibacterial activity of gentamicin and ciprofloxacin against gram negative bacteria: . *Pharmacological Research*, 39(4), 321-323.
- Mokube, M. N., Atashili, J., Halle-Ekane, G. E., Ikomey, G. M., & Ndumbe, P. M. (2013). Bacteriuria amongst pregnant women in the Buea Health District, Cameroon: Prevalence, predictors, antibiotic susceptibility patterns and diagnosis. *PloS one*, 8(8), e71086.
- Mokube , M. N., Atashili, J., Halle-Ekane , G. E., Ikomey, G. M., & Ndumbe , P. M. (2013). Bacteriuria amongst Pregnant Women in the Buea Health District, Cameroon: Prevalence, Predictors, Antibiotic Susceptibility Patterns and Diagnosis. . *PLoS ONE*, 8(8).
- Noel , N. N., Tufon, E. N., Waindim, N. Y., Akwo, T. T. C., & Bong, R. N. (2013). Prevalence and Antimicrobial Sensitivity Pattern Amongst Pregnant Women Attending the Northwest Regional Hospital,Bamenda. *VRI Cell signaling J*, 1(1), 3-6.
- Okalla, C. E., Shu, E. Y., Etoga, M. E., Mengue, E. R., & Adiogo, D. (2018). Bacterial and fungal uropathogens in diabetic patients and their susceptibility patterns: case study of two hospitals in douala. *Asian Journal of Microbiology and Biotechnology*, 3(3), 79-89.
- Okpere, E. E. (2003). *Medical Disorders in Pregnancy* (E. Okpere Ed.): Uniben press.
- Okusanya, B., Aigere, E., Eigbefoh, J., Okome, G., & Gigi, C. (2014). Is a chlorhexidine reaction test better than dipsticks to detect asymptomatic bacteriuria in pregnancy? *Journal of Obstetrics and Gynaecology*, 34(1), 21-24.
- Olamijulo, J. A., Adewale, C. O., & Olaleye, O. (2016). Asymptomatic bacteriuria among antenatal women in Lagos. *Journal of Obstetrics and Gynaecology*, 36(6), 722-725.
- Olusanya , O., & Olutiola , P. O. (1984). Studies on bacteriuria in patients and students in Ile-Ife, Nigeria. . *West Afr J Med.*, 303, 177-182.
- Onanuga, A., Omeje, M. C., & Eboh, D. D. (2018). Carriage of multi-drug resistant urobacteria by asymptomatic pregnant women in Yenagoa, Bayelsa State, Nigeria. *African journal of infectious diseases*, 12(2), 14-20.
- Patterson, T. F., & Andriole, V. T. (1997). Bacteriuria in Pregnancy. *American Academy of Family Physician*, 1, 807-822.

- Senthinath, T. J., Rajalaksmi, P. C., Keerthana, R., Vigneshwari, R., Revathi, P., Prabhu, N., & Susethira, A. (2013). Original Research Article Prevalence of asymptomatic bacteriuria among antenatal women in rural tertiary care hospital, Tamilnadu, India. *Int. J. Curr. Microbiol. App. Sci*, 2(1), 80-85.
- Sharma, J. B. (1990). Prevalence of significant bacteriuria in preterm labour. *Obstet Gynaecol India*, 40, 336-338.
- Sheiner, E., Mazor-Drey, E., & Levy, A. (2009). Asymptomatic bacteriuria during pregnancy. *The journal of maternal-fetal & neonatal medicine*, 22(5), 423-427.
- Shelton, S. D., Boggess, K. A., Kirvan, K., Sedor, F., & Herbert, W. N. (2001). Urinary interleukin-8 with asymptomatic bacteriuria in pregnancy. *Obstetrics & Gynecology*, 97(4), 583-586.
- Sujatha, R., & Nawani, M. (2014). Prevalence of asymptomatic bacteriuria and its antibacterial susceptibility pattern among pregnant women attending the antenatal clinic at Kanpur, India. *Journal of clinical and diagnostic research: JCDR*, 8(4), DC01.
- Turpin, C., Minkah, B., Danso, K., & Frimpong, E. (2007). Asymptomatic bacteriuria in pregnant women attending antenatal clinic at Komfo Anokye Teaching Hospital, Kumasi, Ghana. *Ghana medical journal*, 41, 26.