

**HAEMATOLOGICAL AND HISTOLOGICAL ASSESSMENT OF JUVENILES OF
Chrysichthys nigrodigitatus IN OGBESE RIVER, ONDO STATE, NIGERIA**

ABSTRACT

Chrysichthys nigrodigitatus catfish is a fish of economic importance in sub-sahara Africa. In Ogbese town, and its environs, it constitutes a means of income and food for fisherfolks and community members. Hence, this study was undertaken to assess health status of *Chrysichthys nigrodigitatus* using heamatology and histological assessment of the fish specie due to the anthropogenic activities that takes place around the river body. A total 120 live fish samples of *Chrysichthys nigrodigitatus* were collected by the assistance of fisherfolks using fish cage at Ogbese River from May to August, 2018. Some water parameters measurements were taken: temperature, pH, DO, Turbidity and Conductivity. Morphometric measurement: Weight (g) and length (cm) of fish were taken. Heamatoloty and histology of fish gills, liver and intestine were determined. Mean water temperature ($27.70 \pm 0.18^\circ\text{C}$), pH (7.36 ± 0.22), DO (6.98 ± 0.15 mg/l), Turbidity (78.50 ± 13.53 NTU) and Conductivity (148.35 ± 27.98) of the river determined respectively. Mean body weight of fish was 148.15 ± 36.53 g, and mean length was 25.64 ± 2.86 cm. The of the fish specie were examined to assess the architecture of the organs. result of haematology studies of *C. nigrodigitatus* revealed high values in the parameters measured. Red Blood Cell was higher than the White Blood Cell with mean value of (225.63 ± 10.45 $10^3/\text{mm}^3$) while Eosinophils recorded lowest parameters with mean value of (1.75 ± 0.52 %). Results of histology of gills, liver and intestines showed that the gill filaments are eroded with a deformation of the cartilage core and also hyperplasia of the secondary lamellae. The intestines showed atrophy in a mucosal layer, hemorrhage and dilation within blood vessels and within serosa of mucosa and for liver, picnotic nucleus are shattered, the hepatocytes are ruptured and there is increased kupffer cell as a result of exposure to pollutants. The results indicated pollution level of the environment have significant impact on health status of fish.

KEYWORDS: *Chrysichthys nigrodigitatus*, Ogbese River, Haematology, Histology.

INTRODUCTION

Fish is one of the most important animal protein sources that are widely consumed by all races and classes of people (Abolude and Abdullahi, 2005). It compares favorably with milk, meat, pork and poultry (James, 1984). Fish and fishery products are highly nutritious and are excellent sources of other dietary essentials like vitamins and minerals. Fish fat contains a high proportion of polyunsaturated fatty acids which may help to decrease the incidence of atherosclerosis and heart related diseases (Akande, 2011). Fish also provide an important complement to the predominantly carbohydrate based diet of many people in Nigeria (Akande, 2011).

The silver catfish *Chrysichthys nigrodigitatus* (Lacepede, 1803) is a highly valued food-fish included among the dominant commercial catches exploited in Ogbese river, Ondo State, Nigeria. It is restricted to the bottom of deep water, omnivorous; consume bivalves, detritus, chironomid, crustaceans and vegetable matter (Bankole *et al.*, 2011). This fish can be raised in both fresh and brackish water environments.

43 Fish health can be adversely affected by temperature changes, habitat deterioration and aquatic
44 pollution (Skouras *et al.*, 2003). Hematological parameters are considered an important indicator
45 of fish health status, and provide valuable information to assess the fish welfare (Azevedo *et al.*,
46 2006). Hematology is also used as an indicator of physiological and pathological changes in fish
47 (Chekrabarty and Banerjee 1988, Martins *et al.*, 2008). It can be affected by several factors
48 including gonad maturation (Ranzani-Paiva and Godinho, 1985), dissolved oxygen alterations
49 (Ranzani-Paiva *et al.*, 2000), gender (Lusková, 1998), spawning and water temperature (Joshi
50 1982), lotic or lentic environment (Val *et al.*, 1985), handling stress and transportation (Gbore *et*
51 *al.*, 2006), fish inflammation (Martins *et al.*, 2006), size, feeding and stocking density (Rey
52 Vázquez and Guerrero, 2007), microbial infection and parasitism (Martins *et al.*, 2004, Azevedo
53 *et al.*, 2006. Jamalzadeh *et al.*, 2009).

54 Ogbese region comprises Ogbese community and some neighboring agrarian settlements that
55 sustain it with agricultural produce. The location of Ogbese in the rain forest zone in South
56 Western Nigeria gives it a natural tendency of wood, timber and food production in the region.
57 The community serves as an economic life wire of Akure North Local Government Area of
58 Ondo State that produces food crops in large quantities. With these economic potentials, the
59 town still remains a remote rural settlement in the State.

60 Pollution of the rivers examined in this study is mainly through run-off activities from
61 agricultural practices and commercial activities. Many studies have shown that very large
62 quantities of heavy metals are found in run-off associated with the operation of motor vehicles,
63 atmospheric fallout and road surface materials (Harper, 1985). To the environmental scientists,
64 the ultimate concern of trace metal contaminants in receiving water is their toxic impact on
65 aquatic organisms and fish species (Sutherland and Tolosa, 2000; De Carlo *et al.*, 2004).
66 Assessing pollutants in different components of the ecosystem is an important task in preventing
67 risk to natural life and public health. Pollutants entering these receiving waters by way of run-off
68 conveyance systems, indiscriminate dumping of wastes e.t.c, may adversely impact many of the
69 desired uses. The Ogbese community has undergone great economic development in recent
70 years. In fact, it is notably one of the fastest growing, economically important communities in
71 Ondo State and handles a considerable number of micro- industries. The very popular market
72 (Ogbese market) and the timber business coupled with unequalled agricultural practices have
73 drawn people from several cultural backgrounds in the country to make the settlement inter-
74 tribal. This increase in anthropogenic activities surrounding the area has lead to an increase in
75 environmental degradation. These multiple sources make it especially difficult to identify and
76 isolate the risks associated with this contaminated water. Unfortunately, records of water quality
77 parameters are non-existing and no known monitoring programmes on the water quality have
78 been initiated within the state.

79

80 **MATERIALS AND METHODS**

81 **Study Area**

82 The study site was Ayede, Ogbese River along Akure-Benin expressway in Ondo State. The area
83 lies between E6⁰SE8⁰ and longitude N4⁰N6⁰E. The river has its source from Ayede-Ekiti in Ekiti
84 state and flows through Ogbese in Ondo State to Edo State. The Ogbese community is about
85 10km east of Akure, the Ondo state capital.

86 **Collection of Water Samples**

87 Water samples were collected using water samplers at 10 cm depth at three points locations from
88 the river body, and parameters were determined using multi- parameter machine Model No: for
89 dsissolved oxygen, temperature, turbidity, conductivity, and pH.
90

91 **Collection of Fish**

92 120 live *Chrysichthys nigrodigitatus* fish samples were collected by the assistance of fisherfolks
93 using fish cage at Ogbese River from May to August, 2018. They were then transported alive in
94 buckets containing water to the Marine Biology Laboratory of the Department of Fisheries and
95 Aquaculture Technology, Federal University of Technology, Akure.

96 97 **Length-weight Measurement**

98 The weight in grams (g) of each specimen was taken using a digital weighing balance, which
99 was wiped dry between samples. Standard length was measured in centimeter (cm) using a meter
100 ruler.

101 Condition factor of the fish was assessed to know the state of being of the fish.

$$102 \quad K = \frac{100 \times W}{L^3}$$

103
104 K = Condition Factor

105 W = Body Weight of Fish in gram (g)

106 L = Standard Length of Fish in centimetre (cm)
107

108 **3.4 Haematological Analysis**

109 Blood samples were taken from the caudal vein of each fish using a syringe and transferred to
110 5ml of Ethylene Diamine Tetraacetic Acid (EDTA) bottles. After blood collection in the
111 laboratory, the samples were maintained on ice and sent to the laboratory of Animal Production
112 and Health Technology, Federal University of Technology, Akure for hematological analysis.

113 The haematological parametres analysed were; Erythrocyte Sedimentation Rate Count (ESR),
114 Packed Cell Volume Count (PCV), Red Blood Cell Count (RBC), Haemoglobin Concentration
115 (Hgb), White Blood Cell Count (WBC), Lymphocyte Count, Neutrophils Count, Monocytes
116 Count, Basophils Count, Eusonophils Count. Mean Corpuscular Volume (MCV), Mean
117 Corpuscular Haemoglobin (MCH) And Mean Corpuscular Haemoglobin Concentration (MCHC)
118 were calculated according to (Houston, 1990).

119 The Haemoglobin was calculated as: Hb (g/100ml) = Absorbance of test x Concentration of
120 standard Absorbance of standard Absorbance of standard Total erythrocyte (RBC)

121 Red Blood Cell and White Blood Cell were calculated thus; = C x D x 4000

122 Where;

123 C = dilution factor (20)

124 D = number of cells counted

125 Hematocrit/ PCV = $\frac{\text{Volume of packed red blood cell} \times 100}{\text{Volume of whole blood}}$
126

127

128 White blood cell (WBC) = %WBC X total WBC + thrombocytes counts
129

129

130 The red cell indices – MCHC, MCH and MCV were derived thus;
131

131

132 Mean Cell Hemoglobin Concentration (MCHC) = $\frac{\text{Hemoglobin (g/100ml)} \times 100}{\text{PCV(\%)}}$
 133

134
 135 Mean Corpuscular Haemoglobin (MCH) = $\frac{\text{Hemoglobin (g/100ml)} \times 100}{\text{RBC (x10,000rbc/mm}^3\text{)}}$
 136

137 Mean Cell Volume (MCV) = $\frac{\text{PCV} \times 100}{\text{RBC (x10,000rbc/mm}^3\text{)}}$
 138
 139

140
 141 **3.5 Histological Analysis**

142 The fish specimen was dissected using a dissecting set. The gills, liver and intestines were then
 143 removed and rinsed in distilled water to remove blood stains. The organs were then placed in a
 144 10ml sample bottle with 10% formalin for preservation and transported to the Anatomy and
 145 Veterinary Laboratory at the University of Ibadan for Histological Analysis.
 146

147 **3.6. Statistical Analysis**

148 Data collected were analysed using one-way ANOVA. Further tests were done using Duncan
 149 Multiple Range Test. And test of significance were done at $P > 0.05$.
 150

151 **4.0. Results and Discussion**

152 **4.1. Physico-Chemical Parameters of River Ogbese**

153 The physicochemical properties of water obtained from River Ogbese are presented in

154 Table 1.

155 Table 1: Physicochemical parameters of River Ogbese.

Parameters	Range	Mean±SD
DO (mg/l)	5.80 – 7.99	6.98 ± 0.15
Turbidity (NTU)	67.00– 97.00	78.50 ± 13.53
Temperature (°C)	26.44 – 30.64	27.70 ± 0.18
Conductivity (µohm's/cm)	119.0– 178.0	148.35 ± 27.98
Ph	6.81-8.12	7.36 ± 0.22

156

157 **Length, Weight, Condition Factor (K) and LWR of *Chrysichthys nigrodigitatus***

158 Length (cm), Weight (g), Length / Weight Relationship and Condition factor (K) of *C.*
 159 *nigrodigitatus* obtained at River Ogbese (Table 2). The average body weight of *Chrysichthys*
 160 *nigrodigitatus* used was $148.15 \pm 36.53\text{g}$ which ranged from 106g – 185g, while the average
 161 body length was $25.64 \pm 2.86\text{cm}$ ranging between 23cm – 30cm. The condition factor was 0.88.
 162 The “b” values of the fish were not equal to 3, hence growth in the individual species was
 163 allometric (i.e. b values were less/greater than 3) showing that the rate of increase in body length
 164 is not proportional to the rate of increase in body weight.

165

166

Table 2: Morphometric Characteristic of *Chrysichthys nigrodigitatus*

Length / Weight Relationship	Measurement
Length (cm)	25.64 ± 2.09
Weight (g)	148.15 ± 28.56
Condition Factor (K)	0.88
Intercept (a)	2.08
Slope (b)	2.29
Coefficient of determination (r ²)	0.64

167

168 **Haematological Parameters of *Chrysichthys nigrodigitatus***

169 Tables 3 and 4 showed haematology characteristics of *Chrysichthys nigrodigitatus*. The result
 170 showed high values in parameters measured. Red Blood Cell was higher than the White Blood
 171 Cell with mean value of (225.63±10.45). Eosinophils recorded the lowest parameters with mean
 172 value of (1.75 ±0.52).

173

174 Table 3: Haematological Profile of *Chrysichthys nigrodigitatus* from River Ogbese.

Parameters	MAY	JUNE	JULY	AUGUST
ESR	3.50±0.71 ^a	4.00±0.78 ^a	3.75±0.42 ^a	4.00±0.00 ^a
PCV (%)	24.50±0.71 ^a	22.50±0.41 ^a	23.50±1.41 ^a	24.50±0.28 ^a
RBC (µL)	237.00±8.49 ^a	218.00±4.24 ^b	219.50±9.19 ^b	228.00±11.31 ^c
WBC (µL)	123.00±7.07 ^a	113.50±2.12 ^b	115.50±13.44 ^b	113.50±10.61 ^b
Hb (gdL-1)	8.15±0.21 ^a	7.80±0.42 ^a	8.00±0.28 ^a	8.50±0.21 ^a
Lymphocytes	59.00±1.41 ^a	50.00±0.00 ^a	55.00±1.41 ^a	59.50±2.12 ^a
Neutrophils	25.00±0.00 ^a	34.00±2.83 ^a	22.50±2.12 ^{ab}	23.00±4.24 ^{ab}
Monocytes	12.50±1.41 ^a	12.00±2.83 ^a	13.50±2.12 ^a	13.00±1.41 ^a
Basophils	2.00±0.71 ^a	2.50±0.91 ^a	2.00±0.41 ^a	2.50±0.71 ^a
Eosinophils	1.50±0.71 ^a	1.00±0.71 ^a	2.50±0.71 ^a	2.00±0.00 ^a
MCHC (gdL-1)	33.27±0.09 ^a	33.19±0.21 ^a	33.19±0.29 ^a	33.27±0.16 ^a
MCH	3.44±0.03 ^a	3.58±0.06 ^a	3.56±0.02 ^a	3.50±0.10 ^a
MCV (pg)	10.34±0.07 ^a	10.78±0.11 ^a	10.71±0.13 ^a	10.75±0.23 ^a

175 Values on the same row with the same superscript alphabet are not significantly different. N = 30

176

177

178

179

180
181

Table 4: Range and Mean Haematological Profile of *Chrysichthys nigrodigitatus* from River Ogbese

Parameter	Range	Mean±SD	SR
ESR (mm)	3.00– 4.00	3.81±0.35	4-10
PCV (%)	23.00-25.00	23.75±0.76	21-26
RBC (10 ³ /mm ³)	213.0– 243.0	225.63±0.45	200-250
WBC (10 ³ /mm ³)	106.0-128.0	116.38±8.19	100-150
Hb (g/100ml)	7.60 – 8.30	8.11 ±0.27	5-10
Lymphocytes	58.00– 61.00	55.88±1.19	64-80
Neutrophils (%)	20.00 -26.00	26.13±2.33	25-30
Monocytes (%)	10.00– 15.00	12.75±1.69	10-20
Basophils (%)	2.00– 3.00	2.25±0.53	2-5
Eosinophils (%)	1.00– 2.00	1.75±0.52	1-2
MCHC (gdL-1)	33.04 – 33.33	33.23±0.13	30-45
MCH (pg)	3.40 – 3.60	3.52±0.07	5-10
MCV (pg)	10.20 – 10. 90	10.65±0.22	10-15

182 Data are presented as Means ± S.D. ESR =Erythrocyte Sedimentation Rate, PCV =Packed Cell
183 Volume, HB =Haemoglobin, RBC =Red Blood Cell, WBC =White Blood Cell, MCV =Mean
184 Corpuscular Volume, MCHC =Mean Cell Haemoglobin Concentration, MCH =Mean Cell
185 Haemoglobin. S.R = Standard Range

186

187 **Histology of *Chrysichthys nigrodigitatus***

188 Results of histology of gills, liver and intestines of *Chrysichthys nigrodigitatus* are given in the
189 plates 1 - 13 below. The gill filaments are eroded with a deformation of the cartilage core and
190 also hyperplasia of the secondary lamellae. The intestines show atrophy in a mucosal layer,
191 hemorrhage and dilation within blood vessels and within serosa of mucosa. Liver histology
192 revealed shattered picnotic nucleus, ruptured hepatocytes and increased kupffer cell.



193 **Histology of the Gills**



194

195 **PLATE 2:** The gill filaments are eroded
 196 cartilage Magnification; x 100
 197

195 **PLATE 3:** There is a deformation of the
 196 core Magnification x 100
 197



198

199 **PLATE 4:** The gill arch and gill filaments are
 200 showing visible signs of lesions



201 **Magnification; x400**

202

203 **4.4.2 Histology of the Intestines**

200 **PLATE 5:** There is hyperplasia of the eroded
 201 secondary lamellae

202 **Magnification; x 400**





204

205 **PLATE 6:** shows atrophy in a mucosal layer

PLATE 7: Intestine shows sign of haemorrhage

206 **Magnification; x 100**

Magnification; x 100





207



208 **PLATE 8:** shows hemorrhage and dilation **PLATE 9:** shows severe degeneration and
 209 within blood vessels and within serosa of mucosa. necrosis of mucosal membrane of intestine

210 **Magnification; x400**

Magnification; x400



212 **PLATE 10:** The picnotic nucleus are shattered **PLATE 11:** The hepatocytes are ruptured
213 **Magnification; x 100** **Magnification; x 100**
214



215 **PLATE 12;** There is increased kupffer cell **PLATE 13;** Visible lesions seen
216 **Magnification; x400** **Magnification; x400**

217 **GF= Gill Filaments, CC= Cartilage Core, GA= Gill Arch, GL= Gill Lamellae, ML= Mucosa**
218 **Layer, SL= Serosa Layer, PN= Picnotic Nucleus, KC= Kkupffer Cell.**
219

220 **DISCUSSION**

221 Results of physicochemical parameters of water obtained in this study were within the tolerable
222 range of fish as recommended by WHO (2001 and 2006) except for DO. The result was similar
223 to the reports of Ansa (2004) on the benthic macrofauna of the Andoni flats in the Niger Delta

224 Area of Nigeria, Chindah *et al.*,(1998) on effect of municipal waste discharge on the
225 physicochemical and phytoplankton in a brackish wetland in Bonny Estuary and Ladipo *et al.*,
226 (2011) on seasonal variations in physico-chemical properties of water in some selected locations
227 of Lagos Lagoon who opined that waters with little change in physicochemical parameters are
228 generally more conducive to aquatic life. Most organisms including *C. nigrodigitatus* do not
229 tolerate wide variations in physicochemical parameters and if such conditions persist death may
230 occur. High oxygen demand experienced in the study is in line with Adebayo *et al.*, (2007).

231 Ujjania *et al.*, (2012) opined that condition factor greater or equal to one is good, indicating a
232 good level of feeding, and proper environmental condition. Mean K-values gotten from this
233 study (0.88) were less than one (1) in samples, hence revealing that the species fell slightly from
234 been healthy. This support the report of Gesto *et al.*, (2017) who worked on the Length-Weight
235 Relationship and Condition factor of *C. gariepinus* and *O. niloticus* of Wudil River, Kano,
236 Nigeria, and obtained condition factor less than one (1). Also feeding intensity, availability of
237 food, fish-size, age, sex, season, stage of maturation, fullness of the gut, degree of muscular
238 development and amount of reserved fat (Gupta and Banerjee, 2015) have influence on also K
239 factor of fish

240 The observation of absolute Isometric growth ($b = 3$) in nature is occasional (Bagenal 1978;
241 Basseyy and Ricardo, 2003), and deviation from isometric growth is often observed in most
242 aquatic organisms which changes shape as they grow (Thomas *et al.*, 2003). The differences in
243 the length-weight relationship also agrees with the report of Olurin and Aderibigbe (2006) who
244 stated that the differences may be due to sex and developmental stages of fish.

245 Mean haematocrit value of *C. nigrodigitatus* was $23.75 \pm 0.76\%$ which did not differ considerably
246 from those found by Badawi and Said 1971 and Etim *et al.*, 1999. The Red Blood Cell counts
247 has a mean value of $225.63 \times 10^6 \text{mm}^3 \pm 10.45 \times 10^6 \text{mm}^3$. The Packed cell volume (PCV) has a
248 mean value of $23.75 \pm 0.76\%$. Haemoglobin concentration has a mean value of $8.11 \pm 0.27 \text{g/dl}$.
249 The mean haemoglobin value is low which may be due to the exposure of fish to pollutants
250 resulting in inhibitory effect of those substances on the enzyme system responsible for the
251 synthesis of haemoglobin according to Pamila *et al.*, 1991. The low hb value in the water body
252 may also be associated with less active fishes. Similar results were reported by Engel and Davis,
253 (1964) and Rambhaskar and Rao, (1987). Eisler suggested that there was a correlation between
254 haemoglobin concentration and the activity of the fish. The more active fishes tend to have
255 higher haemoglobin values than the more sedentary ones (Pradan *et al.*, 2012). The high
256 erythrocyte number was associated with fast movement, predaceous nature and high activity
257 with streamlined body (Satheeshkumare*et al.*, 2011). A fall in count, Hb% and PCV%, in the
258 fishes, due to water pollution, has been reported along with acute anemia (Singh,1995).
259 According to Singh *et al.*, 2002), the discharge of waste may cause serious problems as they
260 impart odour and can be toxic to aquatic animals. The organic wastes present in Ogbese river
261 seem to cause stress in the fish and as such seem to be responsible for the changes in the
262 hematological parameters. The PCV or haematocrit is an important tool for determining the
263 amount of plasma and corpuscles in the blood (measurement of packed erythrocytes) and is used
264 to determine the oxygen carrying capacity of blood (Larsson *et al.*, 1985). Hematocrit or PCV in
265 the present study is low compared to the works of (Joshi *et al.*, 2002) and (Banerjee and
266 Banerjee, 1988) have suggested that pollutant exposure decreases the TEC count, Hb content and
267 PCV value due to impaired intestinal absorption of iron.

268 There were variation in WBC quantity and leukocyte cell proportions (neutrophil, monocyte) in
269 the fish specimens. The implication of this result is that the fish has been able to defend itself
270 from invading pathogens both by cell and antibody-mediated responses (Kumar *et al.*, 1999).
271 Similar results were obtained by Sahan and Cengizler, (1894) on carp caught from different
272 regions of Seyhan River. Leukocytosis is directly proportional to severity of stress condition in
273 maturing fish and is a result of direct stimulation of immunological defense due to the presence
274 of pollutants in water bodies. This is in conformity with the report of Saravanan and
275 Harikrishnan, (1999) in freshwater fish, *Sarotherodon mossambicus* , when exposed to sublethal
276 concentration of copper and endosulfan and by Nanda, (1997) in respect of *Heteropneustes*
277 *fossilis* during nickel intoxication. This may be attributed to alteration in blood parameters and
278 direct effects of various pollutants. The lymphocytes are reported to be responsible for immune
279 response (Cazenave *et al.*, 2005), while neutrophils are reported to show the greatest sensitivity
280 to change in the environment. Their characterization and identification is therefore, of
281 significance for assessing the changes in the physiological state of fishes

282 Marked variations like hyperplasia, vacuolation, deformation of cartilage core, bubbling of gill
283 filament, epithelial lifting, lamellar fusion; secondary lamellar damage, shorter secondary
284 lamellae and erosion of secondary lamellae were noticed in the gill tissues of *C. nigrodigitatus*
285 collected from river Ogbese. Similar results were obtained by several works: Fernandes and
286 Mazon, (2003), Simonato *et al.*, (2008), Rajeshkumar *et al.*, (2015), as they revealed alterations
287 like aneurysm, mucous deposition, hypertrophy, fusion of secondary lamellae, ruptured epithelial
288 layer, lifting of primary lamellae, lamellar swelling and necrosis. Through the gills, as the main
289 site of xenobiotic transfer, the toxins are distributed through their bodies accumulating in tissues
290 and organs and may have deleterious effects Vasanthi,*et al.*, (2015).

291 The extent of liver damage observed in the present investigation indicates that chronic exposure
292 always causes impairment to the architecture of the tissue. Since liver is involved in
293 detoxification of pollutants (Lagadic *et al.*, 2000), it is susceptible to a greater degree of
294 disruption in its structural organization due to toxic stress. Some distinct changes like rupture of
295 hepatocytes, melanomacrophages, increased Kupffer cell, increased pycnotic nucleus,
296 vacuolation, ruptured nucleus, Blood congestion, cytoplasmic vacuolation and nucleus
297 disorganization were observed in the liver of fish. Macrophage aggregates have been suggested
298 as potentially sensitive histological biomarkers and or immunological biomarker of contaminant
299 exposure (Schmitt *et al.*, 2000). Histological changes observed in various studies in liver taken
300 from the fishes exposed to pollutants include increased vacuoles in the cytoplasm, changes in
301 nuclear shapes, focal area of necrosis (death of cells in a localized area), ischemia (blockage of
302 capillary circulation), hepatocellular shrinkage, and regression of hepatocytic microvilli at the
303 bile canaliculi, fatty degeneration and loss of glycogen.(Marchand *et al.*, 2012) reported that
304 histopathological changes of fish liver from polluted freshwater system shows structural
305 alterations in hepatic plates or cords, multiple focal areas of cellular alterations leading to a loss
306 of uniform hepatocyte structure, steatosis, cytoplasmic and nuclear alterations (hypertrophic and
307 pyknotic nuclei) of hepatocyte, increase in the size of melanomacrophage centers (MMCs), and
308 focal areas of necrosis. The results from this study also agrees with the result of microscopic
309 examination of liver specimens from Lagos and Ologe Lagoon which were consistent with the
310 findings of Olarinmoye *et al.* (2009) in which liver of *C. nigrodigitatus* from Lagos lagoon
311 showed several alterations including vacuolar hepatocellular degeneration and hepatic necrosis.

312 Histology of the Intestine in the study revealed visible sign of lesions. Although, uptake of
313 metals occurs mainly through gills, it may also occur via intestinal epithelium. Histopathological
314 alterations in the intestine of *C. nigrodigitatus* included severe degenerative and necrotic changes
315 in the intestinal mucosa and sub mucosa, atrophy in the muscularis and sub mucosa and
316 aggregations of inflammatory cells in the mucosa and sub mucosa with edema between them.
317 These findings are in agreement with those of Hanna *et al.*, (2005), Bashir (2010), Yousafzai *et*
318 *al.*, (2010) and Soufy *et al.*, (2007), who opined that pollutants and contaminants affects gills by
319 epithelial lifting, hyperplasia of epithelial cells and blood congestion within filaments and in
320 liver tissue produced hemolysis between hepatocytes, cytoplasmic degeneration and necrosis.
321 Whereas an aggregation of inflammatory cells, edema in an intestinal mucosal layer and
322 hemorrhage between blood vessels were the main alterations observed in the intestine. The
323 changes seemed to be more pronounced in the liver and gills rather than the intestine.

324 **Conclusion**

325 Human activities including industrialization and agricultural practices contributed immensely in
326 no small measure to the degradation and pollution of aquatic environment which adversely has
327 effects on the water bodies that is a necessity for life. Since water pollution has direct
328 consequences on human well beings, an effective teaching strategy in the formal education sector
329 is essential for aquatic health

330 Regulation and monitoring is an effective way of pollution management. There is need to enact
331 legislation to regulate various types of pollution as well as to mitigate the adverse effects of
332 pollution.

333

334 **REFERENCES**

- 335 Abolude, D. S. and Abdullahi, S. A. (2005). Proximate and mineral contents in component parts
336 of *Clarias gariepinus* and *Synodontis schall* from Zaria, Nigeria. *Nigerian Food Journal* 23:1- 8.
- 337 Adams, S.M., Greeley, M.S and Ryon, M.G. (2000), "Evaluating effects of contaminants on fish
338 health at multiple levels of biological organization: extrapolating from lower to higher levels",
339 *Human and Ecological Risk Assessment*, Vol. 6 No. 1, pp. 15-22.
- 340 Adebayo, O. T., Fagbenro, O. A., Ajayi, C. B and Popoola, O.M. (2007). Normal haematological
341 profile of *Parachanna obscura* as a diagnostic tool in aquaculture. *International Journal of*
342 *Zoological Research*. 3(4): 193 – 199.
- 343 Adelegan, M.M. (2004). *Nigerian Petroleum Law and Practice* (Ibadan. Nigeria: Fountain
344 Books).
- 345 Adeniyi, O. (2014). Regional Planning – The Geography of Nigeria Development, in J.S.
346 Oguntoyinbo, Areola, O and M. Filani (eds), Heinemann Thadan. 1983;437–446.
- 347 Adeyemi, J.A. (2014). Oxidative stress and antioxidant enzyme activities in the African catfish,
348 *Clarias gariepinus*, experimentally challenged with *Escherichia coli* and *Vibrio fischeri*.
349 *Fish Physiology and Biochemistry* 40: 347–354.
- 350 Adhikari, and Sarkar, B (2004). Effects of cypermethrin and carbofuran on certain haematological
351 parameters and prediction of their recovery in fresh water teleost, *Labeo rohita* (Ham).
352 *Ecotoxicology and Environmental Safety* 58: 220-226. Ainsworth AJ, Dexiang C, Wterstrat PR

353 (1991). Changes in peripheral blood leucocyte percentage and function of neutrophils in stressed
354 channel catfish. *Journal of Aquatic Animal and Health* 3: 41- 47.

355 Adite, A., Winemiller, K O, and Fiogbe, E D (2006). Population structure and reproduction of
356 the African bony tongue *Silver Catfish* in the So River-floodplain system (West Africa):
357 implications for management. *Ecology of Freshwater Fishes.*,**15**: 30-39.

358 Affonso, E.G., Polez, VLP., Corrêa, CF, Mazon AF, Araújo MRR, Moraes G, and Rantin FT.
359 (2002). Blood parameters and metabolites in the teleost fish *Colossomamacropomum* exposed to
360 sulfide or hypoxia. *ComprehensiveBiochemical Physiology.*; Part C 133:375-382.

361 Agedah, E.C., Ineyougha, E.R., Izah, S.C., and Orutugu, L.A. (2015). Enumeration of total
362 heterotrophic bacteria and some physico-chemical characteristics of surface water used for
363 drinking sources in Wilberforce Island, Nigeria. *Journal of Environmental Treatment*
364 *Techniques.* 3(1), 28 – 34.

365 Ajayi, A. and Ikporokpor, D., (2002). *International Environmental law* (Ardley son:
366 Transnational Publishers).

367 Akande, G. R. (2011). Fish Processing Technology in Nigeria: Challenges and Prospects. In:
368 Aiyelaja, A.A and Ijeomah, H.M. (Eds.). *Book of Reading in Forestry, Wildlife Management*
369 *and Fisheries.* Topbase Nigeria Ltd. New Oko Oba, Lagos, pp. 772-808.

370 Akpan, A.W. (2004). The Water Quality of Some Tropical Freshwater Bodies in Uyo (Nigeria)
371 Receiving Municipal Effluent, SlaughterHouse Washings and Agricultural Land Drainage.
372 *Environmentalist*, 24: 49-55.

373 Al-Kaheem, H. F., Al-Ghanim, K. A., Ahamad, Z., Temraz, T. A., Al-Akel AS, Al-Misned F,
374 and Annazri HA (2008). Threatened fish species (*Aphanius dispar*) in Saudi Arabia, A Case
375 Study. *Pakistan journal of Biological Sciences*, 1-8.

376 Amukam, O., (1997). *Pollution control regulation the Nigerian oil industry* (Lagos:
377 N.I.A.L.S.1997).

378 Angaye, T.C.N., Zige, D.V. and Izah, S.C. (2015). Microbial load and heavy metals properties of
379 leachates from solid wastes dumpsites in the Niger Delta, Nigeria. *Journal of Environmental*
380 *Treatment Techniques*, 3(3): 175 – 180.

381 Ansari, T.M., Marr, I.L. and Tariq, N. (2004). Heavy Metals in Marine Pollution Perspective - A
382 Mini Review. *Journal of Applied Sciences.* 4(1): 1-20.

383 Araoye PA (1999). Spatio-temporal distribution of the fish *Synodontis schall* (Teleostei:
384 Mochokidae) in Asa lake, Ilorin, Nigeria. *Revised Biological Tropics* **47**: 1061-1066.

385 Arimoro, F.O. (2009). Impact of Rubber Effluent Discharges on the Water Quality and
386 Macroinvertebrate Community Assemblages in a Forest Stream in Niger Delta. *Chemosphere*,
387 77: 440-449.

388 Arimoro, F.O. and Ikomi, R.B. (2008). Response of Macroinvertebrate Communities to Abattoir
389 Wastes and Other Anthropogenic Activities in a Municipal Stream in the Niger Delta,
390 Nigeria. *Environmentalist.* 28: 85-98.

391 Authman, M. N., Zaki, M. S., Khallaf, E. A. and Hossam, H. A. (2015). Use of Fish as
392 Bioindicator of the Effects of Heavy Metal Pollution. *Journal of Aquacultural Research and*
393 *Development* 6 (4): 2155-9448.

394 Awoyemi, O.M., Achudume, A.C. and Okoya, A.A. (2014). The Physicochemical Quality of
395 Groundwater in Relation to Surface Water Pollution in Majidun Area of Ikorodu, Lagos State,
396 Nigeria. *American Journal of Water Resources*, 2(5): 126-133

397 Azevedo Tmp, Martins MI, Bozzo Fr And Moraes Fr.(2006). Haematological and gill responses
398 in parasitized tilapia from Valley of Tijucas River, SC, Brazil. *Sci Agric* 63(2): 115-120.

399 Ballarin, L., M. Dall'Oro, D. Bertotto, A. Libertini, A. Francescon and A. Barbaro, (2004).
400 Haematological parameters in Umbrina cirrosa (Teleostei, Sciaenidae): A comparison between
401 diploid and triploid specimens. *Composition of Biochemical Physiology. Part A: Mol.*
402 *Integrated Physiology.*, 138: 45-51

403 Banerjee, V. and M. Banerjee, (1988). Effect of heavy metal poisoning on peripheral hemogram
404 in *H. f o s s ili s* (Bloch) mercury, chromium and zinc chlorides (LC50). *Composition of*
405 *Physiology and Ecology.*, 13: 128-134.

406 Banerjee, T.K., I. Chatterjee, R.K. Jana, P. Chand and S.K. Das, (2009). Detoxification of the
407 effluents generated following recovery of metals from polymetallic sea nodules using
408 phytoremediation technique. *Proceedings of the National Symposium on Functional Biodiversity*
409 *and Ecophysiology of Animals*, Feb. 21-23, Department of Zoology, Banaras Hindu University,
410 India,-pp: 55.

411 Bankole, N. O., Yem, I. Y.and Olowosegun, O. M. (2011). Fish Resources of Lake Kainji,
412 Nigeria. In: Raji A Okaeme N. and Ibeun MO (Eds.). *Forty Years on Lake Kainji Fisheries*
413 *Research*, NIFFR, New Bussa, Nigeria, pp. 20-42.

414 Bashir, N., (2010). Bioaccumulation of heavy metals in organs of *Labeo rohita* and *Cyprinus*
415 *carpio*. BS thesis, Department of Zoology, GC University, Faisalabad.

416 Basorun, J.O. (2013) *Basic Elements of Urban and Regional Planning*, Shalom Publisher,
417 Akure;

418 Bassey, E. A. and Ricardo, P. K. (2003). Seasonality in growth of *Aphyosemiongradneri*
419 (Bolenger) in Mfangmfangpond in Uyo, Nigeria. *The Zoologist* 2: 68-75.

420 Camarago, M.M.P. and Martinez, C.B.R.(2007). Histopathology of gills, kidney and liver of a
421 Neotropical fish caged in an urban stream. *Neotropical Ichthyology*, 5: 327-336.

422 Cazenave, J., D.A. Wunderlin, A.C. Hued and M. de los Angeles Bistoni, (2005).
423 Haematological parameters in a neotropical fish, *C o r y d o r a s p a l e a t u s* (Jenyns,
424 1842) (Pisces, Callichthyidae), captured from pristine and polluted water. *Hydrobiologia*, 537:
425 25-33.

426 Chekrabarth, P and Benerjee, V. (1988). Effects of sublethal toxicity of three organophosphorus
427 pesticide on the peripheral haemogram of the fish, (*Channa punctatus*). *Environmental Ecology*
428 6: 151-158.

429 Chindah, A. C. (1998). The effect of industrial activities on the periphyton community at
430 the upper reaches of New Calabar River, Niger Delta, Nigeria. *Water Resources* 32 (4) 1137 –
431 1143.

432 Collares-Perreira, M. J., Cowx, I.G., Caelho, M. M (2002). *Conservation of Freshwater Fishes:*
433 *Options for Future*. 1st Edn. Iowa State University Press, Ames, IA. ISBN- 0-85238-2863,;pp:
434 472.

435 Craig, J (2000). *Percid Fishes. Sytematics, Ecology and Exploitation*. 1st Edn. Blackwell Science,
436 Oxford pp. 400

437 Davids, CBB., Ekweozor.EAS., Daka, E. R., Dambo, W. B. and Bartimacus EAS (2002). Effects
438 of Industrial Effluents on some Hematological Parameters of *Sarotherodon melenothero* and
439 *Tilapia guineensis*. *Global Journal of Pure and Applied Science* 8: 305-310.

440 Diersing, N. (2009). "Water Quality: Frequently Asked Questions." Florida Brooks National
441 Marine Sanctuary, Key West, FL

442 Dudgeon, D. (2003). The contribution of scientific information to the conservation and
443 management of freshwater biodiversity in tropical Asia. *Hydrobiology*; 500: 295-314.

444 Dudgeon, D., Smith, R. E. W (2006). Exotic species, fisheries and conservation of freshwater
445 biodiversity in tropical Asia. The case of Spike River, Papua New Guinea. *Aquatic Conserv:
446 Marine Freshwater Ecosystem* **16**:203-215.

447 Fakayode, S.O., (2005). Impact assessment of industrial effluent on water quality of the receiving
448 Alaro River in Ibadan, Nigeria. *Ajeam-Ragee*, 10: 1-13

449 Faniran, A., (1991). Water resources development in Nigeria. University of Ibadan, Ibadan, pp:
450 95

451 Fernandes, M. N. and A. F. Mazon. (2003). Environmental pollution and fish gill morphology. In:
452 Val, A. L. & B. G. Kapoor (Eds.). *Fish adaptations*. Enfield, Science Publishers., 203-231.

453 Filani, M. O. (1985). Regional Planning in Nigeria: The Critical Issues of Economic Dependency
454 in Abiodun, J.O. (ed) *Urban and Regional Planning Problems in Nigeria*, Ile-Ife, University of
455 Ife Press; pp-57.

456 Gallardo, M.A., M. Sala-Rabanal, A. Ibarz, F. Padros, J. Blasco, J. Fernandez-Borra and J.
457 Sanchez, (2003). Functional alterations associated with winter syndrome in gilthead sea bream (
458 *Sparus aurata*). *Aquaculture*, 223: 15-27.

459 Gbore, F. A., Oginni, O., Adewole, A. M and Aladetan, J. O. (2006). The effect of transportation
460 and handling stress on hematology and plasma biochemistry in fingerlings of *Clarias gariepinus*
461 and *Tilapia zillii*. *World Journal of Agricultural Sciences* 2(2): 208-212.

462 Getso, B.U., Abdullahi, J.M. and Yola, I.A. (2017). Length-weight relationship and condition
463 factor of *Clarias gariepinus* and *Oreochromis niloticus* of Wudil River, Kano, Nigeria. *Journal
464 of Tropical Agriculture, Food, Environment and Extension*. 16(1): 1-4

465 Gleitsmann, B.A., M.M.Kroma and T. Steenhuis (2007). Analysis of a rural water supply project
466 in three communities in Mali: participation and sustainability. *National Resource Forum*, 31:
467 142-150

468 Gupta S. and Banerjee S. (2015). Length-weight relationship of *Mystus tengara* (Ham.-Buch.,
469 1822), a freshwater catfish of Indian subcontinent. *International Journal of Aquatic Biology*. 3(2):
470 114-118.

471 Hanna, M.I., Shaheed, I.B. and Elias, N.S., (2005). A contribution on chromium and lead toxicity
472 in cultured *Oreochromis niloticus*. *Egypt. J. aquat. Biol. Fish.*, 9: 177- 209.

473 Harper, H. H. (1985). Fate of heavy metals from runoff in stormwater management systems. Ph.D.
474 Dissertation, University of Central Florida, Orlando, Florida.

475 Hinton, D. E., and Lauren, D. J, (1990) Integrative histopathological approaches to detecting
476 effects of environmental stressors of fishes. *American Fish Sociology. Sym.* **8**: 51-66.

477 Hinton, D.E., Baumann, P.C., Gardner G.C., Hawkins, W.E., Hendricks, J.D., Murchelano,
478 R.A., Okihiro, M.S. (1992). Histopathologic biomarkers. In: Huggett R.J. et al. (eds) *Biomarkers:*

- 479 Biochemical, Physiological and Histological Markers of Anthropogenic Stress. Lewis Publishers,
480 Chelsea, pp. 155–210
- 481 Houston, A. H. (1990). Blood and circulation. In: Schreck CB and Moyle PB (Eds), Methods in
482 fish biology. Am Fish Soc, Bethesda, Maryland, p. 273-334.
- 483 Ibrahim, S. A., Authman, M. M., Gaber, H. S. and El-Kasheif, M. A. (2013). Bioaccumulation of
484 Heavy Metals and their Histopathological Impact on Flesh of *Clarias gariepinus* from El-
485 Rahawydrain, Egypt. International Journal of Environmental Science and Engineering 4: 51-73.
- 486 Iqbal, F., Qureshi, I.Z. and Ali, M. (2004) Histopathological changes in the kidney of common
487 carp, *Cprinuscarpio*, following nitrate exposure. *Journal Resources (Science)*, 4, 411–418.
- 488 Ita, E. O. (1993). Inland fishery resources of Nigeria. CIFA Occasional paper No. 20, Rome
489 FAO.;120pp.
- 490 Jamalzadeh, H. R., Keyvan, A., Ghomi, M. R and Gherardi, F. (2009). Comparison of blood
491 indices in healthy and fungal infected Caspian salmon (*Salmo trutta caspius*). African Journal
492 Biotechnology. 8(2): 319-322.
- 493 James, O. (1984). The Production and Storage of Dried Fish. In: FAO Fisheries Report (ITALY),
494 No.279. Supplementary.
- 495 Jiraung, K.W., Upathama, E. S, Kruatrachuea, M., Sahaphongse, S., Vichasrigramsa, S., and
496 Pokethitiyooka, P. (2002). Histopathological effects of round up, a glyphosate herbicide, on Nile
497 tilapia (*Oreochromis niloticus*). *Science, Asia*. 28: 121-127. DOI: 10.2306/scienceasia1513-
498 1874.2002.28.121
- 499 Joshi, B. D. (1982). Circannual fluctuations in some blood components of the fish *Rita rita*, in
500 relation to certain ecophysiological conditions. *Uttar Pradesh J Zoology* 2: 62-66
- 501 Joshi, P.K., M. Bose and D. Harish, (2002). Haematological changes in the blood of *Clarias*
502 *atrachus* exposed to mercuric chloride. *Journal of Ecotoxicology and Environmental*
503 *Monitoring*., 12: 119-122.
- 504 Kalu, V. E., and LL.M, B. L, (2009): Toxic Wastes And the Nigerian Environment; Dept. Of
505 Private & Property Law, University of Benin: An Appraisal.
- 506 Katte, V.Y., M.F. Fonteh and G.N. Guemuh, (2003). Domestic water quality in urban centres in
507 Cameroon: A case study of dschang in the west province of African waters ., 25: 45-56
- 508 Kumar, S., S. Lata and K. Gopal, (1999). Deltamethrin induced physiological changes in
509 freshwater cat fish *Heteropneustes fossilis* . *Bulletin of Environmental Contamination*
510 *Toxicology*.; 62: 254-258.
- 511 Ladipo, M. K., Ajibola, V. O., and Oniye, S. J. (2011). Seasonal Variations in Physicochemical
512 properties of water in some Selected locations of the Lagos Lagoon. *Science World Journal Vol*
513 *6* (No 4).
- 514 Lagadic, L., Amiard, J. C., Caquet, T. (2000) Biomarkers and evaluation of the ecotoxicological
515 impact of pollutants. In: Lagadic L, Caquet T, Amiard JC, Ramade F, Use of Biomarkers for
516 Environmental Quality Assessment. Science Publishers, Enfield (NH) USA.
- 517 Laleye, R. (1995). Climatic and anthropogenic effects on fish diversity and yields in the Central
518 Delta of the Niger River. *Aquatic Living Resources*; 8: 43-58.

- 519 Larsson, A., C. Haux and M.L. Sjobeck, (1985). Fish physiology and metal pollution: Results
520 and experiences from laboratory and field studies. *Ecotoxicology and Environmental Safety*, 9:
521 250-281.
- 522 Lavanya, S., Ramesh, M., Kavitha, C. and Malarvizhi, A. (2011). Haematological, biochemical
523 and ion regulatory responses of Indian major carp *Catla catla* during chronic sublethal exposure
524 to inorganic arsenic. *Chemosphere* 82: 977–985.
- 525 Lusková, V. (1998). Factors affecting haematological indices in free-living fish populations. *Acta*
526 *Vet Brno* 67: 249-255.
- 527 Maheswaran, R., Devapaul, A., Velmurugan, B. and Ignacimuthu, S. (2008). Haematological
528 studies of freshwater fish, *Clarias batrachus* (L.) exposed to mercuric chloride. *International*
529 *Journal for Integrated Biology*. 2(1): 49 - 54.
- 530 Mallatt, J. (1985) Fish gill structural changes induced by toxicants and other irritants: A
531 statistical review. *Canadian Journal of Fisheries and Aquatic Sciences*. 42:630-648.
- 532 Marchand, M. J., Dyk, J. C. V., Barnhoorn, IEJ, Wagenaar GM. (2012). Histopathological
533 changes in two potential indicator fish species from a hypereutrophic Freshwater ecosystem in
534 South Africa: A baseline study.
- 535 Martins, M. I., Moraes, F.R, Fujimoto, Ry, Onaka, Em, Bozzo, Fr And Moraes, Jre. (2006).
536 Carrageenin induced inflammation in *Piaractus mesopotamicus* (Osteichthyes: Characidae)
537 cultured in Brazil. *Bol Inst Pesca* 32(1): 31-39.
- 538 Martins, M. I., Mouriño, J. P., Amaral, G. V., Vieira, F. N., Dotta, G., Jatobá, A. B., Pedrotti, F.
539 S., Jerônimo, G. T., Buglioneneto, C. C. and Pereira-Jr, G. (2008). Haematological changes in
540 Nile tilapia experimentally infected with *Enterococcus* sp. *Braz J Biol* 68(3): 657-661.
- 541 Martins, M. I., Tavares-Dias, M., Fujimoto, R. Y., Onaka, E. M. and Nomura, D. T. (2004).
542 Haematological alterations of *Leporinus macrocephalus* (Osteichthyes: Anostomidae) naturally
543 infected by *Goezia leporini* (Nematoda: Anisakidae) in fish pond. *Arq Bras Medicine, Veterinary*
544 *Zootechnique*. 56(5): 640-646.
- 545 Meyers, T. R., Hendricks, J. D. Histopathology. In: Rand, GM., Petrocelli, SR. (Eds.), (1985)
546 *Fundamentals of Aquatic Toxicology: Methods and Applications*. Hemisphere Publishing
547 Corporation, Washington, USA, 283- 331.
- 548 Nanda, P., 1997. Haematological changes in the common Indian cat fish *Heteropneustes fossilis*
549 under nickel stress. *Journal of Ecobiology*., 9: 243-246.
- 550 Nwani, C. D., Nnaji, M. C., Oluah, S. N., Echi, P. C., Nwamba, H. O., Ikwuagwu, O. E., Ajima,
551 M. O. (2014). Mutagenic and physiological responses in the juveniles of African catfish, *Clarias*
552 *gariepinus* (Burchell 1822) following short-term exposure to praziquantel. *Tissue and Cell* 46:
553 264–273.
- 554 Ogbonna, E.A., and Ekweozor, P., (2000): The Adverse Effects of Crude Oil Spills in the Niger
555 Delta. *Urhobo Historical Society*.
- 556 Olaniran, N.S. (1995). “Environment and Health: An Introduction”, in Olaniran, N.S. et al (Ed)
557 *Environment and Health*. Lagos. Micmillan Nigeria Publication Company for NCF, 34-151.

558 Olarinmoye, O., Taiwo V., Clarke E., Kumolu-Johnson C., Aderinola O., Adekunbi F. (2009).
559 Hepatic pathologies in the brackish water catfish (*Chrysichthys nigrodigitatus*) from
560 contaminated locations of the Lagos lagoon complex. *Appl. Ecol. Environ. Res.* 7: 277–286

561 Olorunfemi, J.F. and H.I. Jimoh, (2000). Anthropogenic activities and the environment. In
562 contemporary issues in environmental studies.

563 Omotoso, T., Lane-Serff G.F., and Young, R. (2015). Issues in River Water Quality, Assessment
564 and Simulation in a West Africa Sub-Region. E-proceedings of the 36th IAHR World Congress
565 28 June – 3 July, 2015, The Hague, the Netherlands, pp 1 – 8

566 Ortiz, J. B., Gonzalez de Canales, M. L. and Sarasquete, C. (2003) Histopathological changes
567 induced by lindane (γ -HCH) in various organs of fish. *Science Marine* 67: 53-61.

568 Orubu, E.A. (2006) The Community of Oil Exporting Countries New York: Cornell University
569 Press).

570 Otuaga, P. M. (2015). Flow pattern of River Ogbese in Akure, Ondo State Nigeria. Proceedings
571 of 2015 international conference on disaster management in civil engineering. Pp 14-20.

572 Oyewo, E.O., and Don-Pedro, E.N. (2003). Influence of salinity variability on heavy metal
573 toxicity of three estuarine organisms. *Journal of Nigeria Environmental Science* 1(2), 141-155.

574 Peebuaa, P., Kruatrachuea, M., Pokethitiyooka, P. and Kosiyachindaa P. (2006). Histological
575 Effects of Contaminated Sediments in Mae Klong River Tributaries, Thailand, on Nile tilapia,
576 *Oreochromis niloticus*. *Science Asia*, 32, 143-150.

577 Poleksić, V., Savić, N., Rašković, B., and Marković, Z. (2006): Effect of different feed
578 composition on intestine and liver histology of trout in cage culture. *Biotechnology in Animal*
579 *Husbandry* 22:359-372.

580 Rajeshkumar, S., Karunamurthy, D., Halley, G., Munuswamy, N. (2015) An integrated use of
581 histological and ultra-structural biomarkers in *Mugil cephalus* for assessing heavy metal
582 pollution in east Berbice- Corentyne, Guyana, *International Journal of Bioassays*. 4(11):4541-
583 4554.

584 Ramesh, F. (2006). Influence of sago effluent on the physiological, biochemical, haematological
585 and histological aspects of the Indian catfish *Clarias batrachus*. Ph.D. Thesis, Bharathiar
586 University, Coimbatore.

587 Ranzani-Paiva, M. J. and Godinho, H. M. (1985). Estudos hematológicos do curimatá,
588 *Prochilodus scrofa* Steindachner, 1881 (Osteichthyes, Cypriniformes, Prochilodontidae). Série
589 vermelha. *Bol Inst Pesca* 12(2): 25-35.

590 Ranzani-Paiva, M. J., Silva-Souza, A. T., Pavanelli, G. T., Takemoto, R. M. and Eiras, AC.
591 (2000). Hematological evaluation in commercial fish species from the floodplain of the upper
592 Paraná River, Brazil. *Acta Science* 22: 507-513.

593 Reed, W., Burchard, J., Hopson, A. J., Jennes, J., and Yam, I. (1967). Fish and Fisheries of
594 Northern Nigeria. Ministry of Agriculture, Northern Nigeria, Gaskiya, Zaria. 226pp.

595 Rey Vázquez, G. and Guerrero, G. A. (2007). Characterization of blood cells and hematological
596 parameters in *Cichlasoma dimerus* (Teleostei, Perciformes). *Tissue and Cell* 39: 151-160.

597 Richter, B.D., Braun, D.P., Mendelson, M.A. and Master, L. L. (1997) Threats to imperiled
598 freshwater fish fauna. *Conserv. Biology*; 11: 1081-1093.

599 Rodrigues, E. L. and Fanta, E. (1998). Liver histopathology of the fish *Brachydanio rerio* after
600 acute exposure to sublethal levels of the organophosphate Dimetoato 500. *Revista Brasileira de*
601 *Zoologia*, 15, 441-450.

602 Sahan, A. and I. Cengizler, (2002). [Determination of some haematological parameters in spotted
603 barb (*Capoeta barroisi* Lortet, 1894) and roach (*Rutilus rutilus* , Linnaeus, 1758) living in
604 Seyhan river (Adana city region)]. *Turkish Journal of Veterinary and Animal Science.*, 26: 849-
605 858.

606 Saravanan, J.S. and R. Harikrishnan, (1999). Effects of sublethal concentration of copper and
607 endosulfan on haematological parameters of the freshwater fish *Sarotherodon mossambicus*
608 (Trewaves). *Journal of Ecobiology.*, 11: 13-18

609 Satheeshkumar, P., D. Senthilkumar, G. Ananthan, P. Soundarapandian and A.B.
610 Khan, (2011). Measurement of hematological and biochemical studies on wild marine
611 carnivorous fishes from Vellar estuary, Southeast coast of India *Composition of Clinical*
612 *Pathology.*, 20: 127-134.

613 Schmitt, C.J., Dethloff, G. M. (2000). Biomonitoring of Environmental Status and Trends
614 (BEST) Program: selected methods for monitoring chemical contaminants and their effects in
615 aquatic ecosystems. Information and Technology Report USGS/BRD-2000--0005. Columbia,
616 (MO):U.S.

617 Selma, k. and Hatice, P. (2004). The Effects of Pollution on Haematological Parameters of Black
618 Goby (*Gobius niger* L., 1758) in Foca and Aliaga Bays, Turkey, E.U. *Journal of Fisheries &*
619 *Aquatic Sciences*, (1- 2):113 -117.

620 Shah, S. L. and Altindag, A. (2004). Haematological parameters of tench (*Tinca tinca* L.) after
621 acute and chronic exposure to lethal and sublethal mercury treatments. *Bulletin of Environmental*
622 *Contamination and Toxicology.* 73: 911 - 918.

623 Shanthi, K., Kiran, Joseph. and Manimeghalai, M.(2009). Studies on the biochemical changes in
624 the liver and Kidney due to Steriling biotech Ltd effluent in freshwater fish, *Labeorohita*. *Indian*
625 *Journal of Environment and Ecoplanning.* 16(1), 145-150.

626 Silva Souza, A. T., Almeida, S.C. and Machado, P. M. (2002). Hematologia: o quadro sanguineo
627 de peixes do rio Tibagi. In: Medri ME, Bianchini E, Shibatta OA and Pimenta JA (Eds), *A bacia*
628 *do rio Tibagi*. Edit. Ed. dos Editores, Londrina, PR, p. 449-471.

629 Simonato, J.D., Guedes, L.B., Martinez, B.R., (2008) Biochemical, physiological and
630 histological changes in the neotropical fish *Prochilodus lineatus* exposed to diesel oil.
631 *Ecotoxicology and Environmental Safety.* 69:112- 120.

632 Singh, M., (1995). Haematological responses in a freshwater teleost *Channa punctatus* to
633 experimental copper and chromium poisoning. *Journal of Environmental Biology*, 16: 339-341.

634 Singh, K.D., B. Srivastava and A. Sahu, (2002). Non-conventional absorbents for fresh water
635 treatment containing phenolic compounds. *Proceedings of the 22nd Annual Meeting American*
636 *Society for Reproductive Immunology*, June 6-9, 2002, Chicago, IL., pp: 73-74.

637 Skeleton, P. H. (2002). An overview of the challenges of conserving freshwater fishes in South
638 Africa. In: *Conservation of Freshwater Fishes: Options for Future*, Collares-Perreira, M.J., I.G.
639 Cowx and M.M Caelho (Eds.). *Fishing News Books, Blackwell Science, Oxford*, ISBN-0-
640 85238-2863, Pp: 221-236.

641 Skouras, A., Broeg, K., Dizer, H., Von Westernhagen, H., Hansen, P. and Steinhagen, D. (2003).
642 The use of innate immune responses as biomarkers in a programme of integrated biological
643 effects monitoring on flounder (*Platichthys flesus*) from the southern north Sea. *Helgol. Marine*
644 *Resources* 57: 190–198

645 Soufy, H., Soliman, M., El-Manakhly, E. and Gaafa, A., (2007). Some biochemical and
646 pathological investigations on monosex *Tilapia* following chronic exposure to carbofuran
647 pesticides. *Global Veterinary.*, 1: 45-52.

648 Srivastava, S.K., Tiwari, P.R. & Srivastav, A.K. (1990). Effects of chlorpyrifos on the kidney of
649 freshwater catfish *Heteropneustes fossilis*. *Bulletin of Environmental Contamination and*
650 *Toxicology*. Vol. 45: 748–751.

651 Sutherland, R. A. and Tolosa, C. A. (2000). Multi-element analysis of roaddeposited sediment in
652 an urban drainage basin, Honolulu, Hawaii. *Environmental Pollution*. 110: 483-495

653 Thomas, J., Venus, S., and Kurup, B.M. (2003): Length-Weight relationship of some deep sea
654 fishes inhabiting the continental slope beyond 250m depth along west coast of India. *Naga*.
655 *ICLARM Q.26* 17-21.

656 Udayakumar, V. (2005). Studies on the influence of treated dairy effluent on certain
657 physiological, biochemical, haematological and histological aspects of freshwater catfish *Clarias*
658 *batrachus*. M.Phil. Dissertation, Bharathiar University, Coimbatore.

659 Ujjania, N.C., Kohli, M.P.S. and Sharma, L.L. (2012). Length-weight relationship and condition
660 factors of Indian major carps (*C. catla*, *L. rohita* and *C. mrigala*) in Mahi Bajaj Sagar,
661 India. *Research Journal of Biology*. 2 (1): 30-36

662 UNESCO-WWAP, (2003). Water for people, water for life

663 USAID Markets (United States Agency for International Development). (2010). *Best*
664 *management practices for fish farmers in Nigeria*. Washington DC: USAID.

665 Val, A.L., Schwantes, A. R., Almeida-Val, V. M. F. and Schwantes, M. L. B. (1985). Hemoglobin,
666 hematology, intraerythrocytic phosphates and whole blood bohr effect from lotic and lentic
667 *Hypostomus regani* populations (São Paulo-Brazil). *Composition of Biochemical Physiology*
668 80B(4): 737-741.

669 Van der Oost, R.; Beyer, J., and Vermeulen, N.P.E. (2003). Fish bioaccumulation and
670 biomarkers in environmental risk assessment: A review of *Environmental Toxicology and*
671 *Pharmacology*. 13, 57-149.

672 Vasanthi, L. A., Revathi, P., Mini, J., Natesan, M. N. (2013) Integrated use of histological and
673 ultrastructural biomarkers in *Mugil cephalus* for assessing heavy metal pollution in Ennore
674 estuary, Chennai, *Chemosphere.*; 91:1156- 1164.

675 Velkova-Jordanoska, L. and Kostoski, G. (2005), “Histopathological analysis of liver in fish
676 (*Barbus meridionalis* Petenyi Heckel) in reservoir Trebenista”, PSI Hydrobiological Institute,
677 Macedonia.

678 Webster.com. (2010). Definition from Webster Dictionary 08-13 Retrieved 2010-08-26

679 WHO, (2001). Water Quality Surveys: A guide for the collection and interpretation of water
680 quality data; Studies and Reports in Hydrology N 23, UNESCO/WHO.

681 WHO, (2006). World Health Organization: Guidelines for drinking water quality; Geneva.

682 Wikipedia, the free encyclopedia. <http://en.wikipedia.org/wiki/pollution>. Retrieved 20-06-2013

683 World Health Organisation (2006). Meeting the MDG drinking water and sanitation target: the
684 urban and rural challenge of the
685 decade. http://www.who.int/water_sanitation_health/monitoring/jmpfinal.

686 Yousafzai, A. M., Douglas, P., Khan, A. R., Ahmad, I. and Siraj, M., (2010). Comparison of
687 heavy metals burden in two freshwater fishes, Wallago attu and Labeo dyccheilus with regard to
688 their feeding habits in natural ecosystem. Pakistan Journal of Zoology., 42: 537-544.

689
690
691

UNDER PEER REVIEW