

2

3 **Common findings in Ultrasound Examination of the Liver**
4 **and Kidneys in Sokoto Metropolis.**

5

6 **ABSTRACT**

7 **Objectives:** *To evaluate and document common findings in ultrasound examination of the liver*
8 *and kidneys in Sokoto metropolis and to reveal gender and age distribution of the findings in*
9 *these organs.*

10 **Materials and methods:** *The data was collected retrospectively from the hospital archives and*
11 *records using a data capture sheet.*

12 **Results:** *Total sample of 322 ultrasound reports of the liver and kidneys in which some patients*
13 *had two findings, one in the liver and another in the kidney, some had liver findings only, some*
14 *kidney findings only, while some had normal ultrasound examination. Liver findings recorded a*
15 *dominant male gender with a frequency of 75 (70.1%). Findings in the liver reported were liver*
16 *cirrhosis, hepatomegaly, fatty liver, primary liver cell carcinoma, hepatic metastasis, chronic*
17 *liver disease (CLD), hepatitis, hepatoma and liver cyst. Liver cirrhosis had the highest frequency*
18 *with 29 (27.1%), hepatomegaly with a frequency of 17 (15.9%) followed by fatty liver and plcc*
19 *with a frequency of 13 (12.1%) each. The age range of 40-49 years had the highest frequency for*
20 *the liver findings with 23 (21.5%), followed by 60-69 years with 18 (16.8%) while 10-19 years*
21 *had the least frequency with 3 (2.8%). For the kidney findings, male gender also dominated the*
22 *findings with a frequency of 107 (64.1%). The age range that had the highest frequency was 20-*
23 *29 years with 43 (25.7%), followed by 30-39 years with a frequency of 27 (16.2%). The age*
24 *range with the least frequency was 80-89 years with 6 (3.6%). The pathological kidney findings*
25 *reported were pyelonephritis, renal parenchyma disease (grades I, II, III, and IV),*
26 *hydronephrosis, renal calculi, renal cyst, ectopic kidney, renal mass, renal failure and nephrotic*
27 *syndrome. Pyelonephritis was recorded as the commonest finding with a frequency of 67*
28 *(40.1%), followed by renal parenchymal disease with 35 (21.0%).*

29 **Conclusion:** *The study revealed that male gender is at higher risk of having both liver and*
30 *kidney diseases in Sokoto metropolis than the female and the age of those at higher risk*
31 *ranged from 20-49 years.*

32 *Liver cirrhosis and pyelonephritis are the commonest diseases of the liver and kidneys*
33 *respectively.*

34 **KEYWORDS:** Ultrasound, Liver, Kidneys, Metropolis, Echogenicity.

35 **1.0 INTRODUCTION**

36 Ultrasound machines use high frequency sound waves to produce images of diagnostic
37 quality [1]). They generate sound waves and receive the reflected echoes from the body
38 tissues to which the sound waves are applied to and these waves are emitted from
39 piezoelectric crystals from the ultrasound transducer. The piezoelectric crystals changes
40 the electrical signals to mechanical vibrations and changes mechanical vibrations to
41 electrical signals [2]. As the ultrasound waves pass through various body tissues, they are
42 reflected back to the transducer creating an image on the ultrasound screen [3].

43 Ultrasonography of the liver and kidneys involves scanning the patient's right upper
44 quadrant and left and right flanks respectively from different planes with patients placed
45 in various positions in order to help the sonographer to visualize and evaluate the normal
46 anatomy of the organs and as well detect abnormalities related to the organs [4].
47 Ultrasound is a good modality for imaging the Liver and the kidneys due to its ability to
48 demonstrate their anatomy which makes it easier to detect pathology, its availability, low
49 cost, non-invasive and do not use ionising radiation.

50 The liver sonographically appears slightly more echogenic than the renal cortex. It has a
51 homogeneous echo texture except for the anechoic (echo free) blood vessels and the gall
52 bladder [1]. The following are common indications for liver Ultrasonography: Hepatitis,
53 liver cirrhosis, fatty liver, Abscess, Liver cyst, Hepatomegaly, Hepatocellular carcinoma,
54 Portal hypertension, Hepatic calcification, etc. [1].

55 Renal ultrasound (US) is a common examination, which has been put to practice in
56 diagnosis of renal pathology and assessment of renal anatomy for decades [5].
57 Sonography is an essential tool in nephrology which is not only used in the diagnosis and
58 management of kidney disease, but also for the guidance of invasive procedures which is

59 non as interventional sonography. For this reason, it is essential for nephrologists to have
60 a thorough understanding of sonography and its uses in nephrology. Technological
61 advances over the past 15 years have resulted in high-quality scanners that are both
62 portable and affordable, which has greatly expanded the use of point-of-care sonography
63 by clinicians. Although nephrologists have been lagging in this area, an increasing
64 number are incorporating sonography into their practice, and training programs are
65 finally starting to meet this need [6].

66 Sonographically, the renal cortex is usually less echogenic than the liver and the spleen,
67 but in young infants is isoechoic or slightly hyperechoic with respect to these organs. The
68 central renal sinus appears hyperechoic than the renal cortex, the liver and the spleen [1].

69 Because of their location, architecture, and limited spectrum of pathology, the kidneys
70 are ideally suited for evaluation by ultrasound. In addition, it is safe, readily available,
71 easily performed at the bedside or in the office, and free of radiation. For these reasons,
72 sonography is the preferred imaging modality and often the only one required. Evaluation
73 includes assessment of the size and shape, the echogenicity, the urinary space (including
74 the lower urinary tract), the presence of masses, and the vasculature [6].

75 **2.0 MATERIALS AND METHOD**

76 A retrospective non experimental study design was adopted for the study. A secondary
77 source of data was used and was recorded using data capture sheet. The data were
78 collected from the archives of Radiology, Urology and Medicine of Usmanu Danfodiyo
79 University Teaching Hospital University (UDUTH) and Specialist Hospital Sokoto
80 (SHS). A non-probabilistic (convenient) sampling technique was used in selecting data.
81 1,651 sonographic reports of patients that underwent liver and kidney sonography from
82 January, 2017 to June, 2018 at UDUTH and Specialist Hospital Sokoto. Yaro Yamane's
83 formula was used to calculate the sample size for this study is 322 Ultrasound reports of
84 the liver and the kidneys with patients' details such as Gender, Age and date of the
85 examination with sonographic reports [7].

86 All ultrasound reports of the liver and kidneys with incomplete patients' details which
87 include gender age and date of examination. All the examinations done outside the scope
88 of the study; before January 2017 and after June 2018 were excluded. The data was
89 collected retrospectively in the Hospital records and archives using data capture sheet.
90 Descriptive statistics was used to analyse the data, Mean, Frequencies and percentages
91 were calculated.

92

93

3.0 RESULTS

94 The total number of data collected or sample size used was 322 with a gender distribution
95 of males, 207 (64.3%) and females, 115 (35.7). It has a mean of 39.2 and a standard
96 deviation of 28.0.

97 Table 3.1 shows gender and age distribution for the subjects with male gender recording
98 the highest frequency and the age range of 20-29 years having the highest frequency

99 Table 3.2 shows the frequency and percentage of the general findings base on the organs
100 affected and those that had normal sonographic examination.

101 Table 3.3 shows gender and age distribution for the pathological liver findings with male
102 gender recording the highest frequency and the age range of 40-49 years recording the
103 highest frequency

104 Table 3.4 shows gender distribution, frequency and percentage of pathological liver
105 findings. Liver cirrhosis was the pathology of the liver with the highest frequency. Male
106 gender had highest frequency.

107 Table 3.5 shows gender and age distribution of pathological kidney findings with the
108 male gender also recording the highest frequency, and the age range of 20-29 years
109 recording the highest frequency.

110 Table 3.6 shows gender distribution, frequency and percentage of pathological kidney
111 findings. Pyelonephritis is seen to be the most common kidney pathology.

112

113 **Table 3.1: Gender and age distribution for the subjects**

GENDER	FREQUENCY	PERCENTAGE (%)
Male	207	64.3
Female	115	35.7
TOTAL	322	100
AGE RANGE (YEARS)		
0-9	29	9.0
10-19	30	9.3
20-29	63	19.6
30-39	48	15.0
40-49	53	16.5
50-59	30	9.3
60-69	37	11.5
70-79	22	6.8
80-89	10	3.1
TOTAL	322	100

114

115 **Table 3.2: Table 4. 2: Frequency and percentage of all the findings**

ORGAN	FREQUENCY	PERCENTAGE (%)
Liver only	90	28.0
Kidneys only	150	46.6
Both liver and kidneys	17	5.3
Normal findings	65	20.2
TOTAL	322	100

116 **Table 3.3: Gender and age distribution for liver findings**

AGE RANGE (YEARS)	GENDER		FREQUENCY	PERCENTAGE (%)
	M	F		
< 20	6	4	10	9.3
20-29	6	2	8	7.5
30-39	12	4	16	15.0
40-49	16	7	23	21.5
50-59	11	6	17	15.9
60-69	11	7	18	16.8
70-79	7	2	9	8.4
80-89	5	1	6	5.6
TOTAL	74	33	107	100

117

118

119

120


121

122

123

124

125



126

127

128 Figure 3.1: A bar chart showing gender and age distribution for liver findings

129

130

131

132

133 **Table 3.4: Gender distribution, frequency and percentage of pathological liver findings**

PATHOLOGY	GEMNDER		FREQUENCY	PERCENTAGE (%)
	M	F		
Liver cirrhosis	18	11	29	27.1
Hepatomegaly	12	5	17	15.9
Fatty liver	7	6	13	12.1
PLCC	11	2	13	12.1
Liver metastasis	6	3	9	8.4
CLD	6	2	8	7.5
Hepatitis	5	2	7	6.5
Hepatoma	3	1	4	3.7
Hepatorenal syndrome	2	0	2	1.9
Sepsis	2	0	2	1.9
Liver cyst	2	0	2	1.9
Liver failure	0	1	1	0.6
TOTAL	74	33	107	100

134

135


136

137

138

139

140



141

142 Figure 3.2: A pie chart showing percentage of pathological liver findings

143

144

145

146

147

148 **Table 3.5: Gender and age distribution for kidney findings**

AGE RANGE (YEARS)	GENDER		FREQUENCY	PERCENTAGE (%)
	M	F		
< 20	18	14	32	19.2
20-29	29	14	43	25.7
30-39	20	7	27	16.2
40-49	11	12	23	13.8
50-59	8	3	11	6.2
60-69	9	3	12	7.2
70-79	9	4	13	7.8
80-89	3	3	6	3.6
TOTAL	107	60	167	100

149

150

151


152

153

154

155

156



157

158 Figure 3.3: A bar chart showing gender and age distribution for pathological kidney
 159 findings

160

161

162

163

164

165 Table 3.6: Gender distribution, frequency and percentage of pathological kidney findings

PATHOLOGY	GENDER		FREQUENCY	PERCENTAGE (%)
	M	F		
Pyelonephritis	47	22	69	41.3
Renal parenchyma disease	20	15	35	21.0
Hydronephrosis	16	9	25	15
Renal cyst	6	6	12	7.2
Renal calculi	6	5	11	6.6
Ectopic kidney	3	0	3	1.8
Renal mass	3	0	3	1.8
Renal failure	1	1	2	1.2
Nephrotic syndrome	1	1	2	1.2
Hepatorenal syndrome	1	1	2	1.2
Sepsis	2	0	2	1.2
Renal agenesis	1	0	1	0.6
TOTAL	107	60	167	100

166

167


168

169

170

171

172



173

174 Figure 3.4: A pie chart showing percentage of pathological kidney findings

175

176

177

178

179 **Discussion**

180 The total of 322 samples were collected and analysed, with a predominant gender of male
181 which carried up to 64.3% of the whole sample, leaving the female gender with the
182 remaining 35.7%. Some patients had findings related to both the liver and kidneys but the
183 predominant requested ultrasound examination and the predominant sonographic findings
184 were those related to the kidneys accounting for up to a frequency of 167 and a
185 percentage of up to 51.9% of all the findings. Liver findings followed with a frequency
186 and percentage of 107 and 33.2 respectively while some had normal sonographic
187 examination (65, 20.2%), in other words; their pathologies were not detected by
188 sonography, though the absence of disease or pathology in them could not be ruled out.
189 The age range of 20-29 years predominated the requests and the findings with a
190 frequency of 63 (19.6%)

191 After analysing the liver findings, the revealed predominated gender was the male
192 gender with a frequency and percentage of 75 and 70% respectively. This was in lined
193 with a similar study conducted in Asaba, Delta State, Nigeria [8]. The gender distribution
194 of dominant male gender is also in line with that of the study conducted on “Pattern of
195 abnormal sonographic findings in patients with clinical suspicion of chronic liver disease
196 in Sokoto and its environs”. The study also recorded a dominant male gender of up to
197 70.5% of the liver pathological findings. A study also recorded a dominant male gender
198 in liver pathology; about 77% of the liver cases were males [9]. The age range of 40-49
199 years had the highest frequency of 23 (21.5%) and of this age range, there were 16 males
200 (69.6%). This was in contrast with that of Ugwu, A. who recorded 51-60 years as the
201 predominant age range, but was in line with the study conducted by Echejoh in 2008
202 because the age range fell within the age range of 21-50 which was recorded. The age
203 range that recorded the lowest frequency was 80-89 years with a frequency of 6 (5.6%)
204 [10]. The predominant pathological ultrasonic liver findings that were recorded are Liver
205 Cirrhosis accounting for up to 29 and 27.1% frequency and percentage respectively. Of
206 this frequency for liver cirrhosis, male gender recorded a frequency of 18 (62.1%). This

207 was followed by Hepatomegaly which recorded a frequency of 17 and a percentage of
208 15.0%. Fatty liver and Primary liver cell carcinoma (PLCC) had the same frequency and
209 percentage which was 13 (12.1%) each. The findings for the liver are in contrast to
210 similar studies conducted in Asabar, Delta State, Nigeria in which Hepatomegaly was
211 recorded as the most common finding liver pathology, with up to 30.8% of the liver
212 findings, followed by Fatty liver which recorded up to 20.8% of the liver findings [8].
213 The pathological findings are also in contrast with a study conducted by Nwokediuko *et*
214 *al*, 2013 who recorded primary liver cell carcinoma (PLCC) and liver cirrhosis as the
215 commonest findings with a percentage of 44.3% and 20.4% respectively.

216 For the kidneys, the male gender also recorded the highest frequency and percentage
217 which was 107 and 64.1% respectively while female recorded 60 (35.9%). This is in line
218 with the study conducted in Asaba, Delta, Nigeria [7]. The age range that dominated the
219 highest findings ranged from 20-29 years with a frequency and percentage of 43 and
220 25.7% respectively. Of this age range there were 29 males (67.4%) and 14 females
221 (32.6%). This was followed by the age range of 30-39 years which recorded a frequency
222 of 27 (16.2%). This is in contrast with that of Ugwu, 2014 who recorded a dominant age
223 range of 31-40 years (21.6%) for kidney findings. The age range that recorded the lowest
224 frequency was 80-89 years which had a frequency of 6 (3.6%) which is in lined with the
225 study in Asaba that had 81-90 years with a frequency of 2 only [8].

226 The pathological finding that dominated the kidney findings after the analysis was
227 Pyelonephritis which had a frequency and percentage of 69 (41.3%). Of the recorded
228 pyelonephritis, male gender were mostly affected, they recorded a frequency of up to 47
229 (68.1%) while female gender recorded 22 (21.9%). This was followed by renal
230 parenchyma disease (grades I, II, III, or IV) which recorded a frequency and percentage
231 of 35 (21.0%). Hydronephrosis was the third common finding with 25 and 15.0%
232 frequency and percentage respectively. This contrasted the study in Asaba which
233 recorded the following in their order of frequency and percentage; Nephropathy 33
234 (23.7%), Hydronephrosis 30 (21.6%) and Renal cyst 30 (21.6%).

235 **Conclusion**

236 The study revealed that male gender is at higher risk of having both liver and kidney
237 diseases in Sokoto metropolis than the female and the age of those at higher risk ranged
238 from 20-49 years.

239 Liver cirrhosis and pyelonephritis are the commonest diseases of the liver and kidneys
240 respectively.

241 Ethical approval was obtained from the Human Research and Ethics Committee of
242 UDUTH and specialist hospital Sokoto before the study commenced.

243 **REFERENCES**

- 244 1. Jane B. 2004 *Abdominal Ultrasound; How, Why And when*, 2nd ed. London,
245 ChurchHill Livingstone. 2004; pp 92, 166.
- 246 2. Rumack CM, Wilson SR, Charboneau JW, Levine D. *Diagnostic Ultrasound*. 4th
247 Ed, Philadelphia, Elsevier. 2011; pp 18-30.
- 248 3. Fikri M A, Ashraf F H and Peter C. Clinical Ultrasound Physics. *Journal of*
249 *Emergencies, Trauma and Shock* 2011 **4**(4): 501–503.
- 250 4. Matthias H and Reihls T. *Ultrasound Teaching Manual; The basics of performing*
251 *and interpreting Ultrasound Scans*, New York, Georg Thieme. 1999; P 23, 37.
- 252 5. Hansen KL, Nielson BM, Ewertsen C. Ultrasonography of the Kidney; A Pictorial
253 review. *Journal of Diagnostics*, 2015; **6**(2) 1-2.
- 254 6. Charles O. Renal Relevant Radiology: Use of Ultrasound in Kidney Disease and
255 Nephrology Procedures. *Clinical Journal of American Society of Nephrology*.
256 2014; **9**(2):37381.
- 257 7. Joseph EN. (2017), Sample size determination in qualitative and quantitative
258 research. Available at [https://educacinfo.com/sample-size-determination-
259 qualitative-quantitative-research/](https://educacinfo.com/sample-size-determination-qualitative-quantitative-research/). Accessed on 13th October 2018.

- 260 8. Ugwu A. Common Findings In Liver And Kidneys Of Patients Undergoing
261 Ultrasound Examination In Asaba, Nigeria [Online]. Available at
262 <https://afribary.com> Accessed on (15th July, 2017).
- 263 9. Nwokediuko SC, Osuala PC, Uduma UV, Alaneme AK, Onwuka C, Mesigo C.
264 Pattern of Liver Disease Admissions in Nigeria Tertiary Institution. *Journal of*
265 *Public Health Africa*. 2013; **16**(3): 339-342.
- 266 10. Echijoh GO, Tanko MN, Manasseh AN, Silas OA, Ogala-Echejoh SE and
267 Mandong BM. Liver Cirrhosis in Jos North-Central Nigeria. *Jos Journal of*
268 *Medicine*. 2008; **3**(1) 4-5.

269

UNDER PEER REVIEW