

14  
15  
16  
17  
18  
19  
20  
21

**Effect of Sorghum-Tigernut *Ibyer* (A Traditional Gruel) on the Fasting Blood Glucose Levels and Body Weight of Alloxan-Induced Diabetic Rats**

---

**ABSTRACT**

**Background:** There is growing interest in the use of natural foods in the management of chronic diseases like diabetes. *Ibyer* is a fibre rich gruel consumed amongst the Tiv people of Benue State made from whole sorghum or millet flours.

**Aim:** The aim of the study was to evaluate the effect of sorghum-tigernut *ibyer* on the fasting blood glucose levels and body weight of alloxan monohydrate-induced diabetic rats.

**Methods:** Sorghum flour (SF) and tigernut flour (TNF) were blended at different proportions (100:00; 90:10; 80:20; 70:30) for the purpose of *ibyer* production. The flour samples were subjected to proximate analysis using standard analytical procedures, the sensory attributes of *ibyer* produced from the different flour samples was evaluated on a 9-point hedonic scale. Thirty (30) male Wistar rats (100–180 g body weight) were grouped into five (1-5) each group containing six rats. They were induced with diabetes by injecting them with 150ml/kg of body weight with alloxan monohydrate dissolved in saline water (0.9% NaCl) except for group 1. Blood samples were collected from the tail of the rats, prior to induction, 48hrs after induction and 72hrs after three days of continuous feeding with test diet. Fasting blood glucose was measured using a standard glucometer and test strips.

**RESULTS:** The proximate composition of the flour samples showed a reduction in moisture (10.12-4.46), crude protein (7.45-6.25), and carbohydrate (76.10-70.53) but with a significant enhancement in the fibre content (1.00-3.00%), crude fat (3.22-12.86%) and crude ash (2.11-2.93%) depending on the level of substitution. The sensory attributes indicated that *ibyer* produced from the flour samples were generally acceptable though that produced from 10% tigernut flour addition was most preferred. Fasting blood glucose levels after 72hrs of feeding ranged from 5.52mmol/L in group through to 5.45mmol/L in group 5 with group 2 having the highest average fasting blood glucose levels of 17.60mmol/L. the body weight changes were irregular in all diabetic rat groups.

**CONCLUSION:** The results indicated that sorghum-tigernut *ibyer* has the potential to lower fasting blood glucose.

14  
15  
16  
17  
18  
19  
20  
21

**Keywords:** *Sorghum, Tigernut, Ibyer, Diabetic rats, Blood glucose level*

**1. INTRODUCTION**

Diabetes is a chronic disease in which food (particularly the nutrient – carbohydrate) is not properly absorbed in the body to be used as energy [1]. Diabetes and hypertension are global health disorders afflicting millions of people worldwide with an ever-increasing incidence and prevalence the upsurge of diabetes in Africa has been linked with rapid urbanization and changing dietary habits [2, 3]. Use of indigenous foods has been advocated

22 to reduce the incidence of chronic, diet-related non-communicable diseases such as obesity,  
23 diabetes, cardiovascular diseases and stroke [4]. The development of diabetes mellitus,  
24 obesity, cancer and cardiovascular disease (CVD) has been reported to be linked to the  
25 intake of high glycemic index (GI) foods while intake of low GI foods has been shown to play  
26 a positive role in the management of these diseases [5].

27  
28 *Ibyer* is an indigenous non-alcoholic gruel made from cereals (maize, sorghum and millet),  
29 consumed in Nigeria especially in Benue state by Tiv people [6]. It is prepared by cooking  
30 reconstituted whole cereal flour or paste in water and is often served with beans products  
31 such as *akpukpa* (native bread) and beans cake. Sour type (*ibyer-i-angen*) arises as result of  
32 the fermentation step that is undertaken before the porridge is prepared. The flour or wet  
33 milled paste is usually reconstituted with water to form slurry or paste after which it is left for  
34 a given time to enable fermentative microbial organisms to act on the product, producing the  
35 characteristic sourness associated with the product. Traditionally, the fermentation time lasts  
36 overnight.

37 Sorghum (*Sorghum Bicolor L. Moench*) is the fifth most important cereal after wheat, rice,  
38 maize and barley in terms of production and utilization. The total world annual sorghum  
39 production is over 60 million tons from cultivated area of 45 mile for food, alcoholic  
40 beverages and the grain is one of the staple foods for poor and rural people. Sorghum is  
41 gluten-free thus can be consumed by people with celiac diseases [7]. Other important  
42 nutrients of sorghum include; dietary fibre, fat-soluble and B-vitamins and minerals [8].  
43 Sorghum flour is used for flours, porridge and side dishes, malted and distilled beverages  
44 and special food such as popped grain, its protein content is higher than many grains.  
45 Sorghum is rich in antioxidant which is believed to help lower the risk of cancer, diabetes,  
46 heart diseases and other neurological diseases.

47 Tigernut has been used extensively mainly for human consumption in Spain [9]. It was an  
48 important food in ancient Egypt [10]. Nowadays tigernuts are cultivated in Northern Nigeria,  
49 Mali, Senegal, Ahana and Togo where they are used primarily uncooked as a side dish [11].  
50 The flour is a good alternative for other flour like wheat flour, as it is gluten free and good for  
51 people who cannot take gluten in their diets. It is also used to make cakes and biscuits and  
52 the oil is used for cooking [12]. The dietary fibre content of tigernut is effective in the  
53 treatment and prevention of diseases such as colon cancer, coronary heart diseases,  
54 obesity, diabetes and gastro-intestinal disorders [13]. This research was aimed at  
55 determining the effect of sorghum-tigernut '*Ibyer*' on fasting blood glucose levels of alloxan  
56 monohydrate induced diabetic rats.

57  
58

## 59 **2.0 MATERIAL AND METHODS**

60  
61

### 62 **2.1 Sample Preparation**

63 The red sorghum grain variety and dried tigernuts were purchased from Wadata Market in  
64 Makurdi, Benue State, Nigeria. Alloxan monohydrate was purchased from Sigma Chemical  
65 Co. (St. Louis, MO, USA).

### 66 **2.2 Sample Preparation**

67 **2.2.1 Sorghum Flour:** The preparation of sorghum flour was done according to the  
68 method described by Sergio [14]. Sorghum grains were dry-cleaned by handpicking, washed  
69 in a large volume of water to remove impurities and sun-dried. The dried sorghum grains  
70 were milled into sorghum flour using hammer mill.

71 **2.2.2 Tigernut Flour:** Tigernut flour was prepared according to the method described by  
72 Adejuyitan [15]. Dry tigernuts (brown variety) were sorted to remove unwanted materials like  
73 stones, pebbles and other foreign seeds before washing with tap water. The cleaned nuts  
74 were sun dried to a moisture content of about 13% then milled and sieved.

76 **2.3 Formulation of Flour Blends:** the two flours (SF and TNF) were blended in a ratio  
77 of 100: 0, 90:10, 80:20 and 70:30 respectively.

79 **2.4 Preparation of *Ibyer-i-angen*:** After the preparation of the sorghum-tigernut flour  
80 blends, the flour samples were mixed with 200ml of water in five different containers with a  
81 cover and kept at ambient temperature ( $30\pm 2^{\circ}\text{C}$ ) for 24hrs to allow natural fermentation take  
82 place. The fermented paste from the two flour blends was diluted with water and cooked in a  
83 tower aluminium pot containing about 80cl of water on a gas cooker. The mixture was stirred  
84 for about 7-8min to achieve desired consistency. For feeding to the animals, the  
85 prepared/cooked *ibyer* was oven dried and milled.  
86 Flow chat for production of *ibyer-i-angen* is shown in figure.1



97 **Fig.1: Traditional method of production of *Ibyer-i-angen***

## 98 **2.5 Determination of Proximate Composition**

### 99 **2.5.1 Determination of Moisture Content**

100 The moisture content was determined by the method of AOAC [16]. Cleaned crucibles were  
101 dried in the oven at  $100^{\circ}\text{C}$  for 1hr to obtain a constant weight and then cooled in the  
102 desiccators. A known quantity of 2.0g of the sample was weighed into the crucible and dried  
103 at  $100^{\circ}\text{C}$  until a constant weight was obtained.

104  
105 Percentage moisture content =  $\frac{\text{weight loss}}{\text{weight of sample}} \times \frac{100}{1}$

### 108 **2.5.2 Determination of Ash Content**

109 The AOAC [16] method was used to determine the ash content. 2g of sample was weighed  
110 into a pre-heated cooled crucible. This sample was charred on a Bunsen flame inside a fume  
111 cupboard. Sample was transferred into a pre-heated muffle furnace at  $550^{\circ}\text{C}$  for 3-5hrs until  
112 a white or light grey ash was obtained.

113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163

$$\text{Ash content} = \frac{\text{weight of ashing}}{\text{weight of sample}} \times \frac{100}{1}$$

### 2.5.3 Determination of Crude Protein

Crude protein was analysed according to the method of AOAC [17]. A portion (2g) of sample was weighed into a cleaned kjeldah digestive flask, 8g of catalyst mixture (anhydrous copper-sulphate mixture)  $\text{CuSO}_4$  and few pieces of anti-bumps glass seed were also put in the flask. 10ml of concentrated  $\text{H}_2\text{SO}_4$  were added and gently swirled until no particles of the sample was adhere to the side of the flask. It was then heated strongly until the solution became a clean light blue-green color. This was heated for one hour. This was allowed to cool quantitatively transferred into a 20ml volumetric flask and up to the mark. Blank determination was carried out. The Markham Semi-micro was used for steam distillation. Digestion sample (5ml) was pipette into the unit, 7ml of 50% NaOH was added and the unit closed and heated the liberated ammonia distilled into 5ml boric indicator mixture. This distillate (20ml) was collected and tide of the delivery tube was rinsed with distillate; was then titrated with 0.1 N HCl until green colour changed to purple.

$$\% \text{ Nitrogen} = \frac{\text{titre} - \text{blank} \times 0.0014 \times \text{normal} (0.4) \times (DF25)}{\text{weight of sample}} \times 100$$
$$\% \text{ crude protein} = \% \text{ Nitrogen} \times 6.25 \text{ (general factor)}$$

### 2.5.4 Determination of Fat

The Soxhlet extraction method of AOAC [16] was used. A Soxhlet extraction with a reflux condenser and a 500ml round bottom flask was fixed. 2g of sample was weighed into a labeled thimble. Petroleum ether (300ml) was filled into the round bottom flask. The extractor thimble was sealed with cotton wool. The soxhlet apparatus was allowed to reflux for 6hrs, the thimble was then removed with care and the petroleum ether collected at the top and drained into a container for re-use. When the flask was free of ether, it was then removed and dried at  $105^\circ\text{C}$  for 1hr in an oven, cooled in desiccators and weighed.

$$\text{Fat content} = \frac{\text{weight of fat}}{\text{weight of sample}} \times \frac{100}{1}$$

### 2.5.5 Determination of Crude Fibre Content

The crude fibre content was determined according to the method of AOAC [16]. 1g of sample was weighed into 250 conical flasks. 100ml  $\text{SO}_2$  (1.25%) was added and heated on hot plate to boil for 30mins. The  $\text{SO}_2$  was drained through Muslim cloth. The filtrate was washed with hot distilled water till it is free from acid and transferred to other 250ml conical flasks. 100ml of 1.25% KOH was added and heated to boil for 30mins. The alkaline was drained and the filtrate washed with hot distilled water and then acetone to remove the KOH. The filtrate was transferred to a crucible and dried in hot air oven at  $105^\circ\text{C}$  to constant weight. Percentage crude fibre was determined using the formula;

$$\text{Crude fibre} = \frac{w_2 - w_1}{w_1} \times \frac{100}{1}$$

### 2.5.6 Determination of Carbohydrate Content

Carbohydrate content of the sample was determined by the different methods described by Ihekoronye and Ngoddy [18].

$$\% \text{ carbohydrate} = 100 - (\% \text{ moisture} + \% \text{ fat} + \% \text{ ash} + \% \text{ protein} + \% \text{ fibre})$$

## 2.6 Sensory Analysis

164 Sensory characteristics of *ibyer-i-angen* prepared from sorghum and tigernut was evaluated  
165 and assessed by 20 semi-trained panelists of Department of Food Science and Technology,  
166 University of Agriculture, Makurdi who are familiar with *ibyer-i-angen*. Fresh samples of  
167 cooked porridge/gruel were assessed for appearance, taste, mouth feel, flavour and general  
168 acceptability. The judges recorded the quality characteristics of each sample using nine-  
169 point hedonic scale, where 1 = dislike extremely, 5 = neither like nor dislike, and 9 = like  
170 extremely as described by Iwe [19].  
171

## 172 **2.7 Animal Treatments**

173 Thirty (30) male Wistar rats (100–180 g body weight) were obtained from the small Animal  
174 Experimental Unit of the National Veterinary Research Institute Vom, Plateau State. The rats  
175 were housed under standard hygienic conditions in metal cages with wood shavings as  
176 bedding. Rats were also kept under natural thermal environmental conditions with ambient  
177 temperature of 24 C–26 C and relative humidity of 70%–80%, and approximately alternating  
178 12hr light/dark cycles. They were given access to a standard pelletized rat chow and water  
179 *ad libitum*. The animal experiments adhered to the Guide for the Care and Use of Laboratory  
180 Animals.  
181

## 182 **2.8 Induction of Experimental Diabetes**

183 Diabetes mellitus (DM) was induced in overnight-fasted rats by a single i.p. injection of  
184 freshly-prepared alloxan monohydrate, dissolved in a cold physiological saline (0.9% NaCl)  
185 solution at the dose rate of 150 mg/kg body weight. The animals were given free access to  
186 5% glucose solution in order to overcome the alloxan-induced hypoglycaemia for the first  
187 one-hour post-treatment with alloxan monohydrate. Blood glucose concentration of the rats  
188 was estimated 48hours after alloxan administration and DM was confirmed by analysis of  
189 blood samples, collected from the vein at the tip of the tail, using a portable blood  
190 glucometer and glucose test strips (On Call®Plus, Hannover, Germany). Animals with blood  
191 glucose concentration equal or more than 14 mmol/L were considered diabetic and used in  
192 the entire experimental group. Animals were weighed and randomly assigned to six groups  
193 and treated as follows:

194 Group 1, Control group: (normal saline only);

195 Group 2, Diabetic + Treated with Sample A (alloxan 150 mg/kg, i.p);

196 Group 3, Diabetic + Treated with Sample B (alloxan 150 mg/kg, i.p);

197 Group 4, Diabetic + Sample C (alloxan 150 mg/kg, i.p);

198 Group 5, Diabetic + Sample D (alloxan 150 mg/kg, i.p);

199 Experimental animals were treated with the samples, after 48hrs of verification of diabetes,  
200 daily for 3 days.  
201

## 202 **2.9 Measurement of Blood Glucose**

203 Experimental animals were rearranged according to the blood glucose concentration, except  
204 the control group, before commencement of treatment. Blood glucose concentration in all  
205 experimental groups were recorded following overnight fasting 72h after commencing the  
206 feed trials, using a portable glucometer (On Call®Plus, Hannover, Germany) and glucose  
207 test strips.  
208

## 209 **2.10 Measurements of Body Weight**

210 Rats were weighed individually at Day 0 (before induction of DM), day 2 (post induction of  
211 DM) and 72 h (post-treatment with feed samples) using a digital precision weighing balance,  
212 and the body weights were recorded to calculate the body weight gains.  
213

214

215

## 216 **2.11 Statistical Analysis**

217 Data collected were subjected to Analysis of Variance (ANOVA). Means were separated with  
 218 Fisher's LSD using SPSS software (2009 model) and judged significantly different at 95%  
 219 confidence level ( $p < 0.05$ ).

220

### 221 3. RESULTS AND DISCUSSION

#### 222 3.1 Results

##### 223 3.1.1 Proximate composition of blends of sorghum and tigernut flours

224

225 Table 1 shows the results obtained for the proximate composition of blends of sorghum and  
 226 tigernut flours. The values of the proximate composition expressed in percentage showed  
 227 significant difference at  $p < 0.05$  between all the samples, however sample B (2.05), C (2.67)  
 228 and also sample D (3.18) had significant difference in the fibre content, there was a  
 229 difference in sample A (1.00) with the lowest value. Some of the parameters had a  
 230 decreasing trend across the samples from A - D with moisture content ranging from 5.49 to  
 231 10.12, crude protein ranged from 6.28 to 7.45, and carbohydrate ranged from 72.11 to 76.10  
 232 whereas crude fat showed an increasing trend of 3.22 to 10.22 and crude ash increased  
 233 from 2.11 to 2.72 having the lowest values of all the parameters with significant difference  
 234 between the samples.

235

236 **Table 1: Proximate composition of Blends of sorghum and tigernut flours**

237 *Mean with different superscript within the same column are significantly different at*  
 238 *( $P < 0.05$ ).*

239 LSD: Least Significant Difference

240 SF: Sorghum Flour; TNF: Tigernut flour.

241 A: 100% SF. B: 90% SF and 10% TNF C: 80% SF and 20% TNF D: 70% SF and 30%

242 TNF

243

Parameters (%)	Samples				LSD
	A	B	C	D	
Moisture	10.12 <sup>a</sup> ±0.02	8.23±0.01 <sup>b</sup>	6.23 <sup>c</sup> ±0.01	5.49 <sup>d</sup> ±0.01	0.02
Protein	7.45 <sup>a</sup> ±0.00	7.19 <sup>b</sup> ±0.00	7.01 <sup>c</sup> ±0.00	6.28 <sup>d</sup> ±0.01	0.02
Fat	3.22 <sup>e</sup> ±0.01	4.84 <sup>d</sup> ±0.00	8.43 <sup>c</sup> ±0.00	10.22 <sup>b</sup> ±0.00	0.01
Fibre	1.00 <sup>d</sup> ±0.03	2.05 <sup>c</sup> ±0.03	2.67 <sup>b</sup> ±0.01	3.18 <sup>a</sup> ±0.02	0.04
Ash	2.11 <sup>d</sup> ±0.01	2.39 <sup>c</sup> ±0.00	2.65 <sup>b</sup> ±0.01	2.72 <sup>a</sup> ±0.00	0.01
Carbohydrate	76.10 <sup>a</sup> ±0.02	75.32 <sup>b</sup> ±0.01	73.01 <sup>c</sup> ±0.02	72.11 <sup>d</sup> ±0.01	0.03

##### 244 3.1.2 Sensory properties of lbyer produced from blends of sorghum and tigernut 245 flours.

246 The results of the sensory evaluation of lbyer produced from blends of sorghum and tigernut  
 247 flours are shown in table 2. The sensory qualities measured were appearance, mouth feel,  
 248 taste, flavour and general acceptability. The results showed that the appearance ranged  
 249 from 6.20 to 7.15 across the samples with sample D having the lowest sample A, B and C  
 250 having similar mean rating. Flavour properties were ranged from 6.05 to 6.75, taste ranged  
 251 from 6.50 to 6.90 and general acceptability ranged from 6.45 to 7.40 with no significant  
 252 difference in all the samples. There was significant difference in sample C, and D but no  
 253 difference in sample A and B in the mouth feel across the samples. Taste was highest in  
 254 sample B. All the samples were generally acceptable though sample B was most preferred.

255  
256  
257

**Table 2: Results of the Sensory scores of “lbyer” produced from blends of sorghum and tigernuts flour.**

Sample	A	B	C	D	LSD
Appearance	7.150 <sup>a</sup>	7.050 <sup>a</sup>	6.850 <sup>a</sup>	6.200 <sup>ab</sup>	0.943
Flavour	6.650 <sup>a</sup>	6.750 <sup>a</sup>	6.750 <sup>a</sup>	6.050 <sup>a</sup>	1.171
Mouth feel	7.150 <sup>a</sup>	6.950 <sup>a</sup>	6.300 <sup>ab</sup>	5.600 <sup>bc</sup>	0.958
Taste	6.900 <sup>a</sup>	6.750 <sup>a</sup>	6.650 <sup>a</sup>	6.500 <sup>a</sup>	1.168
General acceptability	7.400 <sup>a</sup>	7.150 <sup>a</sup>	6.750 <sup>a</sup>	6.450 <sup>a</sup>	1.015

258 *Mean with different superscript within the same column are significantly different at (P<0.05).*

259 LSD: Least Significant Difference

260 **Key:**

261 SF: Sorghum Flour; TNF: Tigernut flour.

262 A: 100% SF. B: 90% SF and 10% TNF C: 80% SF and 20% TNF D: 70% SF and 30%

263 TNF

264

265

### **3.1.3 Effect of Sorghum-tigernut lbyer on the Fasting Blood glucose levels of alloxan induced diabetic rats.**

266

267

268

269

270

271

272

273

274

275

The results of the fasting blood glucose levels of alloxan induced diabetic rats before and after feeding with test diets are presented in figure 2. The results showed that the average fasting glucose levels of the diabetic rats before feeding with the test diets ranged from 22.30mm/L to 32.00mm/L and 3.15mm/L to 6.05mm/L in the non-diabetic rats. However, there was considerable reduction in average fasting blood glucose levels of the diabetic rats 0hrs through to 72hrs post treatment. The results ranged from 5.52mm/L from group 1 through to 5.45mm/L in group 5 with rats' group 2 having the highest average fasting blood glucose levels of 17.60mm/L

276

### **3.1.4 Body Weight of Rats before and after Feeding with Test Diet**

277

278

279

280

281

282

283

284


285

286

287

288

Figure 3 presents results of body weight changes during the experimental period. Results showed steady weight increase in the non-diabetic rat group from 98.717 to 118.450g from 0hrs to 72hrs post feeding. However, the diabetic rat groups showed slight variations in body weight changes with groups 2 and 4 showing steady decrease in weight ranging from 151.075g to 175.017 and 154.100g to 171.700 in groups 2 and 4 respectively. However, groups 3 and 5 had body weight changes in no particular order. Group 4 showed an initial average body weight decrease from 165.94 to 147.72g 48hrs after induction and a subsequent increase from 147.72 to 153.660g 72hrs after feeding. While rat groups 5 showed an initial average weight gain from 166.925g at 0hrs to 175.500g 48hrs after induction but decreased after 72hrs to 155.025g.



289  
 290  
 291  
 292  
 293  
 294  
 295  
 296  
 297  
 298

**Fig. 2. Fasting blood glucose levels of alloxan induced diabetic rats before and after feeding with test diet.**

Group 1(control; normal saline) Group 2(sample A at 100%SF) Group 3(sample B at 90%SF and 10%TNF) Group 4(sample C at 80%SF and 20% TNF) and Group 5 (sample D at 70%SF and 30%TNF)

key:  
 SF; Sorghum flour  
 TNF: Tigernut flour

UNDER PUBLICATION



299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

**Fig. 3. The average body weight of the rats (grams).**

Group 1(control; normal saline) Group 2(sample A at 100%SF) Group 3(sample A at 90%SF and 10%TNF) Group 4(sample C at 80%SF and 20% TNF) and Group 5 (sample D at 70%SF and 30%TNF)

Key: SF; Sorghum flour

TNF: Tigernut flour

### 3.2 Discussion

#### 3.2.1 Proximate composition of sorghum and tigernut flour blends.

There was a significant ( $P<.05$ ) difference in the moisture content of all the samples. The moisture content of the sample flour decreased with increase in the tigernut flour substitution. The gradual decrease in moisture content might be due to the low moisture content found in tigernuts. The result is in agreement with those of Abiose *et al* [20] who reported a decrease in moisture content in flour. Low moisture content is an indication of good keeping quality of any food product; it also reduces microbial growth thereby extending the shelf life of the product.

The protein content of sample A, B, C, and D were determined to be 7.45%, 7.19%, 7.01%, and 6.28% respectively. The study also showed that there was significant ( $P<.05$ ) difference in protein content of the samples. The protein content decreased with increase in tigernut flour. This may be attributed to the low protein content of tigernut as reported by Ade-Omowaye *et al* [21]. Protein content may also vary due to the genotype, temperature, soil and fertility and environmental conditions during plants development. The protein content of the four samples meets the recommended dietary allowance (RDA) of 0.8g of protein per kilogram of body weight per day [22].

The flours of A, B, C, and D contained fat that ranged from 3.22%, to 10.22%, With sample A having the lowest fat content and sample D having the highest fat content. It has been reported that the sorghum flour contained about 3% fat [18]. Ibrahim [23] also reported to have 4.25% fat in sorghum flour. There was significant ( $P<.05$ ) difference in the fat content

328 of all the samples. The increase in the fat content is due to the increase in the tigernut  
329 substitution [21, 24]. Zahras and Ahmed [25] reported fat content in tigernut flour about  
330 38.6%. Tigernut tubers contains up to 30% of non-drying oil which is used for cooking and  
331 making soap [26]. Hence, defatting the nuts before utilization may yield better result.  
332 The fibre content of sample A, B, C, and D were found to be 1.00%, 2.05%, 2.67%, and  
333 3.18%. These results are in accordance with Ade-Omowaye *et al* [21] who reported an  
334 increase in the fibre content of tigernut-wheat flour with increasing tigernut flour substitution.  
335 There was a significant ( $P<.05$ ) difference in the fibre content of all the samples indicating  
336 that tigernut which has been reported to be a rich source of dietary fibre had a huge impact  
337 on the fibre content of the product. The increased fibre content is a good source of dietary  
338 fibres which lowers serum cholesterol and prevents gastrointestinal problems. Crude fibre  
339 also adds bulk to diet and is advantageous in bowel movement.  
340 The values of ash content of samples A, B, C, and D were found to be 2.11%, 2.39%,  
341 2.65%, and 2.72 with sample A having the lowest value and D the highest value. Ihekoronye  
342 and Ngoddy [18] reported an ash composition of 2% in sorghum flour. There was a  
343 significant ( $P<.05$ ) difference in the ash content of all the samples as the level of tigernut  
344 substitution increased. The ash content result is in agreement with the reports of Adebowale  
345 [27] and comparable with those of Aremu *et al* [28] who reported ash content of 2.38 to 2.57.  
346 Ash content is an indication of the mineral content in flour.  
347 Carbohydrates decreased from 76.10 to 70.53 which is within the range of those reported by  
348 Ihekoronye and Ngoddy [18]. There was a significant ( $P<.05$ ) difference in carbohydrate  
349 content of all the samples. The decrease in carbohydrate content is attributed to an increase  
350 in tigernut flour. The powdered products however, are higher in carbohydrates and this may  
351 be attributed to reduced moisture content. Carbohydrates are extremely important as  
352 component of foods which they are sources of energy, flavour and bulk.

353

### 354 **3.2.2 Quality characteristics on the sensory attributes of lbyer produced from** 355 **blends of sorghum and tigernut flour.**

356 There was no significant ( $P<.05$ ) difference in appearance of the sample A, B and C but  
357 there was a significant ( $P<.05$ ) difference in sample D. This could be attributed to the fact  
358 that tigernut has a colour that is distinct from that of sorghum and so as the amount of  
359 tigernut in each sample increased across the rows there would be a considerable change in  
360 the appearance of the products. It was however noted that the sensory scores for  
361 appearance were considerable high even though with slight differences. This means all  
362 products were appealing to the eyes.

363 There was no significant ( $P<.05$ ) difference in the mouth feel of sample A and B but there  
364 was a significant difference in sample C and D. The differences in mouth feel could be  
365 attributed to higher amounts of fibre in samples C and D. *lbyer* is a gruel made from whole  
366 cereals, particularly sorghum and millet, due to this it has a coarse feel when consumed.

367 There was also no significant ( $P<.05$ ) difference among the samples in terms of taste,  
368 flavour and general acceptability. All the samples were generally acceptable indicating that  
369 *lbyer* of acceptable eating qualities can be produced from flour blends and sorghum and  
370 tigernuts. These results are in agreement with the reports of Kure and Wyasu [6] who  
371 reported general acceptability of *lbyer* produced from sorghum and soy addition meaning  
372 addition of other legumes to the product may not affect the eating qualities of the product.

373

### 374 **3.2.3 Effect of sorghum-tigernut lbyer on the Fasting Blood Glucose Levels Alloxan** 375 **Induced diabetic rats.**

376 *lbyer* made from flour blends of sorghum and tigernut at different levels of substitution. The  
377 results showed that the average fasting blood glucose of the rats in the various groups  
378 differed significantly. After three days of consuming approximately 230g of the test diets  
379 daily, their fasting blood glucose levels dropped significantly. The results showed that rat  
380 groups fed with samples containing higher amounts of tigernut had lower fasting blood

381 glucose levels than those of the control group. This could be attributed to the fact that  
382 tigernut is a rich source of fibre [29, 30]. Fibre has been reported to exert some  
383 hypoglycemic effects in subjects with type II diabetes. It is also reported that arginine an  
384 amino acid (not determined) found in tigernut has the potential of stimulating the release of  
385 insulin thereby ameliorating the effect of diabetes [31]. This could be one of the reasons for  
386 the lowered blood glucose levels after three days of consecutive feeding with the test diet.  
387 Several studies have shown that sorghum extracts and sorghum rich diets exert  
388 hypoglycemic effects in either human subjects or lab animals, thus agreeing with the findings  
389 of this research [32, 33]. The average fasting blood glucose levels of the non-diabetic rats  
390 ranged from 22.30mm/L to 32.00mm/L and 3.15mm/L to 6.05mm/L. However, there was  
391 considerable reduction in average fasting blood glucose levels of the diabetic rats 0hrs  
392 through to 72hrs post treatment. The results ranged from 5.52mm/L from group A through to  
393 5.45mm/L in group E with rats group B having the highest average fasting blood glucose  
394 levels of 17.60mm/L.

395

#### 396 **3.2.4 Effect of Sorghum-tigernut lbyer on the weight of alloxan induced diabetic rats.**

397 Body weight changes during the experimental period showed steady weight increase in the  
398 non-diabetic rat group from 98.717 to 118.450g from 0hrs to 72hrs post feeding. However,  
399 the diabetic rat groups showed slight variations in body weight changes with groups 2 and 4  
400 showing steady decrease in weight ranging from 151.075g to 175.017 and 154.100g to  
401 171.700 in groups 2 and 4 respectively. However, groups 3 and 5 had body weight changes  
402 in no particular order. Group 4 showed an initial average body weight decrease from 165.94  
403 to 147.72.g 48hrs after induction and a subsequent increase from 147.720 to 153.660g  
404 72hrs after feeding. While rat groups E showed an initial average weight gain from 166.925g  
405 at 0hrs to 175.500g 48hrs after induction but decreased after 72hrs to 155.025g. This trend  
406 could be attributed to the fact the diabetic rats were fed a carbohydrate and fibre-rich diet  
407 only consecutively for three days (72hrs). The mono diet could have effect on normal weight  
408 gain and growth rate resulting to the decrease in weight of the animals.

409

410 **4. CONCLUSION**

411 The results of the proximate composition revealed that, inclusion of tigernut flour in to  
412 sorghum flour at levels of up to 30% resulted in notable increase in the fat, fibre and ash  
413 with the substitution of both samples while protein, moisture and carbohydrate decreased.  
414 The significant increase in fibre content could be nutritionally advantageous in diabetic diets.  
415 Results of the sensory evaluation of *ibyer* produced from sorghum and tiger flour blends  
416 showed that all the samples were generally acceptable, though sample b produced from  
417 90:10% sorghum and tigernut flour blends was most preferred. From the results, it is evident  
418 that nutritious diets can be formulated by complementing unexploited tuber and cereals like  
419 tigernut and sorghum respectively, such blends could be used to diversify their uses to  
420 develop new products.

421 The fasting blood glucose levels of the diabetic rats have shown that tigernut inclusion in the  
422 sorghum flour for *ibyer* production has a significant role to play in lower blood glucose levels.  
423 Although the post prandial glucose levels in the test animals was not studied. Results  
424 showed that this product can be used in the management of diabetes. Although the  
425 mechanism of action of in achieving these results was not studied and so remains unclear.  
426 The effect of the diet on the weight changes showed that, the diet had properties of reducing  
427 weight, or maintaining weight in subjects. This could be insight to weight management  
428 amongst the diabetes and even obesity.

429 **REFERENCES**

- 431 1. Bellows L and Nichols K (2012). Position Statement of the American Diabetes  
432 Association. Nutrition Recommendation and intervention for diabetes: Diabetes  
433 Care, vol.31,suppl.1.
- 434 2. Jervel J (1995). Poor Nutrition and Chronic diseases. United Nations administrative  
435 Committee on coordination sub- committee on Nutrition. (Acc/SCN) news Vol.13:  
436 Pg. 10-14.
- 437 3. SCN (2006): Diet related chronic diseases and the double burden of malnutrition  
438 committee on Nutrition News (Development in international Nutrition) Vol.33: Pg.13-  
439 19.
- 440 4. Odenigbo UM (2001). Incorporation of Afzelia Africana in dietary management of  
441 type-2 diabetes in Nnewi, Anambra State, Nig. Ph.D. Thesis. Home Sci. Nutrition  
442 and Dietetics, University if Nig. Nsukka.
- 443 5. Ludwig-Wasserman NU (2009): New Diabetes Nutrition Therapy Recommendations;  
444 what you need to Know.
- 445 6. Kure OA. and Wyasu G. (2013). Influence of natural fermentation, malt addition and  
446 soya- fortification on the sensory and physico-chemical characteristics of *Ibyer*-  
447 Sorghum gruel. *Advances in Applied Science Research*. Vol 4(1): Pg.345-349
- 448 7. Taylor JRN, Achoch TJ and Bean SR (2006). Novel Food and Non-food uses for  
449 Sorghum and Millets. Articles in press *Journal of Cereal Science*, Vol.44: Pg.252-  
450 271.
- 451 8. Waniska RO, Poe JH and Bandy O.R. (2004). Effects of growth conditions on grain  
452 molding and phenols in sorghum Caryopsis, *Journal of cereal Science*, Vol. 10, Pg.  
453 217-225.
- 454 9. Gambo, A. (2012). Sensory evaluation of yoghurt produced from blend of Cowmilk,  
455 Soymilk and Tigernut milk. In: Proceedings of 36th Annual Conference/AGM of  
456 Nigerian Institute of Food Science and Technology, Lagos October. Pp.440- 441.
- 457 10. Neigbi OA (1992): Quality characteristics of Candies Produced from Tigernuts  
458 Tubers (*Cyperusesculentus*) and Melon Seeds (*Colocynthiscitrulus*) Milk Blends.  
459 *Global journal of Science Frontier Research: Agriculture and veterinary*. Vol.15 Issue  
460 2 Version

- 461 11. Omode MO, Ibiom OF and Okoi A. (2009): Studies on the fungi and phytochemical  
462 and proximate composition of dry and fresh tigernuts (*Cyperuseculentus*. L).  
463 *International Research journal of Biotechnology*. Vol.4 (1) pp. 11-14.
- 464 12. Wise GE (2009): Tigernut-Soy Milk: Preparation, Proximate Composition and  
465 Sensory Qualities. *International Journal of food and Nutrition Science*. Vol.1. No 4.
- 466 13. Anderson KN and Helena D (1994): Dietary Treatment of Diabetes: A Systematic  
467 Review medtex International.
- 468 14. Sergio Serna-Saldivar (2010). Cereal Grains: Properties, Processing, and Nutritional  
469 Attributes CRC Press, ISBN 97814398156012010
- 470 15. Adejuyitan JA, Otunolam ET, Akandem ET, Bolarinwa IF and Oladokun FM (2009):  
471 Some physicochemical properties of flour obtained from fermentation of tigernut  
472 (*Cyperuseculentus*) sourced from a market u Ogbomoso, Nigeria. *African Journal of*  
473 *Food Science*. Vol.3(2) pp.051-055.
- 474 16. AOAC (2010): Official methods of Analysis, Association of Official Analytical  
475 Chemist Washington, D. C.
- 476 17. AOAC (2005): Official methods of Analysis, Association of Official Analytical  
477 Chemist Washington, D. C.
- 478 18. Ihekoronye and Ngoddy (1985): Integrated Food Science and Technology for the  
479 Tropics.
- 480 19. Iwe, M.O (2002). Hand book of sensory Evaluation methods and analysis Rojoint  
481 communication services Ltd. Enugu. Pp 72-75.
- 482 20. Abiose SH and Ikujenlola AV (2014): Comparison of chemical composition,  
483 functional properties and amino acids composition of quality protein maize and  
484 common maize (*Zea may* L.)
- 485 21. Ade-Omomowaye BIO, Akinwande BA, Bolarinwa IF and Adebisi AO (2008).  
486 Evaluation of tigernut (*cyperuseculentus*) -wheat composite flour and bread. *African*  
487 *Journal of Food Science* Vol.(2) pp.087-091.
- 488 22. Yemi B. (2010): Malnutrition; the case of Death In Nutrition, Pg. 785.
- 489 23. Ibrahim, F.S., E.E. Babiker, N.E. Yousif and A.H. El-Tinay, (2005). Effect of whey  
490 protein supplementation and/or fermentation on biochemical and sensory  
491 characteristics of sorghum flour. *J. Food Technol.*, 3: 118-125
- 492 24. Estoshola E and Oraedu AC (2012). Fatty acid composition of Tigernut tubers,  
493 Baobab seeds (*Adansoniadigitata* L.) and their mixtures. *American Journal of*  
494 *Chemical Sciences*. Vol.73: Pg. 255-267.
- 495 25. Zahras A and Ahmed MS (2014). Explaining the suitability of incorporating tigernut  
496 flour as novel ingredient in gluten-free biscuit. *Journal of Food and Nutritional*  
497 *Sciences*. Vol. 64, No1, pp 27-36.
- 498 26. Nwainikpe R (2010). The phytochemical proximate and amino acid composition of  
499 the extracts of two varieties of tigernut (*Cyperuseculentus*) and their effects on  
500 sickle cell hemoglobin polymerization. *J.med. Med.sci*, 2010.1, 543-549.
- 501 27. Adebowale, A.A., Adegoke, M.T., Sanni, S.A., Adegunwa M.O., and Fetuga,G.O.  
502 (2012). Functional Properties and Biscuit Making Potentials of Sorghum-wheat Flour  
503 Composite. *American Journal of Food Technology*, 7: 372-379.
- 504 28. Aremu M. O, Bamidele TO, Agere H, Ibrahim H and Aremu SO (2015): Proximate  
505 Composition and Amino Acid Profile of Raw and Cooked Black Variety of Tigernut  
506 (*Cyperuseculentus* L.) Grown in Northern Nigeria. *Journal of Biology Agriculture*  
507 *and Health care*. Vol.5. No.7.
- 508 29. Sánchez ZE, Fernández LJ and Angel Pérez AJ (2012). Tigernut (*Cyperus*  
509 *esculentus*) Commercialization: Health Aspects, Composition, Properties and Food  
510 Applications. *Comprehensive Reviews in Food Science and Food Safety*. 11(4),  
511 pp.366–377.
- 512 30. Gambo A and Da' (2014). Tigernut (*cyperus esculentus*): composition, products,  
513 uses and health benefits – a review. *Bayero journal of pure and applied sciences*.

- 514 Vol. 7(71), Pg.56– 61.  
515 31. Wu GY and Meininger CJ (2000). Arginine nutrition and cardiovascular function.  
516 *Journal of Nutrition*. Vol. 130(11), Pg. .2626–2629.  
517 32. Chung IM, Kim EH, Yeo MA, Kim SJ, Seo M and Moon HI (2011). Antidiabetic  
518 effects of three Korean sorghum phenolic extracts in normal and streptozotocin-  
519 induced diabetic rats. *Food Research International*. Vol. 44(1), pp.127–132.  
520 33. Park, J. H., Lee, S. H., Chung, I. M., & Park, Y. (2012). Sorghum extract exerts an  
521 anti-diabetic effect by improving insulin sensitivity via PPAR-?? in mice fed a high-fat  
522 diet. *Nutrition Research and Practice*, 6(4), 322–327.  
523 <https://doi.org/10.4162/nrp.2012.6.4.322>

524  
525  
526

UNDER PEER REVIEW