

# Comparative evaluation of antioxidant potential in thermally processed, underutilized food grains of the Himalayan region

## ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

**Aims:** To determine the bioactive components and total antioxidant capacity (TAC) in selected underutilized crops of the Himalayan region viz. Barnyard millet, Grain amaranth, Rice bean, Black soybean and Horsegram.

**Study design:** Experimental design (Lab experiment).

**Place and Duration of Study:** Department of Foods and Nutrition, College of Home Science, G. B. Pant University of Agriculture and Technology, Pantnagar, Uttarakhand, in the year 2016-18.

**Methodology:** We applied different processing techniques (covered pan cooking and pressure cooking) in the underutilized crops and analyzed the total phenol, total flavonoids and total antioxidant capacity (Ferric reducing antioxidant power (FRAP) assay and (2, 2-Diphenyl-1-picryl hydrazyl) (DPPH) by using standard methods.

**Results:** It was found that black soybean had highest phenolic content after both thermal treatments (3233.76 mg GAE/100g for pan cooked and 1883.11 mg GAE/100g for pressure cooked samples) and TAC by both FRAP (6423.76 mg TE/100g for pan-cooked and 4415.58 mg TE/100g for pressure cooked) and DPPH (536.41 and 453.98 mg TE/100g for pan and pressure cooked samples, respectively) method. Among raw samples, rice bean contained the highest flavonoid content and TAC by FRAP assay. In contrast raw grain Amaranth showed the lowest phenolic content. Further, pressure cooking was found to be better for barnyard millet, while in pulses, pan cooking yielded the best results (in terms of increased value/lower losses).

**Conclusion:** It can be concluded that among pulses pan cooked black soybean was found to have a good store of bioactive compounds as compared to rice bean and horsegram. The pressure cooking method was found to be suitable for millet like barnyard.

*Keywords: Underutilized, thermal processing, rice bean, phenolics, flavonoids, total antioxidant capacity*

## 1. INTRODUCTION (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

Underutilized cereals, millets and pulses are an important group of crops which have special significance in subsistence farming and nutritional security of resource poor masses in developing countries. The underutilized food grains have a vast scope for not only supporting the commercially grown crops by reducing pressure on their availability, but they are a cheap source of nutrients and can be raised at low management costs [1]. Underutilized crop species which are rich in micro-nutrients can contribute effectively in making diets more balanced; hence they play an important role in combating silent hunger. Biodiversity International and International Centre for Underutilized crops listed 200 such underutilized crop species for different eco-geographical region of the world. Minor millets like barnyard millet, pseudo cereals like grain amaranth, and pulses like rice bean, black soy bean, horse gram are important under-utilized crops of the Himalayan region.

28 Minor millets are a group of grassy plants with short slender stalks and small grains  
29 possessing remarkable ability to survive under severe drought conditions. The nutritional  
30 significance of minor millets lies in their richness in micro nutrients like calcium, iron,  
31 phosphorus, vitamins and sulphur containing amino acids. Barnyard millet is the fastest  
32 growing of all millets, and produces a crop within 80 days of sowing. In India, Uttarakhand is  
33 the major producer of barnyard millet. The crop occupies 74, 000 hectares area with a  
34 production and productivity of 87, 000 tonnes, and 857 kg/ha, respectively [2].It is an  
35 important source of vital minerals like niacin, magnesium, phosphorus, manganese, iron and  
36 potassium. It contains high amounts of protein, fiber, essential amino acids like methionine,  
37 lecithin, and vitamin E [3].

38 Besides millets, among pseudo cereals amaranth is a highly nutritious crop with higher  
39 protein content than other cereal species. Amaranth seeds are not true cereals. They are  
40 dicotyledonous in contrast to cereals (e.g. wheat, rice, barley) which are monocotyledonous.  
41 They are referred to as pseudo cereals, as their seeds resemble in function and composition  
42 to those of the true cereals. Amaranth seeds are small (1-1.5 mm diameter), they are  
43 lenticular in shape and weight per seed ranges between 0.6-1.3 mg [4]. Many health  
44 benefits are attributed to amaranth seeds, such as decreasing plasma cholesterol  
45 levels, stimulating the immune system, antitumorigenic activity, reducing blood glucose levels  
46 and improving the condition of hypertension and anaemia patients [5]. Concerning the  
47 antioxidant activity, researchers reported that amaranth, as well as other pseudocereals,  
48 have an antioxidant capacity comparable to that of soybean and rice. Also these authors  
49 reported that the main antioxidant compounds of pseudocereals are polyphenols [6]. Some  
50 researchers studied distinct amaranth cultivars and identified polyphenols (isoquercetin and  
51 rutin) and phenolic acids (syringic and vanillic acids) with antioxidant capacity [7].  
52 Among underutilized pulses and legumes, black soya bean, rice bean and horsegram  
53 are nutritionally superior to traditionally consumed pulses. Black soybean has long  
54 been consumed in the Far East and Southeast Asia as an important source of natural  
55 antioxidants which is attributed to the anthocyanin content in its seed coat [8]. Despite the  
56 possible health benefits of colored soybean, there is limited information related to its  
57 chemical constituents of pharmacological and nutritional importance. Another  
58 underutilized legume ricebean (*Vigna umbellata* L.), also known as climbing mountain bean,  
59 mambi bean and oriental bean, is native to Southeast Asia [9,10]. Rice bean has exhibited  
60 excellent antioxidant capacity and anti-diabetic potential of the sixteen species of beans [11].  
61 Rice bean is also known as one of the promising pulses of mountain regions due to  
62 its high yield potential.

63 Horsegram that belongs to the family *Fabaceae* is a grain legume having excellent  
64 nutritional and remedial properties with excellent climate resilience making it  
65 adaptable to harsh environmental conditions [12]. It is one of the most important under  
66 exploited food legumes being grown almost all over the world. Temperate and sub-tropical  
67 regions, encompassing countries of East and Northeast Africa, Asian countries particularly,  
68 India, China, Philippines, Bhutan, Pakistan, Sri Lanka and Queensland in Australia are  
69 known to grow horse gram [13,14].

70 Polyphenols are the primary antioxidants that donate hydrogen atoms to the free radicals  
71 [15]. They delay or prevent oxidation of lipids, proteins and DNA by reactive oxygen species  
72 that is produced in cells during oxidation [16]. The measurement of total antioxidant capacity  
73 (TAC) is a useful tool in evaluating of the anti-oxidative role of the investigated compounds.  
74 The total antioxidant power as an 'integrated parameter of antioxidants present in a complex  
75 sample' is often more meaningful to evaluate health beneficial effects because of the  
76 cooperative action of antioxidants [17]. The health benefits of whole grain products are in part  
77 attributed to their unique antioxidant activities [18]. It is common knowledge that most grains  
78 (cereals and pulses) have to be processed/ cooked in some way before being eaten.  
79 Cooking increases digestibility and palatability of grains. Normal hydrothermal process such  
80 as boiling, pressure cooking, or canning are utilized to make these foods edible. Dry heat

81 processes such as puffing, roasting and oil frying can also be used. Processing leads to  
82 inactivation of anti-nutritional factors as well as changing the level of bioactive compounds.  
83 In the study area pressure cooking and covered pan cooking are commonly used processing  
84 methods for pulses and cereals. Thus these two processing techniques were adopted to  
85 assess their impact on the bioactive potential of the samples.

86  
87 Information regarding antioxidant activity of underutilized crops will help in understanding  
88 their role in combating chronic degenerative diseases. Therefore, the present study was  
89 planned to estimate and compare the total phenol, total flavonoids and total antioxidant  
90 capacity by FRAP and DPPH in five underutilized crops of the state of Uttarakhand,  
91 India. Among grains dehusked barnyard millet and amaranth seeds were taken, while  
92 among pulses, black soybean, rice bean and horse gram were studied. All evaluations were  
93 done for the sample in their raw state, as well as after thermally processing them in covered  
94 pan, pressure cooker


## 95 96 **2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY** 97 **(ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)**

### 98 99 100 **2.1 Sample collection**

101 Samples of dehusked barnyard millet, grain amaranth, black soybean, rice bean and  
102 horsegram were procured from villages of Tehri and Nainital districts of Uttarakhand, India.

### 103 104 **2.2 Sample preparation**

105 The samples were prepared by using the method shown in Fig1.  
106



107  
108 **Fig 1: Methodology for sample preparation**

109  
110 **2.3 Thermal Processing of samples**

111 Five gram of sample was taken for Thermal Processing of each underutilized grain. Amount  
112 of water used in covered pan cooking was 15 ml. For pressure cooking 20 ml water was  
113 used. Water after processing was not drained. Glass test was used to check the completion  
114 of cooking of the samples.

115  
116 **2.4 Determination of total phenolic content**

117 Total phenolic content was determined using the standard method [19]. Known aliquot of  
118 sample was taken and volume was made up to 1.5 ml with distilled water. To this 0.5ml of  
119 Folin ciocalteau reagent was added. After that 10 ml of 7.5% Na<sub>2</sub>CO<sub>3</sub> was added and  
120 incubation was done at 37° C for 60 minutes. Absorbance of the developed blue color  
121 complex was measured at 750nm wavelength. For blank, 1.5 ml of distilled water was  
122 treated in the same way as sample.

123 Total phenolic content was expressed as mg gallic acid equivalents (GAE)/100g. Gallic acid  
124 (5-20 µg) was used as standard.

125  
126 **2.5 Determination of total flavonoids**

127 Total flavonoids content was estimated by using the standard method [20]. A known aliquot  
 128 of sample was taken and volume was made up to 5 ml with distilled water. Then 0.3 ml of 5%  
 129 NaNO<sub>2</sub> was added. After 5 minutes 0.6 ml of 10% of AlCl<sub>3</sub> was added. After another 6  
 130 minutes 2.0 ml of 1N NaOH was added and contents were mixed. Then 2.1 ml of distilled  
 131 water was added to make volume up to 10 ml. The absorbance of developed pink color was  
 132 measured at 510 nm wavelength against distilled water as blank and expressed as mg rutin  
 133 equivalents (RE)/100g. Rutin (50-200 µg) was used as standard.

134

### 135 **2.6 Determination of total antioxidant capacity (TAC) by ferric reducing** 136 **antioxidant power (FRAP) assay**

137 Two methods were used to determine the total antioxidant capacity of samples. Total  
 138 antioxidant capacity by FRAP assay was determined by using standard method with some  
 139 modifications [21, 22]. Different aliquots of sample were taken and the volume was made up  
 140 to 0.3 ml with distilled water. Then 1.8 ml of FRAP working reagent was added and the  
 141 contents were incubated at 37° C for 10 minutes. Absorbance of developed blue coloured  
 142 complex was measured at 593nm wavelength against 0.3 ml of distilled water as blank and  
 143 expressed as mg trolox equivalents (TE)/100g. Trolox (5-20µg) was used as standard.

144

### 145 **2.7 Determination of total antioxidant capacity by DPPH (2, 2-Diphenyl-1-picryl** 146 **hydrazyl)**

147 Total antioxidant capacity using DPPH radical was measured by the standard method with  
 148 some modification [23,24]. Different sample aliquots were taken and the volume was made  
 149 up to 1 ml with methanol. Then 3 ml of DPPH reagent was added to the sample and  
 150 contents were mixed properly. Incubation was done for 20 minutes at 37°C. Absorbance of  
 151 the developed colour of oxidized solution was measured at 517 nm wavelength against  
 152 methanol as blank and expressed as mg trolox equivalents (TE)/100g. Trolox (10-40 µg)  
 153 was used as standard.

154

### 155 **2.8 Statistical analysis**

156 Data from this study were reported as mean ± SD from three replicates for each test. One  
 157 way ANOVA test was applied using IBM SPSS statistics version 20 programme to observe  
 158 any difference among the treatments with respect to bioactive potential.

159

## 160 **3. RESULTS AND DISCUSSION**

161

### 162 **3.1 Thermal processing/cooking**

163 Barnyard millet took 15 minutes and 5 minutes to be completely cooked in covered pan and  
 164 pressure cooker (Table 1). Grain Amaranth was not processed by these methods as it is not  
 165 boiled or pressure cooked for usual consumption. The pulses, rice bean, black soybean and  
 166 horse gram were fully cooked by 26 minutes, 23 minutes, and 25 minutes respectively, when  
 167 cooked in covered pan. Cooking time for the three pulses was 8 minutes, 6 minutes and 7  
 168 minutes respectively when cooked in a pressure cooker (Table 1). Thus among the three  
 169 pulses, black soybean took less time to cook in comparison to the other 2 pulses in both  
 170 covered pan and pressure cooking methods. It is thought that high pressure rice cooker is  
 171 more energy-intensive than a normal rice cooker during the cooking process [25].

172

**Table 1: Processing of selected samples**

Sr. No.	Samples	Processing time (in minutes)		Amount of water used for processing (ml)	
		Covered pan	Pressure	Covered pan	Pressure

		cooking	cooking	cooking	cooking
	<b>Millet</b>				
1	Barnyard millet (dehusked) ( <i>Echinochloa frumentacea</i> )	15	5	15	20
	<b>Pulses</b>				
1	Rice bean ( <i>Vigna umbellata</i> )	26	8	15	20
2	Black soybean ( <i>Glycine max</i> )	23	6	15	20
3	Horsegram ( <i>Macrotyloma uniflorum</i> )	25	7	15	20

173

174

### 3.2 Bioactive compounds and total antioxidant capacity

175

Each of the five samples, barring grain amaranth, were given two heat treatments and subjected to four assays to establish their bioactive potential. Results were statistically analysed to see the impact of different cooking methods on the bioactive compounds, as well as establish significant differences between samples. Comparison was done among the treatments and among the selected samples. Following were the findings of the study:

176

Total phenolic content of barnyard millet ranged from 21.32 (pressure cooked) to 33.08 (raw)

177

mg GAE/ 100g (Table 2). Similar results for total phenolic content (22.905 to 36.90 mg

178

GAE/100g) of raw dehusked barnyard were reported [26]. However, pan cooking led to

179

lesser losses of total phenolic content (29.41 mg GAE/100g). Total Flavonoid content was

180

found to be minimum (31.42 mg RE/100g) in covered pan cooked barnyard millet and

181

maximum (157.14mgRE/100g) in raw barnyard millet, while in pressure cooked barnyard

182

millet, total flavonoid content was found to be 60.75mgRE/100g (Table 2). The results

183

indicate that loss of flavonoids was less in pressure cooked barnyard millet samples as

184

compared of pan cooked sample. This has also been statistically established. Regarding

185

results of total antioxidant capacity by FRAP; raw and pressure cooked barnyard millet

186

showed a similar value of 60.75, while covered pan cooking increased the total antioxidant

187

capacity (TAC) to 80.51 mg TE/100 g. Total Antioxidant capacity (TAC) by DPPH

188

method of raw barnyard millet exhibited the highest significant ( $p \leq 0.05$ ) value (59.81

189

mg TE/100 g) as compared to the thermally processed samples. In covered pan cooking

190

and pressure cooking treatment TAC was reduced to 46.506 and 53.517 mg

191

TE/100 g, respectively. However pressure cooking treatment was found better than pan

192

cooking as the latter method caused higher loss of TAC (by DPPH).

193

194

195

196

197 In raw amaranth seeds, total phenolic and total flavonoid contents were found to be 4.41 mg  
 198 GAE/100g and 60 mg RE/100g, respectively. While total antioxidant capacity by FRAP and  
 199 DPPH was found to be 59.13 and 67.83mgTE/100gm respectively (Table 2). Some  
 200 researchers reported lower values than that of the present study of total phenol in  
 201 *Amaranthus cruentus* v. *Aztek* i.e. 2.95 mg GAE/g, further they also reported the antioxidant  
 202 activity by FRAP in amaranth seeds as 3.37 mmol Fe<sup>2+</sup> per kg dry weight[26].  
 203 Regarding raw and thermally processed pulses, a significantly high (p≤0.05) phenolic  
 204 content of 1029 mg GAE/100g was found in raw rice bean (Table 2). These values were  
 205 higher than that of 16 cultivars of raw rice bean (1.63-1.82 percent) [27]. In raw Horse gram  
 206 the phenolic content was 156.06 mg GAE/100g which was significantly higher (p≤0.05) than  
 207 both processed samples of the pulse. In contrast covered pan cooked soybean exhibited  
 208 the highest significant (p≤0.05) phenolic content (3233.76 mg GAE/100g.) as compared to  
 209 the raw and pressure cooked samples.  
 210 Total Flavonoid content of raw rice bean showed statistically significant and highest  
 211 value of 857.14mgRE/100g as compared to the two processed samples (Table 2).  
 212 Similarly pan cooked black soybean and horse gram showed significant (p≤0.05) highest  
 213 value of 512.98 and 356.09 mg RE/100grespectively, when compared to their raw and  
 214 pressure cooked forms.  
 215 Total Anti-oxidant capacity (TAC) measured by FRAP method had significant (p≤0.05)  
 216 maximum value in raw rice bean (4818.18mgTE/100gm), pan cooked soybean  
 217 (6423.76) and pan cooked horse gram (2542.60), similarly TAC analysed by DPPH  
 218 method exhibited significant (p≤0.05) highest value in raw rice bean(204.626), pan  
 219 cooked black soybean (536.41) and pan cooked horse gram(145.5 mg TE/100gm) (Table 2).  
 220 A study showed similar values for total phenolic content and total flavonoid content in raw  
 221 black soya bean as 613.00 mg GAE/100g and 219 mg RE/100g respectively [28].  
 222

223 **Table2: Bioactive components and total antioxidant capacity of samples**  
 224

Sr.No.	Sample	Raw/processed	Total phenolic content(mg GAE/100g)	Total Flavonoids (mg RE/100g)	TAC by FRAP assay (mg TE/100g)	TAC by (DPPH) Mg TE/100g
<b>Millet and Pseudocereal</b>						
1.	Barnyard millet (dehusked) ( <i>Echinochloa frumentacea</i> )	Raw	33.08±0.001	157.14±0.004	60.75±0.004	59.891±0.002
		Pan cooked	29.41±0.003	31.42±0.001	80.51±0.002	46.506±0.003
		Pressure cooked	21.32±0.002	54.29±0.002	60.75±0.003	53.517±0.002
		F value	42.609	544.517	170.855	1182.932
		S/NS	S	S	S	S

2.	Amaranth seeds ( <i>Amaranthus cadatus</i> linn)	Raw	4.41±0.004	60±0.004	59.13±0.008	67.83±0.003
<b>Pulses and legumes</b>						
1	Rice bean( <i>Vigna umbellata</i> )	Raw	1029.41±0.005	857.14±0.000	4818.18±0.003	204.626±0.004
		Pan cooked	544.117±0.003	642.86±0.003	3428.57±0.003	186.678±0.002
		Pressure cooked	352.94±0.001	514.28±0.001	3798.70±0.002	178.092±0.004
		F value	168.557	15.967	102.415	45.038
		S/NS	S	S	S	S
2.	Black soybean( <i>Glycine max</i> )	Raw	200±0.010	422.07±0.003	2204.30±0.001	206.45±0.001
		Pan cooked	3233.76±0.003	512.98±0.004	6423.76±0.001	536.41±0.0013
		Pressure cooked	1883.11±0.001	428.57±0.003	4415.58±0.002	453.98±0.005
		F value	6179.570	5.441	390.901	34813.938
		S/NS	S	NS	S	S
3.	Horse gram( <i>Macrotyloma uniflorum</i> )	Raw	156.06±0.030	238.96±0.028	340.36±0.002	33.74±0.002
		Pan cooked	146.96±0.012	356.09±0.002	2542.60±0.004	145.55±0.004
		Pressure cooked	107.35±0.007	100.83±0.005	2097.40±0.005	114.81±0.005
		F value	14.616	7.339	5731.097	1232.177
		S/NS	S	NS	S	S

225 S- Significant, NS- Non significant

226

227

228

229

230

### 3.3 Proximate composition

Proximate composition of the five underutilized grains as reported in literature (Table 3). reveals moisture content in the range of 7.10 to 11.39g%, ash content ranged from 3.05% to 4.27%, crude protein ranged from 6.93 to 35.35g %, crude fat ranged from 0.62 to 18.1g%,

231 carbohydrate ranged from 24.45 to 71.87g% and energy was ranged from 302.63 to 356.22  
 232 Kcal in the selected samples (Table 3).  
 233  
 234  
 235  
 236  
 237  
 238  
 239  
 240  
 241  
 242  
 243

**Table 3: Proximate composition of selected samples**

Sr. No.	Samples	Moisture %	Ash%	Crude protein %	Crude fat %	Crude fibre %	Carbohydrate(By difference)	Energy (Kcal)
	<b>Millet and Pseudo cereal</b>							
1	Barnyard millet (dehusked) ( <i>Echinochloafrumentacea</i> )*	11.39 ±0.05	4.27± 0.02	6.93± 0.13	2.02± 0.06	2.98± 0.12	71.87±0.10	333±0.35
2.	Amaranth seeds ( <i>Amaranthescadatuslinn</i> )**	9.20± 0.40	3.05± 0.30	13.27 ±0.34	5.56± 0.33	-	61.46±0.6	356.22 ±0.10
	<b>Pulses</b>							
1	Rice bean ( <i>Vigna umbellata</i> )**	11.12	3.54	19.97	.74	-	51.26	302.63
2	Black soybean ( <i>Glycine max</i> )***	7.10	4.25	35.35	18.15	10.70	24.45	-
3	Horsegram ( <i>Macrotyloma uniflorum</i> )**	9.28	3.24	21.73	0.62	-	57.24	329.90

\*-Verma et al (2015), \*\*- Longvah et al(2017), \*\*\*-Ciabotti et al (2016)

#### 4. CONCLUSION

244  
 245  
 246  
 247  
 248  
 249 It was concluded from the study that bioactive components were reduced due to processing  
 250 in barnyard millet. However pressure cooking was found better than covered pan cooking as  
 251 it led to greater losses. This may be due to the prolonged cooking time in the latter method.  
 252 Thermal processing (both covered pan cooked and pressure cooked) of the samples  
 253 showed black soybean had the highest phenolic content and total antioxidant capacity by

254 both FRAP and DPPH methods. Thermally processed (both covered pan and pressure  
255 cooked) rice bean had the highest flavonoid. In contrast raw grain Amaranth showed the  
256 lowest phenolic content and lowest total antioxidant capacity by FRAP assay. Thus it may be  
257 concluded that black soybean is the richest source of bioactive compounds among  
258 all selected samples. This reinforces the health benefits of black soybean followed by rice  
259 bean whereas barnyard millet and amaranth have lower bioactive potential as  
260 compared to pulses. Further, it may also be concluded that in case of pulses covered pan  
261 cooking yielded the better results as compared to pressure cooking in terms of lesser losses  
262 of bioactive compounds and total antioxidant capacity.

UNDER PEER REVIEW

263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
  
275  
276  
277  
  
278  
279  
  
280  
281  
  
282  
283  
284  
  
285  
286  
287  
  
288  
289  
290  
291  
292  
  
293  
294  
295  
  
296  
297  
  
298  
299  
300

## COMPETING INTERESTS

“Authors have declared that no competing interests exist.”

## REFERENCES

1. Sankhala A, Chopra S, Sankhala A K. Effect of processing on tannin, phytate and invitro iron in underutilized millets-Bajra, The Indian J Nutr And Dietetics.2004;41: 55-61.
2. Gupta A, Sood S, Agrawal PK, Bhatt JC. Under-Utilized Food Crops Of Himalayan Region: Utilization And Prospective. Newer Approaches To Biotechnology.2013.
3. Chandrasekhar U. Appropriate technology for utilization of nutrient dense local plant foods. ProclXth Asian Congress Nutrition.2003; 48: 12-16.
4. Bressani R. Amaranth. In: B. Caballero, editors. Encyclopaedia of food sciences and nutrition (pp. 166e173). Oxford: academic press; 2003.
5. Caselato-Souza VM, Amaya-Fafran J. State of knowledge on amaranth grain: a comprehensive review. Journal of food science. 2012; 77(4): r93-r104.
6. Gorinstein S. The total polyphenols and the antioxidant potentials of some selected cereals and pseudocereals. European food research and technology.2007; 225( 3-4): 321-328.
7. Rosa B D L, Fomsgaard I S, Laursen B, Mortensen A G, Olvera-Martínez L, Silva-Sanchez C, Mendoza-Herrera A, González-Castaneda J, Leon-Rodríguez A. Amaranth (*AmaranthusHypocondriacus*) as an alternative crop for sustainable food production: phenolic acids and flavonoids with potential impact on its nutraceutical quality. J Cereal Sci. 2009; 49(1):117-121.
8. Astadi IR, Astuti M, Santoso U, Nugraheni PS. In Vitro Antioxidant Activity OfAnthocyanins Of Black Soybean Seed Coat In Human Low Density Lipoprotein (LDL). Food Chem. 2009; 112: 659–663.
9. Bolivar A, Luis CZ. Impact of germination on phenolic content and antioxidant activity of 13 edible seed species. Food Chem. 2010; 119: 1485–1490.
10. Saikia P, Sarkar CR, Borua I. Chemical Composition, Antinutritional Factors And Effect Of Cooking On Nutritional Quality Of Rice Bean [*Vigna Umbellate* (Thunb; Ohwi And Ohashi)]. Food Chem. 1999; 67:347–352.

- 301 11. Yao Y, Cheng XZ, Wang LX, Wang SH, Ren GX. Biological potential of  
302 sixteen legumes in china. *Int J Mol Sci.* 2011; 12: 7048–7058.
- 303 12. Kumar D. Horsegram Research: An Introduction. In: Kumar D. (Ed.),  
304 Horsegram In India. 2006. Pp1-10.
- 305 13. Durga KK. Variability and Divergence in horsegram (*Dolichos Uniflorus*). *J*  
306 *Arid Land.* 2012; 4(1): 71–76.
- 307 14. Krishna KR. Legume agro ecosystems of south India: nutrient dynamics,  
308 ecology and productivity. Brown Walker Press, Florida, USA. 2010; Pp 372-  
309 382
- 310 15. Singh B, Singh JP, Kaur A, Singh N. Bioactive compounds in banana and  
311 their associated health benefits—a review. *Food Chem.* 2016; 206:1–11.
- 312 16. Amarowicz R, Estrella I, Hernández T, Robredo S, Troszyńska A, Kosińska  
313 A, Pegg RB. Free radical-scavenging capacity, antioxidant activity, and  
314 phenolic composition of green lentil (*Lens Culinaris*). *Food Chem.* 2010;  
315 121(3):705-11.
- 316 17. Ghiselli A, Serafini M, Natella F, Scaccini C. Total antioxidant capacity as a  
317 tool to assess redox status: critical view and experimental data. *Free Radical*  
318 *Bio Med.* 2000; 29(11):1106-14.
- 319 18. Tian L, Liu J, Ma L, Zhang L, Wang S, Yan E, Zhu H. Isolation and  
320 Purification of Antioxidant and ACE-Inhibitory Peptides from Yak  
321 (*Bosgrunniens*) Skin. *J Food Processing Preservation,* 2017; 41(5), e13123  
322
- 323 19. Singleton VI, Orthofer R, Lamuela-Raventos RM. Analysis of total phenol and  
324 other oxidation substrates and antioxidants by means of Folin-Ciocalteu  
325 reagent. *Methods of Enzymology.* 1999; 299:152-158.  
326
- 327 20. Zhishen J, Mengcheng T, Jianming W. The determination of flavonoid  
328 contents in mulberry and their scavenging effects on superoxide radicals.  
329 *Food Chem.* 1999; 64(4):555-559.  
330
- 331 21. Benzie I, Strain JJ. Ferric reducing/ antioxidant power assay: direct measure  
332 of total antioxidant activity of biological fluids and modified version for  
333 simultaneous measurement of total antioxidant power and ascorbic acid  
334 concentration. *Methods in Enzymology.* 1999; 299:15-27.

- 335 22. Tadhani MB, Patel VH, Subhash R. In vitro antioxidant activities of stevia  
336 rebaudiana leaves and callus. J Food Compos Ana.2009;20(3):323-329.
- 337 23. Brand-Williams W, Cuvelier ME, Berset CLWT. Use Of A Free Radical  
338 Method To Evaluate Antioxidant Activity. LWT-Food Sci Tech.1995;  
339 28(1):25-30.
- 340 24. Lee JH, Kim HJ, Lee BW, Lee YY, Lee BK, Ko JY, Woo KS.  
341 Physicochemical and antioxidant properties of rice cooked with different  
342 proportions of black soybeans and cooking methods. J Food bioch. 2018;  
343 42(6):E12671.
- 344 25. Goudar G, Sathisha GJ. Effect of extrusion and flaking on the retention of  
345 nutrients and phenolic compounds in millet grains. Int J Food SciNutr.2016;  
346 1(4):08-11.
- 347 26. Pasko PA, Barton H , Zagrodzki P, Gorinstein S , Fołta M , Zachwieja Z.  
348 Anthocyanins, total polyphenols and antioxidant activity in amaranth and  
349 quinoa seeds and sprouts during their growth. Food Chem.2009;115 :994–  
350 998.
- 351 27. Katoch R.Nutritional potential of rice bean (*VignaUmbellata*): an  
352 underutilized legume. J Food Sci Tech. 2013; 78(1):C8-16.
- 353 28. Malencic D,Cvejic J, Miladinovic J. Polyphenol content and antioxidant  
354 properties of colored soybean seeds from central Europe J Medicinal  
355 Food.2012; 15(1):89-95.