

1 *In vitro* study of anti-Salmonella activities of *Boerhaavia diffusa* Leaf extract

2

3

4 **Abstract**

5 Various strategies have been employed in the treatment and management of *Salmonella*
6 infection however, *Salmonella* strains have gained resistance to antibiotics. This study was to
7 determine *in vitro* anti-Salmonella activity of *Boerhaavia diffusa* leaf extract against clinical
8 isolate of *Salmonella typhi* and *Salmonella typhi* ATCC 14028. The aqueous and ethanol
9 extracts of *B. diffusa* was studied for their antibacterial activity against pathogenic *Salmonella*
10 *typhi*. The *in vitro* antibacterial activity was performed by agar well diffusion method and broth
11 dilution using spectrophotometric method and the results were expressed as the average
12 diameter of zone of inhibition of bacterial growth around the well and optical density
13 respectively. It was observed that aqueous extract exerted slightly higher activity than ethanolic
14 extract as revealed by mean diameter of zone of inhibitions at concentration of 200 mg/ml,
15 aqueous extract had 35.21 ± 0.47 mm (*Salmonella typhi* ATCC 14028) compared with ethanol
16 extract 26.41 ± 0.32 mm (clinical). However, in broth dilution method, ethanol extract
17 significantly ($p < 0.05$) reduced the cell, at 48 hours, the optical density of clinical isolate of *S.*
18 *typhi* treated at concentration of 200 mg/ml of extract was 0.47 ± 0.02 nm while at the same
19 concentration of extract, aqueous extract had optical density of 0.52 ± 0.11 nm respectively.
20 Phytochemical assay revealed that tannin (5.18 ± 0.02 mg/g) and quinone (8.45 ± 0.13 mg/g) in
21 ethanol extract was significantly ($p < 0.05$) higher than aqueous extract while saponin
22 (14.18 ± 0.06 mg/g) was higher in aqueous extract. The ethanol and aqueous extracts of leaves of
23 *B. diffusa* whole plant exhibited significant antibacterial activity against both clinical and typed
24 *Salmonella typhi*. Therefore, the plant extract could be used for the treatment of Salmonellosis,
25 however, the *in vivo* studies is needed to ascertain the safety of the extract.

26 **Key words:** Anti-Salmonella activity, plant extracts, agar well diffusion, broth dilution,
27 *Salmonella* strains

28 **Introduction**

29 The bacterium *Salmonella typhi* causes typhoid fever (Prescott *et al.* 2005 and Doughari *et al.*,
30 2007). The bacterium is a gram-negative, motile, non-spore, non-capsulated bacillus that can
31 be contracted through contaminated water, milk, food or fruits and vegetables or via

32 convalescent or chronic carriers (Doughari, 2005). It has also been linked with zoonotic
33 transmission via reptiles and common domestic pets (Birgitta *et al.*, 2005). *Salmonella enterica*,
34 which is a group of Gram-negative bacterial pathogens capable of infecting humans and animals,
35 cause significant morbidity and mortality worldwide (Christenson, 2013). Certain serotypes
36 adapted to human, such as *Salmonella typhi* (*S. typhi*) and *Salmonella paratyphi* (*S. paratyphi*),
37 usually cause severe diseases in humans, such as enteric fevers (typhoid and paratyphoid fevers).
38 In most endemic areas like Africa, Asia, and Latin America, approximately 90% of enteric fever
39 is typhoid. This disease is an important global health problem with an estimated 16 million cases
40 and 600 000 deaths each year.

41 Various strategies have been employed in the treatment and management of *Salmonella*
42 infection. Fluoroquinolones and tetracyclines are most commonly used to treat *Salmonella*
43 infections. However, *Salmonella* strains resistant to these antibiotics have been reported in Korea
44 and other countries (Wangari, 2017). One major concern to public health has been the global
45 dissemination of *S. typhimurium* Definitive Type 104, which is resistant to cotrimoxazole,
46 nalidixic acid and ampicilin (Perron *et al.*, 2008; Kariuki *et al.*, 2010). The rise in antibiotic-
47 resistant strains has led to increased interest in use of plant materials to develop new effective
48 drugs (Wangari, 2017). Moreover, conventional antityphoid drugs are becoming more and more
49 unavailable to the common man in Africa due to increased cost (Donald *et al.*, 2015).

50 The rise in antibiotic-resistant strains has led to increased interest in use of plant materials to
51 develop new effective drugs (Wangari, 2017). It has been reported that 80% of the world
52 population are rural dwellers and rely on medicinal plants for their daily medications, also,
53 plants have been reported to have minimal or no side effects compared to antibiotics (Bibitha
54 *et al.*, 2002; Maghrani *et al.*, 2005). *Boerhaavia diffusa* is traditionally known in Nigeria as
55 *Etiponla* in Yoruba, *Azeigwe* in Igbo and *Babba-juju* in Hausa. *B. diffusa* is a perennial
56 creeping weed, prostrate or ascending herb, up to 1 m long or more, having spreading
57 branches (Nayak and Thirunavoukkarasu, 2016).

58 The root, leaves, aerial parts and the whole plant of *B. diffusa* (L. syn) are used worldwide for
59 the treatment of a number of disorders e.g. liver complaint, kidney disorders, rheumatism
60 e.t.c. (Katampe *et al.*, 2018). The quest to identify and isolate novel phytochemicals from *B.*
61 *diffusa* has led many researchers to discover various compounds such as flavonoids, alkaloids,
62 glycosides, steroids, triterpenoids, lipids, lignans, carbohydrates, proteins, and glycoproteins

63 from its leaves, stems, seeds and roots (Girach *et al.*, 2006). Sourav (2012) explored the Anti
64 bacterials from *Boerhaavia diffusa*. In his study, the chloroform and alcohol extracts of the
65 plant were screened against six bacteria viz *Staphylococcus aureus*, *Escherichia coli*, *Proteus*
66 *mirabilis*, *Salmonella typhimurium*, *Pseudomonas aeruginosa* and *Klebsiella aerogenosa*.
67 Chloroform extract showed activity against *E. coli*, *S. typhimurium* and *P. aeruginosa* while
68 the alcohol extract was active against *P. mirabilis* and *S. typhimurium*. The present study was
69 undertaken to further investigate antibacterial activity of *Boerhaavia diffusa* on typed and
70 clinical strains of *Salmonella typhi* with the view to provide scientific evidence for its
71 application as a medicinal plant.

72 **Materials and methods**

73 **Collection of leaves of *Boerhaavia***

74 Fresh leaves of *Boerhaavia diffusa* were collected from the School of Health
75 Technology, Oda Road, Akure, and identified in the Department of Crop, Soil and Pest
76 Management, Federal University of Technology, Akure Ondo State.

77 **Preparation of plant extract**

78 **Aqueous extraction**

79 The aqueous extractions of the water soluble ingredient were carried out using the filter method.
80 A 2g of each of the grounded leaves were extracted by successive soaking for 2days using 50ml
81 of distilled water in a 250ml sterile conical flask. The extracts were concentrated in vacuum at
82 60°C and stored in universal bottles and refrigerated at 4°C prior to use (Ogoti *et al.*, 2015).

83 **Ethanol extraction**

84 The organic solvent leaf extract was prepared by 2g of plant mixture with ethanol and kept for
85 two days. The extract was concentrated to one-fifth volume, filter sterilized and stored at 4°C
86 (Ogoti *et al.*, 2015).

87 **Test organism**

88 The clinical bacterial strains were obtained from the Department of Microbiology, Federal
89 University of Technology Akure. Clinical *Salmonella typhi* and typed (ATCC 14028) *Salmonella*
90 *typhi* were used. The isolates were confirmed based on cultural, morphological and biochemical
91 characteristics following standard methods of identifying *Salmonella typhi* (Cheesbrough, 2004).
92 The bacterial strain was grown in nutrient broth for 12-18 hours at 37°C on rotary shaker. Cells
93 were grown at 37°C for 18 hours and cultures were kept at 4°C.

94 **Antimicrobial susceptibility tests**

95 **Standardization of the inoculum**

96 The inoculum was prepared by inoculating colonies of fresh test cultures into sterile
97 distilled water. The turbidity was compared to 0.5McFarland standard prepared according to
98 method of Cheesbrough, (2004).

99 **Antibacterial susceptibility assay**

100 The extracts were dissolved and diluted using 50 % v/v dimethylsulphoxide (DMSO) to
101 obtain different concentrations (50, 100 and 200 mg) in 1 mL. The 50 mg/ml, 100 mg/ml and
102 200 mg/ml of the extracts of *B. diffusa* leaves were introduced into the wells of Muller Hinton
103 agar plate. The plates were incubated aerobically at 37°C and examined after 24 hours. The
104 plates were examined for microbial growth inhibition and the Inhibition Zone Diameter (IZD)
105 was measured to the nearest millimeter and compared with those produced by the commercial
106 antibiotic ciprofloxacin which was used as control. Effect of extract on anti-*Salmonella* efficacy
107 of the extract in broth was also assayed using spectrophotometric method, the absorbance of the
108 tube was read at 620 nm (Cheesbrough, 2004; Marcelin *et al.*, 2016).

109 **Antibiotics sensitivity test using commercial**

110 Antibiotics sensitivity test of the bacterial isolates were determined by disc diffusion
111 method as described by Cheesbrough (2004). Standard inoculum of 18 hours broth was spread
112 on Muller Hinton agar using sterile swab in triplicate. The antibiotic discs were placed on the
113 plate at equidistance. The plates were then incubated for 24 hours at 37°C and diameter of zone
114 of inhibition were measured and recorded. The commercial antibiotics discs (Fondoz
115 Laboratories Ltd, Nigeria) used were; Chloramphenicol (CH) 30 µg, Sparfloxacin (SP) 25 µg,
116 Ciprofloxacin (CPX) 10µg, Amoxicillin (AM) 25µg, Augmentin (AU) 30µg, Gentamycin (CN)
117 10µg, Pefloxacin (PEF) 5µg, Ofloxacin (OFX) 5µg, Streptomycin (S) 10 µg and Septra (SXT)
118 30µg.

119 **Minimum Inhibitory Concentration (MIC) and Minimum Bacteriocidal Concentration** 120 **(MBC) of *Boerhaavia diffusa* Extracts**

121 The Minimum Inhibitory Concentration (MIC) and Minimum Bacteriocidal Concentration
122 (MBC) of the extracts were determined using the broth (tube) dilution technique (Anibijuwon
123 and Udeze, 2009). Dilutions of the extract in Mueller Hinton broth were prepared in tubes. The
124 concentration of inoculum was also standardized to 0.5 McFarland's turbidity, The Mueller

125 Hinton broth in tubes containing the different concentration of plant extract, 50, 100, and 200
126 mg/ml were then inoculated with 0.5 ml of the standardized culture. The tubes were then
127 incubated at 37°C for 24 hours. MIC and MBC values were recorded.

128 **Screening of phytochemical compounds**

129 The various solvent extracts of the powder of leaves of *Boerhaavia diffusa* were subjected to
130 phytochemical tests for the identification of various action constituents using the method of
131 Marcelin *et al.* (2016). The following major pharmaceutical valuable phytochemical compounds
132 were analyzed qualitatively and quantitatively; alkaloids, phenols, tannins, flavonoids, quinones,
133 saponins, terpenoids, sterols and cardiac glycosides.

134 **Statistical analysis of data**

135 Data obtained were subjected to analysis of variance and means were compared using Duncan's
136 New Multiple Range Test (DNMRT) with the aid of SPSS software at $p \leq 0.05$ level of
137 significance.

138 **Results**

139 The test organisms used for this study were identified based on biochemical characteristics
140 common to *Salmonella typhi*. The result is presented in Table 1. The antibiotic sensitivity
141 patterns of commercial antibiotics on the two strains of *S. typhi* are presented in Figure 1. The
142 result revealed that the zones of inhibition of antibiotics against typed isolates was higher than
143 that of clinical isolates however, chloramphenicol had highest inhibition against the isolates
144 (STC=24.30±0.42 mm, STT=24.36±0.07 mm). The results of antibacterial activity of both
145 water and ethanol crude extracts of *B. diffusa* showed anti*Samonella* activity on the two strains
146 of *S. typhi* tested at different concentrations, with aqueous extract exerting slightly higher
147 activity than ethanolic extract as revealed by mean diameter of zone of inhibitions, 200 mg/ml
148 of aqueous extract had highest (35.21±0.47) zone of inhibition (Figure 2). Minimum inhibitory
149 concentration (MIC) and minimum bactericidal concentrations (MBC) of the extracts is shown
150 in Tables 2. The ethanol and aqueous extract had the same MIC (100 mg/ml) on typed isolate,
151 also, there was no difference in the MIC and MBC of ethanol and aqueous extract on typed
152 isolate.

153 The anti-*Salmonella* efficacy of *Boerhaavia diffusa* extracts in broth was assayed and were
154 shown in Figure 3, 4, 5 and 6. The result presented in Figure 3 revealed the effect of ethanol
155 extract on clinical isolate of *S. typhi*, it was noted that the extract significantly ($p < 0.05$) reduced


156 the cell, at 48 hours, the optical density of clinical isolate of *S. typhi* treated with 50, 100, 200
 157 mg/ml of extract were 0.52±0.03, 0.50±0.10, 0.47±0.02 nm respectively while at the same
 158 concentration of extract, aqueous extract had optical density of 0.64±0.21, 0.54±0.03, 0.52±0.11
 159 nm respectively (Figure 4). Also, the anti-*Salmonella* efficacy of *B. diffusa* ethanol extracts on
 160 typed isolate of *S. typhi* is shown in Figure 5. It was observed that the extract significantly
 161 (p<0.05) reduced the cell, at 48 hours, the optical density were of typed isolate of *S. typhi*
 162 treated with 50, 100, 200 mg/ml of extract were 0.49±0.00, 0.48±0.01 and 0.37±0.12 while at
 163 the same concentration, aqueous extract had optical density of 0.62±0.03, 0.53±0.11 and
 164 0.49±0.21 nm respectively (Figure 6).

165 Both plant extracts (ethanolic and aqueous) were subjected to preliminary qualitative
 166 phytochemical evaluation. The phytochemical profiles of the two solvent extracts from plant
 167 used in this study are presented in Table 2. The analysis revealed the presence of alkaloids,
 168 phenol, glycosides, steroids, carboxylic acid, reducing sugar, flavonoids, saponins, tannins,
 169 proteins, triterpenoids, quinines, carbohydrates and sterols. Also, tannin (5.18±0.02 mg/g) and
 170 quinone (8.45±0.13 mg/g) in ethanol extract was significantly (p<0.05) higher than aqueous
 171 extract while saponin (14.18±0.06 mg/g) was higher in aqueous extract.

172 **Table 1: Biochemical characteristics of *Salmonella* strains**

Biochemical characteristics	<i>Salmonella typhi</i> (Clinical isolate)	<i>Salmonella typhi</i> (ATCC 14028)
Gram reaction	-ve	-ve
Shape	Rod	Rod
Motility	+ve	+ve
Catalase	+ve	+ve
Coagulase	-ve	-ve
Citrate	+ve	+ve
H ₂ S	+ve	+ve
Lactose	-ve	-ve
Glucose	+ve	+ve
Fructose	+ve	+ve
Sucrose	-ve	-ve
Galactose	+ve	+ve
Indole	-ve	-ve
Methyl red	+ve	+ve
Voges-Proskauer	-ve	-ve
Oxidase	-ve	-ve

173 **Key:** -ve = negative +ve = positive




175

176 **Figure 1: Antibiotic sensitivity pattern of commercial antibiotic discs on *S. typhi* strains**

177 Key: STC – *Salmonella typhi* (clinical isolate), STT – *Salmonella typhi* (typed isolate),
 178 Chloramphenicol (CH) 30 µg, Sparfloxacin (SP) 25 µg, Ciprofloxacin (CPX) 10µg, Amoxicillin
 179 (AM) 25µg, Augmentin (AU) 30µg, Gentamycin (CN) 10µg, Pefloxacin (PEF) 5µg, Ofloxacin
 180 (OFX) 5µg, Streptomycin (S) 10 µg and Septra (SXT) 30µg.

181

182



183

184 **Figure 2: Anti-Salmonella activity of Boerhaavia diffusa extracts**

185 **Key:** AQ = Aqueous extracts of *Boerhaavia diffusa*, ET= Ethanolic extracts of *Boerhaavia*
 186 *diffusa*, Cpx= Ciprofloxacin

187


188

189 **Table 2: Minimum Inhibitory Concentration (MIC) and Minimum Bacteriocidal**
 190 **Concentration (MBC) of Boerhaavia diffusa extracts on Salmonella**

<i>Boerhaavia diffusa</i> extracts	Ethanolic extract		Aqueous extract	
	S1	S2	S1	S2
MIC (mg/ml)	100	100	50	100
MBC (mg/ml)	50	100	50	100


191 Key: S1 - *S. typhi* (Clinical isolate), S2 – *S. typhi* (ATCC 14028)

192




193

194 **Figure 3: Effect of *Boerhaavia diffusa* ethanol extract on Clinical isolate**




195

196 **Figure 4: Effect of *Boerhaavia diffusa* aqueous extract on clinical isolate**



197

198 **Figure 5: Effect of *Boerhaavia diffusa* ethanol extract on typed isolate**



199

200 **Figure 6: Effect of *Boerhaavia diffusa* aqueous extract on typed isolate**

201 **Table 3: Qualitative analysis of phytochemicals in *Boerhavia diffusa* leaf extracts**

Phytochemical	Ethanolic extract	Aqueous extract
203 Alkaloids	-	+
204 Tannins	-	+
205 Flavonoids	+	+

206	Quinones	+	-
207	Saponins	+	+
208	Terpenoids	+	-
209	Sterols	+	-
210	Cardiac Glycosides	-	+
211	Phenols	+	+

212 Key : + = Present, - = Absent

213
214 **Table 4: Quantitative phytochemical screening of aqueous and ethanol extracts of *B.***

215 *diffusa*

216	Phytochemicals	Ethanollic extract	Aqueous extract
	Tannins (mg/g)	5.18±0.02 ^a	3.90±0.22 ^a
	Quinones (mg/g)	8.45±0.13 ^b	6.60±0.31 ^a
	Saponins (mg/g)	6.36±0.24 ^a	14.18±0.06 ^b
	Triterpenoids (mg/g)	8.56±0.08 ^a	8.89±0.31 ^a
	Steroid (mg/g)	9.03±0.11 ^a	6.73±0.14 ^a
	Glycosides (mg/g)	30.39±0.06 ^b	28.29± 0.03 ^a
	Flavonoids (mg/g)	9.98±0.61 ^a	11.26±0.33 ^a

217 Discussion

218 *Salmonellosis* and enteric fever are always a public health concern in most developing countries,
219 which are mostly low or middle-income countries with inadequate sanitation and hygiene,
220 particularly, regarding food, water and disposal of human excreta (Marcelin *et al.*, 2016).
221 Different plants and their parts (flowers, buds, leaves, stem, bark, fruits, skin, pulp and root) have
222 been used for thousands of years to enhance the flavour and aroma of food. In addition, plants
223 are rich in a wide variety of second metabolites such as Alkaloids, Flavonoids, Phenols, which
224 were found in vitro to have antimicrobial properties (Abbas *et al.*, 2007; Marcelin *et al.*, 2016).
225 In this study, extracts of *Boerhaavia diffusa* leaves were investigated for antibacterial activity
226 against *Salmonella typhi*. Plant extracts were used to investigate antibacterial activity against two
227 bacterial strains (Clinical *Salmonella typhi* and *Salmonella typhi* ATCC 14028). In this study,
228 antibacterial activity of *B. diffusa* leaf extracts was compared against the test bacteria with
229 activities of model antibiotics. The higher antibacterial activity of model antibiotics is not
230 surprising, since the antibiotics are in a refined state. The standard antibiotics (ampicillin,

231 amoxicillin, ciprofloxacin, ofloxacin, chloramphenicol) used in this study are first line drugs
232 employed in the treatment of typhoid fever (Prescott *et al.*, 2005).

233 The aqueous and ethanol extracts exhibited different zones of inhibition against the isolates,
234 however, aqueous extract had higher zones of inhibition than ethanol extract. This is consistent
235 with other studies (Ujowundu *et al.*, 2008). Antimicrobial action may be due to the synergistic
236 action of different chemical constituents, some of which probably are lost upon extraction with
237 solvent (Shahina, 2007; Ogoti *et al.*, 2015; Marcelin *et al.*, 2016). Water could be a better
238 extraction solvent than ethanol for *B. diffusa* leaf, also, the demonstration of higher activity by
239 the aqueous solvent may be an indication that the phytoconstituents in the plant leaves are more
240 soluble in water than the organic solvent (Marjorie, 1999). The antimicrobial potential of *B.*
241 *diffusa* and other plants sourced from traditional healers through an ethnobotanical survey of
242 anti-infective plants in Egbado South in Ogun State, Nigeria was previously reported by Abo and
243 Ashidi (1999). This study also corroborates the findings of Madani and Jain (2008) who reported
244 higher anti-*Salmonella* activity in aqueous extract of *Terminalia belerica* than chloroform and
245 acetone extracts. It has been reported that different phytoconstituents have different degrees of
246 solubility in different types of solvents depending on their polarity. In a traditional setting, water
247 is the solvent largely used to prepare these concoctions.

248 It was noted from this study that plant extracts tested by microdilution technique and the optical
249 density was measured after 48 hours showed that ethanol extract had higher anti-*Salmonella*
250 activity compared to aqueous extract which was higher in values obtained from agar well
251 diffusion technique. It could be that the bioactive components in ethanol extract did not diffuse
252 into agar in agar well but was able to inhibit microbial cells directly in broth. This was
253 previously reported by Olila *et al.* (2001) and Wangari (2017) that the active components of the
254 extract does not diffuse into Muller Hinton agar, however, they were able to cause inhibition of
255 microbial cells in broth microdilution.

256
257 The preliminary qualitative phytochemical screening carried out showed that the leaf extracts of
258 *B. diffusa* contain vital secondary metabolites such as alkaloids, saponins, tannins and
259 glycosides. The bioactive compounds in medicinal plants have been reported to be the active
260 principles responsible for the pharmacological potentials of medicinal plants (Edeoga *et al.*,
261 2005). The presence of these chemicals in the leaves and root of these plants justifies the local

262 uses of these plants for the treatment of various ill conditions. Phytoconstituents such as
263 saponins, phenolic compounds and glycosides have been reported to inhibit bacterial growth
264 and to be protective to plants against bacterial and fungal infections (Okwute, 1992; Wangari,
265 2017). Ethanol extract of *Boerhavia diffusa* leaves possess some phytochemicals like Alkaloids,
266 Anthraquinone, Glycoside, Flavanoids and Tannins. Saponins are natural glycosides that act as
267 hypoglycemic, antifungal and serum cholesterol lowering agents in animals (Desai *et al.*, 2009).
268 Saponins are essential elements in ensuring hormonal balance and synthesis of sex hormones
269 (Okwu, 2004). Tannins are bitter polyphenolic compounds that hasten the healing of wounds.
270 They also possess anti-diuretic and anti-diarrhea properties (Okwu, 2004). Terpenoids was
271 present in both ethanolic extract of and aqueous extracts of AOU and AFU. Terpenoids have
272 been found to be useful in the prevention and therapy of several diseases, including cancer.
273 Terpenoids are also known to possess antimicrobial, antifungal, anti- parasitic, antiviral, anti-
274 allergic, antispasmodic, antihyperglycemic, anti-inflammatory and immunomodulatory
275 properties (Shah *et al.*, 2008). The presence of these compounds promises its potential
276 application in the treatment of microbial ailment. However, tannins were present in aqueous
277 extract of but not in the ethanolic extract. Saponin and flavanoid is higher in the aqueous extract
278 of the leaf (14.18 and 11.26 mg/g) than the ethanolic extract (6.36 and 9.98 mg/g).

279 **Conclusion**

280 Most of the antibiotics used nowadays have lost their effectiveness due to development of
281 resistant genes in microbes. The antibiotics are sometimes associated with side effects such as
282 hypersensitivity, immune suppression and allergic reaction.

283 More interest is being shown in developing alternative antimicrobial drugs for the treatment of
284 infectious diseases without side effects. The results of our present study demonstrates anti-
285 *Salmonella* activity of aqueous and ethanol extract of *Boerhavia diffusa*, tannin and quinone
286 were higher in ethanol extract while saponin was higher in aqueous extract, using agar well
287 diffusion, the aqueous extract showed higher anti-*Salmonella* efficacy while the broth
288 microdilution examined by spectrophotometer revealed that ethanol extract had higher anti-
289 *Salmonella* efficacy. In the present study, the anti-salmonella activity of *Boerhaavia diffusa*
290 may be attributed to individual or synergistic effect of phytoconstituents present in it. The
291 ethanol and aqueous extracts of leaves of *B. diffusa* whole plant exhibited significant
292 antibacterial activity against both clinical and typed *Salmonella typhi*. Therefore, the plant

293 extract could be used for the treatment of Salmonellosis, however, the in vivo studies is needed
294 to ascertain the safety of the extract.

295 **Recommendation**

296 Based on our findings, it is therefore recommended that both agar well diffusion and broth
297 dilution method should be used to affirm the antimicrobial efficacy of the plant extracts.

298 **References**

- 299 Abbas A. M. Noor M. A. and Sarmina Y. (2007). Antimicrobial screening of different extracts of
300 *Piper longum* Linn. *Res. J Agri and Biol sci.*, **3**(6): 852-857.
- 301 Abo K.A and Ashidi J.S. (1999). Antimicrobial screening of *Bridelia micrantha*, *Alchornea*
302 *cordifolia* and *Boerhavia difussa*. *African Journal of Medicine and Medical Sciences*
303 **28**(34):167-169.
- 304 Anibijuwon I.I and Udeze O.A (2009). Antimicrobial activity of *Carica papaya* (Pawpaw Leaf)
305 on some pathogenic organisms of clinical origin from South-Western Nigeria.
306 *Ethnobotanical Leaflets*, **13**(7): 850-864.
- 307 Bibitha B., Jisha V.K., Salitha C.V., Mohan S., Valsa A.K. (2002). Antibacterial activity of
308 different plant extracts. *Indian Journal of Microbiology*, **42**:361-363.
- 309 Birgitta J., Yvonne A., Carl E (2005). Effect of regulation and education and reptile associated
310 salmonellosis. *Emerging Infectious Diseases*, **11**:398-403.
- 311 Christenson J.C. *Salmonella* infections. *Pediatr Rev* 2013; **34**(9): 375-83.
- 312 Desai S.D., Desai D.G., Kaur H. (2009). Saponins and their Biological Activities. *Pharma*
313 *Times*, **41**(3):88-92
- 314 Donald S. T., Donatien G., Siméon P. C. F., Charles F., Fabrice K. and Merline N. D. (2015). *In*
315 *vivo* anti-salmonella activity of aqueous extract of *Euphorbia prostrata* Aiton
316 (Euphorbiaceae) and its toxicological evaluation. *Asian Pacific Journal of Tropical*
317 *Biomedicine*. **5**(4): 310-318
- 318 Doughari J.H (2005). A comparative study on effects of crude extracts of some local medicinal
319 plants and some selected antibiotics on *Salmonella typhi* pp 1-15. M. Sc. Thesis Federal
320 University of Technology, Yola; Adamawa State, Nigeria. Yola. pp. 1-4.
- 321 Doughari J.H, Pukuma M.S., De N. (2007). Antibacterial effects of *Balanites aegyptiaca* L. Drel.
322 and *Moringa oleifera* Lam. on *Salmonella typhi*. *African Journal of Biotechnology*,
323 **6**:2212-2215.
- 324 Edeoga H.O, Okwu D.E, Mbaebie O. (2005). Nutritional values of some known conventional
325 leafy vegetables of Nigeria. *African Journal of Biotechnology*, **4**:685-688.
- 326 Girach R.D. and Singh V.K. (2006). Notable plants in ethnomedicine of Jajpur District, Orissa,
327 India. In: Recent Progresses in Medicinal Plants. *Biopharmaceuticals*, **14**:495-501.
- 328 Kariuki S, Revathi G, Kiiru J, Mengo DM, Mwituria J, Muyodi J, Munyalo A, Yik YT, Kathryn
329 EH, Robert AK, Gordon D (2010). Typhoid in Kenya is associated with dominant
330 multidrug-resistant *Salmonella enterica* Serovar Typhi haplotype that is also widespread
331 in Southern Asia. *Journal of Clinical Microbiology* **48**(6):2171-2176.
- 332 Katampe I., Uwem M., and Uwem J. (2018). Antioxidant and Antimicrobial Effects of Aqueous
333 and Ethanolic Extracts of Selected Medicinal Plants from Southern Nigeria. *Journal o*
334 *Advances in Medical and Pharmaceutical Sciences*, **17**(2): 1-12.

335 Madani A. and Jain S. K. (2008). Anti-Salmonella activity of Terminalia bellerica: In vitro and in
336 vivo studies. *Indian Journal of Experimental Biology*, **46**: 817-821

337 Maghrani M, Zeggwagh N, Michel J, Eddouks M (2005). Antihypertensive effect of *Lepidium*
338 *sativum* L. in spontaneously hypertensive rats. *J. Ethnopharm.* **100**(102): 193-197.

339 Marcelin U., Bharathidasan R. and Prince L. (2016). Antisalmonella Activity of Selected
340 Medicinal Plants. *World Journal of Pharmacy and Pharmaceutical Sciences*, **5**(4): 2512-
341 2519

342 Marjorie M.C. (1999). Plant products as antimicrobial agents. *Clin. Microbiol. Rev.* **12**(4):564-
343 582.

344 Nayak P. and Thirunavoukkarasu M. (2016). A review of the plant *Boerhaavia diffusa*: its
345 chemistry, pharmacology and therapeutical potential. *The Journal o Phytopharmacology*,
346 **5**(2):83-92

347 Ogoti P., Magiri E., Magoma G., Kariuki D. and Bii C. (2015). In vitro anti-Salmonella activity
348 of extracts from selected Kenyan medicinal plants. *Journal of Medicinal Plants*
349 *Research*, **9**(8): 254-261

350 Okwu D.E. (2004). Phytochemical and vitamin content of indigenous species of southeastern
351 Nigeria. *Journal of Sustainable Agriculture and the Environment*, 6:30-37.

352 Okwute S.K. (1992). Plant derived pesticidal and antimicrobial agents for use in Agriculture. A
353 review of phytochemical and biological studies on some Nigerian plants. *J. Agric. Sci.*
354 *Technol.* **2**(1):62-70.

355 Olila D, Olwa-Odyek A, Opuda-Asibo J (2001). Antibacterial and antifungal activities of
356 extracts of Zanthoxylum chabeum and Wurburgia ugandensis, Uganda medicinal plants.
357 *Afr. Health Sci.* **1**(2):66-72.

358 Perron GG, Bell G, Quessy S (2008). Parallel evolution of multidrug-resistance in *Salmonella*
359 enteric isolated from swine. *FEMS Microbiol. Lett.* **281**(1):17-22.

360 Prescott LM, Harley JP, Klein DA (2005). Microbiology, McGraw-Hill, Paris. pp. 937-960.

361 Shah A. S., Wakade A. S. and Juvekar A. R. (2008). Immunomodulatory activity of methanolic
362 extract of *Murraya koenigii* spreng leaves. *Indian J Exp Biol* **46**: 505-9.

363 Shahina N., Samia A. and Sheikh A. R (2007). In vitro antimicrobial activity of the extracts
364 derived from *Terminalia catappa*. *Res. J Microbial.*, **2**(2): 180-184.

365 Sourav Das (2012). Antimicrobial activity study of ethanolic extract of *Boerhaavia diffusa* whole
366 plant. *International Journal Of Pharmacy & Life Sciences*, **3**(10): 2006-2009

367 Ujowundu C.O., Igwe C.U., Enemor V.H.A., Nwaogu L.A., Okafor O.E.(2008). Nutritive and
368 anti-nutritive properties of *Boerhavia diffusa* and *Commelina nudiflora* leaves. *Pak. J.*
369 *Nutr.* **7**(1):90-92.

370 Wangari D. U. (2017). In vitro study of anti-Salmonella activities of medicinal plants. *Global*
371 *Journal of Medicinal Plants Research.* **3** (1): 301-308