

1  
2

3 **LIMING EFFECTS OF SAWDUST ASH AND LIME ON SUNFLOWER YIELD IN ACIDIC SOIL**  
4 **OF SOUTHEASTERN NIGERIA**

5 **ABSTRACT**

6 A pot experiment was carried out to determine the effect of sawdust ash and lime ( $\text{Ca}(\text{OH})_2$ ) on soil  
7 characteristics and yield of sunflower in acidic soil of southeastern Nigeria. The experiment was laid out in  
8 split-plot design, using sawdust ash (0, 1, 2, 3, 4  $\text{t ha}^{-1}$ ) as the sub plot and lime (0, 0.5, 1.0, 1.5  $\text{t ha}^{-1}$ ) as  
9 the main plot. Results showed that with the exception of organic carbon there was significant effect of  
10 treatments on all soil chemical properties. Lime and sawdust ash (SDA) as single and combined  
11 treatments significantly increased total nitrogen ( $P=0.05$ ), available phosphorus ( $P<0.010$ ), and base  
12 saturation ( $P<0.012$ ). The interaction between SDA and lime significantly ( $P=0.05$ ) increased total  
13 exchangeable bases and effective cation exchange capacity, while soil pH was significantly increased  
14 ( $P=0.05$ ) by single applications. The increases in soil chemical properties led to significant positive  
15 response of the sunflower. With the exception of number of leaves, other plant parameters (Plant height,  
16 stem diameter, head weight, 50 seed weight, head diameter) had significant increases for sawdust ash  
17 alone at  $P=0.05$ . Correlation studies showed positive significant relationship between soil pH and  
18 sunflower yield. The study showed that sunflower performed best at the combination of 3  $\text{t ha}^{-1}$  SDA and  
19 1.5  $\text{t ha}^{-1}$  lime producing a mean head weight of 45.4 g.

20 Keywords: sawdust ash, lime, soil acidity, sunflower

21

22 **1. INTRODUCTION**

23 Sunflower (*Helianthus annuus* L.) is the third most important oil seed crop next to soybean and groundnut  
24 as a source of edible oil in the world. It is cultivated globally [1] due to its adaptability to wide range of soil  
25 and climatic conditions. It is tolerant to high temperature and high humidity and will grow well in any light

26 textured, well drained sandy loam soil [2]. Sunflower is not highly drought tolerant; however its highly  
27 branched tap root system allows it to extract more soil moisture. In Nigeria, sunflower is commonly  
28 cultivated in the savannah regions having maximum annual rainfall of about 1500 mm [3]. Interestingly,  
29 studies have shown encouraging yield results in humid high rainfall area with application of phosphorus  
30 fertilizer [4], adjustments in planting time [5], and planting density [6, 7]. However production could be  
31 greatly hindered by soil acidity in the highly weathered soils of this region.

32 Soil acidity is a limiting factor in sunflower production. A good example is a 10% decrease in sunflower  
33 yield between soil pH of 4.7 – 5.3 [8]. One of the critical effects of soil acidity in highly weathered soils of  
34 humid southeastern Nigeria is the unavailability of Phosphorus [9, 10, 11]. Phosphorus deficiency in  
35 sunflower could affect physiological development such as leaf area and photosynthetic rate per unit of  
36 leaf area [12] consequently affecting yield. In the attempt to improve yield in humid region of Nigeria,  
37 Adebayo et al. [4] found good responses in sunflower yield from the application of phosphorus fertilizer. In  
38 addition, significant responses of sunflower yield to liming have been reported in acid soils. Kovacevic et  
39 al [13] recorded about 49 % increase in sunflower yield by liming. Similar finding was previously reported  
40 by Blamey and Nathanson [14].

41 The use of lime for amelioration of acidic soils functions by increasing the availability of nutrients and  
42 reducing toxicity of Al and Fe ions in such soils. Sawdust ash is a source of lime, but due to the  
43 complementary qualities of mineral nutrients contained in sawdust ash and high calcium carbonate  
44 equivalent of commercial lime, Clapham and Zibilske [15] suggest that sawdust ash be used as a  
45 supplement rather than replacement for lime. Several studies have been conducted on sawdust/wood ash  
46 for liming in the study region, but the effect of sawdust ash in combination with lime on sunflower has not  
47 been previously investigated. Other studies on sunflower in the study area have focused on agronomic  
48 practices such as plant population, intercropping and fertilizer management. The objective of this study  
49 was to determine the effect of liming on sunflower yield in acidic soil of southeastern Nigeria.

50

51

## 52 2. MATERIALS AND METHOD

53 **2.1 Experimental site:** This study was conducted outdoors at Michael Okpara University of Agriculture,  
54 Umudike, Abia State (Lat. 5° 29' N and Long.7° 32' E, 122 m.a.s.l), Southeastern Nigeria. Climate type is  
55 generally humid tropic with rainforest vegetation type. It is characterized by uniform high temperatures  
56 which changes slightly during the year. The average annual temperature and rainfall are 26.9 °C and  
57 2046 mm respectively.

58 **2.2 Pot experiment:** Bulk topsoil samples were collected from the Eastern farm of same institution, air  
59 dried and sieved through 2 mm sieve. The bulk soil was thoroughly mixed and 10kg of soil weighed into  
60 each of the sixty 12-litre plastic buckets. Sunflower (*Helianthus annus* L.) seeds were obtained from  
61 National Institute for Horticulture Research and Training, Ibadan. Sawdust was collected from timber  
62 shed, Umuahia Abia State and burnt to obtain ash. Factor levels comprised four rates (0, 0.5, 1.0, 1.5  
63 tonnes per hectare) of commercial lime [Ca(OH)<sub>2</sub>] and five rates (0, 1, 2, 3, 4 tonnes per hectare) of  
64 sawdust ash. They were arranged in a split- plot design and replicated three times. The different  
65 treatment combinations were applied to the 60 buckets containing soil, mixed thoroughly and watered  
66 adequately. After 1 week of treatment application, the seeds were planted. Two seeds were planted per  
67 bucket and later thinned to one seedling per bucket. Uniform watering and weeding were carried out as  
68 required throughout the growing season.

69 **2.3 Statistical analysis:** Data collected were subjected to Analysis of Variance (GENSTAT) and their  
70 means compared using least significant difference at 5% level of probability, while Pearson correlation  
71 was performed using SPSS 13.0.

72

73

74

75

76

77 **3. RESULTS AND DISCUSSION**

78 **3.1 Properties of soil and sawdust ash used for the study**

79 Textural class of the soil is loamy sand with low clay content. Clay play important roles in water and  
80 nutrient retention capacity of soils. Soils low in clay content could lead to heavy leaching of soil nutrients,  
81 even when fertilizer is applied [16].

82 The pH values ranged from 5.36 - 5.64 and 4.37 - 4.45 in water and CaCl<sub>2</sub> respectively. The low pH also  
83 resulted to high exchangeable acidity. Soil pH values in salts are generally lower than those measured in  
84 water. The indication is that these soils at their natural pH are negatively charged [17]. Soil pH  
85 requirement for sunflower ranges between 6.5 and 7.5 (in water). The low pH values could generally  
86 reduce crop yield [18] and in particular, the performance and yield of sunflower [19].

87 Organic carbon content falls within the range for ultisols found in this region as indicated by Eswaran [20].  
88 The soil organic carbon reflects soils' fertility status [21].

89 Available phosphorus was lower than the critical level of 15 mg kg<sup>-1</sup> for most crops [22, 23]. In acidic soils,  
90 phosphorus is known to be associated with secondary minerals such as Fe and Al (hydrus) oxides  
91 through sorption [24], however low effective cation exchange capacity obtained suggests small presence  
92 of secondary minerals having high specific surface [25]. Deficiency in available phosphorus results to  
93 stunted growth, purplish discoloration of leaves. It also affects flowering, fruit formation and seed  
94 production [2]. Uptake of major nutrients elements by sunflower has also been reported to be facilitated  
95 by phosphorus application in the forest zone [26].

96 Total nitrogen was also low and below the critical level of 0.15 % for optimum crop production [27]. This  
97 may be attributed to heavy leaching caused by high rainfall pattern experienced in this region.

98 Basic cations were low however calcium exceeded the critical level of 2 Cmol kg<sup>-1</sup> for most crops [28].  
99 Base saturation had mean of 79.97 %, while exchangeable acidity was between 1.34 and 1.40 Cmol kg<sup>-1</sup>.

100

101

102

103

104

105 **Table 1: Soil and sawdust ash physico-chemical properties**

Properties	Soil	Sawdust ash	Method
Sand	73.0 %	-	Hydrometer method [29]
Clay	12.8 %	-	
Silt	14.2 %	-	
pH (1:2.5 sample:H <sub>2</sub> O)	5.48	10.7	Glass electrode pH meter [30]
pH (1:2.5 sample:CaCl <sub>2</sub> )	4.40	10.5	
Organic Carbon	1.37 %	-	Wet oxidation [31]
Available Phosphorus	11.00 µgg <sup>-1</sup>	17.33 µgg <sup>-1</sup>	Bray 1 [32]
Total Nitrogen	0.10 %	0.30 %	Micro kjeldahl [33]
Exchangeable Acidity	1.36 Cmolkg <sup>-1</sup>	-	KCl extraction [34]
Potassium	0.09 Cmolkg <sup>-1</sup>	34.78 %	NH <sub>4</sub> <sup>+</sup> -acetate extraction
Calcium	2.80 Cmolkg <sup>-1</sup>	5.53 %	
Magnesium	2.00 Cmolkg <sup>-1</sup>	2.13 %	
Sodium	0.10 Cmolkg <sup>-1</sup>	0.12 %	
Total exchangeable bases	4.99 Cmolkg <sup>-1</sup>	42.56 %	
ECEC	6.24 Cmolkg <sup>-1</sup>	-	
Base Saturation	79.97 %	-	

\*ECEC: Effective cation exchange capacity

Sawdust ash had pH values of 10.7 and 10.5 in water and CaCl<sub>2</sub> respectively (Table 1). It also contained 0.30 % of total nitrogen, 17.33 µgg<sup>-1</sup> available phosphorus, and 5.53, 2.13, 0.12, 34.78 Cmol kg<sup>-1</sup> of Ca, Mg, Na and K respectively. Data indicate alkalinity and higher concentrations of nutrients compared to initial soil sample. Hence, plant ash is a potential improver of acidic soils for better crop yield. Similar results have been obtained by several authors [35, 36, 37, 38, 39].

106

107

### 108 3.2 Effect of sawdust ash and lime on soil chemical properties

109 Effects of sawdust ash and lime on soil chemical properties are shown in Table 2. Although there was no  
110 significant effect of sawdust ash (SDA) and lime on organic carbon as several researchers [37, 36, 35]  
111 have previously reported, there were increases relative to control. The highest mean value of 1.55 %  
112 was obtained at treatment 1  $\text{tha}^{-1}$  SDA  $\times$  1  $\text{tha}^{-1}$  lime.

113 There were significant increases in total N as a result of the application of treatments. Increase in soil pH  
114 encourages increases in microbial activities which are responsible for the breakdown of crop residues,  
115 contributing to the availability of nitrogen, phosphorus and sulphur in soils.

116 Main and interaction effects of SDA and lime had positive significance ( $P=0.05$ ) on available  
117 phosphorus. This may be attributed to the fact that lime [ $\text{Ca}(\text{OH})_2$ ] contains  $\text{Ca}^{2+}$  which increases pH  
118 thereby releasing adsorbed P, as well as the high presence of P and Ca in SDA. Odedina *et al.*[35]  
119 found that SDA and other plant derived ashes increased soil N, P, K, Ca, Mg contents of soils.

120 Basic cations (K, Mg, Na and Ca) significantly increased ( $P=0.05$ ) with all treatments. This finding is in  
121 line with that of several other researchers [40, 38, 37]. This confirms the positive effect of ash and lime  
122 on cationic nutrients. The implication of this observation is that the nutrient contained in the treatments  
123 were mineralized by microbial activities [41] and made available in soil solution for subsequent plant  
124 uptake. The following treatment combinations had the highest levels of K, Mg, Na and Ca respectively: 4  
125  $\text{tha}^{-1}$  SDA  $\times$  1.5  $\text{tha}^{-1}$  lime, 2  $\text{tha}^{-1}$  SDA  $\times$  1  $\text{tha}^{-1}$  lime, 0  $\text{tha}^{-1}$  SDA  $\times$  0.5  $\text{tha}^{-1}$  lime and 1  $\text{tha}^{-1}$  SDA  $\times$  0  
126  $\text{tha}^{-1}$  lime. From these results, it could be suggested that ash be added as a supplement to lime and not  
127 as a replacement. This corroborates the finding of Clapman and Zibilske [14].

128 Significant increases ( $P<0.012$ ) were recorded for total exchangeable bases, base saturation and ECEC.  
129 Increases ranged from 9.26 to 56.74 % for total exchangeable bases, 4.94 to 10.32 % for base  
130 saturation, and 1.48 to 50.50 % for ECEC. These results indicate that application of lime and SDA on  
131 acidic soil could have positive influence on both soil and crops being grown.

132 Expected reductions were observed for exchangeable acidity relative to control. This could be as a result  
133 of replacement of  $H^+$  and  $Al^{3+}$  by the basic cations present in SDA and lime occurring at the exchange  
134 sites. This consequently increased the pH of soil solution. Increase in soil pH upon SDA application  
135 affirms that ash has a liming effect. Several researchers [42, 37, 40] have successfully used plant ash as  
136 liming material.

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

UNDER PEER REVIEW

**Table 2: Effect of sawdust ash and lime on soil chemical properties**

		SAW DUST ASH (tha <sup>-1</sup> )						
	LIME(tha <sup>-1</sup> )	0	1	2	3	4	LSD (0.05)	
pH (water)	0	5.510	5.577	5.727	5.870	5.927	Lime NS	
	0.5	5.573	5.827	5.780	6.007	6.087	SDA	0.1595
	1.0	5.823	5.790	5.913	6.070	6.167	L×S	NS
	1.5	5.953	5.917	5.587	6.220	6.063		
pH (CaCl <sub>2</sub> )	0	4.073	4.093	4.377	4.663	4.637	Lime	0.2107
	0.5	4.320	4.563	4.587	4.780	4.987	SDA	0.2070
	1.0	4.513	4.727	4.677	5.023	5.450	L×S	NS
	1.5	4.620	4.790	4.290	5.117	5.383		
Exchangeable Acidity	0	6.64	20.19	15.47	8.15	7.24	Lime	NS
	0.5	11.78	11.30	11.59	12.99	12.20	SDA	0.2254
	1.0	13.71	11.59	17.60	6.84	10.87	L×S	NS
	1.5	14.76	14.58	9.92	19.66	10.02		
ECEC	0	6.64	20.19	15.47	8.15	7.24	Lime	NS
	0.5	11.78	11.30	11.59	12.99	12.20	SDA	2.756
	1.0	13.71	11.59	17.60	6.84	10.87	L×S	5.635
	1.5	14.76	14.58	9.92	19.66	10.02		
Base Saturation	0	79.10	94.01	91.93	92.68	91.84	Lime	2.483
	0.5	90.39	92.92	93.59	95.22	95.48	SDA	2.136
	1.0	92.23	94.32	94.32	91.42	95.10	L×S	4.323
	1.5	94.27	92.58	92.58	97.31	94.02		
Exchangeable bases	0	5.24	18.99	14.53	7.55	6.64	Lime	NS
	0.5	10.64	10.50	10.58	12.39	11.67	SDA	2.756
	1.0	12.64	10.93	16.66	6.31	10.34	L×S	5.635
	1.5	13.96	13.85	9.19	19.13	9.49		
Available Phosphorus	0	20.67	20.67	24.00	24.00	24.67	Lime	2.404
	0.5	23.67	24.67	18.67	27.00	23.67	SDA	3.514
	1.0	27.00	21.00	20.00	26.00	24.00	L×S	6.564
	1.5	20.67	26.67	26.60	32.67	20.67		
Total Nitrogen	0	0.080	0.107	0.080	0.067	0.080	Lime	0.01152
	0.5	0.100	0.080	0.097	0.047	0.077	SDA	0.01063
	1.0	0.067	0.060	0.080	0.053	0.080	L×S	0.02117
	1.5	0.137	0.067	0.067	0.097	0.080		
Organic Carbon	0	1.335	1.493	1.453	1.397	1.353	Lime	NS
	0.5	1.500	1.493	1.340	1.540	1.377	SDA	NS
	1.0	1.493	1.547	1.387	1.343	1.370	L×S	NS
	1.5	1.493	1.413	1.417	1.490	1.397		

- ECEC = effective cation exchange capacity
- SDA = sawdust ash
- L×S = lime and sawdust ash interaction

### 3.3 Effect of sawdust ash and commercial lime on growth and yield of sunflower

Growth parameters (Plant height, number of leaves, stem diameter), were obtained at the 10<sup>th</sup> week after planting. As expected, these parameters had increases relative to control. These increases were not significant for number of leaves (Fig. 2). However, there were significant ( $P<0.05$ ) increases in plant height and stem diameter. Similar finding was observed by Patterson *et al* [40].

The main effect of SDA resulted to significant increases in plant height (Fig. 1). The tallest plants of mean value, 123.70 cm were obtained at treatment combination 3  $\text{tha}^{-1}$  SDA  $\times$  1.5  $\text{tha}^{-1}$  lime. Presumably, the highest increase in plant height at this treatment may be due to the increases observed in available phosphorus and other nutrients. Phosphorus promotes cell division [43] which is manifested in plant height during vegetative growth.

For stem diameter, significant differences were observed in the main effects of SDA and lime. The highest mean value of 2.03 cm was recorded at the application rates of 2  $\text{tha}^{-1}$  SDA  $\times$  1  $\text{tha}^{-1}$  lime (Fig. 3).




Fig.1. Effects of sawdust ash and lime on plant height





Fig.2. Effects of sawdust ash and lime on number of leaves



**Fig. 3: Effects of sawdust ash and lime on stem diameter**

After harvesting, yield parameters were obtained and analyzed. The control had the least yield, indicating increases with treatment application. Only main effects of SDA had positive significant impact on these yield parameters. This suggests that SDA was able to release nutrients for improved sunflower yield. These increases in yield may be attributed to the increases in soil pH and nutrients provided by SDA.

The yield mean values ranged from 12.6 to 45.4 g, 1.73 to 5.33 g and 6.27 to 14.10 cm, with the highest yields recorded at application rates of 3  $\text{tha}^{-1}$  SDA  $\times$  1.5  $\text{tha}^{-1}$  lime, 3  $\text{tha}^{-1}$  SDA  $\times$  1.5  $\text{tha}^{-1}$  lime and 4  $\text{tha}^{-1}$  SDA  $\times$  1.5  $\text{tha}^{-1}$  lime, for head weight (Fig. 4), 50 seed weight (Fig. 5) and head diameter (Fig. 6) respectively. The positive response of sunflower to SDA and lime is similar with the earlier results obtained with several crops such as Cowpea [44, 37], Tomato [36], Amaranthus [45], Okra [36] and maize [38]. These studies attributed crop responses to SDA application which increased soil pH and nutrients such as N, P, K, Ca, Mg.




Figure 4: Effects of sawdust ash and lime on head weight





Figure 5: Effects of sawdust ash and lime on 50 seed weight



**Figure 6: Effects of sawdust ash and lime on head diameter**

### **3.4 Relationships between growth and yield parameters of sunflower and soil chemical properties**

Pearson correlation study was used to determine relationships between growth and yield parameters of sunflower and some soil chemical properties (Table not shown). Results showed that soil pH had positive relationship with all parameters obtained from pot experiment, indicating that increase in soil pH led to increases in these parameters. This finding corroborate earlier studies that grain yield of sunflower was highly positively correlated with soil pH [19]. The results ascertain that in soils of low pH, increment in pH through liming enhanced sunflower yield.

There was an expected inverse relationship between plant parameters and exchangeable acidity. However, this negative relationship was not significant for number of leaves. Data also showed that increases in plant height, head diameter, head weight and 50 seed weight were dependent on available Phosphorus. However, their dependence was not significant for plant height and head diameter.

## Conclusion

In southeastern Nigeria, where soil acidity poses a threat to sustainable crop production, acid sensitive and heavy feeder crop such as sunflower cannot be successfully grown without soil amelioration. The imbalance created by the use of chemical fertilizers alone has resulted to a quest for alternative sources of nutrients such as sawdust ash. The combined use of commercial lime and sawdust ash is expected to advance both economic and environmental management strategies in agriculture.

The present study showed that the soil was acidic, nutrient deficient and available phosphorus was below the optimum level required by sunflower. However, results indicated that application of sawdust ash and commercial lime based on agronomic principles such as lime requirement or for improving soil nutrient status have the potential to increase yield of sunflower in acidic, nutrient deficient soils of southeastern Nigeria.

Treatment combination of  $3 \text{ tha}^{-1}$  SDA  $\times$   $1.5 \text{ tha}^{-1}$  lime gave the most satisfactory yield of sunflower with regards to head weight and 50 seed weight by supplying soil nutrients and increasing pH.

## References

- [1] Groove JH, Summer ME. Yield and leaf composition of sunflower in relation to NPK, and lime treatments. *Nutrient Cycling in Agroecosystems*. 2005; 3/4: 367-378.
- [2] Aduayi EA, Chude VO, Adebusuyi BA, Olayiwola SO. Fertilizer use and management practices for crops in Nigeria. Abuja, Nigeria: Federal Ministry of Agriculture and Rural Development; 2002.
- [3] Aladodo SS, Mathew DE, Otu JO, Najib Y, Akoshile CO. Diurnal and interannual variability of rainfall in Nigeria climatic zones as seen from in-situ and satellite measurements. *Journal of Environment and Earth Science* 2018; 8(2): 9-18

- [4] Adebayo AG, Akintoye HA, Olatunji MT, Shokalu AO, Aina OO. Growth response and flower yield of sunflower to phosphorus fertilization in Ibadan, southwestern Nigeria. Report and Opinion. 2010; 2(4): 29-33
- [5] Lawal BA, Obigbesan GO, Akanbi WB, Kolawole GO. Effect of planting time on sunflower (*Helianthus annuus* L.) productivity in Ibadan Nigeria. African Journal of Agricultural Research. 2011; 6(13): 3049-3054
- [6] Ogunremi EA . The effect of plant population on sunflower seed yield in south western Nigeria. Ife Journal of Agriculture. 1979; 1: 51-56
- [7] Olowe VIO. Effect of plant population density on growth and yield of sunflower (*Helianthus annuus* L.) in the transition zone of south west Nigeria. Tropical Agricultural Research & Extension. 2005; 8: 37-44
- [8] Sutradhar A, Lollato RP, Butchee K, Arnall DB. Determining Critical Soil pH for Sunflower Production. International Journal of Agronomy. 2014; Accessed 25 June 2019. Available: <https://doi.org/10.1155/2014/894196>
- [9] Ibia TO, Effiong GS, Ogban PI, Obi JC. Relationships between phosphorus forms and parent materials in soils of southeastern Nigeria. Acta Agronomica Nigeriana. 2008; 8(2): 104-112
- [10] Osodeke VE. Determination of the residual value of applied phosphorus in some soils of southern Nigeria. J. Sustain, Agric. Environ. 2000; 2(1):139-143
- [11] Ohaeri JE, Eshett ET. Phosphorus forms and distribution in selected soils formed over different parent materials in Abia state of Nigeria. Agro-Science Journal of Tropical Agriculture, Food, Environment and Extension. 2011; 10(3): 28 - 37
- [12] Rodriguez D, Zubillaga MM, Ploschuck E, Keltjens W, Goudriaan J, Lavado R. Leaf expansion and assimilate prediction in sunflower growing under low phosphorus conditions. Plant Soil. 1998; 202: 133-147
- [13] Kovacevic V, Kovacevic J, Lalic A, Jurkovic, Krizmanic M. Influences of liming on maize, sunflower and barley. Cereal Research Communications. 2006; 34(1): 1-13

- [14] Blamey FPC, Nathanson K. Relationships between aluminum toxicity and sunflower yields on an Avalon medium sandy loam *Agrochimica*. 1977; 9: 59-65
- [15] Clapham WM, Zibilske LM. Wood ash as liming amendment. *Communications in Soil Science and Plant Analysis*. 1992; 23(11): 1209-1227
- [16] Brad NC, Weil RR. *The nature and properties of soils*. 12<sup>th</sup> ed. New Jersey: Prentice Hall Inc.; 1999.
- [17] Villapando RR, Graetz DA. Phosphorus sorption and desorption properties of the spodic horizon from selected Florida Spodosols. *Soil Sci. Soc. Am. J.* 2001; 65:331-339
- [18] Stevenson FJ, Cole MA. *Cycles of soils: Carbon, Nitrogen, Phosphorus, Sulfur, Micronutrients*. 2<sup>nd</sup> ed. Canada: John Wiley and Sons; 1999.
- [19] Ungaro MG, Ouaggio JA, Gallo PB, Dechen SCF, Neto FL, Melo de Castro O. Sunflower response to soil acidity. *Bragantia*. 1985; 44(1): 41-48
- [20] Eswaran H. Assessment of global resources: Current status and future needs. *Pedologie*. 1993; 43(1): 19-39.
- [21] Woomer PL, Ingram JSJ. *The biology and fertility of tropical soils. The Tropical Soil Biology and Fertility*. Nairobi, Kenya: Report; 1990
- [22] Enwezor WO, Ohiri AC, Opuwaribo EE, Udo EJ. *Literature review on soil fertility investigations in Nigeria*. Lagos: Federal Ministry of Agriculture and Natural Resources; 1990.
- [23] Osodeke VE, Uba AF. Determination of phosphorus fraction in selected soils of southeastern Nigeria. *International Journal of Natural and Applied Sciences*. 2005; 1(1): 10-14.
- [24] Bridgham SD, Johnston CA, Schubauer-Berigan JP, Weishampel P. Phosphorus sorption dynamics in soils and coupling with surface and pore water in riverine wetlands . *Soil Sci. Soc. Am. J.* 2001; 65: 577-588
- [25] Lin C, Banin A. Effect of long-term effluent recharge on phosphate sorption by soils in a wastewater reclamation plant. *Water, Air and Soil Pollution*. 2005; 164: 257-273.

- [26] Fagbayide JA, Adeoye GO. Sunflower growth and mineral uptake as influenced by fertilizer phosphorus. *Tropical oilseeds Journal*. 1999; 4: 21-28.
- [27] Adeoye GO, Agboola AA. Critical level for soil pH, available P, K, Zn and Mn and maize content of P, Cu, and Mn in the sedimentary soils of southeastern Nigeria. *Fertilizer Research*. 1984; 6: 65-71
- [28] Agboola AA, Corey RB. The relationship between soil pH, organic matter, available phosphorus, exchangeable calcium, magnesium and nine elements in the maize tissue. *Soil Sci*. 1973; 115(5): 367-375.
- [29] Bouyoucos GH. A recalibration of the hydrometer method for mechanical analysis of soil. *Agron. Journ*. 1951; 43: 434-438
- [30] Mclean EO. Exchangeable aluminium. In: Black CA, editor. *Methods of soil plant II. Chemical methods*. Madison: ASA; 1965.
- [31] Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter and proposed modification of the chromic acid and titration method. *Soil sci*. 1934; 37: 29-38
- [32] Bray RH, Kurtz LT. Determination of total organic and available forms of phosphorus in soils. *Soil Sci*. 1945; 59: 39-45
- [33] Jackson ML. *Soil chemical analysis*. Englewood cliffs, NY: Prentice Hall; 1962
- [34] Kamprath EJ. (1967). *Soil acidity and response to liming*. Int. Soil Testing Series N. C. State University: Tech Bull. No. 4; 1967
- [35] Odedina SA, Odedina JN, Ayeni SO, Arowojolu SAA, Adeyeye SD, Oyeniyi SO. Effect of types of ash and soil fertility nutrient availability and yield of tomato and pepper. *Nig. J. Soil Sci*. 2003; 13: 61-67
- [36] Owolabi O, Ojeniyi SO, Amodu AO, Hassan K. (2003). Response of cowpea, okra and tomato to sawdust ash manure. *Moor Journal of Agricultural Research*. 2003; 4(2): 178-182
- [37] Awodun MA. Influence of sawdust ash on soil chemical properties and cowpea performance in southwest Nigeria. *Intl. J. Soil Sci*. 2007; 2(1) :78-81

- [38] Onwuka MI, Osodeke VE, Okolo NA. Amelioration of soil acidity using cocoa husk ash for maize production in Umudike area of south east Nigeria. *Tropical and Subtropical Agroecosystems*. 2007; 7: 41-45.
- [39] Ewulo BS, Babadele OO, Ojeniyi SO. Sawdust ash and urea effect on soil and plant nutrient content and yield of tomato. *Am-Eurasian J. Sustain. Agric.* 2009; 3(1): 88-92
- [40] Patterson SJ, Acharya SN, Thomas JE, Bertschi AB, Rothwell RL. Barely biomass and grain yield and canola seed yield response to land application of wood ash. *Agron. J.* 2004; 96: 971-977
- [41] Baath E, Arnebrant K. Growth rate and response of bacteria communities to pH in ash treatment forest soils. *Soil Biol. Chem.* 1994; 26: 995-1001.
- [42] Ayeni LS, Ayeni OM, Oso OP, Ojeniyi SO. Effect of sawdust and wood ash applications in improving soil chemical properties and growth of cocoa (*Theobroma cacao*) seedlings in the nurseries. *Agric. J.* 2008; 3: 323-326.
- [43] Ugese FD and Avav T. Response of mungbean (*Vigna radiata*) to phosphorus application in a southern guinea savanna location of Nigeria. *Nigeria Journal of Sustainable Tropical Agricultural Research*. 2005; 13: 77-80
- [44] Obi O, Ekperigin J. Effects of wastes and soil pH on growth and yield of crops. *African Soils*. 2001; 33: 3-5.
- [45] Ojeniyi SO, Adejobi KB. Effect of ash and goat dung manure on leaf nutrient composition growth and yield of *Amaranthus*. *Niger Agric J.* 2002; 33: 46-57

UNDER PEER REVIEW