

GENETIC VARIABILITY AND CHARACTER ASSOCIATION FOR OIL YIELD AND ITS COMPONENTS IN PHYSIC NUT (*Jatropha curcas* L)

ABSTRACT

The need to find suitable alternative to fossil fuel necessitated the search for non-food oil crops like physic nut (*Jatropha curcas*). Improvement of the crop requires adequate knowledge of extent of genetic variability present. Therefore, a study was carried out to determine genetic variability and character association among forty local and exotic accessions of physic nut. The accessions were evaluated in two years in randomized complete block design with two replications. Results indicated highly significant variation among the accessions for all characters studied except plant height and number of seeds per fruit. High estimates of heritability were observed in number of leaves (84.10%) and seed oil (94.97%) while heritability was low in number of seeds (11.30%) and plant height (13.36), other characters had moderate heritability. High heritability estimates were accompanied by relatively low genetic advance in all the traits except kernel oil, an indication that the character has high selection value with less environmental influence. Correlation analysis showed that tall plants produced bigger seeds because of the positive association between plant height and seed weight. Therefore, seed weight can be used as selection index.

INTRODUCTION

The global demand for energy is ever increasing. Fossil fuels have been the major energy source in the world over time. However, the environment is currently challenged by the continuous use of fossil fuels as it causes serious environmental problems such as air pollution, global warming, and environmental degradation amongst others. In recent times, attention has been shifted to the use of biofuel which has less environmental problems. Majority of oil crops used in biofuel production are food crops. This can greatly impact on food production negatively, arising from competition with food supply and can lead to food shortage especially in the developing countries. Thus, there is need to focus on non-edible oil plants to serve as feedstock for biofuel. *Jatropha curcas* is one of many of these promising non-edible oil plants under consideration.

Jatropha curcas, also known as the physic nut, is a shrub which can grow up to 5-7 meters [1, 2]. It originates from Mexico or Central America, and was introduced to Africa by the Portuguese sailors in the 16th century [1]. The physic nut is one of the leading non-edible oil plants [4] that produces a substantial quantity of oil from its seeds. The oil content has been reported to range between 25 and 50% [4, 5, 6]. The use of *J. curcas* oils has increased in importance and has, in some cases, replaced the use of fossil fuels. Apart from the prominence gained in oil production, *J. curcas* is important in carbon sequestration. [7]. Nahar and Sunny [8] showed that a mature plant absorbs CO₂ at a rate of 18lbs per year, while Becker *et al.* [9] in their study spanning over a 20-year period revealed that one hectare of *J. curcas* could sequester up to 25 tons of CO₂ from the atmosphere every year.

An important plant like *J. curcas* needs genetic improvement in order to alter its status from wild to a cultivable crop with higher yield and oil content. One of the major constraints in achieving genetic improvement is absence of information about genetic variability. Genetic variability is a

pre-requisite for improvement of any crop because variability precedes for selection. According to Jubera [11] without the presence of sufficient genetic variability of the desired traits that are of economic interest, an attempt to improve this crop will be unsuccessful or a failure.

Currently in Nigeria crop improvement work in this plant is very limited. To fully exploit this multiple utility biofuel crop, work is required to select the best genotypes for oil content and quality. The genetic diversity in the natural population appears narrow since no remarkable genetic variability has been observed. Therefore, in commencing any crop improvement programme, there is need to determine the amount, cause and nature of variation that is present in the species of interest. In view of the foregoing, the main objective of this study is to evaluate the magnitude of genetic diversity, heritability and correlation between yield and its contributing characters in thirty two accessions of *J. curcas* from six states in Nigeria and one exotic genotype from Brazil.

MATERIALS AND METHODS

This experiment was conducted at the Laboratory of the Department of Crop, Soil and Pest Management and Teaching and Research Farm of the Federal University of Technology Akure (7° 16' N, 5° 12' E) located in the rainforest area of Southwestern Nigeria. The location is characterized by a bimodal pattern of rainfall with an annual mean of about 1300mm with a mean temperature of 27° C and the climate is of the sub-humid type. The vegetation of the experimental plot was mixture of *Panicum maximum* and *Chromolaena odorata* plants.

Mature fruits were collected from thirty-two accessions of *Jatropha* from six states in Nigeria [Edo (6), Osun (5), Kogi (6), Ekiti (4), Kwara (4) and Ondo (7)] and fruits from eight trees from

Brazil. Experimental plot was manually cleared and seeds were planted directly on flat ground. The experimental design was a randomized complete block design. The experimental plot was divided into three replications each measuring 3m by 33m. Alleys of 1m were created between replications. Seven seeds were selected randomly from the fruits of each accession and planted on the plot allocated to each accession and replicated three times. Planting was done on 18th August, 2010 for the first experiment and 1st October, 2011 for the second. Glyphosate was used to control newly germinated weeds 2 weeks before planting. Thereafter manual weeding was carried out at 3 weekly intervals. The following quantitative data were collected on each genotype for agronomic and yield characters from 12 weeks after planting (WAP):

Plant height (cm): measured from base of plant at soil level to the tip of the plant with a meter tape

Stem girth (cm): measured at the base of the plant with the aid of vernier caliper

Number of branches: the total number of branches per plant was counted.

Number of leaves: the total number of leaves per plant was counted.

Leaf length (cm): measured from base of the leaf to the tip of the leaf. The mean length of matured leaves (20 leaves/plant) was recorded.

Petiole length (cm): measured from point of attachment at the stem to the base of leaf. The mean length of matured leaves (20 leaves/plant) was recorded.

Number of seeds: the total number of seeds harvested per plant throughout the fruiting period was counted.

Seed weight (g): 100 seeds per plant were weighed using electronic balance.

Percentage oil content: percentage seed, kernel and shell oil was estimated using a Soxhlet extraction method.

The data obtained were subjected to analysis of variance using PROC GLM of SAS [11]. Genotypic and phenotypic coefficients of variation were computed according to the procedure of Singh and Chaudhary [13] as follows:

$$\text{Genotypic coefficient of variation (GCV)} = \frac{\sqrt{\sigma_g^2}}{\bar{x}} \times 100$$

$$\text{Phenotypic coefficient of variation (PCV)} = \frac{\sqrt{\sigma_p^2}}{\bar{x}} \times 100$$

GCV and PCV values obtained were categorized as low, moderate and high according to Siva-Subramanian and Menon [14],

0 – 10%:	Low
10 – 20%:	Moderate
21% and above:	High

Broad sense heritability was estimated as the ratio of genotypic variance to the phenotypic variance and expressed in percentage by Singh and Chaudhary [12] as:

$$\text{Heritability (H)} = \frac{\sigma^2_g}{\sigma^2_g + \sigma^2_e} \times 100$$

Where: σ^2_g = genotypic variance
 σ^2_p = phenotypic variance
 σ^2_e = environmental variance
 $\sigma^2_g + \sigma^2_e = \sigma^2_p$

Heritability percentage was categorized as low, moderate and high as indicated by Elrod and Stanfield [15] as follows,

0 – 20%:	Low
20 – 50%:	Moderate
50% and above:	High

Genetic advance (GA) was computed according to the formula given by [16] Johnson *et al.* (1955)

$$GA = \frac{\sigma^2_g}{\sqrt{\sigma^2_p}} \times K$$

Where:

$K = 2.06$ (selection differential at 5%)

σ^2_g = Genotypic variance

σ^2_p = Phenotypic variance

\bar{x} = Sample mean of the character

Analysis of covariance was carried out on pairs of variables which exhibited significance ($P \leq 0.05$) in the regression analysis. The generated components of the covariance were used to estimate the phenotypic and genotypic correlation coefficients as suggested by Singh and Chaudhary [12]:

$$\text{Phenotypic Correlation Coefficient } (r_p) = \frac{COV_p(X_1, X_2)}{\sqrt{\sigma^2_{p(X_1)} \times \sigma^2_{p(X_2)}}}$$

$$\text{Genotypic Correlation Coefficient } (r_g) = \frac{COV_g(X_1, X_2)}{\sqrt{\sigma^2_{g(X_1)} \times \sigma^2_{g(X_2)}}}$$

Where:

$COV_g(X_1, X_2)$ = Genetic covariance among traits X_1 and X_2 .

$COV_p(X_1, X_2)$ = Phenotypic covariance among traits X_1 and X_2 .

$\sigma^2_{g(X_1)}$ and $\sigma^2_{g(X_2)}$ = Genetic variance for traits X_1 and X_2 , respectively.

$\sigma^2_{p(X_1)}$ and $\sigma^2_{p(X_2)}$ = Phenotypic variance for traits X_1 and X_2 , respectively.

Test of significance of correlation was done by comparing the computed values against tabular 'r' values given by Fisher and Yates [16].

RESULTS

The mean squares from the analysis of variance for the eleven plant characters that were investigated are shown in Table 1. Mean square for replicate was significant in most characters tested including number of branches, plant height, stem girth, number of seed and seed oil. Highly significant differences for genotype were observed in all the characters except plant height and number of seeds. The means, ranges and standard deviation of eleven characters among the forty genotypes of *J. curcas* are presented in Table 2. Plant height ranged from 110cm to 279cm with mean value of 180.29 cm. Wide variation was observed for number of leaves, number of branches and number of Seed as they varied from 82 to 632, 3 to 45 and 2 to 513 respectively. Moderate variation was observed in stem girth, percentage kernel oil and percentage seed oil. Low variation was recorded for leaf length and seed weight as they ranged from 12 cm to 18cm and 62.08g to 79.38g respectively.

Estimates of genotypic coefficient of variation, phenotypic coefficient of variation, genotypic variance, phenotypic variance, heritability, genetic advance and expected genetic gain for eleven agronomic characters of forty genotypes of *J. Curcas* are presented in Table 3. Number of branches, number of seeds and seed weight showed high genotypic and phenotypic coefficients of variation (21.19%, 45.96, 27.40%, 81.54 % and 21.22%, 39.82%) respectively. Number of leaves, leaf length, kernel oil and seed oil had low genotypic and phenotypic coefficients of variation. Low genotypic coefficient of variation were observed for petiole length, plant height and stem girth. Phenotypic coefficients of variation for these characters were moderate. Phenotypic coefficient of variation was higher than genotypic coefficient of variation in all the

characters evaluated. High estimates of heritability were observed in number of leaves (84.10%) and seed oil (94.97%), while heritability was low in number of seeds (11.30%) and plant height (13.36). Other characters had a moderate heritability. Genetic advance and expected genetic gain were observed to be low for all the characters studied.

The phenotypic (r_p) and genotypic (r_g) correlation coefficients among the eleven characters in the forty *J. curcas* genotypes are presented in Table 4. Number of branches showed significant and positive phenotypic and genotypic correlations with stem girth (0.70, 0.72), number of seeds (0.33, 0.49) and leaf length (0.31, 0.29), respectively but correlated significantly and positively with plant height (0.36) only at the genotypic level, and with kernel oil (-0.34) at the phenotypic level only. Petiole length had significant and positive r_p and r_g with plant height (0.34, 0.40), stem girth (0.37, 0.53) and leaf length (0.36, 0.70) respectively but also with shell oil (0.29) only at the genotypic level. It had significant negative correlation at phenotypic level with number of seeds (-0.76). Number of leaves had significant and positive r_g with number of seeds (0.61) but significant and negative r_g with seed weight (-0.40). At genotypic level, plant height showed significant positive correlation with number of seeds (0.29) and seed weight (0.62), but correlated significantly and negatively with leaf length (-0.35). At the phenotypic level, plant height correlated significantly and positively with stem girth (0.36). Stem girth had significant positive r_p and r_g with leaf length (0.35, 0.66), and also showed significant positive r_g with seed weight (0.41), but negative with number of seeds (-0.47). Leaf length showed significant and negative r_g with kernel oil (-0.34). Kernel oil had significant and positive genotypic correlations with shell oil (0.38), seed oil (0.92) and seed weight (0.31), and significant positive phenotypic correlations with shell oil (0.35), seed oil (0.91) only. Shell oil also showed a significant and

positive r_p and r_g with seed oil (0.71, 0.71) and r_g with seed weight (0.57). Seed oil showed significant positive r_g with seed weight (0.48).

DISCUSSION

Crop improvement is achieved when there is significant genetic variability among genotypes for the desired traits. The observed differences among the genotypes evaluated in this study differed in yield characters tend to suggest that breeding work can be initiated through selection or hybridization. The high level of variability among the genotypes suggests that heterosis could be utilized to produce superior hybrids which can further be used in breeding programme to develop superior genotypes [17]. Development of such genotypes requires that overall variability should be partitioned into heritable and non-heritable components with the aid of genetic parameters such as genotypic and phenotypic coefficients of variation and heritability [19]. In this study, phenotypic variance was greater than genotypic variance in all the characters for yield parameters. Similar findings were reported in okra [18], in cucumber [20] and in the fluted pumpkin [20]. The high genotypic and phenotypic coefficients of variation observed in number of seeds and seed weight suggests that greater potential might be expected in selection for these characters. The observed high heritability values in number of leaves and seed oil content indicate that the characters were under genetic control and less influence of the environment on these characters. According to Ojo and Amanze [22], high heritability strongly suggests that there is potential for large genetic determination for these characters which can be exploited for improvement of oil yield in this crop.

Since high heritability does not always indicate a high genetic gain, heritability with genetic advance considered together should be used in predicting the ultimate effect of selecting superior genotypes [23]. In this study, high heritability estimates were accompanied by relatively low genetic advance in all the traits except kernel oil, an indication that the character has high selection value with less environmental influence. Therefore, improvement by direct phenotypic selection is possible. Whenever a high heritability value is accompanied by high genetic advance, it may be governed by additive gene action and improvement with respect to these characters could be brought about by phenotypic selection [23; 20]. Moderate heritability with low genetic advance were observed for some characters in this study. They include petiole length, stem girth, kernel oil, shell oil and seed weight. This indicates the existence of intermediate additive and non-additive gene action. Consequently improvement of these characters can be achieved through careful selection. The low heritability and low genetic advance recorded for plant height and number of seeds suggest that these characters governed by non-additive gene action. The correlation coefficients among characters were determined at the phenotypic and genotypic levels. In general, most genotypic correlation coefficients were higher in magnitude than phenotypic correlation coefficients. This can be interpreted as a strong inherent genotypic relationship between characters studied, though their phenotypic expression was influenced by environmental influence. Direct selection for improvement some of these traits may not be possible since variation among the genotypes is attributed largely to the environment.

Significant and positive correlation between two characters suggests that these characters can be improved simultaneously in a selection programme [24; 20]. This is because it shows mutual relationships among characters and selection for one will translate to selection of the other [20].

For the improvement of oil yield in this crop, it was necessary to determine the magnitude and the direction of relationship between the oil yield and its components. Studies on the character associations showed that tall plants might produce bigger seeds because of the positive association between plant height and seed weight. The leaves of tall plants can easily intercept more light than short plants. High light interception will lead to greater accumulation of photosynthate. Accumulation of photosynthate in leaves has been reported to increase seed weight of okra [18]. The reduction in oil yield caused by big branches and broad leaves could be attributed to diversion of photosynthate for seed yield to vegetative growth (branch and leaf). This is in agreement with the findings of Patil *et al.* [25]. Therefore, it can suggest that to ensure good yield pruning should be carried out on this crop as it is done in other tree crops. Seed weight was observed to have significant genotypic relationship with seed oil and shell oil also. An increase in weight may therefore lead to an increase in the latter two traits. As such, seed weight may be used as selection index. Prasanthi *et al.* [26] reported that the weight of seed per plant and numbers of seeds per plant were important selection indices. This evaluation may be useful in developing reliable selection indices for important agronomic traits in *Jatropha curcas*. It is recommended that future research explore molecular means to further confirm the outcome of this research and establish the relationship between the two methods.

Table 1. Mean squares from analysis of variance (ANOVA) for eleven agronomic characters of forty *J. curcas* genotypes

Source	DF	Number of Branches	Petiole Length (cm)	Number of Leaves	Plant Height (cm)	Stem Girth (cm)	Number of seeds per plant	Leaf Length (cm)	Kernel Oil (%)	Shell Oil (%)	Seed Oil	Seed Weight (g)
Rep	2	135.36*	2.71ns	957.59ns	3223.80*	36.60**	34193.89**	0.40ns	0.54ns	1.75ns	4.04*	0
Genotype	42	60.79*	11.66**	36128.71**	1463.77ns	12.78**	6034.32ns	1.67**	40.05**	12.57**	69.16**	70.99**
Error	84	33.59	3.43	490.68	1000.9	6.63	4366.73	0.78	0.36	0.70	1.20	0

*Significant at $p < 0.05$, **Significant at $p < 0.01$

NBR = number of branches LFLT = leaf length (cm) PLT = petiole length (cm)

PLHT = plant height (cm) SHOL = shell oil (%) KNOL = kernel oil (%)

STGT = stem girth (cm) SDOL = seed oil (%)

NOSD = number of seeds SDWT = seed weight (g)

NOLV = number of leaves

Table 2. Mean, range and standard deviation of eleven characters among forty genotypes of *J.*

curcas

Characters	Mean	Minimum	Maximum	Standard deviation
Number of branches	14.21	3.00	45.00	6.64
Petiole length (cm)	19.02	13.90	30.70	2.47
Number of leaves	293.22	82.00	632.00	110.42
Plant height(cm)	180.29	110.00	279.00	34.46
Stem girth (cm)	23.54	16.70	31.70	3.02
Number of seeds	86.05	2.00	513.00	73.35
Leaf length(cm)	14.88	12.00	18.00	1.03
Kernel oil (%)	61.87	51.88	71.88	3.66
Shell oil (%)	11.53	3.75	15.00	2.15
Seed oil (%)	73.40	61.25	83.76	4.80
Seed weight (g)	68.97	62.08	79.38	4.83

Table 3. Estimates of Genetic coefficient of variation, Phenotypic Coefficient of variation, Genotypic variance, Phenotypic variance, Heritability, Genetic advance and Expected genetic gain for eleven agronomic characters in *J. curcas* genotypes

Characters	Genotypic coefficient of variation (%)	Phenotypic coefficient of variation (%)	Genotypic variance σ_g^2	Phenotypic variance σ_p^2	Environmental variance σ_e^2	Heritability (%)	Genetic Advance	Expected genetic gain (%)
Number of branches	21.19	45.96	9.07	42.66	33.59	21.26	2.86	20.12
Petiole length (cm)	8.72	13.08	2.74	6.17	3.43	44.44	2.27	11.97
Number of leaves	1.35	1.47	15.66	18.62	490.68	84.10	7.48	2.55
Plant height (cm)	6.89	18.85	154.29	1155.18	1000.89	13.36	9.35	5.19
Stem girth (cm)	6.08	12.52	2.05	8.68	6.63	23.62	1.43	6.09
Number of seeds	27.40	81.54	555.86	4922.59	4366.73	11.30	16.32	18.97
Leaf length (cm)	3.66	6.97	0.30	1.08	0.78	27.55	0.59	3.96
Kernel oil (%)	0.54	1.11	0.11	0.47	0.36	23.94	0.34	0.55
Shell oil (%)	12.11	24.22	1.95	7.80	5.85	25.00	1.44	12.47
Seed oil (%)	6.48	6.65	22.65	23.85	1.2	94.97	9.55	13.02
Seed weight (g)	21.22	39.82	22.08	77.71	55.63	28.42	5.16	23.31

Table 4. Genetic correlation (r_g) and phenotypic correlation (r_p) for eleven characters of 40 *J. curcas* genotypes.

		Petiole length (cm)	Number of leaves	Plant height (cm)	Stem girth (cm)	Number of seeds	Leaf length (cm)	Kernel oil (%)	Shell oil (%)	Seed oil (%)	Seed weight (g)
Number of branches	r_g	0.21	0.04	0.36*	0.72**	0.49**	0.29*	-0.17	-0.02	-0.14	0.14
	r_p	0.28	0.07	0.26	0.70**	0.33*	0.31*	-0.34*	-0.05	-0.28	0.14
Petiole length (cm)	r_g		-0.06	0.40**	0.53**	-0.76**	0.70**	-0.19	0.29*	-0.03	0.18
	r_p		-0.01	0.34*	0.37**	-0.14	0.36*	-0.13	0.23	0.01	0.06
Number of leaves	r_g			0.23	-0.12	0.61**	0.03	0.05	-0.21	-0.05	-0.40**
	r_p			0.16	-0.03	0.24	0.06	0.05	-0.18	-0.04	-0.2
Plant height (cm)	r_g				0.11	0.29*	-0.35*	-0.02	0.04	0.01	0.62**
	r_p				0.36*	0.12	0.28	-0.02	0.06	0.01	0.11
Stem girth (cm)	r_g					-0.47**	0.66**	-0.21	0.04	-0.14	0.41**
	r_p					0.22	0.35*	-0.09	0.03	-0.05	0.1
Number of seeds	r_g						-0.27	-0.41**	-0.98**	-0.79*	-0.79**
	r_p						0.03	-0.11	-0.32*	-0.23	-0.13
Leaf length (cm)	r_g							-0.34*	0.1	-0.21	0.07
	r_p							-0.18	0.06	-0.11	0.02
Kernel oil (%)	r_g								0.38**	0.92**	0.31*
	r_p								0.35*	0.91**	0.15
Shell oil (%)	r_g									0.71**	0.57**
	r_p									0.71**	0.26
Seed oil (%)	r_g										0.48**
	r_p										0.23

*, ** significant at 0.05, 0.01 probability level respectively.

REFERENCES

1. Heller, J. (1996). Physic nut, *Jatropha curcas*. Promoting the conservation and use of under-utilized and neglected crops. International Plant Genetic Resources Institute, IPGRI, Rome, Italy, 66 pp.
2. Atabani, A. E., Silitonga, A. S. and Ong, H. C. (2013). Non-edible vegetable oils: A critical evaluation of oil extraction, fatty acid compositions, biodiesel production, characteristics, engine performance and emissions production. *Renewable and Sustainable Energy Reviews* 18: 211-245.
4. No, S. Y. (2011). Inedible vegetable oils and their derivatives for alternative diesel fuels in CI engines: A review. *Renewable and Sustainable Energy Reviews* 15(1): 131-149.
5. Pramanik, K. (2003). Properties and use of *Jatropha curcas* oil and diesel fuel blends in compression ignition engine. *Renewable Energy* 28(2): 239-248.
6. Pant, K. S., Kumar, D. and Gairola, S. (2006). Seed oil content variation in *Jatropha curcas* L. in different altitudinal ranges and site conditions in H.P. India Lyonia.11: 31–34.
7. Pandey, V. C., Singh, K. and Singh, J. S. (2012). *Jatropha curcas*: A potential biofuel plant for sustainable environmental development. *Renewable and Sustainable Energy Reviews* 16(5): 2870–2883
8. Martin, G. and Mayeux, A. (1985).Curcas oil (*Jatropha curcas* L.): a possible fuel. *Agr.Trop.*9:73- 75.
9. Nahar K and Sunny SA (2014) *Jatropha curcas* L: A sustainable feedstock for the production of bioenergy and by products. *Journal of Energy and Natural Resources* 3: 51–57.
10. Becker, K, Wulfmeyer, V, Berger, T (2013) Carbon farming in hot, dry coastal areas: An option for climate change mitigation. *Earth System Dynamics* 4(2): 237–251.

11. Jubera, M. A. (2008). Morpho-Physiological and Molecular Diversity in *Jatropha curcas*(L.) Msc. Thesis submitted to the University of Agricultural Sciences, Dharwad.
12. Statistical Analysis System (SAS). 2000. Online Doc. Version 9.2. Cary,NC: SAS Institute Inc.
13. Singh, R.K. and Chaudhary, B.D. (1985). Biometrical Methods in Quantitative Analysis. Kalyani Publishers. New Delhi. 318pp.
14. Siva-Subramanian, S. and Menon, M. (1973). Heterosis and inbreeding depression in rice. *Madras Agricultural Journal*, 60:1139.
15. Elrod, S. and Stanfield, W. (2002). Genetics. 4th Ed. Tata McGraw Hill Co., New Delhi.
16. Johnson, H.E., Robinson, H.F. and Comstock, R.E. (1955). Estimates of genetic and environmental variability in soybean. *Agronomy Journal* 47:314-318.
17. Fisher, G. and Yates, M. (1963). Statistical tables for biological, Agricultural and Medical Research, 6: 62-63.
18. Makinde, S.C. (1988). Genetic characterization of okra (*Abelmoschus esculentus* L.) Moench cultivars. Unpublished M. Sc. Thesis, Faculty of Science, University of Ilorin, Ilorin. Nigeria.
19. Ariyo, O.J. (1987). Variation and heritability of fifteen Characters in okra (*Abelmoschus esculentus* L.). *Tropical Agriculture*, 34: 67-71
20. Afangideh, U. and Uyoh, E.A. (2007). Genetic variability and correlation studies in some varieties of cucumber (*Cucumis sativus* L.). *Jordan J. Agric. Sci.* 3(4): 376-384.
21. Fayeun, L.S., Odiyi, A.C., Makinde, S.C.O. and Aiyelari, O.P. (2012). Genetic Variability and Correlation Studies in the Fluted Pumpkin (*Telfairia occidentalis* Hook.F.) *Journal of Plant Breeding and Crop Science* 4(10): 156-160.

22. Ojo, D.K. and C.O. Amanze. (2001). Prediction of grain yield through heritability and genetic advance yield parameters in soybean (*Glycine max. L.*) Merr). *Nigerian Journal of Ecology* 3: 10-13.
23. Ali, A., Khan, S., Asad, M.A., 2002. Drought tolerance in wheat: Genetic variation and heritability for growth and ion relations. *Asian Journal of Plant Science*, 1: 420-422.
24. Hayes, H. K., Immer, F. R. and Smith, D. C. (1955). *Methods of Plant Breeding*, 2nd ed,; McGraw-Hill Book Co.: New York, NY, USA. P 551.
25. Patil, S.J., Mutanal, S.M., Shahpurmath, G.B. and Mannikeri, I. M. (2006). Effect of pruning and nutritional levels on growth of *Jatropha (Jatropha curcas)* under rainfed conditions, *Indian J. Agroforestry*, 8(2): 5-9.
26. Prasanthi, L., Maheswara, P., Reddy, P. S., Sudhakar, B., Babu, B., and Reddy, R., (2009). Genetic variability and path analysis for yield and yield attributes in *Jatropha (Jatropha curcas L.)* *J. Oilseeds Res.*, 26: 230-233.