

# Fertilization and the impact of the soil elements supply on the production of plums in organic farming

## Abstract

Organic farming takes up more and more areas in agricultural production. It is based on the principles of preserving human and animal health and maintaining the balance in agrobiocenosis and therefore prohibits the use of a majority of water-soluble mineral fertilizers and phytopharmaceuticals of chemical origin (except those from the list of allowed products). In ecological farming of plums, optimization of fertilization is very important, which must comply with ecological principles and regulations which included fertilization with a relatively narrow range of allowed organic fertilizers (manure, compost, manure, slurry, peat, guano, sawdust) and mineral additives (calcium carbonate, crude phosphates, basic slag, raw potassium salt, potassium sulphate, gypsum, wood ash, calcium chloride, sodium chloride, etc.).

Nutrition methods in organic plum production is different from nutrition in conventional production, primarily due to limited selection of available fertilizers, and the most important difference is the inability to use water soluble individual and complex fertilizers. It is precisely for this reason that in organic agriculture, the emphasis on maintaining humidity, optimum pH and soil moisture is considerably more important than necessary prerequisites for sufficient availability of plant nutrients in the soil.

*Keywords: Ecological plum production, fertillization optimization, keeping humility, organic fertilizers*

## 1. INTRODUCTION

Plum in Bosnia and Herzegovina is a national fruit. In the last century, ex-Yugoslavia was one of the largest plum producers with 18% of world production, but over time, plum lost the economic importance it once had. Nevertheless, it is today the most widespread fruit culture on the territory of Bosnia and Herzegovina. According to the National Directorate of Statistics (2006), in Bosnia and Herzegovina there are 10.9 million plum trees with an average yield of 9.3-9.5 kg per tree. The fruit of plum is rich in vitamins and abundance of minerals, pectin and organic acids, and occupies a very important place in human nutrition. It is a good antioxidant, and its anthocyanins can provide human protection against oxidative stress. Fruit plums contain 75 to 87% water and 13 to 25% dry matter. The taste of the fruit depends on ratio of sugar and organic acids. In the fruit of the plum predominates glucose, while it is less fructose. Among the acids most are malic and citric. The total amount of organic acids expressed as malic acid ranges from 0.39 to 2.28%. The percentage of acids and sugars varies with the same variety, depending on climatic and soil conditions. In the mature fruit of plums the pH value ranges from 3.3 to 3.6.

The fleshy part of the plum of the plum contains up to 0.69% of pectic substances, which are important in the technology of jellies, marmalades and similar products. Nitrogenous substances participate in the chemical composition of the fruit with 0.6 to 0.8% (amino acids, amides, ammonia compounds and nitrogen bases), and minerals with only 0.5%. Of the mineral substances, the highest is potassium (54.59%) and phosphorus (17.7%). Plum fruit contain large amounts of fat (Mitrović Olga et al., 2009).

Bosnia and Herzegovina fulfills the basic geographical and climatic precondition for plum cultivation, and the remaining attention is paid to the technology of breeding, and most of the nutrition and protection of plums. The plum lives and yields on the same land for 30-50 years, and in some cases even longer. Through this relatively long period, plum with its root system from the soil, take out large amounts of nutrients are available in the available form that are necessary for the growth and development of vegetative and generative organs. Particular accent is given to the high yielding as a necessary factor for the production process. A very important factor of cost-effective production is the maintenance of soil fertility, but sometimes the soil does not have the optimal amount of nutrients for the plant, so it is necessary to provide agricultural land with the help of agro-technical measures by fertilizing the nutrients in the soil or plant nutrients by foliar application.

Of course, the repair of chemical, physical and biological characteristics of the land should not be done by the heart, but only with the analysis of the soil, in compliance with the rules of the agronomic profession in accordance with modern plum cultivation technologies. Applying modern knowledge about plum cultivation technology, while adhering to already known principles of sustainable agricultural production and processing technology, plums can result in growing plum production and

61 revitalizing the status of an economically significant culture. A study of agroecological and  
62 technological characteristics of production with the soil conditioning plan and fertilization plan prior to  
63 planting and the fertilization plan for the future plum planting, significantly contributes to the  
64 sustainable ecological production of plums in Bosnia and Herzegovina, which could quantitatively and  
65 qualitatively realize the great potential of placing on the market of organic products.

66 Organic production is a complete food production management system that combines the best  
67 ecological practices, a high degree of biodiversity, the conservation of natural resources, the  
68 application of high animal welfare standards and the way of production in line with consumer  
69 expectations, using natural substances and processes. Organic production has a dual social role. On  
70 the one hand, it provides public goods that contribute to the protection of the environment and animal  
71 welfare, and on the other, contributes to the development of rural areas (EC Regulation 834/2007).  
72 Organic agriculture is a specific agricultural production system that represents a systematic and  
73 comprehensive approach to sustainable survival, taking into account factors that influence sustainable  
74 development and vulnerability at the physical, economic and socio-cultural level (Eyhorn, 2007).  
75 Therefore, organic production can significantly contribute to the development of rural communities  
76 (Lobley et al., 2005; El-Hage Scialabba and Hattam, 2002; Eyhorn, 2007).

77 At the same time, organic food can respond to increasingly lucrative consumer demands for healthy,  
78 quality and safe food, whose production will not impair water and soil quality, biodiversity and animal  
79 welfare. Therefore, it is classified as a higher quality food. According to Little and Frost (2008), the  
80 main elements of organic production systems are:

- 81 • Good soil management leading to good fertility, maintaining high content of organic matter, high  
82 microbiological activity and good land structure,
- 83 • Well-designed crop rotation that are important for balanced crop nutrition, weed control and  
84 minimize disease and pest problems,
- 85 • preventive, not chemical, approach to controlling weeds, diseases and pests,
- 86 • profitable contribution of organic cover crops and livestock.

## 91 2. MATERIAL AND METHODS

92 For this work the data of the Federal Hydrometeorological Institute of BiH were used for: temperature,  
93 precipitation and insolation for 2017. Data were processed for the whole of BiH, but the accent was  
94 placed on the territory of the Tuzla Canton. For temperature it is described average air temperatures,  
95 average annual temperatures, temperature deviations and based on this is given a conclusion on  
96 whether and how much it affects plum cultivation. According to the distribution of percentile, rainfall  
97 conditions in 2017 were classified into categories (10-25 percentiles), normal (26-74 percentiles), rain  
98 (75-90 percentiles) and very rainy (> 98 percentile). Months with the highest precipitation are shown,  
99 as are the months with the least amount of precipitation and extreme values. Measured both the  
100 number of hours of sunshine and the number of sunny days. On the basis of the obtained data from  
101 the Federal Hydrometeorological Institute as well as the soil conditions in BiH and the Tuzla Canton  
102 area, the state and possibilities of ecological production of plums in BiH as well as the orientational  
103 standards of fertilization of plums in organic farming are taken into consideration in the principles of  
104 ecological agriculture.

## 106 3. RESULTS AND DISCUSSION

### 107 3.1. Temperature

108 During deep winter hibernation plum tolerate absolute minimum temperatures even below -30 ° C. In  
109 the opening phase of the flower buds, the flowers can be damaged at -1 ° C to -5 ° C, and in the full  
110 flowering phase, frost is expected at temperatures from -0.5 ° C to -2.2 ° C. As with other fruiting  
111 species, small fruits are still sensitive, and they suffer at temperatures of -0.5 ° C to -2 ° C. The quality  
112 of the fruit most depends on the average temperature during June, July and August. If the average  
113 temperatures are 18 - 20 ° C these months then these areas are excellent for the production of plums  
114 (Ivo Krpina et al., 2004). Average annual air temperatures in 2017 were considerably above the  
115 standard normal value (1961-1990) throughout the observed territory of Bosnia and Herzegovina. The  
116 average air temperatures ranged between 10.4 ° C in Bugojno and 16.0 ° C in Mostar, on the  
117 mountains between 1.8 ° C on Bjelašnica and 8.2 ° C on Ivan Sedlo. Temperature deviations ranged  
118 from 0.6 ° C to Bjelašnica to 2.1 ° C in Gradačac (Table 1.). The spatial distribution of average annual  
119 temperatures in Bosnia and Herzegovina is shown in Figure 1. As for temperature conditions in  
120 Bosnia and Herzegovina, we can conclude that the average temperatures do not indicate bigger

121 problems for plums or deviations of 0.6 ° C on Bjelašnica to 2, 5 ° C in Gradacac are not a major  
 122 problem in the plum production.  
 123




Figure 1. Spatial distribution of average air temperature

124  
 125  
 126  
 127  
 128  
 129

Table 1. The deviation of the average annual air temperature related to the climatic standard normal and the associated percentiles

STATION	ANOMALY	PERCENTILES	STATION	ANOMALY	PERCENTILES
BIHAĆ	1.8	100	Livno	1.3	100
BJELAŠNICA	0.6	91	Mostar	1.5	100
BUGOJNO	1.6	100	Neum	///	///
DRVAR	1.4	100	Sanski Most	1.6	100
GRADAČAC	2.1	100	Sarajevo	1.4	100
IVAN SEDLO	1.0	99	Stolac	1.2	100
JAJCE	1.1	100	Tuzla	1.5	100
KALESIJA	1.5	100	Zenica	1.7	100

130  
 131  
 132  
 133  
 134  
 135  
 136  
 137  
 138  
 139  
 140  
 141  
 142  
 143  
 144  
 145  
 146  
 147  
 148

### 3.2. Precipitation

For the cultivation of plums, the annual amount of precipitation, as well as their distribution during the year, are a very important factor. For normal growth, development and fruiting of the plum it is necessary that the soil be sufficiently humid during the entire vegetation period. Most plum varieties are best managed in regions with an annual rainfall of 700 to 1000 mm (with a fall of 350-600 mm in the vegetation period) and relative humidity of 75-85% (Mišić, 2006). The critical boundary for plum cultivation is 600 mm precipitation per year. However, the amount of this quantity depends on several factors, primarily on the layout of precipitation, soil properties, temperatures in the course of vegetation, etc. (Ivo Krpina et al., Voćarstvo, Zagreb, 2004). In BiH, in the so-called plum area (areas where there is a tradition of plum cultivation in which favorable agronomic conditions for plum cultivation such as Gradacac, Čelić, Srebrenik, etc.) fall more than 800 mm of precipitation. The analysis of annual rainfall amounts expressed in % of the average values shows that in 2017, deviations in relation to the perennial average (1961-1990) ranged between 83.7% in Mostar and 116.2% in Bihać. Total rainfall in 2017 was in the range of 825.4 mm at the meteorological station Gradacac to 1522.6 mm at the meteorological station Bihać (Table 2).

Table 2. The deviation of the average precipitation sum related to the climatic standard normal and the associated percentiles

STATION	ANOMALY	PERCENTILES	STATION	ANOMALY	PERCENTILES
BIHAĆ	116.2	89	Livno	92.2	33
BJELAŠNICA	108.7	58	Mostar	83.7	17

BUGOJNO	110.1	74	Neum	///	///
DRVAR	94.2	38	Sanski Most	107.5	70
GRADAČAC	94.3	32	Sarajevo	100.6	47
IVAN SEDLO	94.8	38	Stolac	80.0	12
JAJCE	107.8	60	Tuzla	113.1	94
KALESIJA	139.5	90	Zenica	113.0	87

149

150

151 December and September were the months with the highest rainfall in 2017, while August is the  
 152 month with the smallest amount. At the meteorological station Mostar in August, 4.8 mm of  
 153 precipitation was measured, while in Bihać in September it was measured 335.0 mm, which is the  
 154 largest monthly sum in 2017 at one station. In the inhabited areas, the highest height of the snow  
 155 cover was measured in Bihać 54 cm (18.01.), While in the mountainous areas (Bjelašnica) 113 cm  
 156 (11.03.) Was recorded.

156

157

158 The number of precipitation days  $\geq 1.0$  mm in 2017 ranged from 76 in Mostar to 131 days at the Ivan  
 159 Sedlo meteorological station. Most days with rainfall  $\geq 1.0$  mm was registered in December and April  
 160 and at least in August. Most days with precipitation  $\geq 10.0$  mm were recorded at the weather station  
 161 Ivan Sedlo (48), and at least at the meteorological station Gradačac (26). In September, most days  
 162 were recorded with precipitation  $\geq 10.0$  mm, and in August on most station there were no precipitation  
 163 days  $\geq 10.0$  mm. As for Tuzla, there were 96 days with precipitation  $\geq 1.0$  mm, with precipitation  $\geq 10$   
 164 mm 35 days and precipitation  $\geq 20$  mm for 16 days. (Federal Hydrometeorological Institute of BiH)  
 165 (Table 3).

164

165

166

167

168

**Table 3. Number of days with characteristic rainfall, snow cover and maximum snow cover height**

169

170

STATION	$\geq 1.0$ MM	$\geq 10.0$ MM	$\geq 20.0$ MM	NUMBER OF DAYS > 1 CM	MAX HEIGHT
	2017. number of days	2017. number of days	2017. number of days	2017. number of days	cm
SARAJEVO	113	30	15	70	39
MOSTAR	76	40	22	0	0
BIHAĆ	110	45	26	45	54
TUZLA	96	35	16	49	26
BJELAŠNICA	128	37	17	211	113

171

172

173

174

### 3.3. Insolation

175

176

177


178

179

180

181

The number of hours of sunshine (insolation) in 2017 was higher in all analyzed stations compared to the thirty year average (1961-1990). Less hours of sunshine (1986) were recorded at the meteorological station Bjelašnica, which is 342 hours more sunshine than the perennial average. Most hours (2694) were recorded at the meteorological station Mostar, or 340 hours more sunshine from the perennial averages (Figure 2). The largest deviation was recorded at the meteorological station Bihać, where during 2017 there were 590 hours of sunshine more than average. The lowest deviation of the sunshine relative to the perennial average was recorded in Mostar (Table 4).



182  
183 **Figure 2. Spatial distribution of insolation (h) in 2017**

184 The most pronounced monthly positive deviation of the sunshine expressed in% of the average  
185 values relative to the reference series (1961-1990) was registered in January at the meteorological  
186 station Sarajevo. The number of hours of sunshine at the meteorological stations in Sarajevo was  
187 198% higher than the average, and 113 hours of sunshine were recorded. The most pronounced  
188 monthly negative deviation of the sunshine was recorded in December at the meteorological station  
189 Bjelašnica. Bjelašnica recorded 60 hours of sunshine, which amounts to about 72% of the average  
190 value.

191 **Table 4. Number of hours of sunshine in 2017**

STATION	NUMBER OF HOURS OF SUNSHINE COMPARED TO PERENNIAL AVERAGES
MOSTAR	340 hours more compared to perennial averages
BIHAĆ	590 hours more compared to perennial averages
BJELAŠNICA	342 hours more compared to perennial averages

192  
193  
194 **3.4. Necessary amounts and concentrations of nutrients in plum production**

195 For normal life of the plums, as well as other green plants, 17 biogenic elements are necessary  
196 (Nešković et al., 2003). Biogenic macroelements plants are adopt and accumulate in large quantities,  
197 and other biogenic elements (Fe, B, Zn, Mn, Mo, Cu, Cl, Ni), i.e. microelements are used in smaller  
198 quantities. Biogenic macroelements primarily are used as building materials, and biogenic  
199 microelements as biocatalysts in the plant. Plum is supplied with carbon and partly oxygen from the  
200 air, while other biogenic elements come from the soil. However, the above-ground organs of the plant,  
201 especially the leaf, can absorb biogenic elements with the so-called foliar or leaf nutrition. In this way,  
202 plants can provide the necessary nutrients. Plum leaves are the richest mineral substances, followed  
203 by buds and fruits (Table 5 and 6).

204  
205 **Table 5. Quantities of macroelements expressed in% of dry matter in various plum organs**  
206 **(Sarić, 1975. i**  
207 **Childers, 1969.)**  
208

NUTRITION	FRUIT	LEAF	SPROUT	BUDWOOD	TREE AND BRANCHES	ROOT
N %	0,71-1,20	1,80-2,10	0,43	0,97	0,37	0,34
P <sub>2</sub> O <sub>5</sub> %	0,22-0,30	0,33-0,60	0,10	0,27	0,09	0,10

K <sub>2</sub> O %	1,41-2,27	1,80-3,00	0,23	0,43	0,21	0,21
CAO %	0,11	2,80-5,60	0,84	2,31	0,63	0,59

209  
210  
211  
212  
213

**Table 6. Optimal quantities of biogenic elements in normal plum leaves (in mg per kg of dry substance) in August (Childers, 1966.; Westwood, 1978.; Faust, 1989.)**

ELEMENT	CHILDERS (1966.)	WESTWOOD (1978.)	FAUST (1989.)
N	25.300-30.000	22.000	25.000
K	8.400-46.600	14.000	25.000
CA	27.800-51.600	10.000	25.000
P	550-770	1.200	2.000
MG	26.900-29.400	2.400	4.000
S	2.200-4.700	-	-
FE	-	50	120
MN	53-93	25	50
ZN	-	18	30
B	33	35	35
CU	7-10	4	10

214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226

The amount of biogenic elements in the normal plum leaves (Table 6) ranges from 4 mg for copper to 51,600 mg per kilogram of dry calcium. For normal growth and fruiting, the same amount of copper is as important as 12,900 times more calcium. The plum root receives mineral matter dissolved in water in the form of ions (anions and cations) by diffusion, osmosis and contact ion exchange. The absorption and content of mineral substances are influenced by: the selective permeability of protoplasm in the absorption root zone, temperature, pH, antagonism and synergism between anion and cation, aeration, stage of individual plant development, phenofase in the annual plum cycle, and the organ of the plant receiving mineral substances. When the concentration of nutrients is above the level of good supply, there is no need for fertilization to increase availability, as economically justified results will not be obtained. In that sense it is very important to interpret the results of soil analysis well.

### 3.5. Organic fertilization and soil

227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253

Some studies (Blanco-Canqui et al., 2017, Hondebrink et al., 2017, Shah et al., 2017) show that eco-systems of plant production function normally or in some aspects even better than conventional systems involving the use herbicides and other agrochemicals. The necessary soil stability in ecological agroecosystems is improved by the addition of organic matter through organic manure, compost, green manure, organic by-products on organic farms and well-designed crop rotation that largely involve the sowing of cover crops and shamrock. The above mentioned agro-technical measures in practice mitigate the negative effects of soil treatment on the structure and quantity of organic matter, i.e. humus. Yagodin (1984) states that the continuous application of solid organic fertilizers improves the chemical properties of soil, increases the humus content, microbiological activity, and there is also a positive influence on water-air relations in the soil. Furthermore, the capacity to replace the bases and the saturation level of the adsorption complex with bases (Ca, Mg, K) increases, while the acidity slowly decreases, as is the amount of mobile aluminum, iron and manganese. Butorac and sar. (1988) point out that the application of organic fertilizers is one of the ways of removing the harmful effect of excessive acidity of the soil, primarily thanks to the mechanism of antitoxic activity. This mechanism is activated to a greater degree only during the many years of intensive application of organic fertilizers, when organic acids contained in organic manure block aluminum, copper and other elements in the form of heavily soluble organic compounds, and organic acids form chelates with metals. Along with the mentioned effect of organic acids on reducing the toxicity of aluminum, manganese and other elements, and other substances from organic fertilizers, such as antagonistic elements (P, Ca, Mg, S, Si), they also have an impact on the suppression of toxicity. Continuous application of organic fertilizers allowed in organic agricultural production increases the amount of soil organic matter, reduces erosion, improves soil infiltration and soil aeration, improves the soil's biological activity during the decomposition of organic soil, and promotes long-term yield of cultures acting residually to release nutrients in the soil Hati and Bandyopadhyay, 2011).

### 3.6. Plan for fertilizing a young plantation to full fertility

Young orchards are plantation at the stage of intense growth and fruit development. This is the stage from planting till it develops the basic skeletal branches, with the characteristics of the intended breeding form. This phase is shorter in intensive than extensive plantations. Young fruit trees should be regularly fertilized in order to foster faster growth and development. The correct fertilization depends largely on the rate of growth and the development of fruit trees. In the first and second year fruit is fertilized individually, and the fertilization zone around the fruit trees should be slightly wider than the crochets (Kantoci, 2012). The result plums fertilization depends predominantly on the biogenic element or other factor that is at a minimum (Mišić, 2006). The selection of fertilizers depends on the soil, climate and weather conditions, variety and plum substrate, crown shape and breeding system, growth and yield, nutrient consumption of plums, quality, composition and physiological activity of fertilizers (Mišić, 2006). When in the agromelioration all biogenic elements are brought to the proper level, the fertilization of plum potassium and phosphorus may be absent during the first few years of the care of a young plantation (Mišić, 2006). Otherwise, fertilization with nitrogen, phosphorus and potassium is carried out from the first year. The orientation norms of certain nutrients during the first three years of young sponge care are shown in Table 5 (Mišić, 2006).

**Table 7. Orientation norms of individual nutrients (in g per tree) for fertilization of young plums**

YEAR PLANTATION	AFTER N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
FIRST	30	10	40
SECOND	60	20	80
THIRD	90	30	120

Depending on the soil properties, a greater or lesser part of nutrients from the fertilizer binds (immobilized) in forms that are hard to reach by fruit trees. For example, easily available phosphate (monoclonation phosphate) is often bind to iron or aluminum in the soil in a heavily accessible or inaccessible form (tetra and penta phosphate iron or aluminum). Nitrogen in the soil is easily subjected to rinsing processes, resulting in large losses if fertilization is not done gradually on several occasions (Kantoci, 2012).

Fertilization for medium high yield:

Average nutrient requirements for medium high yield are nitrogen (N) 90 kg / ha, phosphorus (P<sub>2</sub>O<sub>5</sub>) 50 kg / ha and potassium (K<sub>2</sub>O) 140 kg / ha. In the autumn after harvest, it is necessary to give from 390 kg / ha of NPK 6-18-36 (or 1 kg / tree), in March 100 kg / ha urea (or 0.4 kg / tree), and in May 60 kg / ha of KAN (or 0.3 kg / tree), (Table 8).

**Table 8. Average nutrient requirements for medium high yield**

PERIOD OF YEAR	KIND AND QUANTITY OF FERTILIZER
AUTUMN	NPK 6-18-36 390 kg/ha
MARCH	UREA 100 kg/ha
MAY	KAN 60 kg/ha

Foliar nutrition should be carried out on two occasions (or more in case of a more expressed deficit of microelements):

- before blooming: 0.5% solution foliar preparation on Boric (B) base
- after blooming: 1 to 2% solution of foliar preparation with macro and micro elements (Pavlović, 2016).

Fertilization for high yield:

Average nutrient requirements for high yield are nitrogen (N) 160 kg / ha, phosphorus (P<sub>2</sub>O<sub>5</sub>) 90 kg / ha and potassium (K<sub>2</sub>O) 250 kg / ha. In the autumn after the harvest, it is necessary to give 680 kg / ha NPK 6-18-36 (or 1.7 kg / tree), in March 180 kg / ha urea (or 0.5 kg / tree) and in May with 100 kg / ha KAN (0.4 kg / tree). (Table 9.)

**Table 9. Average nutrient requirements for high yield**

PERIOD OF YEAR	KIND AND QUANTITY OF FERTILIZER
AUTUMN	NPK 6-18-36 680 kg/ha
MARCH	UREA 180 kg/ha
MAY	KAN 100 kg/ha

306

307 On lime soil it is recommended that in the autumn after harvest you give 830 kg / ha NPK 5-20-30S  
 308 (or 2.1 kg / tree), in March 175 kg / ha urea (0.5 kg / tree), and in May 100 kg / ha of AN (0.3 kg /  
 309 tree).

310 For organic fertilizers can be used compost, vermicompost, slurry, manure, rigid cow manure, solid  
 311 chicken manure, rigid horse manure, rigid pig manure and chicken dry manure . Each of them has  
 312 different ranges of nutrient concentration. Range of nutrient concentrations are different due to the  
 313 different proportions of the litter of these solid and liquid excretions. The litter has a lower nitrogen  
 314 content, but absorbs a large amount of liquid secretions and increases the concentration of N and K,  
 315 which are mainly excreted in the urine. The extracellular extracts, however, contain mostly  
 316 phosphorus, which is why the low concentration of phosphorus in the slurries (Lončarić, 2015). Since  
 317 the plum requires a large amount of nutrients, and the use of mineral fertilizers is not allowed in  
 318 organic production, alternative solutions are used. Dry chicken manure is imposed as one of favorable  
 319 solutions for soil with a lower level of available phosphorus because in 1 ton of dry chicken manure  
 320 there is 30-38 kg N, 20-30 kg P<sub>2</sub>O<sub>5</sub> and 20-22 kg K<sub>2</sub>O (Lončarić Z. , 2015). In solid chicken manure  
 321 (in fresh state) the concentrations are slightly lower: 12-20 kg / t N, 12-14 kg / t P<sub>2</sub>O<sub>5</sub> and 5-7 kg / t  
 322 K<sub>2</sub>O (Lončarić, 2015). On the other hand, for soil with lower potassium level is more suitable for  
 323 organic fertilizer mixtures containing liquid manure (liquid manure) and plant waste (green mass,  
 324 straw, corn) that are richer in potassium (eg composted mixtures of plant waste and manure).  
 325

326

#### 4. CONCLUSION

327 The content of organic matter is an important factor in fertilizing plums. On the one hand, it provides  
 328 the necessary nutrients, and now on the other hand it affects the physical and chemical properties of  
 329 the soil. Organic matter of soil (humus) is the most important factor of soil elasticity. The amount of  
 330 organic matter directly affects the fertilization needs. In the first year of breeding, in the basic  
 331 fertilization of fruit with a thick assembly, it is fertilized by one fruit with 0.15 to 0.2 kg with some of the  
 332 complex fertilizers (depending on the availability of P and K in soil) as NPK 7-20-30, NPK 6-18-36,  
 333 NPK (SO<sub>3</sub>) 7-14-21 (24) or 0.1 kg NPK 8-26-26, NPK 10-30-20, NPK (MgO, SO<sub>3</sub>) 7-14-21 (2 , 18). In  
 334 the second year of cultivation fertilization is done at the same time and with the same types of  
 335 fertilizers, with the amount increasing by 30 to 50%. In terms of nutrition in the fullfruiting phase, plum  
 336 is significantly different from other types of fruit trees, because in addition to nitrogen, it requires large  
 337 amounts of potassium in the soil. Fertilization is best done on the basis of chemical soil analyzes,  
 338 nutritional status of plums determined by the analysis of leaves and the expected yield. The specificity  
 339 of fertilization in organic agriculture is necessarily omitted from water-soluble mineral fertilizers, and  
 340 the necessary nutrients are introduced into the soil by organic fertilizers and allowed additives  
 341 (organic and / or mineral origin) from the list of permitted fertilizers in organic production. Plum  
 342 annually yields 40 to 60 kg / ha N. Orientation of this amount of available nitrogen in the soil is done  
 343 by organic fertilizers with different nitrogen concentration ranges. It is used cow, chicken, pig, horse  
 344 manure and dry chicken manure with the highest nitrogen content (30 to 38 kg / t N). Phosphorus and  
 345 potassium are provided partly from the already mentioned organic fertilizers, and partly by the use of  
 346 natural minerals or by-products such as phosphorites, Tomas phosphate, potassium sulphate,  
 347 potassium chloride. Microelements are provided by fertilizing organic fertilizers, as well as micro-  
 348 fertilizers and preparations that are used by foliar or soil application in organic farming.  
 349

350

351

#### 5. LITERATURE

352 Alföldi et al., (2002.). In: El-Hage Scialabba, N.; Hattam, C. (Eds) Organic agriculture, environment  
 353 and food security. Environment and Natural Resources Management Series, N°4. FAO, Rome  
 354 Anastasijevic, V., Sisakovic, N.: Composting of urban green spaces' organic waste as a must of  
 355 sustainable urban greenspace management. Proceeding of the 8th Symposium of flora of  
 356 Southeastern Serbia and Neighbouring Regions, Nis, 2005.  
 357 Benaković D. (2016.). Principi utvrđivanje optimalne gnojidbe i mjera kondicioniranja tla  
 358 Blanco-Canqui, H., Francis, C. A., Galusha, T. D. (2017.). Does organic farming accumulate carbon in  
 359 deeper soil profiles in the long term?. Geoderma, 288, 213-221.  
 360 Butorac, A. (1999.). Opća agronomija, Zagreb

- 361 Ćosić, T., Mesić M. (2008). Fertilizers and fertilizers, Faculty of Agriculture, University of Zagreb  
362 Kantoci, D. (2012). Fertilizing fruit trees. Bulletin of the Protection of Bilja, 35 (5), 40-45.  
363 Kastori, R. (1983). The Role of Elements in Plant Nutrition. Matica srpska, Novi Sad  
364 Kordic A. (2014). Application of visual diagnostics and analysis of plant material in horticulture.  
365 Krpina, I. (2004). Fruit production.  
366 Kurtović, M. et al. (2008). Integral production of bone fruits, Faculty of Agriculture in Sarajevo,  
367 Sarajevo  
368 Little T, Frost D. (2008). Organic fruit and vegetable production  
369 Lončarić, Z. (2010). Internal materials for the module "Fertilization in organic agriculture" and  
370 presentations from lectures. Osijek.  
371 Lončarić, Z., Parađiković, N., Popović, B., Lončarić, R., Kanisek, J. (2015). Fertilization of fertilizers,  
372 organic fertilizers and composting. Faculty of Agriculture, University of Osijek.  
373 Lončarić, Z., Popović, B., Ivezić, V., Karalić, K., Manojlović, M., Čabovski, R., Lončarić, R. (2014).  
374 Mineral and organic fertilization on family farms in the border region of Croatia and Serbia.  
375 Proceedings of the 49th Croatian and 9th International Symposium of Agronomists. Marić, S.,  
376 Lončarić, Z., editor (s). Faculty of Agriculture, University of Osijek, Osijek. 77-81.  
377 Marschner, H. (1995). Mineral nutrition of higher plants. Academic Press Ltd.  
378 Mišić, P. (2006). Plum.  
379 Mitrović Olga, Zlatković Branislav, Kandić Miodrag, Gavrilović-Damnjanović Jelica, Popović Branko,  
380 Milinković Vladan: Fruit growing, Technological characteristics of fruits of some new varieties of plums  
381 for drying, vol. 43, no. 167-168 (2009) 101-106  
382 Mratinić, E. (2004). Plum. Partenon, Belgrade  
383 Shah, A., Askaard, M., Rasmussen, I. A., Jimenez, E. M. C., Olesen, J. E. (2017). Productivity of  
384 organic and conventional arable cropping systems in long-term experiments in Denmark. European  
385 Journal of Agronomy, 90, 12-22.  
386 Vukadinović V., Bertić B. (2013). Fertilization philosophy: everything you need to know about  
387 fertilization.  
388 Vukadinović, V. and Bertić, B. (1989). Practicum from agrochemistry and nutrition of plants, Faculty of  
389 Agriculture in Osijek, Osijek.  
390 Vukadinović, V. and Lončarić, Z. (1998). Plant nutrition, Faculty of Agriculture in Osijek, Osijek  
391 Vukadinović, V. and Lončarić, Z. (1998). Plant nutrition. Faculty of Agriculture in Osijek.  
392 Yagodin, B.A. (1984). Agricultural chemistry, Mir Publishers, Moscow.  
393 Znaor, D. (1996). Organic farming. Nakladni zavod Globus. Zagreb.