

**Capacity for parasitism of *Trichogramma* spp. in  
tomato fruit borer under different temperatures**

**Abstract:**

---

The parasitoid in eggs of *Trichogramma* genre is the most studied in the world, being bred widely and used for flooding releases. This study aimed to evaluate the capacity for parasitism of *Trichogramma galloi* Zucchi, 1988 (Hymenoptera: Trichogrammatidae) in *Neoleucinodes elegantalis* (Guenée) (Lepidoptera: Crambidae) eggs at different temperatures. The experiment was developed at the Nucleus for Scientific and Technological Development in Phytosanitary Management (NUDEMAFI) in which the daily and accumulated biological parameters were assessed, as well as total parasitized eggs by female, sexual ratio (number of females/number of males + females), viability of the eggs (number of eggs with orifice/number of parasitized eggs x 100) and number of individuals per egg at temperatures 18, 21, 24, 27 and 30 °C. The eggs of the fruit borer were offered daily to each *T. galloi* female at each temperature until the death of the parasitoid could be verified. The larger number of parasitized eggs was found in the first 24h, at temperatures 24 and 27 °C with 17 parasitized eggs. The accumulated parasitism in *N. elegantalis* eggs reached 80% of total parasitized eggs for each thermal range (18, 21, 24, 27 and 30 °C) on the 2nd, 3rd, 3rd, 1st and 2nd days respectively. The ideal parasitism conditions for this lineage vary between 24 and 27°C. Therefore, it is conclude that the studied *T. galloi* lineage has the adequate biological parameters in *N. elegantalis* eggs, demonstrating promise in phytosanitary management of this pest.

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

*Keywords: Egg parasitoid, Neoleucinodes elegantalis, Phytosanitary management, Tomato fruit, Oviposition, Trichogrammatidae,*

**1. INTRODUCTION**

Among the pests that attack the tomato culture, the tomato fruit borer *Neoleucinodes elegantalis* (Guenée) (Lepidoptera: Crambidae) is considered one of the main pests for its preference for this culture and the damages caused directly in the fruit, making them inadequate for consumption and industrial processing, with significant loss (Miranda et al., 2005; Picanço et al., 2007; Fornazier et al., 2010; Pratisoli, 2015; Carvalho et al.; 2017; Silva et al., 2017; Moraes & Foerster, 2015).

23 Since this is a culture of high risk, with high intensity for pest attack, it is important to  
24 implement practices that aim to manage these pests. Among management methods, biological  
25 control is a viable technique especially when using parasitoids of the *Trichogramma* genre  
26 since it acts on the eggs avoiding the larvae to penetrate the fruit, reducing the loss caused by  
27 caterpillar feeding in its interior (Plaza et al., 1992, Oliveira et al., 2017).

28 The egg parasitoid *Trichogramma* is the most often studied in the world, being greatly bred  
29 and used in flooding releases (Hassan, 1997; Davies et al., 2009). The advantage of its use is  
30 its capacity to control pests from different cultures. Moreover, they are highly specialized and  
31 efficient (Haji et al., 2002; Wang et al., 2007; Wang et al, 2018; Arruda et al. 2014).

32 In Brazil, studies aiming at the use of *Trichogramma* were initiated over 30 years ago, with  
33 excellent results in many cultures, more recently *Trichogramma galloi* Zucchi, 1988  
34 (Hymenoptera: Trichogrammatidae) being the most often used species, released in about  
35 500,000 hectares every year in sugar cane to control the cane borer *Diatraea saccharalis*  
36 *Fabricius, 1794* (Lepidoptera: Crambidae) (Parra, 2010, Arruda et al. 2014, Zago et al. 2007,  
37 Oliveira et al., 2017).

38 For *N. elegantalis* studies have demonstrated its potential to use *Trichogramma* in its  
39 management (Blackmer et al., 2001). Nonetheless, other studies must be conducted for better  
40 reliability on the use of these parasitoids in the management of *N. elegantalis*. These studies  
41 must involve the efficacy of the species, biological characteristics, thermal demands, ideal  
42 release numbers and dispersion capacity (Oliveira et al., 2017).

43 Thus, the aim of this study was to evaluate the potential for parasitism of *T. galloi* in *N.*  
44 *elegantalis* eggs at different temperatures.

45

## 46 **2. MATERIAL AND METHODS**

47

48 The experiment was conducted in the Entomology Department at the Nucleus for Scientific  
49 and Technological Development in Phytosanitary Management (NUDEMAFI) at the Agronomic  
50 Sciences Center at the Federal University of Espírito Santo (CCAUE-UFES), Alegre, ES  
51 (Brazil). A lineage Tg1 of *T. galloi* species was used, provide by BUG Biological Agents.

52

### 53 **Breeding of the alternative host *Anagasta kuehniella* Zeller (Lepidoptera:Pyralidae)**

54 The alternative host *Anagasta kuehniella* Zeller (Lepidoptera: Pyralidae) was bred in  
55 homogenized diet of whole wheat flour (60%), corn (37%) and beer yeast (3%). The diet was  
56 disposed into plastic bins (30 x 25 x 10 cm) with corrugated cardboard strips (25 x 2 cm) on  
57 the inside, with the host eggs randomly selected for the diet. The adults were collected daily,  
58 with an adapted vacuum and transferred into PVC tubes (150 mm diameter x 25 cm height)  
59 with nylon strips folded in its interior for oviposition (Pratissoli, 2010).

60

61 **Breeding of *T. galloi***

62 For the maintenance of parasitoids, *Anagasta kuehniella* eggs were inactivated in germicide  
63 lamp during 50 minutes and fixated in rectangles of sky blue cardboard (8.0 x 2.0 cm), with  
64 arabic gum diluted to 20%. Those cards were inserted in glass tubes (8.5 x 2.4 cm),  
65 containing adult parasitoids recently emerged. Furthermore, the tubes were sealed with PVC  
66 plastic film to avoid parasitoid escape. The cards were kept in the tubes for 24 hours and later  
67 stored in clean glass tubes (9 x 3 cm) in an acclimatized room at  $25 \pm 1$  °C, relative humidity  $70$   
68  $\pm 10\%$  and photoperiod of 14h.

69

70 **Breeding of *N. elegantalis***

71 Breeding of pests was conducted in an acclimatized room ( $25 \pm 2$  °C, RH  $70 \pm 10\%$  and photoperiod  
72 phase of 12h). Adults were kept in acrylic cages and fed with a solution of 10% honey. For  
73 oviposition, tomato fruit from the F1 wire were conditioned in the cages. Daily, the eggs were  
74 removed from the tomato fruit and distributed in african eggplant fruit (mean 5 eggs/fruit)  
75 which remained in plastic containers covered in non-woven fabric serving as places for  
76 pupation of caterpillars. Once this phase is finished, pupae were transferred into plastic  
77 containers or Petri dishes and stored in acclimatized chambers in the above mentioned  
78 conditions until adults emerge, then again taken to the acrylic cages.

79

80 **Capacity of parasitism**

81 *N. elegantalis* eggs with up to 12h of age were collected from the tomato fruit with the help of a  
82 scalpel and glued to sky blue cardboard (0.5 x 2.0 cm) with a brush and arabic gum at 20%.  
83 For each temperature of the study, 20 recently emerged females were isolated in *eppendorf*  
84 tubes (2.0 ml), containing drops of honey for feeding and sealed with the tubes' own lid. The  
85 cards with the 20 tomato fruit borer eggs were offered daily to each one of the *T. galloi*  
86 females at each temperature (18, 21, 24, 27 and 30 °C) until the death of the parasitoid was  
87 confirmed. The cards removed daily were identified and bagged (23.0 x 4.0 cm) and kept at its  
88 respective temperatures.

89 The following biological parameters were assessed: daily and accumulated parasitism, total  
90 parasitized eggs per female, sexual ratio (number of females/number of males + number of  
91 females), viability of the eggs (number of eggs with orifice/number of parasitized eggs x 100)  
92 and number of individuals per egg at different temperatures.

93 The experiment was conducted with a completely casual design, with five treatments  
94 (temperatures) and 20 repetitions, each repetition represented by a *T. Galloi* female. For data  
95 analysis, a regression with test F was used at 5% probability level.

96

97 **3. RESULTS**

98

99 Daily parasitism decreased at all temperatures with the advance of age of the *T. galloi* female.

100 At the different studied temperatures, higher rates of parasitism occurred in the first 24 hours,  
101 presenting between 13 and 17 parasitized eggs. The higher rates of parasitism were at  
102 temperatures 24 °C and 27 °C with around 17 eggs parasitized (Figure 1).

103 In terms of longevity of the females, it was noted that lower temperatures (18 °C to 24°C)  
104 females were able to live longer due to a reduction in physiological activity of females when  
105 exposed to lower temperatures. At higher temperatures (27 °C and 30 °C), there is higher  
106 energy expenditure and, consequently, females lived for a shorter period of time (Figure 1).

107 The parasitism period for *T. galloi* females was increased in the thermal range of 18 to 24°C  
108 (5, 7 and 8 days) and in the 27 to 30°C range there was a decrease (5 and 4 days). Therefore,  
109 the ideal conditions for survival of *T. galloi* vary between 24 °C and 27 °C where better  
110 performance was observed.

111 Accumulated parasitism in *N. Elegantalis* eggs in the studied thermal range reached 80% of  
112 total parasitized eggs in a maximum of three days. At extreme temperatures (18 and 30°C)  
113 this condition was reached in two days. At milder temperatures (21 and 24°C) the  
114 accumulated parasitism reached 80% in three days and at 27 °C was reached in the first day  
115 of parasitism (Figure 1). Due to the 80% parasitism, it was noted that the potential for  
116 parasitism of this lineage occurs within the first days of parasitism, independent from  
117 temperature.

118 For this lineage, the 24°C temperature highlights the total amount of eggs parasitized per  
119 female reaching an average of 30 eggs. In terms of viability, as the temperature increased  
120 there was reduction in viability, coming to 50% at 30°C. For sexual ratio, it was observed that  
121 when temperature increased there was a higher number of males in the population, but the  
122 number of individuals per egg were constant (Figure 2).


123

124

125

126

127



129 **Figure 1** – Daily and accumulated parasitism for *T. galloi* in *N. elegantalis* eggs at different  
 130 temperatures.

131


132

133

134

135

136



158

159 **Figure 2** – Total *N. elegantalis* eggs parasitized by *T. galloi*, viability of parasitism, sexual ratio and  
 160 of descendents per eggs at different temperatures.

161

162 **4. DISCUSSION**

163 We verified that the temperature interferes in the potential for parasitism and biological  
 164 characteristics of *T. galloi*.

165 Among the main factors affecting biological characteristics of species in the *Trichogramma*  
 166 genre, temperature is highlighted since with its increase, there is lower performance by  
 167 females causing metabolism to increase and, therefore, reducing parasitism (Hansen &  
 168 Jensen, 2002; Pratisoli et al., 2003; Rahimi-Kaldehy et al 2018, Pratisoli et al. 2004). This  
 169 was demonstrated in the present study since extreme temperatures showed a decrease in  
 170 parasitism with only 11 eggs parasitized on average at 18°C, 10 eggs at 21°C and 14 eggs  
 171 parasitized at 30°C (Figure 2).

172 The potential for parasitism in the first days may be directly connected to the instinct of animal  
173 preservation once all species in the *Trichogramma* genre present this behavior (Pratissoli at  
174 al., 2004; Zago et al.,2007; Paes et al., 2018). This behavior may be related to the parasitism  
175 of 80% of the eggs as studies have confirmed this rate to be, in most cases, in the first few  
176 days of life in females (Pratissoli at al., 2004; Zago et al.,2007).

177 Parasitism period may vary according to temperature and within each temperature. This fact  
178 may be related to the capacity of adaptability in each species and/or lineage of *Trichogramma*  
179 to the habitat in which it was collected (Hansen & Jensen, 2002; Pratissoli at al., 2004; Zago et  
180 al.,2007; Arruda et al., 2014; Paes et al., 2018).

181 Accumulated parasitism is another factor that may be related to the capacity of adaptability of  
182 each species and/or lineage of *Trichogramma* to the habitat in which it was collected since the  
183 necessary time to reach total percentage is variable (Pratissoli at al., 2004; Zago et al.,2007).  
184 The range of temperature in which species and/or lineage of *Trichogramma* present their  
185 higher potential for parasitism (number of parasitized eggs) is between 24 and 27°C (Hansen  
186 & Jensen, 2002; Pratissoli at al., 2004; Zago et al.,2007; Arruda et al., 2014).

187 Through viability there seems to also be direct interference from temperature. It is possible to  
188 verify that there is an inverse relationship between the percentage of emergence of  
189 descendents and the increase in temperature. However, this was not found in any other  
190 studies.

191 The variation in sexual ratio has been reported as influenced specially by temperature  
192 (Vinson, 1997, Rahimi-Kaldehy, et al, 2018). This was verified in extreme temperatures once  
193 humidity, female age, and host were constant for all temperatures.

194 On the number of descendents per egg, it was verified that it was constant, that is, one  
195 individual per egg independent from temperature. The variation in this factor is directly related  
196 to nutritional and morphological characteristics of the egg such as size, shape, thickness,  
197 corion stiffness and lay behavior (Hassan, 1997; Bakthavatsalam, et al., 2013, Paes et al.,  
198 2018).

199 It was verified that the lineage studied for *T. galloi* holds true the adequate biological  
200 parameters for parasitism in *N. elegantalis* eggs, proving to be promising in phytosanitary  
201 management of this pest.

202

203

204

## 5. CONCLUSION

205 The studied *T. galloi* strain presents the appropriate biological parameters for parasitism in *N.*  
206 *elegantalis* eggs, showing promise in the phytosanitary management of this pest.

207

208

209

210

## COMPETING INTERESTS

211

“Authors have declared that no competing interests exist”

212

213

214

## REFERENCES

215

Arruda LA, Leite R C, Tonquelski GV, Leal AF, Borges FSP, Rodrigues LA. Eficiência do

216

parasitismo de três espécies de *Trichogramma* (*T. galloi*, *T. atopovirilia* e *T. bruni*) sobre ovos

217

da praga *Datraea saccharalis*. Global Science Technology. 2014;07:67 – 75.

218

Bakthavatsalam N, Tandon PL, Bhagat D. Trichogrammatids: Behavioural Ecology In.

219

Sithanantem CR, Ballal, Jajali SK, Bakthavatsalam N. (Ed). Biological Control of Insects Pests

220

using Egg Parasitoids. 2013;77-103. New Delhi-India: Springer

221

Blackmer JL, Eiras AE, Souza CLM. Oviposition preference of *Neoleucinodes elegantalis*

222

(Guenee) (Lepidoptera: Crambidae) and rates of parasitism by *Trichogramma pretiosum* Riley

223

(Hymenoptera: Trichogrammatidae) on *Lycopersicon esculentum* in São José de Ubá, RJ,

224

Brazil. Neotropical Entomology. 2001;30:89-95.

225

Carvalho GS, Silva LB, Reis SS, Veras MS, Carneiro E, Almeida MLS, Silva AF, Lopes GN.

226

Biological parameters and thermal requirements of *Trichogramma pretiosum* reared on

227

*Helicoverpa armigera* eggs. Pesquisa agropecuária brasileira. 2017;52:961-968.

228

Davies AP, Pufke US, Zalucki MP. *Trichogramma* (Hymenoptera: Trichogrammatidae)

229

Ecology in a Tropical Bt Transgenic Cotton Cropping System: 18 Sampling to Improve

230

Seasonal Pest Impact Estimates in the Ord River Irrigation Area, Australia. Journal Economic

231

Entomological. 2019;102:1018-1031.

232

Fornazier M, Pratisoli D, Martins D S. Principais pragas da cultura do tomateiro estaqueado

233

na região das montanhas do Espírito Santo. In: Incaper(Ed.). Tomate; 2010;185-226. Vitória:

234

Incaper.

235

Haji FNP, Prezotti L, Carneiro JS, Alencar JA *Trichogramma pretiosum* para controle de

236

pragas no tomateiro industrial, Vol. 1: Controle biológico no Brasil: Parasitoides e predadores

237

(ed. Parra, JRP, Botelho, SM, Ferreira, BSC, Bento JMS) Manole, São Paulo, SP, Brazil,

238

2002, pp. 477-494.

239

Hassan, AS. Seleção de espécies de *Trichogramma* para o uso em programas de controle

240

biológico., Vol. 1: Trichogramma e o Controle Biológico Aplicado (ed. Parra JRP & Zucchi

241

RA) FEALQ, Piracicaba, São Paulo, Brazil, pp. 183- 205. 1997.

242

Hansen LS, Jensen KMV. Effect of Temperature on Parasitism and Host-Feeding of

243

*Trichogramma turkestanica* (Hymenoptera: Trichogrammatidae) on *Ephestia kuehniella*

244

(Lepidoptera: Pyralidae). J. Econ. Entomol. 2002;95:50-56.



245 Miranda MMM, Picanço MC, Zanuncio JC, Bacci L, Silva EM. Impact of integrated pest  
246 management on the population of leafminers, fruit borers, and natural enemies in tomato.  
247 *Ciência Rural*. 2005;35:204-208.

248 Moraes CP, Foerster LA. Thermal requirements, fertility, and number of generations of  
249 *Neoleucinodes elegantalis* (Lepidoptera: Crambidae). *Neotropical Entomology*. 2015;44:338-  
250 344.

251 Oliveira CM, Oliveira JV, Silva Barbosa DR, Breda MO, França SM, Duarte BLR. Biological  
252 parameters and thermal requirements of *Trichogramma pretiosum* of the management of the  
253 tomato fruit borer (Lepidoptera: Crambidae) in tomatoes. *Crop Protection*. 2017;99:39-44.

254 Paes JPP, Lima VLS, Pratisoli D, Carvalho JR, Bueno RCOF. Selection of parasitoids of the  
255 genus *Trichogramma* (Hymenoptera: Trichogrammatidae) and parasitism at different eggs  
256 ages of *Duponchelia fovealis*. *Acta Scientiarum Biological Sciences*. 2018;40:1-9.

257 Parra JRP. Egg parasitoid commercialization in the New World, Vol. 1: Egg parasitoids in  
258 agroecosystems with emphasis on *Trichogramma* (ed. Cònsoli FL, Parra JRP, Zucchi RA)  
259 Springer, Dordrecht, Holland: Springer, 2010;373-388.

260 Picanço MC, Bacci L, Crespo ALB, Miranda MMM, Martins JC. Effect of integrated pest  
261 management practices on tomato production and conservation of natural enemies. *Agricultural  
262 and Forest Entomology*. 2007;9:327-355.

263 Plaza AS, León EM, Fonseca JP, Cruz J. Biology, behavior and natural enemies of  
264 *Neoleucinodes elegantalis*. *Revista Colombiana de Entomologia*. 1992;18:32-37.

265 Pratisoli D, Fornazier MJ, Holtz AM, Gonçalves JR, Chioramital AB, Zago HB. Ocorrência de  
266 *Trichogramma pretiosum* em áreas comerciais de tomate, no Espírito Santo, em regiões de  
267 diferentes altitudes. *Horticultura Brasileira*. 2003;21:73-76.

268 Pratisoli D, Oliveira, HN de, Vieira SMJ, Oliveira RC de, Zago HB. Efeito da disponibilidade  
269 de hospedeiro e de alimento nas características biológicas de *Trichogramma galloi*. *Revista  
270 Brasileira de Entomologia*. 2004;48:101-104.

271 Pratisoli D. Guia ilustrado de pragas da cultura do tomateiro. Alegre, Unicopy, 2015. 45p.

272 Rahimi-Kaldehy S, Ashouri A, Bandani A, Ris N. Abiotic and biotic factors influence  
273 diapause induction in sexual and asexual strains of *Trichogramma brassicae* (Hym:  
274 Trichogrammatidae). *Scientific Reports*. 2018;8:1-6.

275 Silva RS, Kumar L, Shabani F, Silva EM, Galdino TVS, Picanço MC. Spatio-temporal dynamic  
276 climate model for *Neoleucinodes elegantalis* using CLIMEX. *International Journal  
277 Biometeorology*. 2017;61:785-795.

278 Vinson SB. Comportamento de seleção hospedeira de parasitoides de ovos com ênfase na  
279 família Trichogrammatidae, Vol. 1: Trichogramma e controle biológico aplicado (ed. PARRA,  
280 JRP, ZUCCHI RA) FEALQ, Piracicaba, SP, Brazil, 1997. pp. 67-120.

281 Wang Z, He K, Bai S. Use of *Trichogramma* in plant protection achievement, challenge and  
282 opportunity. Entomological Research. 2007;37:1-73.

283 Wang Z, Lui Y, Shi M, Huang J, Chen X. Parasitoids wasps as effective biological control  
284 agents. Science Direct. 2018;17:60345-603457.

285 Zago HBD, Pratissoli D, Barros MGC, Gondim JR. Capacidade de parasitismo de  
286 *Trichogramma pratissoli* Querino & Zucchi (Hymenoptera: Trichogrammatidae) em  
287 hospedeiros alternativos, sob diferentes temperaturas. Neotropical Entomology. 2007;36:084-  
288 087.

289

290

UNDER PEER REVIEW