

**Breast cancer awareness, knowledge and beliefs among
Libyan women**

ABSTRACT

Background: Breast cancer (BC) is the most frequent cancer of women. The high mortality in developing countries is associated with late detection, and lack of knowledge and adequate screening programmes.

Objectives: To assess the current level and determinants of knowledge and beliefs regarding BC, risk factors and various screening methods among Libyan women.

Study Design: A cross-sectional descriptive study

Place and Duration of Study: between September and October 2016 among a sample of adult women in western Libya.

Methods: 1091 woman aged between years were asked to fill a validated questionnaire to investigate their knowledge about the risk factors as well as their awareness and screening behaviours of BC.

Results: The majority of women participated in the study were aware of BC early warning signs and symptoms with over 90% of the women were able to list at least one symptom of breast cancer correctly. The most frequent warning sign identified was breast lump (91.0 %), followed by discharge from the nipples (80.6%). Also, 565 (52.7%) of those surveyed were aware that increasing age was associated with a higher incidence of breast cancer and 747 (68.3%) of the respondents identified positive family history as a risk factor. Moreover, 62% know how to perform self-examination (BSE), and only 59% ever performed BSE. The majority (92%) would seek medical advice if they discovered a mass in the breast whereas, about half of those (59%) would consult a male doctor.

Conclusions: Women participated in this study were fairly informed about BC risks and warning signs; the results appear to reflect growing awareness of women regarding BC screening methods. Health education message should be presented and delivered in a culturally-sensitive manner and tailored to provide simple and clear information and avoid false beliefs and misconceptions about the disease, its screening methods and management options.

Keywords: Breast cancer, Breast cancer self-examination, awareness, Libya.

INTRODUCTION:

Breast cancer is the most common cancer in women both in the developed and less developed world. It is estimated that worldwide over 508 000 women died in 2011 due to breast cancer. Although breast cancer is thought to be a disease of the developed world, almost 50% of breast cancer cases and 58% of deaths occur in less developed countries [1].

Breast cancer (BC) represents 10% of all cancers diagnosed annually and the second principal cause of cancer deaths in women worldwide [2,3]. The incidence of new cases is expected to rise from 10 million in 2002 to 15 million by 2025, with 60% of those cases occurring in developing countries. Data from the Arab world have placed breast cancer at the number one position with almost half of cases occurs in women under the age of 50 [4].

In Libya, breast cancer is accounting for more than 25% of all cancer in females with age-standardised mortality rate of 10.9 per 100,000 [5]. Breast cancer survival rates vary greatly

44 worldwide, ranging from 80% or over in North America, Sweden, Japan and Australia to around 60%
45 in Brazil and Slovakia and below 40% in Algeria [6] .

46 The low survival rates in less developed countries can be explained mainly by the lack of early
47 detection programmes, resulting in a high proportion of women presenting with late-stage disease, as
48 well as by the lack of adequate diagnosis and treatment facilities [1].

49 WHO promotes breast cancer control within the context of comprehensive national cancer control
50 programmes that are integrated to non-communicable diseases and other related problems.
51 Comprehensive cancer control involves prevention, early detection, diagnosis and treatment,
52 rehabilitation and palliative care. Raising general public awareness on the breast cancer problem and
53 the mechanisms to control as well as advocating for appropriate policies and programmes are key
54 strategies of population-based breast cancer control [1].

55 Screening and early detection is widely recognized as being a principal factor in reducing the
56 mortality from breast cancer [7]. However, previous studies reported that most breast cancer patients
57 present at advanced stages of the disease which emphasise the need for increasing awareness and
58 improved screening programmes including self-examination, clinical breast examination and
59 mammography [8-9].

60 In 2011, a comparison of the clinic-pathological and epidemiological features of breast cancer in
61 Libya to corresponding data from patients from Nigeria and Finland has reported that approximately
62 51% of Libyan patients were classified in stages 3 and 4 [10,11]. Also, research has shown poor levels
63 of knowledge towards risk factors awareness and screening methods even among young and educated
64 women [12-18].

65 Knowledge deficiency may lead to delayed presentation with advanced stages when little or no
66 benefit is derived from any form of therapy. For presentation at an early stage, women must be “breast
67 aware”; they must be capable of identifying symptoms of BC through routine practice of screening
68 [19].

69 Assessment of the current level of breast cancer awareness and knowledge toward risk factors and
70 screening methods is crucial for the development of awareness campaigns and programmes for
71 women to decrease the burden of the disease and mortality. Therefore, the study will be conducted to
72 assess the current level and determinants of knowledge and beliefs regarding breast cancer, risk
73 factors and various screening methods among Libyan women.

74

75 **MATERIALS AND METHODS:**

76 A cross-sectional descriptive study carried out between September and October 2016 among a sample
77 of adult women in western Libya. Women participating in the study were interviewed using pre-tested
78 validated questionnaire. The questionnaire included 49 questions pertaining three sections: 1) socio
79 demographic characteristics of women participating in the study; 2) knowledge of breast cancer risk

80 factors and warning signs; 3) Knowledge and awareness of women towards breast cancer screening
81 methods knowledge (BSE, CBE, and mammography).

82 Knowledge scoring

83 The questionnaire consisted of 23 items that assessed students' knowledge related to breast cancer (13
84 questions related to BC risk factors and 10 questions related to BC warning signs and 20 items that
85 assessed students' knowledge regarding breast cancer screening methods. These questions were then
86 scored; each correct response was scored one (1) point and each wrong or "don't know" was scored
87 zero (0). A correct response was based on literature and current practice. The knowledge index was
88 calculated for each participant by summing the number of correct answers. The total score of the
89 participants' knowledge regarding breast cancer is 23 (100%). The knowledge level was categorised
90 as "low" for scores within 0-49%, "moderate" for scores within 50-79% and "high" for scores within
91 80-100% [20]. These scores were then used to assess the relationship between socio-demographic
92 factors and level of breast cancer knowledge and warning signs.

93 Statistical analysis:

94 Data were translated to English and analysed using SPSS version 17 (SPSS Inc., Chicago, IL).
95 Descriptive statistics including means, standard deviation, frequencies, and percentages were obtained
96 for all continuous and categorical variables as appropriate. Chi-square test was used to examine the
97 association between the respondents' socio-demographic variables and knowledge of breast cancer.

99 RESULTS

100 The mean age of the 1091 women enrolled in the study was 33.2 (SD 9.6; age range: 18-61) years.
101 Most participants (71.8%) were aged less than 40 years and there were 501 (45.8%) single
102 respondents; 463 (42.2) were students; 996 (79.6%) had a university degree and 930 (93.4%) had no
103 family history of breast cancer (Table 1).

105 **Table 1.** Socio-demographic Characteristics of the Participants
106

CATEGORY	NUMBER	(%)
Age group (years)		
18-25	293	26.9
26-35	384	35.2
36-45	276	25.3
46-55	121	11.1
≥56	17	1.5
Level of education		
Primary	102	9.1
Secondary	306	27.9
University	683	79.6
Marital status		
Single	503	46.3
Married	523	48.1
Divorced	46	4.1
Widowed	19	1.6

Family history of breast cancer in a first degree relative		
Yes	92	8.4
No	985	90.3
No answer	14	1.3
History of breast problem	93	8.5

107

108

109 Section A of Table 2 shows respondents' knowledge of risk factors for breast cancer; The women
 110 surveyed had a fair knowledge of BC risk factors; 565 (52.7%) of those surveyed were aware that
 111 increasing age was associated with a higher incidence of breast cancer and 747 (68.3%) of the
 112 respondents identified positive family history as a risk factor for breast cancer. However, only third of
 113 the participants answered correctly about the effect of early menarche (31.9%) and late menopause
 114 (37.5%).

115 The majority of women participated in the study were aware of BC early warning signs and symptoms
 116 with over 90% of the women were able to list at least one symptom of breast cancer correctly. The
 117 most frequent warning sign identified was breast lump (91.0 %), followed by discharge from the
 118 nipples (80.6%). Only 566 person (52.4%) acknowledged that weight loss could be a warning sign of
 119 breast cancer (Table 2).

120

121 **Table 2.** Knowledge regarding Breast cancer risk factors and warning signs among study participants.
 122

Item (correct answer)	Number	%
A. Risk factors:		
Old age (yes)	565	52.7
Family history of breast cancer (yes)	747	69.2
Hormone replacement therapy (yes)	687	63.7
Alcohol consumption (yes)	604	56.2
Obesity (yes)	514	47.8
High fat diet (yes)	485	45.2
Smoking (yes)	640	59.7
Having children later on in life (yes)	458	42.9
Early menarche (yes)	341	31.9
Late menopause (yes)	399	37.5
Anxiety	536	50.0
Large breast size	399	40.1
Working	267	25.0
B. Early warning signs		
Lump under armpit	851	78.9
Breast lump	988	91.0
Bleeding or discharge from the nipple	874	80.6
Changes in the nipple	826	76.5
Redness of the breast skin	806	74.9
Changes in the size of breast	830	76.6
Changes in the shape of breast	844	78.4
Pain in the breast	784	72.7
Ulceration of the breast skin	824	76.5
Weight loss	566	52.4


123

124

125

126 The results showed that 62% of female participants know how to perform SBE, and only 59% ever
 127 performed BSE. The majority of women in the study (92%) would seek medical advice if they
 128 discovered a mass in the breast whereas, about half of those (59%) would consult a male doctor.
 129 Regarding screening methods, women were more familiar with BSE. Only 20% of participants were
 130 not aware of BSE, compared to 40% of women who were not aware of ultrasound as a BC screening
 131 method. In total, women who didn't know any screening methods constituted only 4%. While 45% of
 132 women were familiar with the five screening methods. Figure 1 shows the distribution of the
 133 knowledge scores amongst the respondents. The median score was 15 with 782 (71.5%) of women
 134 scoring >50% and 240 (22%) had a good score of o 80% or more. Age of the participants, marital
 135 status and their level of education did play a significant role in determining the knowledge attitude,
 136 while positive family history of breast cancer in a first degree relative as well as a history of breast
 137 problem were not significantly associated with BC knowledge (Table 3).

138 **Figure 1.** Distribution of breast cancer risk factors and warning signs knowledge scores.
 139
 140



141

142

143

144

145

146

147

148

Table 3. Relationship between knowledge scores and demographic variables of the respondents

Variable	Knowledge scores			P -value
	Good N (%)	Moderate N (%)	Poor N (%)	
Age group (years)				0.004
16-25	57 (19.5%)	167 (57.0%)	69 (23.5%)	
26-35	116 (30.2%)	152 (39.6%)	116 (30.2%)	
36-45	47 (17.0%)	152 (55.1%)	77 (27.9%)	
46-55	19 (15.7%)	49 (40.5%)	40 (33.1%)	

≥56	0 (0.0%)	9 (52.9%)	7 (41.1%)	
Level of education				
Primary	11 (12.7%)	45 (44.2%)	44 (43.1%)	0.000
Secondary	50 (16.4%)	156 (50.9%)	100 (32.7%)	
University	176 (25.8%)	341 (49.9%)	166 (24.3%)	
Marital status				
Single	116 (23.1%)	258 (51.3%)	129 (25.6%)	
Married	110 (21.1%)	250 (47.8%)	163 (31.1%)	0.036
Divorced	10 (21.7%)	23 (50.0%)	13 (28.2%)	
Widowed	1 (5.3%)	11 (57.9%)	7 (36.8%)	
Positive family history of breast Cancer in a first degree relative	17 (18.5%)	47 (51.1%)	28 (28.6%)	0.27
History of breast problem	19 (20.4%)	48 (51.6%)	26 (28.0%)	0.45

149

150

Chi Square test was performed. Level of significance is at $p < 0.05$.

151

152 Almost 50% of those with good knowledge score aged between 26-35 years and 75% had a university
153 degree. Whereas only 8% of participants with good knowledge score had a positive family history of
154 breast cancer or a previous breast problem.

155 **DISCUSSION**

156 Breast cancer is the most common of all female cancers in Libya [5]. In this study, the knowledge and
157 practice among general population in western Libya was explored. The main findings were that level
158 of awareness of risk factors and early warning signs of BC was moderate, with 71.5 % having good
159 knowledge, also the study showed that 59.2% of women participated in the study perform BSE.

160 The level of knowledge about breast cancer and the screening behaviour is generally poor in Arabic
161 region compared to the developed world [6,10,17]. In the present study, respondents answered
162 correctly that the commonest symptom of breast cancer is a breast mass. Our results are consistent
163 with those of similar studies carried in Saudi Arabia [21-22] and Kuwait [23].

164 The present study showed that women demonstrated higher knowledge of breast cancer screening and
165 risk factors and were more likely to perform BSE compared to other recent studies in neighbouring
166 countries [24-26]. Similar to previous studies [23, 25, 27] the most familiar methods was BSE
167 followed by CBE and mammography.

168 As it was expected, satisfactory knowledge scores were more common among younger participants
169 and those with higher educational levels. However, unlike other studies in the region [24-25] and
170 worldwide [28], the anticipated fact that women who had a breast problem or positive family history
171 of BC would have better knowledge scores could not be demonstrated in the present study. The study
172 revealed that about two thirds of the participants shared a misconception that early menarche and late
173 menopause were not risk factors for BC. This finding was supported by a previous study [29].

174 The results of this study may be influenced by the young age and the relatively high educational level
175 of the surveyed women which may reflect selection bias. A second limitation of the current study is
176 the use of convenience sampling to recruit participants which may limit the generalisability of the
177 findings. Nevertheless, convenience sampling considered a valid data collection method and has been

178 widely used in health education research [30]. In spite of these limitations, the study yielded
179 significant findings that could have implications reorganise the national health education strategy.

180

181 **CONCLUSIONS**

182 Women participated in this study were fairly informed about BC risks and warning signs; the results
183 appear to reflect growing awareness of women regarding BC screening methods. However, health
184 education message should be presented and delivered in a culturally-sensitive manner and tailored to
185 provide simple and clear information and avoid false beliefs and misconceptions about the disease, its
186 screening methods and management options.

187

188

189 **CONSENT**

190

191 All authors declare that written informed consent was obtained from the participants for
192 publication. A copy of the written consent is available for review by the Editorial office/ of this
193 journal.

194

195

196 **ETHICAL CONSIDERATIONS**

197 Permissions were obtained from the local health directorate and prior orientation of participants was
198 carried out. The data collection tools were anonymous, and data confidentiality was maintained
199 throughout the study.

200

201 **REFERENCES**

202 1-WHO. Breast cancer: Prevention and control. [cited2/4/2015]. Available at:
203 <http://www.who.int/cancer/detection/breastcancer/en/>

204 2-Wilson CM, Tobin S, Young RC. The exploding worldwide cancer burden: the impact of cancer on women.
205 *Int J Gynecol Cancer*. 2004;14:1-11.

206 3-WHO, [2006] EMRO. Non communicable diseases. [cited 08/04/2015]. Available from:
207 http://www.emro.who.int/ncd/cancer_breast.htm.

208 4-El Saghier NS, Khalil MK, Eid T, El Kinge AR, Charafeddine M, Geara F *et al*. Trends in epidemiology and
209 management of breast cancer in developing Arab countries: a literature and registry analysis. *Int J Surg*,
210 2007;5:225-33.

211 5-El Mistiri M, Pirani M, El Sahli N, El Mangoush M, Attia A, Shembesh R, Habel S, El Homry F, Hamad
212 Sand Federico M. Cancer profile in Eastern Libya: incidence and mortality in the year 2004. *Annals of*
213 *oncology*, 2010;21:1924-26

214 6-Coleman MP, Quaresma M, Berrino F, Lutz JM, De angelis R, Capocaccia R *et al* . Cancer survival in five
215 continents : a worldwide population based study [CONCORD]. *Lancet Oncol*, 2008;9(8):730-560.

- 216 7-Mai V, Sullivan T and Chiarelli AM.. Breast Cancer Screening Program in Canada: Successes and
217 Challenges. *Salud Publica Mex.* 2009;51 Suppl 2:s228-35.
- 218 8-Abdel-Fattah M, Zaki A, Bassili A, el-Shazly M, Tognoni G. Breast self-examination practice and its impact
219 on breast cancer diagnosis in Alexandria, *Egypt.East Mediterr Health J*, 2000;6:34-40.
- 220 9-Bener A, Alwash R, Miller CJ, Denic S, Dunn EV. Knowledge, attitudes, and practices related to breast
221 cancer screening: a survey of Arabic women. *J Cancer Edu.* 2001;16:215-20.
- 222 10-Boder JME, Abdalla FBE, Elfageih MA, Abusaa A, Buhmeida A, and Collan Y. Breast cancer patients in
223 Libya: comparison with European and central African patients,” *Oncology Letters*,2011;2: 323–330.
- 224 11-Ermiah E, Abdalla F, Buhmeida A, Larbesh E, Pyrhönen S and Collan Y. Diagnosis delay in Libyan female
225 breast cancer, *BMC Research Notes* 2012;5:452.
- 226 12-Milaat WA. Knowledge of secondary-school female students on breast cancer and breast self-examination in
227 Jeddah, *Saudi Arabia.East Mediterr Health J.* 2000;6:338-44.
- 228 13-Madanat H, Merrill RM . Breast cancer risk factors and screening awareness among women nurses and
229 teachers in Amman, *Jordan.Cancer Nursing.* 2002;25:276-82.
- 230 14-Leslie NS, Deiriggi P, Gross S, et al . Knowledge, attitudes, and practices surrounding breast cancer
231 screening in educated Appalachain women. *Oncol Nursing Forum.* 2003;30:659-67.
- 232 15-Amin TT, Al Mulhim AS, Al Meqihwi. A Breast Cancer Knowledge, Risk Factors and Screening Among
233 Adult Saudi Women in a Primary Health Care. *Asian Pac J Cancer Prev.* 2009;10(1):133-8.
- 234 16-Parsa P, Kandiah M, Zulkefli NA, Abdul Rahman H. Knowledge and behavior regarding breast cancer
235 screening among female teachers in Selangor, Malaysia. *Asian Pacific J Cancer Prev.* 2008;9: 221-7.
- 236 17-Ziuo FY, Twoier A, El-khewisky FS .Awareness of women about breast self-examination and risk factors for
237 breast cancer in Benghazi, Libya. *European Journal of Cancer.* 2014;50:e60-e61.
- 238 18- Noreen M, Murad S, Furqan M, Sultan A, Bloodsworth P. Knowledge and Awareness about Breast Cancer
239 and its Early Symptoms among Medical and Non-Medical Students of Southern Punjab, Pakistan. *Asian Pac J*
240 *Cancer Prev*, 2015;16 (3):979-984
- 241 19- Latif R. Knowledge and attitude of Saudi female student towards breast cancer: A cross-sectional study.
242 *Journal of Taibah university medical sciences.* 2014;9(4):328-334.
- 243 20- Akhtari-Zavare M, Juni MH, Said SM, Ismail IZ. Beliefs and behavior of Malaysia undergraduate female
244 students in a public university toward breast self-examination practice. *Asian Pacific J Cancer Prev.*
245 2013;14(1): 57-61.
- 246 21- Radi SM. Breast Cancer awareness among Saudi females in Jeddah. *Asian Pac J Cancer Prev.*
247 2013;14(7):4307-12.
- 248 22- Mohamed H A, Ibrahim Y M, Lamadah S M and Abo El-Magd M H. Application of the Health Belief
249 Model for Breast Cancer Screening and Implementation of Breast Self-Examination Educational Program for
250 Female Students of Selected Medical and Non-Medical Faculties at Umm al Qura University. *Life Sci J.*
251 2016;13(5):21-33.
- 252
253 23- Alharbi N, Alshammari M, Almutairi B, et al. Knowledge, awareness, and practices concerning breast
254 cancer among Kuwaiti female school teachers. *Alexandria J Medicine*, 2011;48:75–82.
- 255 24- Jahan S, Al-Saigul M, Abdelgadir H. Breast cancer: Knowledge, attitudes and practices of breast self-
256 examination among women in Qassim region of Saudi Arabia. *Saudi Med J.* 2006; 27:1737-41.

- 257 25- Abu-Shammala B I, Abed Y. Breast Cancer Knowledge and Screening Behavior among Female School
258 Teachers in Gaza City. *Asian Pac J Cancer Prev.* 2015;16 (17):7707-7711.
- 259 26- Oluwatosin OA, Oladepo A. Knowledge of breast cancer and its early detection measures among rural
260 women in Akinyele Local Government Area, Ibadan, Nigeria. *BMC Cancer* 2006;6:271-5,
- 261 27- Temiz M, Aslan A, Inand T, et al. Knowledge, attitudes, and behaviors of female teachers related to breast
262 cancer and breast examination in southern Turkey. *Breast Care*, 2002;3:55-60.
- 263 28- Grunfeld EA, Ramirez AJ, Hunter MS, Richard MA. Women's knowledge and beliefs regarding breast
264 cancer. *Br J Cancer* 2002;86:1373-8.
- 265 29- Manas K, Duangjai P, Chaowanee C, Suwit D. Knowledge, attitudes and practice of breast cancer screening
266 among female personnel of Walailak University. *Health Expect.* 2015;2014;1-10.
- 267 30- Bowling A. Research methods in health: Investigating health and health services. 3rd ed. ed. Maidenhead:
268 Open University Press.2009.
- 269
- 270
- 271
- 272
- 273

UNDER PEER REVIEW