

# Review Paper

## Anti-bacterial and anti-fungal properties of garlic extract (*Allium sativum*)

---

### ABSTRACT

Garlic (*Allium sativum*) contains various biologically active components that play significant role in treatment of bacterial and fungal infection. It contains sulfur compounds like allicin, ajoene, allylmethyltrisulfide, diallyltrisulfide, diallyldisulphide and others which exhibit various biological properties like antimicrobial, anticancer, antioxidant, immunomodulatory, anti-inflammatory, hypoglycemic and cardiovascular effects. The objective of the current review was to relate various literatures and assess the anti-microbial potential of garlic extract. The antimicrobial potency of garlic can be maximized by increasing the concentration of the extract.

**Keywords:** Antimicrobial activity, Bacteria, Fungi, Garlic extract

### INTRODUCTION

Medicinal plants have been used as traditional treatments for numerous human diseases [1]. In rural areas of the developing countries, they continue to be used as the primary source of medicine [2]. A wide range of microorganisms including bacteria, fungi, protozoa and viruses have been shown to be sensitive to crushed garlic preparations [3-4]. Garlic antimicrobial activities have been recognized for centuries being many of its therapeutic properties first mentioned in 1500 BC in an Egyptian recipe named *Papirus ebers*. Currently, it is used in folk medicine for the treatment of many diseases [5].

Microbial infections are the major cause of morbidity and mortality in the developed and developing country, although a number of antimicrobial agents are available for the treatment and management of infectious diseases [6]. Historically, garlic has been used for centuries worldwide by various societies to combat infectious disease. Garlic can be provided in the form of capsules and powders, as dietary supplements, and thus differ from conventional foods or food ingredients [7].

Garlic (*Allium sativum* L.) exhibit a broad antibiotic activity against both gram negative and gram positive bacteria including species of *Escherichia*, *Salmonella*, *Staphylococcus*,

29 *Streptococcus, Klebsiella, Proteus, Bacillus, Clostridium, Helicobacter pylori* [8-10]. One of  
30 the most potent and active component in garlic is the Sulphur compound called allicin which  
31 is a chemical compound produced when garlic is chopped, chewed or bruised. It is a  
32 powerful antibiotic and an agent that helps the body to inhibit the growth and development of  
33 pathogenic microbes [11].

#### 34 **ANTI-BACTERIAL EFFECTS OF GARLIC EXTRACT (*Alliums sativum*)**

35 Garlic (*Allium sativum*) has antibacterial, antifungal and antiviral properties. Aqueous,  
36 ethanol and chloroform extracts of garlic inhibited the growth of the pathogenic bacteria,  
37 though with varying degrees of susceptibility [9, 12].

38 Disease causing gram positive *bacteria S. mutans* were sensitive to garlic extract. The  
39 diameter of non-growth zone increased as the concentration of garlic extract increase.  
40 Similarly, *Lactobacillus acidophilus* was sensitive to different concentrations of garlic extract  
41 on blood agar medium. Again the diameter of inhibition zone increased as the concentration  
42 of garlic extract increased [13]. Garlic extract tested on *Bacillus subtilis, Staphylococcus*  
43 *aureus, Escherichia coli* and *Salmonella typhi* showed zone of inhibition 29mm, 26mm,  
44 31mm and 25mm respectively. The inhibition zone with streptomycin antibiotics was found to  
45 be 35mm, 33mm, 29mm and 31mm respectively [14].

46 The antibacterial effect of garlic extract was highest against skin pathogenic bacteria *S.*  
47 *aureus* with diameter zones of inhibition ranging from 4.40 – 3.80 cm followed by *S.*  
48 *epidermidis* ranging from 4.13 - 3.57cm, and then *Strep. pyogenes* with zones of inhibition  
49 ranging from 3.40 – 2.67cm. The least inhibition zones recorded about 2.32 – 1.55 cm were  
50 observed in *P. aeruginosa* [15]. Fresh domestic and imported garlic extract are known to  
51 play bacteriostatic activity against the methicillin-resistant *Staphylococcus aureus* (MRSA)  
52 with 31mm inhibition zone [16]. The most sensitive bacteria to garlic extract have been  
53 determined as *S. aureus* of the 26 mm diameter zone of inhibition, followed by *S. enteritidis*  
54 and *B. cereus* was the most resistant bacteria against garlic extract [17]. Garlic extract can  
55 be utilized for the development of broad spectrum antibiotics as it has wide spectrum  
56 antibacterial activity [18].

57 Garlic collected from Tamil Nadu town showed a better zone of inhibition in both gram  
58 negative and gram positive bacterial isolates but maximum diameter was obtained in  
59 *Staphylococcus aureus* and *V. cholera* 30mm, followed by  $\beta$  hemolytic streptococcus and  
60 *Escherichia coli* 26mm [19]. Garlic extract of 100% concentration showed a maximum zone  
61 of inhibition against *B. subtilis* and *Proteus mirabilis* bacteria. This indicates that garlic  
62 extract has the potential of a broad spectrum of activity against both gram-positive and  
63 gram-negative bacteria [20].

64 The concentration (100 mg/ml) of garlic extract had the highest inhibitory effect about  
65 (36mm) inhibition zone for *S. aureus* and about (33mm) inhibition zone for *E. faecalis*. The  
66 zone of inhibition was increased with increasing the concentration of garlic extracts. The  
67 lowest concentrations (5 and 10 mg/ml) were inhibited the bacteria weakly [21]. Another  
68 report also indicated that the effectiveness of 50% garlic extract against *S. aureus* is more  
69 (10 mm zone of inhibition) when compared to (ampicillin) antibiotic at 50% concentration (6  
70 mm zone of inhibition). Similarly 50% of garlic extract against *Pseudomonas aeruginosa*  
71 showed better zone of inhibition (9 mm) compared to 50% of antibiotic (ciprofloxacin) (7mm)  
72 zone of inhibition. But 20% of both garlic extract and antibiotics showed no result against  
73 *Pseudomonas aeruginosa* [22]. The alcoholic extract of garlic against *Staphylococcus*  
74 *aureus* showed that the zone of inhibition was 9 mm recorded for the concentration of 10  
75 mg/ml and 23 mm for the concentration 100 mg/ml. The concentrations 10-20 mg/ml were  
76 rather low active in preventing the growth of *S. aureus*, the concentrations 40-60 mg/ml were  
77 moderate active, while the concentrations 80-100mg/ml was highly active [23].

#### 78 **ANTIFUNGAL EFFECTS OF GARLIC EXTRACT (*Allium sativum*)**

79 The slow pace of newer antibiotic development coupled with the availability of fewer  
80 antifungal agents with fungicidal actions centered on inhibition of ergosterol synthesis has  
81 provided the need to explore nature in search of phytotherapeutic agents with novel targets  
82 and mode of actions [24]. It was revealed from the study of [25], the aqueous and ethanol  
83 diluted garlic extracts showed complete inhibition (100%) of *Botrytis cinerea* at the higher  
84 concentrations (80% and 60%) respectively. At a concentration of 80%, the aqueous and  
85 ethanol diluted extracts inhibited *Penicillium expansum* by 96.21% and 99.21% respectively.  
86 Ethanol diluted extracts seemed to be more effective against *Neofabraea alba* with 80%  
87 extract showing 79.63% inhibition. [26], also reported that the ethanolic extract of garlic  
88 tested against *A. flavus*, *A. niger* and *C. herbarum* showed inhibition diameter of (3.766cm,  
89 4.750cm and 1.883cm) respectively after seven-day incubation on PDA. With inorganic  
90 water base extraction medium garlic extract had inhibition diameter growth (0.934cm) on *A.*  
91 *flavus*, (0.884cm) on *A. niger* and (5.750cm) on *C. herbarum* at day seven of incubation.  
92 A report of [27], indicated that the mean zone of inhibition of *Candida albicans* was found to  
93 be  $28.0 \pm 1.0$  mm with aqueous garlic extract and  $27.5 \pm 0.5$  mm with clotrimazole. This result  
94 indicated that that garlic juice has an equal or a better effect against *Candida albicans*  
95 compared with a commercial antifungal drug. The aqueous garlic extract exhibit higher zone  
96 of inhibition against *Aspergillus niger* was reported to be  $41.0 \pm 4.0$  mm and  $22.5 \pm 1.5$  mm with  
97 clotrimazole. Both aqueous and methanolic extracts of *A. sativum* exhibited antifungal  
98 activities against almost all the tested *C. albican* [28]. Petroleum ether and aqueous extracts

99 of garlic were found to inhibit the growth of the standard organism of *Candida albicans* and  
100 *Aspergillus niger* and the clinical isolates of *A. flavus*, *Curvularia lunata*, *Microsporum*  
101 *audouinii*, *Trichophyton soudanense* and *Trichophyton mentagrophytes* at a concentration of  
102 10mg/ml [29].

103 Antidermatophytic activity of aqueous extract of garlic tested on dermatophytic fungi showed  
104 that zones of inhibition against *Trichophyton mentagrophytes* was (18.00 mm), *Trichophyton*  
105 *rubrum*, (18.33 mm), *Microsporum gypseum* (16.53mm), *Trichophyton verrucosum*  
106 (15.67mm) and *Epidermophyton floccosum* (6.00mm). Inhibition zone of Nystatin used as  
107 control for cultures at a concentration of 1mg/ml varies between 25-31 mm [30]. The extract  
108 of garlic at highest concentration proved highly to be effective in reducing the spore  
109 germination of *Alternaria alternata* and *Rhizopus stolonifera* [31].

## 110 CONCLUSION

111 Garlic extracts exhibit broad spectrum antibiotics against both gram positive and gram  
112 negative bacteria and fungi. The anti-bacterial and anti-mycotic activity of garlic was  
113 increased with increasing the concentration of garlic extract. Garlic extract enhances the  
114 antibacterial activity and protect individual from bacteria invasion. Thus, consuming of garlic  
115 may be utilizing as an economic way for patients or hospital workers and have been  
116 proposed as novel treatments of bacterial infectious diseases also to reduce the problem of  
117 multi-drug resistant pathogenic bacteria.

## 118 REFERENCES

- 119 1. Ross ZM, O'gara EA, Hill DJ, Sleightholme HV, Maslin DJ. Antimicrobial Properties of  
120 Garlic Oil against Human Enteric Bacteria: Evaluation of Methodologies and  
121 Comparisons with Garlic Oil Sulfides and Garlic Powder. Appl.enviro.n.micr.obial.2001;  
122 67(1):475–480.
- 123 2. Gaherwal S, Johar F, Wast N, Prakas MM. Anti-Bacterial Activities of *Allium sativum*  
124 against *Escherichia coli*, *Salmonella Ser. typhi* and *Staphylococcus aureus*. International  
125 Journal of Microbiological Research. 2014; 5(1):9-22.
- 126 3. Ankri S, Mirelman D. Antimicrobial properties of allicin from garlic. Microbes and  
127 Infection.1999; 2:125–129.
- 128 4. Hannan A, Rauf K, Ullah MI, Naeem T, Raja M, Qamar MU, Tahir R, Saba M. Inhibitory  
129 effect of aqueous garlic (*Allium sativum*) extract against clinical isolates of *Salmonella*  
130 *typhi*. African Journal of Microbiology Research. 2012; 6(21):4475-4480.
- 131 5. Groppo FC, Ramacciato JC, Motta RH, Ferraresi PM, Sartoratto A. 2007. Antimicrobial  
132 activity of garlic against oral *Streptococci*. Int J Dent Hygiene.2007; 5:109–115.

- 133 6. Anitha M, Hemapriya J. Efficacy of antibacterial activity of garlic cloves from Tamil Nadu  
134 and Jowai region. Int J Res Med Sci.2014; 2(2):434-438.
- 135 7. Hindi NK. In vitro Antibacterial Activity of Aquatic Garlic Extract, Apple Vinegar and  
136 Apple Vinegar - Garlic Extract combination. American Journal of Phytomedicine and  
137 Clinical Therapeutics.2013; 1(1):042-051.
- 138 8. Ranjan S, Dasgupta N, Saha P, Rakshit M, Ramalingam C. Comparative study of  
139 antibacterial activity of garlic and cinnamon at different temperature and its application  
140 on preservation of fish. Adv. Appl. Sci. Res.2012; 3(1):495-501.
- 141 9. Mikaili P, Maadirad S, Moloudizargari M, Aghajanshakeri S, Sarahroodi S. Therapeutic  
142 Uses and Pharmacological Properties of Garlic, Shallot, and Their Biologically Active  
143 Compounds. Iran J Basic Med Sci. 2013; 16 (10):20-25.
- 144 10. Baljeet SY, Simmy G, Ritika Y, Roshanlal Y. Antimicrobial activity of individual and  
145 combined extracts of selected spices against some pathogenic and food spoilage  
146 microorganism. International Food Research Journal.2015; 22(6):2594-2600.
- 147 11. Chuku EC. Anti-fungal properties of ginger (*Zingiber officinale*) and garlic (*Allium*  
148 *sativum*) on smoked cat fish mycoflora. Niger. J. Mycol.2015; 6:48 – 57.
- 149 12. Bakri M, Douglas CW. Inhibitory effect of garlic extract on bacteria. Arch Oral Biol .2005;  
150 50:645-651.
- 151 13. Motamayel FA, Hassanpour S, Alikhani MY, Poorolajal J, Salehi J. Antibacterial effect of  
152 Eucalyptus (*Globulus labill*) and garlic (*Allium sativum*) extracts on oral Cariogenic  
153 bacteria. Journal of Microbiology Research and Review.2013;1(2):12-17.
- 154 14. Ahmed AM, Ravi S, Ghogare P. Studies on Antimicrobial Activity of Spices and Effect of  
155 Temperature and Ph. on Its Antimicrobial Properties. Journal of Pharmacy and  
156 Biological Sciences.2015;10(1):99-102.
- 157 15. Abdulrahman DM, Daskum AM, Abdulrahim KM, Dadile AM, Amma H. Antibacterial  
158 potency of garlic extract against certain skin pathogenic bacteria. Novel Research in  
159 Microbiology Journal.2017;1(1):3-1.
- 160 16. Strika IA, Basic A.B, Halilović N. Antimicrobial effects of garlic (*Allium sativum* L.).  
161 Bulletin of the Chemists and Technologists of Bosnia and Herzegovina.2017;47:17-20.
- 162 17. Taş E, Erginkaya Z, Kalkan S, Turhan E. Determination of Antimicrobial Effects of  
163 Probiotic Lactic Acid Bacteria and Garlic Extract against Some Foodborne Pathogenic  
164 Bacteria. Turkish Journal of Agriculture - Food Science & Technology.2017; 5(2):125-  
165 131.
- 166 18. Chand B. Antibacterial effect of garlic (*Allium sativum*) and ginger (*Zingiber officinale*)  
167 against *Staphylococcus aureus*, *Salmonella typhi*, *Escherichia coli* and *Bacillus cereus*.  
168 JMBFS.2013; 2(4):2481-2491.

- 169 19. Anitha M, Hemapriya J. Efficacy of antibacterial activity of garlic cloves from Tamil Nadu  
170 and Jowai region. Int J Res Med Sci.2014; 2(2):434-438.
- 171 20. Durairaj S, Srinivasan S, Lakshmanaperumalsamy P. In vitro Antibacterial Activity and  
172 Stability of Garlic Extract at Different pH and Temperature. Electronic Journal of  
173 Biology.2009; 5(1):5-10.
- 174 21. Salih JM, Monawer AT, Abdulkahar IM. Antibacterial activity of garlic against multi-drug  
175 resistant. *Staphylococcus aureus* and *Enterococcus faecalis* in Duhok city. J. Univer.  
176 Duhok.2016;19(1):114-122.
- 177 22. Praba SK, Kumaresan R. Efficacy of antimicrobial activity of aqueous garlic (*Allium*  
178 *sativum*) extract against different bacterial species. J. Chem. Pharm. Res.2014;  
179 6(10):677-679.
- 180 23. Khashan AA. Antibacterial activity of garlic extract (*Allium sativum*) against  
181 *Staphylococcus aureus* in vitro. G.J.B.B.2014; 3(4); 346-348.
- 182 24. Iwalokun BA, Ogunledun A, Ogbolu DO, Bamiro SB,Omojola JJ. In Vitro Antimicrobial  
183 Properties of Aqueous Garlic Extract Against Multidrug-Resistant Bacteria and Candida  
184 Species from Nigeria. J Med Food. 2004;7(3):327-333.
- 185 25. Daniel CK, Lennox CL, Vries FA. In-vitro effects of garlic extracts on pathogenic fungi  
186 *Botrytis cinerea*, *Penicillium expansum* and *Neofabraea alba*. South African Journal of  
187 Science. 2015;111(7):1-8.
- 188 26. Tagoe D, Baidoo S, Dadzie I, Kangah V, Nyarko H. A Comparison of the Antimicrobial  
189 (Antifungal) Properties of Garlic, Ginger and Lime On *Aspergillus flavus*, *Aspergillus*  
190 *niger* and *Cladosporium herbarum* Using Organic and Water Base Extraction Methods.  
191 The Internet Journal of Tropical Medicine.2009;7:1-8.
- 192 27. Abdallah EM. Potential Antifungal Activity of Fresh Garlic Cloves (*Allium sativum* L.) from  
193 Sudan. J. Biotech. Research.2017; 3(11):106-109.
- 194 28. Hailu G, Bitew M, Temesgen M. In vitro antifungal activity of *Croton macrostachys* and  
195 *Allium sativum* extracts against *Candida albicans* and *Trichophyton mentagrophytes*  
196 isolates. Int. J. Biotechnol. Food Sci.2017; 5(3):42-47.
- 197 29. Suleiman EA, Abdallah WB. In vitro Activity of Garlic (*Allium sativum*) on Some  
198 Pathogenic Fungi. European Journal of Medicinal Plants.2013; 4(10):1240-1250.
- 199 30. Mercy KA, Jeoma I, Emmanuel KJ. Anti-dermatophytic Activity of garlic (*Allium sativum*)  
200 extracts on some dermatophytic fungi. Inter. L. N. Scien. 2014; 19:34-40.
- 201 31. Nisa TU, Wani AH, Mir RA. Antimycotic activity of plant extracts on the spore  
202 germination of some pathogenic fungi. Mycopath.2010;8(2):65-69.