

Field Survey of cassava susceptibility to termite infestation as influenced by time, scale and management strategy in selected Owerri Agricultural zones, Southeast Nigeria.

ABSTRACT

The field survey was conducted in 2016 in four selected Agricultural zones of Owerri, Southeastern Nigeria namely Ezinnihite Mbaise (Zone 1), Owerri North (Zone 2), Mbaitoli (Zone 3) and Owerri West (Zone 4). A random selection of thirty cassava farmers from each of the study areas was made. The sample size was made up of a total of one hundred and twenty (120) respondents. Data were collected through structural questionnaire administration to the respondents on the four selected zones and were analyzed using Descriptive Statistics such as the use of Percentages, Frequencies and Means. Result shows that cassava stems were more susceptible to termite attack than tubers especially during dry periods. Equally cassava devastation by termite occurred mostly at planting where newly planted cuttings suffered most. Also, the study reveals that termite infestation on cassava was more between January and May as well as between October and December. On the other hand, respondents submitted that the use of chemical control either singly or in synergy with other **nonchemical** means to control termites appears to be very popular amongst respondents in the study area. Result also shows that youths and mature adults were actively engaged in cassava farming but **the** majority of them were evidently unskilled by virtue of their low literacy levels. Farmer education is recommended to enable the integration of pest/termite avoidance principle into termite control strategy across the zones under study.

Keywords: ?????

1 .INTRODUCTION

Cassava (*Manihot esculenta* Crantz.) is a perennial woody shrub with an edible root which grows in tropical and sub-tropical areas of the world. It has the ability to grow on marginal lands and can tolerate long dry spell (IITA, 2000) [1]. However, cassava does well on well-drained, rich and friable loamy soils (Akinsanmi, 1987) [2].

In Africa, cassava provides a basic daily source of dietary energy and has gained popularity as one of the most important root crops in Nigeria especially in the Southern States (Nwokoma, 1998) [3]. Cassava is the second most important staple food in sub-Saharan Africa and accounts for more than 100 calories per day in the diet of an individual (IITA, 1988) [4].

Cassava roots are processed into a wide variety of granules, pastes, flour, etc. or consumed freshly boiled or raw. It is used in the production of starch, garri, 'foo-foo', wet and dry chips (Nwokoma,

35 1998) [3]. The fresh cassava tuber can be used considerably as a source of feed for livestock (sheep,
36 goats, cattle, pigs, etc.). In many rural households, cassava peel is fed to domestic animals
37 (Ihekoronye and Ngoddy, 1985) [5].

38 In the traditional farming systems where cassava is usually one of the many crops being grown, pest
39 control is often given a low priority and so cassava receives minimal pesticide application. Under
40 such conditions yields are often low (Henry, 1995) [6]. Arthropod pests and diseases are major
41 factors causing this yield reduction (Belloti *et al.*, 1999) [7]. In the humid lowlands, the predominant
42 diseases of cassava include cassava mosaic virus (CMV), cassava bacterial blight (CBB), cassava
43 anthracnose disease (CAD) and root rots. The major insect pests are cassava green mite (CGM:
44 *Mononychellus* spp.), elephant grasshopper (*Zonocerus elegans* L. and *Zonocerus variegatus* Thumb.),
45 cassava mealybug (CM: *Phenacoccus manihotis*), a wide range of rodents and termites (Hillocks and
46 Thresh, 2002) [8].

47 A survey of the distribution of termites in the country by Malaka (1973) [9] has revealed that certain
48 species are restricted to a particular vegetation zone while some are distributed all over the zones.
49 For instance, rainforest appears to have a more dominant species than other vegetation zones. About
50 26 species have been recorded from the Guinea Savanna (ODM, 1997) [10] and of which only 10 are
51 dominant. Altogether 120 species of termites have been identified in Nigeria (Logan *et al.*, 1992) [11]
52 out of which only 20 damage crops and building

53 However, a thorough survey on cassava susceptibility to termite infestation in relation to time and
54 management strategy has not received attention by our farmers in Owerri , therefore, the need to
55 carry out field survey of cassava susceptibility to termite infestation as influenced by time, scale and
56 management strategy in selected Owerri Agricultural zones of Southeastern Nigeria forms the
57 objective of this study.

58 1. Materials and Methods

59 The study was conducted in 2016 cropping season. It was carried out in Owerri Agricultural zone
60 located at the South-western part of Imo State. Owerri is located between Latitude 4° 40' and 8° 15' N
61 and Longitude 6° 40' and 8° 15' E (FDALR, 1985) [12]. It is of the humid tropics. It records means
62 annual rainfall of about 18000 -2190 mm which spans from early March to October. The minimum
63 and maximum mean annual temperatures were 22.5°C and 31.9°C respectively with relative humidity
64 of about 82.6%. (Nwosu and Adeniyi, 1980) [13]. The zone comprises ten Local Government Areas,
65 namely; Aboh Mbaise, Ahiazu Mbaise, Ezinihite Mbaise, Mbaitoli, Ikeduru, Ngor-okpala, Ohaji/
66 Egbema, Owerri Municipal, Owerri North and Owerri West. Farmers in the zone are mainly
67 **smallholders** known for growing such arable crops as maize, melon, yam, cassava etc. (ISADP. 2000)
68 [14].

69 Four out of the ten Local Government Areas were randomly selected for the study. The selected areas
70 are Ezinihite Mbaise in Owutu Community, Owerri North in Azaraubo Community, Mbaitoli in
71 Obinnoha Community, and Owerri West in Obinze Community. These areas were selected based on
72 the quantum of cassava cultivation that was being carried out by farmers. A random selection of
73 thirty cassava farmers from each of the study areas who had admittedly been producing cassava for
74 the past ten years was made. The sample size was made up of a total of one hundred and twenty
75 (120) respondents.

76 Data was collected through structured questionnaire administration to the respondents on the four
77 selected local Government Areas.

78 All Data collected were analyzed using Descriptive Statistics such as the use of Percentages,
79 Frequencies and Means.

80 3. Results and DISCUSSION

81 Table (1) indicates the degree of susceptibility of different parts of the cassava plant to termites
82 infestation in the field. 50% (zone 4) and 43% (zone 1) proportion of the respondents under study
83 claimed that cassava stems tend to exhibit high susceptibility to termites infestation. Also, 33% (zone
84 2) and 20% (zone 1) of them accepted that cassava leaves were attacked by termites, while 6.7%
85 (zone 4), 3.3% (zones 1 and 2) and 0% (zone 3) agreed that cassava tubers were susceptible to
86 termites infestation. In the same Table, 60% multiple responses from zone 3 claimed that optimal
87 termites infestation in cassava field was recorded on different parts of the plant.

88 Also, the distribution of the different levels of termite damage to cassava at various growth phases
89 was presented in Table (2). Result reveals that 73.3% (zone 4) and 56.7% (zone 1) proportion of the
90 respondent sampled in the study area claimed that cassava incurred more damage from termites
91 infestation at planting (establishment) period. This was upheld by 50% (zone 2) and 36.7% (zone 3)
92 proportion of them. Equally, 16.7% (zones 1 and 4) and 23.3% (zone 1) of the respondents agreed
93 that termites cause economic damage to cassava at maturity and harvest periods respectively. Also,
94 the table shows the distribution of respondents by their regular observation in the study area
95 (multiple responses) where 53.3%(zone 3) and 36.7% (zone 2) agreed that attack on cassava by
96 termites occurs at any period of its growth phase.

97 The high susceptibility of cassava stems to termites attack as well as the plant's prone to attack at
98 planting period as claimed by respondents was in line with Onwueme (1978) [15] that termite
99 activities in cassava field are more devastating on the stems and at early stages of their development
100 resulting to poor stand establishment.

101 Table (3) presents the distribution of termite infestation in cassava field by time. 43.3% (zone 4),
102 33.3% (zone 1), 30% (zone 3) and 16.7% (zone 2) proportion of the respondents agreed that
103 termites infestation occur from January to May, while 46.7% (zones 1 and 2), 40% (zone 3) and 50%
104 (zone 4) proportion of them claimed that infestation takes place more in October to December.
105 However, 20% (zone 1) and 13.3% (zone 2) of the respondents accepted that termites infestation
106 occur in June to September.

107 **The result shows** that termite infestation on cassava was more extensive between January to May and
108 between October to December, implies that optimal periods of termites infestation in cassava field
109 coincides with periods of prolonged dry spell. This is in conformity with Taylor (1977) [16] who
110 reported that termites attack the roots, tubers, young seedlings, shoots and stems of crops and their
111 damage is more extensive during the dry season when the crops are weakened by water stress or
112 drought.

113 Table (4) shows the distribution of economic loss from termites in the cassava field by plant part.
114 Majority of the cassava farmers sampled under the study area across the zones admitted that the
115 greatest economic loss from termites infestation occurs on the stems. 50% (zone 4) and 46.67%
116 (zone 1) proportion of them believed that the greatest economic loss on cassava by termites comes
117 from the stems. On the other hand, none of the respondents in zone 1 (0%) and zone 3 (0%) agreed
118 that cassava leaves exhibited any evidence of economic loss from termites infestation. However,
119 6.67% (zone 4) and 3.33% (zone 2) proportion of them accepted that cassava leaves suffered
120 economic loss from termites. Equally, 26.67% (zone 2) and 16.67% (zone 4) of the respondents
121 maintained that cassava tubers also showed marked evidence of economic loss from termites. In the

122 same vein, 56.67% (zone 3) and 40% (zone 1) from multiple responses claimed that huge economic
123 losses were recorded on different parts of the plant, while 26.67% (zones 2 and 4) proportion of
124 them concurred to this claim.

125 Higher economic loss from cassava stems as a result of termite infestation was described by Nweke *et al.* (1994), [17] that cassava field planted early or late in the rainy season often have poor
126 establishment record because termites feed on the planted sticks (cuttings).
127

128 Result in Table (5) shows different methods of termites control measures employed by farmers in
129 their cassava field. Majority of the respondents in zone 4 (30%) and zone 2 (60%) claimed to employ
130 cultural and chemical methods of control respectively. On the other hand, 3.33% (zone 2) of the
131 respondents employed Biological method. In the same Table, 73.33% (zone 3) and 43.33% (zone 1)
132 of the farmers sampled agreed to employ two or more different methods (multiple responses) of
133 control. This claim was upheld by 33.33% (zone 4) and 10% (zone 2) proportion of them

134 Application of chemical control and other non-chemical means at different periods of time either
135 singly or in synergy in the cassava field to control termites appears to be very popular amongst
136 respondents in the study area. Though chemical control is effective but most chemical control
137 measures rely principally on the use of organo-chlorine insecticides such as aldrin, dieldrin, lindane
138 etc. (Umeh, 2002) [18]. Unfortunately this type of control measure is no longer popular due to the
139 associated environmental contamination and health hazards (PAN, UK, 2003) [19]. Any control
140 measure that ensures adequate synergy of these methods and which promote the rapid growth of
141 healthy crop is a suitable means for avoiding termites damage (Schmutterer *et al.*, 1978) [20].

142 The Socio-economic characteristics of respondents were described in Tables 6, 7, 8 and 9 below. In
143 Table 6, 33.33% (zone 2) and 16.67% (zone 1) of the respondents sampled under the study area were
144 between the age group of 25-40 years, while 56.67% (zone 1), 46.67% (zone 2) and 36.67% (zones 3
145 and 4) were between 41-55 years. Also, 56.67% (zone 3) and 50.00% (zone 4) were between the age
146 group of 56-70 years and 6.67% (zone 3) and 3.33% (zone 1) were between 71-100 years. However,
147 the mean age group of respondents were 50.17% (zone 1), 45.83% (zone 2), 59.00% (zone 3) and
148 53.43% (zone 4).

149 In Table 7, larger proportion of the cassava farmers in zone 2 (100%), zone 1 (90%), zone 4 (83.34%)
150 and zone 3 (80%) sampled cultivated under farm size of 1-5 hectares (ha) while, 10.00% (zone 1),
151 20% (zone 3) and 13.33% (zone 4) of them had farm size of 6ha and above. The mean land area were
152 3.5ha (zone 1), 3.00 (zone 2), 4.00ha (zone 3) and 3.57ha (zone 4). On the other hand, Table 8 shows
153 that majority of respondents in zones 1 and 2 (70%) and zone 4 (60%) sampled were males,
154 while 56.67% (zone 3) were females.

155 Apart from that, Table (9) indicates that 93.33% (zone 3), 70% (zone 1), 56.67% (zone 2) and 50%
156 (zone 4) of the respondents sampled acquired non-degree educational training. However, 36.67%
157 (zone 2) and 26.67% (zone 1) of them obtained Bachelor of Science Degree, while 30% (zone 4) had
158 Master of Science Degree.

159 Different ages and mean age of respondents sampled found to fall between the range of 25-70 years
160 and 59 years respectively, implies that majority of them were at their productive age group. It also
161 shows that youths and mature adults are actively involved in cassava production. Equally, the
162 indication that majority of the respondents cultivated under land area of 1-5 ha and on mean land
163 area of 4.00 ha across the zones, showed that land as a productive resource was not a constraint in
164 the study area. Also, on the gender balancing and participation, majority of the respondents were
165 males. This is probably because traditionally, men have right to land than women. Quisumbing

166 (1994) [21] opined that there has been a great disparity between women and men in the size of
 167 landholdings. Apart from that, majority of the respondents sampled were found to be literate but
 168 obtained certificates other than Degree. This implies that larger proportion of them were primarily
 169 un-skilled.

170 **Conclusion/Recommendation**

171 Termite infestation in cassava field and their subsequent attack were more severe during dry periods
 172 than in wet season. However, cassava stems appeared to be more susceptible to attack than the
 173 tubers. On the other hand, farmers in the study areas combined chemical and cultural means a
 174 method of controlling termites in their cassava field.

175 Equally, the study indicated that youths and mature adults actively engaged in cassava farming.
 176 However, a greater proportion of the respondents agreed that land as a productive resource was not
 177 a limiting factor. Finally, the majority of the farmers sampled were primarily unskilled evidenced in
 178 their low literacy level.

179 Farmers in a termite endemic area such as Owerri, Imo State, Nigeria are advised to be conscious of
 180 the two extremes of heavy termite infestation by ensuring that planting of cassava is not carried out
 181 between January and May. Also, harvesting of cassava tubers should not be delayed up to October
 182 through December in the season. However, the study strictly recommends that farmers under this
 183 condition should adopt late planting and early harvesting options.

184 Also, Farmers through the acquisition of qualitative education can aptly adopt the use of pest
 185 avoidance strategy in such a manner that sound and sustainable peculiar termite control package
 186 across the zones under study can be achieved. This will not only be efficacious in reducing termite
 187 load and damage on cassava but also economical in an application.

188 **Table 1: Frequency and Percentage Distribution of Plant Parts Most Susceptible to Attack by**
 189 **Termites According to Zones**

Plant Part	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)
Stems	13	43.3	9	30	10	33.3	15	50
Leaves	6	20	1	33.3	2	6.67	1	3.33
Tubers	1	3.3	1	3.3	0	0	2	6.7
All Parts	3	10	3	10	0	0	5	16.7
Multiple rep.	7.0	23.3	7.0	23.3	18.0	60.0	7.0	23.3
Total	30	100	30	100	30	100	30	100

190 **Source: Computed from Field Survey Data (2016).**

191 **Table 2: Frequency and Percentage Distribution of Damage at Different Plant Growth Phases**
 192 **According to Zone**

Growth phase	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)
planting	17.00	56.67	15.00	50.00	11.00	36.67	22.00	73.33
maturity	5.00	16.67	4.00	13.33	3.00	10.00	5.00	16.67
harvest	7.00	23.33	0.00	0.00	0.00	0.00	3.00	10.00
multiple rep	1.00	3.33	11.00	36.67	16.00	53.33	0.00	0.00
Total	30	100	30	100	30	100	30	100

193 **Source: Computed from Field Survey Data (2016).**

194
 195 **Table 3: Frequency and Percentage Distribution of Termites Infestation by Time According to**
 196 **Zones**

Time (months)	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)
Jan – May	10.00	33.33	5.00	16.67	9.00	30.00	13.00	43.33
June – Sept	6.00	20.00	4.00	13.33	0.00	0.00	2.00	6.67
Oct – Dec	14.00	46.67	14.00	46.67	12.00	40.00	15.00	50.00
Multiple rep.	0.00	0.00	7.00	23.33	9.00	30.00	0.00	0.00
Total	30	100	30	100	30	100	30	100

197 **Source: Computed from Field Survey Data (2016).**

198
 199 **Table 4: Frequency and Percentage Distribution of Economic Loss from Termites by Plant Part**
 200 **According to Zones**

Plant part	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)
Stems	14.00	46.67	13.00	43.33	10.00	33.33	15.00	50.00
Leaves	0.00	0.00	1.00	3.33	0.00	0.00	2.00	6.67
Tubers	4.00	13.33	8.00	26.67	3.00	10.00	5.00	16.67
Multiple	12.00	40.00	8.00	26.67	17.00	56.67	8.00	26.66

rep.

Total	30	100	30	100	30	100	30	100
--------------	-----------	------------	-----------	------------	-----------	------------	-----------	------------

201 **Source: Computed from Field Survey Data (2016)**

202

203 **Table 5: Frequency and Percentage Distribution of Control Measures in Use According to**
 204 **Zones**

Control Method	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)
Cultural	8.00	26.67	5.00	16.67	5.00	16.67	9.00	30.00
Chemical	6.00	20.00	18.00	60.00	1.00	3.33	7.00	23.34
Biological	0.00	0.00	1.00	3.33	0.00	0.00	0.00	0.00
All	2.00	6.67	0.00	0.00	0.00	0.00	4.00	13.33
Multiple								
rep.	13.00	43.33	3.00	10.00	22.00	73.33	10.00	33.33
Others	1.00	3.33	3.00	10.00	2.00	6.67	0.00	0.00
Total	30	100	30	100	30	100	30	100

205 **Source: Computed from Field Survey Data (2016).**

206

207

208 **Table 6: Frequency and Percentage Distribution of Farmers by Age According to Zones**

Age of Farmers	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)	Frequency	Percent (%)
25 - 40	5.00	16.67	10.00	33.33	0.00	0.00	4.00	13.33
41 - 50	17.00	56.67	14.00	46.67	11.00	36.67	11.00	36.67
56 - 70	7.00	23.33	6.00	20.00	17.00	56.66	15.00	50.00
71 - 100	1.00	3.33	0.00	0.00	2.00	6.67	0.00	0.00
Total	30	100	30	100	30	100	30	100
Mean		50.17		45.83		59		53.43

209 **Source: Computed from Field Survey Data (2016).**

210

211 **Table 7: Frequency and Percentage Distribution of Farmers by Farm Size According to Zones**

Farm Size (ha)	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	
1 - 5	27.00	90.00	30.00	100.00	24.00	80.00	25.00	83.34
6 - 10	3.00	10.00	0.00	0.00	6.00	20.00	4.00	13.33
11 - 15	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
16 - 20	0.00	0.00	0.00	0.00	0.00	0.00	1.00	3.33
Total	30	100	30	100	30	100	30	100
Mean		3.5		3		4		3.57

212 **Source: Computed from Field Survey Data (2016).**

213

214 **Table 8: Frequency and Percentage Distribution of Farmers by Gender According to Zones**

Gender	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	Frequency Percent (%)	
Male	21.00	70.00	21.00	70.00	13.00	43.33	18.00	60.00
Female	9.00	30.00	9.00	30.00	17.00	56.67	12.00	40.00
Total	30	100	30	100	30	100	30	100

215 **Source: Computed from Field Survey Data (2016).**

216

217

218

219

220

221

222

223

224

225

226

227 **Table 9: Frequency and Percentage Distribution of Farmers by Level of Education According to**
 228 **Zones**

Education	Zone 1		Zone 2		Zone 3		Zone 4	
	Frequency		Frequency		Frequency		Frequency	
	Percent	(%)	Percent	(%)	Percent	(%)	Percent	(%)
B.Sc	8.00	26.67	11.00	36.67	2.00	6.67	5.00	16.67
M.Sc	1.00	3.33	1.00	3.33	0.00	0.00	9.00	30.00
Ph.D	0.00	0.00	1.00	3.33	0.00	0.00	1.00	3.33
Others	21.00	70.00	17.00	56.67	28.00	93.33	15.00	50.00
Total	30	100	30	100	30	100	30	100

229 **Source: Computed from Field Survey Data (2016).**

230 **COMPETING INTERESTS** Authors have declared that no competing interests exist.

231

232 **REFERENCES**

- 233 1. IITA(International Institute of Tropical Agriculture) 2000. IITA research priorities and
 234 strategies 2000-2005, IITA Ibadan Nigeria.<http://www.iita.org/crop/cassava.htm>
- 235 2. Akinsanmi,O. 1987. Certificate in Agricultural Science. William Clowes and sons limited,
 236 London. Beccles and Colchester Pp. 94-95.
- 237 3. Nwokoma, S.N. 1998. Nigeria's staple foods Nigeria. Spring Field Publishers. Pp. 10-13
- 238 4. IITA (International Institute of Tropical Agriculture) 1988. IITA Research: Priorities and
 239 Strategies, 1988-2000. IITA Ibadan, Nigeria. In: Product Development for Root and Tuber
 240 Crops Vol. 3-Africa. Pp. 193-195.
- 241 5. Ihekoronye, A. I. and Ngoddy, P. O. 1985. Integrated food science and technology for the
 242 tropics. London Macmillan ed. Ltd. Pp. 270-272 Pp.41-54.
- 243 6. Henry, C. 1995. Global cassava sector. Constraints and estimated future. R and D
 244 benefits CIAT, Cali Columbia.
- 245 7. Belloti, A.C., Smith, L., and Lapointe, S.L. 1999. Recent Advances in cassava pest management.
 246 Annual Review of Entomology (44): 343-370.
- 247 8. Hillocks, R. J.and Thresh, J. M. 2002. Cassava. Biology, production and
 248 utilization. Natural resources institute, University of Greenwich, Kent UK
- 249 9. Malaka, S.L.O. 1973. Observations on termites in Nigeria. The Nigerian field. 38(1):24-40.
- 250 10. ODM 1977. Ecology and importance of termites in crops and pastures in Northern Nigeria.
 251 Project Report, 1973-1976. Ministry Overseas Development/Institute of Agricultural
 252 Research, ABU, Zaria, Nigeria. Pp. 131

- 253 11. Logan, J. W., Cowie, R.H. and Wood, T.G. 1992. Termite (isopteran) control in agriculture and
254 forestry by non-chemical methods: a review Bulletin of Entomological Research, 80: 309-330
255 12. FDALR (Federal Department of Agricultural Land Resources) 1985. The reconnaissance soil
256 survey of Imo State (2: 250). Soils report. Pp. 133
257 13. Nwosu, A.C. and Adeniyi, E.O. 1980. Imo State. A survey of resources for development. NISER
258 Ibadan. Pp. 310
259 14. ISADP (Imo State Agricultural Development Programme) 2000. An Assessment Study of the
260 performance of the National Agricultural Technology Support Project in ImoState with focus
261 on Farmers Adoption of Technology and their Socio-economic Improvements. Cochita Nig.
262 Ltd.
263 15. Onwueme, I. C. 1978. Strategies for increasing cocoyam. In: Nigeria food basket Pp. 35-42 in
264 cocoyam in Nigeria. Tropical root crops in a developing Economy. Proceedings of the symp. of
265 the intern. Soc. for Trop. Root Crops Accra Ghana 1991. Pp. 52.
266 16. Taylor, T. A. 1977. Crop Pests and Diseases: Studies in the Development of African Resources
267 4. Oxford University Press, Ibadan. Pp. 70.
268 17. Nweke, F.I., Hahn, S.k. and Ugwu, B. O. 1994. Circumstances of rapid spread of improved
269 cassava varieties in Nigeria. Journal for farming systems. Research extension, 4 (3): 93-119.
270 18. Umeh, V. C. 2002. The need for an integrated management programme for termites in West
271 Africa. Occasional publication of the Entomological Society of Nigeria, 9th-13th October 2000.
272 Nigeria Institute for oil palm research (NIFOR) Benin city, Nigeria. Pp. 71-76
273 19. Pesticide Action Network(PAN,UK) 2003. Eurolink centre, 49 Effra Road London. SW218Z UK.
274 Pp. 9-10.
275 20. Schmutterer,H., Krantz, J., Koch, W. 1978. Diseases, Pests and Weeds in tropical crops. Pp.
276 283-285.
277 21. Quisumbing, A. 1994. Gender differences in Agricultural productivity: A Survey of empirical
278 evidence. Discussion paper series No. 36, Educational and Social Policy Department, World
279 Bank Washington DC.

280

281

282

283

284

285

286

287

288

289

290

291 **Appendix**

292 **SECTION A (Biodata)**

293 **1. Name of Respondent:**.....

294 **2. Age:**

295 **3 Gender:**.....

296 **4 Marital Status:**.....

297 **5 Educational Attainment:**

298 **(i) B.Sc. (ii) M.Sc. (iii) Ph.D (iv) Any other**

299 **6 Occupation**

300 **(i) Farming(ii) Trading (iii) Civil Servant (iv) Any other**

301 **7 Farm location**

302 a. TOWN

303 b. VILLAGE

304 **8 Farm Size**

305 **9 Number of Farm Organization you belong:**.....

306 **10 ANY OTHER INFORMATION:**.....

307 **SECTION B (Specific Objectives)**

308 **a. CASSAVA CULTIVATION**

309 **i. Do you grow Cassava in your Farm?**

310 Yes ()

No ()

311 **ii. If yes, what variety (ies)**

312 - TMS 30555

313 - TMS 30572

314 - TMS 4(2)1425

315 - NR 8083

316 - ANY OTHER

317 **iii. What problems do you often encounter in your cassava Farm?**

318 **b.PESTS**

319 **i.Do you encounter pests problems in your Farm?**

320 Yes ()

No ()

321 **i. If yes , name the common pests that attack the crop**

322 (a) Grasshopper (b) Mealybug (c) Green spider mites (d) Termites (e) Any other

323 **ii. Indicate the parts of the plant that are mostly affected by the named pests**

324

325 (a) Stems (b) Leaves (c) Tubers (d) All of the above

- 326 **iv. Do Termites pose serious problem to cassava cultivation in your locality?**
- 327 Yes () No ()
- 328 **v. If yes, what time of the year do Termites become more prevalent?**
- 329 **vi. Which part of the plant show more visible signs of attack in the field?**
- 330 (a) Stems (b) Leaves (c) Tubers (d) All of the above
- 331 **vii. Which stage of the plant development is more susceptible to termites attack?**
- 332 **ix. Do you recognize more than one kind of termites in your field?**
- 333 Yes () No ()
- 334 **x. If yes, specify names**
- 335 **xi. Specify the major losses that you experience from termites attack**
- 336 **xii. which part of the plant record more economic loss**
- 337 (a) Tubers (b) Stems (c) Leaves
- 338 **xiv. What is the degree of damage caused by termites**
- 339 a. 0% No Infestation
- 340 b. 1-20% Slight Infestation
- 341 c. 21-40% Moderate Infestation
- 342 d. 41-60% Extensive Infestation
- 343 e. 61-80% Very Extensive Infestation
- 344 f. 81-100% Plant completely Infested
- 345 **Quantify the economic loss from termites**
- 346 (a) Readily (b) Significantly (c) Difficult (d) Not at all
- 347 **xv. What method(s) do you use to prevent or control termites?**
- 348 (a) Cultural (b) Chemical (c) Biological (d) All of the above
- 349 (e) None/ any other method.

350

351