

# Physiological responses and nutritional qualities of tomato fruits to chitosan coating during postharvest storage

## ABSTRACT

An experiment was conducted at the Department of Agricultural Chemistry, Bangladesh Agricultural University, Mymensingh, during the period from February to September, 2017 to evaluate the effect of chitosan coating on physiological responses and nutritional qualities of tomato fruits at postharvest storage. There were four treatments of chitosan viz. control, 0.1, 0.2 and 0.3% solution in two storage conditions viz. 4°C and room temperature. The tomato fruit samples were collected at 10, 20, 30 and 50 days after postharvest storage to assess physiological parameters viz. shelf life and weight loss as well as to determine lycopene and mineral constituents viz. Ca, Mg, P, S, Na and K. The mean weight loss of tomato fruits were 0.64, 1.28, 1.59 and 2.28% at 4°C, while it was 0.88, 1.84, 2.60 and 4.80% at room temperature at 10, 20, 30 and 50 days after postharvest storage, respectively. The shelf life of tomato fruits ranged between 58.3-100.0, 50.0-100.0, 33.3-75.0 and 16.7-66.8% at 4°C, while the ranges were 66.8-100.0, 50.0-100.0, 33.3-75.0 and 0.0-41.8% at room temperature at 10, 20, 30 and 50 days after postharvest storage, respectively. As regards to weight loss and shelf life, the study results inferred that chitosan coating with 0.2% solution is useful at postharvest storage of fruits. The study results revealed that storage conditions (4°C and room temperature) did not affect on nutrient contents of tomato fruits but significantly reduced lycopene content at refrigerated condition. But the effect of chitosan coating on different nutrient contents of tomato fruits at different days after postharvest storage were highly significant at both conditions. Finally, the study results concluded that 0.2% chitosan based coatings in tomato fruits proved to extend the shelf life by decreasing the decay incidence and weight loss, and refrigerated condition is better than that of room temperature.

**Keywords:** Chitosan coating, postharvest storage, tomato, nutritional quality

## 1. INTRODUCTION

Tomato (*Lycopersicon esculentum*) is one of the most important supplementary sources of minerals, phenolics and vitamins in human diet. The estimated annual production of tomato in Bangladesh was 385 thousand metric tons in 2017-2018 fiscal year [1], which is not enough to meet up local demand for the country, thus Bangladesh government has been imported several thousand metric tons from foreign countries in every year. Tomato is highly perishable, it encounters several problems in its transportation, storage and marketing [2]. Hence, postharvest losses make its production in most parts of the world unprofitable. According to Rehman et al. [3] postharvest losses in tomatoes can be as high as 25-42% globally. Thousands of tons of vegetables and fruits go to waste annually in Bangladesh due to a lack of sufficient technologies and knowledge on postharvest handling, packaging, storage and transportation. Bangladesh Bureau of Statistics report showed that postharvest

**Comment [A1]:** This is your selling point, always be mindful of your first statement

**Comment [A2]:** Remove this, this is not needed in an abstract, simply tell us what u did

**Comment [A3]:** What informed the choice of this concentrations, has anyone used this before

**Comment [A4]:** State the temperature

**Comment [A5]:** State the colour of tomato used; matured green or ripe?

**Comment [A6]:** Reframe for easy understanding

**Comment [A7]:** What are the parameters?

**Comment [A8]:** Please revisit these two statement as they look contradictory

**Comment [A9]:** -ing

**Comment [A10]:** remove

29 loss of tomato was 27.64% while the national level loss of tomato was 64252 tons in 2015-  
30 2016 [4].

31  
32 Chitosan is commercially produced from shells of crabs, shrimp and lobsters, and coastal  
33 areas of Bangladesh produce huge amount of shrimps. Thus the raw materials for chitosan  
34 production is abundant in Bangladesh, which has a wide scope of use in agricultural field. In  
35 the meantime, Department of Agricultural Chemistry of Bangladesh Agricultural University  
36 (BAU) has extracted chitosan from shells of crabs and shrimp using local techniques.  
37 Chitosan is soluble in dilute organic acids, and its coating is non-toxic and safe, and could  
38 theoretically be used as a preservative for coating fruits [5]. Chitosan exhibits antifungal  
39 activity against several fungi [6]. Meanwhile, it has been well documented that chitosan has  
40 broad-spectrum antimicrobial activity [7, 8] and *in vivo* studies showed that chitosan  
41 treatment could control or delay postharvest decay of fruits and vegetables [9]. Owing to lack  
42 of information on appropriate postharvest treatments, packaging, temperature, etc. the  
43 tomato fruits not only lose their quality like consumer acceptability, nutrient status of fruits,  
44 and financial income to producers but also encounter a substantial postharvest loss.  
45 Considering the facts stated above, this study was undertaken to assess the physiological  
46 effects of chitosan application at postharvest storage, and to determine nutritional qualities of  
47 tomato fruits at different stages of storage.

## 48 2. MATERIAL AND METHODS

### 49 2.1 Collection and Screening of Tomato Fruits

50  
51  
52 To conduct this experiment 15.0 kg of fully matured and partially ripen tomato fruits (cv.  
53 Ruposhi) were collected from farmer's field and immediately brought to the laboratory of the  
54 Department of Agricultural Chemistry, BAU, Mymensingh. After collection, tomato fruits were  
55 screened on the basis of their uniformity in shape, size and level of maturity (colour). Almost  
56 similar shape, size and matured fruits were selected for the experiment. Damaged and  
57 disease infected fruits were removed at the beginning.

### 58 2.2 Treatments of Chitosan

59  
60 Chitosan used in the experiments was collected from the Department of Agricultural  
61 Chemistry, BAU, Mymensingh, which has been extracted from shells of shrimp. There were  
62 4 (four) treatments of chitosan used for the experiment viz. T0 (control/ no chitosan), T1  
63 (0.10% chitosan solution), T2 (0.20% chitosan solution) and T3 (0.30% chitosan solution).

### 64 2.3 Preparation of Chitosan Coating Solutions

65  
66 To prepare 1.0 L of 0.10, 0.20 and 0.30% chitosan solutions, at first exactly 1.0, 2.0 and 3.0  
67 gm of chitosan, respectively were dissolved in three different beakers containing about 25  
68 mL of glacial acetic acid. Then the content was shaken well until chitosan dissolved  
69 completely. After then dissolved chitosan solution was transferred into a litre volumetric flask  
70 containing about 800 mL of distilled water and shaken well. Finally, the volume was made up  
71 to the mark with distilled water. Acid solution without chitosan was used as control. The pH  
72 of the solution was adjusted to 5.0 with 0.1 M NaOH solution.

### 73 2.4 Postharvest Application of Chitosan

74  
75 Previously selected 7-8 tomato fruits were dipped for 30 seconds in each treatment of  
76 chitosan, and same number of fruits were also dipped similarly in the distilled water having  
77 pH 5.0 (control). All treated fruits were allowed to air dried for 1 hr at 20°C. One group was  
78 regarded as a replicate, and there were three replications and two conditions (room temp.  
79 and refrigerated temp.) for the experiment. Thus, there were 24 (4×3×2) groups of tomato

**Comment [A11]:** Has any authri tried this before, if yes what was their findings and why is the current research important.

**Comment [A12]:** Did you use any colour guide for this if yes state it

**Comment [A13]:** Rephrase to " fully matured and partially ripen timato fruits (cv."

**Comment [A14]:** What method was used to extract the chitosan?

**Comment [A15]:** Please be uniform in presenting this, it was presented as 0.1 in the abstract, check and correct for others

**Comment [A16]:** What was the pH of the chitosan solution?

80 fruits in this experiment. The treated and control fruits were packaged in zip-lock bags, to  
81 maintain the relative humidity (RH) about 90-95%, and finally, the samples were stored at  
82 room (20°C) and refrigerated (4°C) temperature.  
83

## 84 2.5 Data Recorded at Postharvest Storage

85 Data on shelf life and weight loss of tomato fruits were measured and recorded at 10, 20, 30  
86 and 50 days after storage. One tomato fruit from each replication was also collected  
87 randomly at the same interval for chemical analyses.  
88

## 89 2.6 Nutritional Quality of Tomato Fruits

90 One tomato fruit sample from each replication was collected at 0 (fresh), 10, 20, 30 and 50  
91 days interval for the determination lycopene and nutrient contents (Ca, Mg, P, K, Na and S).  
92 Lycopene is responsible for the red colour of tomato. The carotenoids in the sample are  
93 extracted in acetone and then taken up in petroleum ether following the method described by  
94 Sadasivam and Manickam [10]. To determine different nutrient elements, collected fruit  
95 samples were cut into small pieces using a sharp stainless steel knife and dried in an electric  
96 oven at 50°C temperature for about 72 hrs. Then the samples were ground by a grinding mill  
97 and used to prepare tomato fruit extract by wet oxidation method using di-acid mixture as  
98 described by Singh et al. [11]. Among the nutrient elements, Ca and Mg were determined by  
99 titrimetrically, P and S were measured by spectrophotometrically, and Na and K were  
100 estimated by flame photometrically as mentioned by Singh et al. [11].  
101  
102

## 103 3. RESULTS AND DISCUSSION

### 104 3.1 Weight Loss of Tomato Fruits

105  
106 Weight losses of tomato fruits in storage at 4°C (in refrigerator) and room temperature are  
107 presented in Fig. 1. At 4°C temperature, the ranges of weight loss of tomato fruits were 0.44-  
108 0.92, 0.97-1.74, 1.13-2.24 and 1.58-3.45% at 10, 20, 30 and 50 days after postharvest  
109 storage (DAPS), respectively. It is apparent from Fig. 1 that the rate of weight loss was  
110 higher in control treatment with the storage time at both temperature. While postharvest  
111 chitosan coating treatment significantly decreased weight loss with increasing  
112 concentrations. But there was very little difference in weight loss of tomato fruits at different  
113 storage time between the treatments T2 and T3. The study results inferred that chitosan  
114 coating with T3 (0.3% solution) is the best to retard water loss of tomato fruits in storage at  
115 4°C temperature.  
116

117 At room temperature, the ranges of weight loss of tomato fruits were 0.70-1.24, 1.40-2.70,  
118 1.75-4.48 and 3.12-8.54% at 10, 20, 30 and 50 DAPS, respectively. Present study revealed  
119 that the weight losses of tomato fruits were almost twice at different storage time, when they  
120 were stored at room condition. Finally, the study results inferred that chitosan coating may  
121 be used to prevent water loss of tomato fruits at postharvest storage and refrigerated  
122 condition is better than that of room temperature. Similar observation was also reported by  
123 Meng et al. [12] in case of table grape fruit stored at 20°C and 0°C temperature. Chien et al.  
124 [13] also reported that coating of citrus fruits with low molecular weight chitosan significantly  
125 decrease weight loss. They also stated that postharvest water retention prevents rapid  
126 deterioration by shriveling of fruits and before shriveling becomes apparent, postharvest  
127 water loss may also alter metabolism and, in some instances, accelerate fruit ripening.  
128 Therefore, reducing water loss from fruit during storage or ripening helps to maintain the  
129 quality of fruit.  
130

**Comment [A17]:** Did you perform any statistical analysis, if yes which one did you use

**Comment [A18]:** Change to "initial"


**Comment [A19]:** Change to mineral

**Comment [A20]:** remove

**Comment [A21]:** state the specification of the equipment used

**Comment [A22]:** same as above

**Comment [A23]:** Has anyone gotten this type of result, if yes compare your result with theirs




131  
132 **Fig. 1 Effects of different doses of chitosan coating on weight loss (in %) of tomato**  
133 **fruits at different days after post-harvest storage (DAPS) at room temperature**  
134 **(a) and 4°C temperature (b).**  
135

**Comment [A24]:** this figures are not well presented, what does T0-T4 and error bars represent, no legend, please adjust for clarity

### 136 3.2 Shelf life of Tomato Fruits at Storage

137  
138 Shelf lives of tomato fruits in storage at 4°C (in refrigerator) and room temperature are  
139 presented in Fig. 2. At 4°C temperature, the ranges of shelf life of tomato fruits were 58.3-  
140 100.0, 50.0-100.0, 33.3-75.0 and 16.7-66.8% at 10, 20, 30 and 50 DAPS, respectively. It is  
141 apparent from Fig. 2 that the shelf life of tomato fruits decreased significantly in control  
142 treatment with the storage time at both conditions. But postharvest chitosan coating  
143 treatment significantly increased shelf life of tomato fruits with increasing concentrations. It is  
144 also prominent from Fig. 2 that the treatment T3 (0.3% chitosan solution) could maintain  
145 shelf life of tomato fruits 100% upto 20 days after storage. Furthermore, the shelf lives of  
146 tomato fruits at storage were 75 and 66.8% at 30 and 50 days, respectively with the same  
147 treatment. So, T3 treatment can be used for long time storage of tomato fruits at postharvest  
148 storage at 4°C temperature.  
149

150 At room temperature, the ranges of shelf lives of tomato fruits were 66.8-100.0, 50.0-100.0,  
151 33.3-75.0 and 0.0-41.8% at 10, 20, 30 and 50 DAPS, respectively. Present study results  
152 revealed that there was no significant difference for shelf life of tomato fruits in between the  
153 treatments T2 and T3. So, it can be inferred from this study that chitosan coating may be  
154 used to extend shelf life of tomato fruits at postharvest storage and refrigerated condition is  
155 better than that of room temperature, which might be due to controlling effect of chitosan on  
156 postharvest diseases of tomato fruits caused by different organisms. Similar observation was  
157 also reported by Liu et al. [14] and they stated that chitosan at 0.5 and 1% could significantly  
158 decrease gray mould and blue mould caused by *Botrytis cinerea* and *Penicillium expansum*  
159 in tomato fruit stored at 25 and 2°C temperature, respectively. Furthermore, Romanazzi et  
160 al. [15] reported that chitosan application had shown promising disease control, at both  
161 preharvest and postharvest stages. According to their report, chitosan showed a dual mode  
162 of action on the pathogen and on the plant, as it reduces the growth of decay-causing fungi  
163 and food borne pathogens and induces resistance responses in the host tissues.  
164



165  
166  
167 **Fig. 2 Effects of different doses of chitosan coating on shelf-life (in %) of tomato fruits**  
168 **at different days after post-harvest storage (DAPS) at room temperature (a) and**  
169 **4°C temperature (b).**

### 170 3.3 Lycopene Content of Tomato Fruits

172 Lycopene is one kind of carotenoids responsible for the red colour of tomato. The amount of  
173 lycopene in tomato fruits at postharvest storage at 4°C (in refrigerator) and room  
174 temperature are presented in Fig. 3. Epidemiological, as well as cell culture and animal  
175 studies suggest that lycopene and the consumption of lycopene containing foods may  
176 reduce cancer or cardiovascular disease risk [16]. At room temperature, the amount of  
177 lycopene present in tomato fruits ranged between 4.07-6.86, 3.76-5.01, 2.64-3.12 and 0.0-  
178 3.08 mg in 100 gm tomato fruits at 10, 20, 30 and 50 DAPS, respectively. The amounts of  
179 lycopene were higher compared to fresh tomato (3.55 mg in 100 gm tomato fruits) at 10 and  
180 20 DAPS, which might be due to extend physiological process during postharvest storage at  
181 room temperature.


182  
183 At 4°C temperature, the amount of lycopene present in tomato fruits ranged between 1.27-  
184 2.32, 0.78-1.54, 0.51-1.14 and 0.39-1.15 mg in 100 gm tomato fruits at 10, 20, 30 and 50  
185 DAPS, respectively. These amounts were smaller compared to fresh tomato (3.55 mg in 100  
186 gm tomato fruits), which might be due to low temperature during postharvest storage (4°C).  
187 It is evident from Fig. 3 that coating of chitosan at different doses did not affect on the  
188 lycopene content of tomato fruits at both temperatures. However, present study revealed  
189 that in most cases, the amount of lycopene in tomato fruits decreased with postharvest  
190 storage time. After bringing the fruit from room temperature to refrigerator temperature, the  
191 abundance of most volatiles was greatly reduced within 3 to 5 hrs [17]. Exposure to storage  
192 temperatures below 13°C may induce significant chilling injury in tomato fruit. Severity of  
193 chilling injury is dependent on the length of the exposure to cold temperature as well as on  
194 the ripening stage of the tomato fruit [18, 19]. They also stated that refrigerator storage at  
195 around 4-6°C temperature may cause a severe alteration in fruit quality of tomato including  
196 fruit discolouration and lycopene degradation. Following prolonged storage at chilling  
197 temperature, a decrease in lycopene content was observed due to a decreased synthesis  
198 and/or an increased breakdown. However, present study revealed that in most cases, the  
199 amount of lycopene in tomato fruits decreased and/or remained unchanged with postharvest  
200 storage time. Lycopene in fresh tomato fruits occurs essentially in the all-trans configuration.

**Comment [A25]:** please correct as instructed in figure 1 and same goes for other figures

**Comment [A26]:** references

**Comment [A27]:** had no effect

201 The main causes of tomato lycopene degradation during processing are isomerization and  
 202 oxidation [20]. Isomerization converts all-trans isomers to cis-isomers due to additional  
 203 energy input and results in an unstable, energy-rich station.  
 204



205  
 206 **Fig. 3 Effects of different doses of chitosan coating on lycopene content (mg in 100**  
 207 **gm sample) in tomato fruits at different days after post-harvest storage (DAPS)**  
 208 **at room temperature (a) and 4°C temperature (b).**  
 209

### 210 3.4 Nutrient Contents of Tomato Fruits

#### 211 3.4.1 Calcium (Ca) content

212  
 213 Effect of chitosan application on Ca content of tomato fruits at different days after  
 214 postharvest storage at both temperatures were highly significant at 1% level of probability  
 215 (Tables 1 and 2). At refrigerated condition, the highest amounts of Ca were recorded from  
 216 10, 30 and 50 DAPS at T3 (0.387%), T2 (0.514%) and T3 (0.518%) treatments, respectively.  
 217 But the lowest amounts of Ca were found from 10 and 50 DAPS at T1 treatment and 30  
 218 DAPS at control treatment. On the other hand, at room temperature, the maximum amounts  
 219 of Ca were recorded from 10, 30 and 50 DAPS at T3 (0.421%), T2 (0.340%) and T2  
 220 (0.624%) treatments, respectively. Instead, the minimum amounts of Ca were found from  
 221 control treatments at different DAPS at room temperature. The amounts of Ca in tomato  
 222 fruits at different DAPS both at 4°C and room temperatures were comparatively higher than  
 223 the fresh tomato fruits (Tables 1 and 2). So, in context of Ca, it may be inferred that the  
 224 treatment T2 (chitosan application at 0.2% solution) can be recommend for postharvest  
 225 storage of tomato fruits. It is also evident from the present study that storage condition (4°C  
 226 and room temperature) did not affect Ca content in postharvest storage of tomato fruits. Paul  
 227 and Shaha [21] obtained 27.0±1.2 mg% Ca in tomato fruits collected from the northern  
 228 region of Bangladesh. According to Parvin et al. [22], the tomato variety *Roma VF* contained  
 229 0.32 to 0.69% Ca, which is almost at par with the present study.

230 **Table 1 Effects of different doses of chitosan coating on nutrient elements (Ca, Mg, P, S, Na and K) of tomato fruits at different**  
 231 **days after post-harvest storage (DAPS) at 4°C temperature**

**Comment [A28]:** Mineral composition

Treatments	Ca (%)			Mg (%)			P (%)			S (%)			Na (%)			K (%)		
	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS
T <sub>0</sub>	0.312b	0.233d	0.402c	0.070c	0.070c	0.129c	0.004b	0.007b	0.001c	0.145b	0.226a	0.191c	0.214c	0.243c	0.230c	0.273c	0.372b	0.365c
T <sub>1</sub>	0.297b	0.463b	0.367d	0.094b	0.048d	0.192b	0.002c	0.008b	0.005b	0.152b	0.199b	0.215b	0.286ab	0.282b	0.248c	0.225d	0.328c	0.398b
T <sub>2</sub>	0.386a	0.514a	0.495b	0.181a	0.165a	0.224a	0.010a	0.012a	0.005b	0.187a	0.202b	0.261a	0.305a	0.209d	0.307a	0.404a	0.307c	0.432a
T <sub>3</sub>	0.387a	0.321c	0.518a	0.094b	0.145b	0.139c	0.004b	0.007b	0.007a	0.202a	0.239a	0.168d	0.268b	0.310a	0.281b	0.343b	0.404a	0.244d
LSD <sub>0.05</sub>	0.0197	0.0146	0.0178	0.0103	0.0168	0.0119	0.0008	0.0020	0.0013	0.0168	0.0188	0.0157	0.0197	0.0157	0.0188	0.103	0.231	0.215
Level of significance	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**
CV (%)	2.98	2.01	2.10	4.69	8.60	3.92	10.48	12.11	17.21	5.27	4.53	4.04	3.97	3.17	3.82	1.85	3.52	3.11
Average content in fresh fruit	0.273 ± 0.036			0.066 ± 0.018			0.003 ± 0.0002			0.158 ± 0.017			0.234 ± 0.021			0.381 ± 0.063		

232 \*\* = Significant at 1% level of probability

**Comment [A29]:** Please define what T0-T4 are

233  
 234  
 235  
 236

237 **Table 2 Effects of different doses of chitosan coating on nutrient elements (Ca, Mg, P, S, Na and K) of tomato fruits at different**  
 238 **days after post-harvest storage (DAPS) at room (25°C) temperature**

**Comment [A30]:** Correct as indicated above

**Comment [A31]:** This does not correlate with the temperature you gave in your methodology(20°C), you may need to cross check please

Treatments	Ca (%)			Mg (%)			P (%)			S (%)			Na (%)			K (%)		
	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS	10 DAPS	30 DAPS	50 DAPS
T <sub>0</sub>	0.233c	0.295b	0.318c	0.093c	0.179c	0.071c	0.008b	0.001c	0.008	0.221b	0.198c	0.262b	0.309b	0.259b	0.356b	0.296c	0.379ab	0.393b
T <sub>1</sub>	0.269b	0.296b	0.478b	0.094c	0.179c	0.093b	0.008b	0.002b	0.008	0.161c	0.239b	0.228c	0.251d	0.267ab	0.384a	0.294c	0.344c	0.365c
T <sub>2</sub>	0.266b	0.340a	0.624a	0.207b	0.252a	0.072c	0.009ab	0.008a	0.008	0.253a	0.291a	0.358a	0.353a	0.266ab	0.393a	0.422a	0.359bc	0.435a
T <sub>3</sub>	0.421a	0.308b	0.441b	0.235a	0.210b	0.176a	0.011a	0.002b	0.007	0.202b	0.229b	0.289b	0.272c	0.281a	0.394a	0.312b	0.395a	0.418a
LSD <sub>0.05</sub>	0.0198	0.0197	0.0963	0.0168	0.0084	0.0133	0.0021	0.0006	0.0017	0.027	0.013	0.029	0.017	0.018	0.025	0.119	0.231	0.238
Level of significance	**	**	**	**	**	**	**	**	ns	**	**	**	**	**	**	**	**	**
CV (%)	3.51	3.42	11.01	5.72	2.24	6.58	12.37	9.56	12.19	6.73	2.84	5.38	3.04	3.51	3.40	2.01	3.34	3.14
Average content in fresh fruit	0.273 ± 0.036			0.066 ± 0.018			0.003 ± 0.0002			0.158 ± 0.017			0.234 ± 0.021			0.381 ± 0.063		

239 \*\* = Significant at 1% level of probability; ns = not significant

**Comment [A32]:** Reduce spacing, define your legends

240

241

#### **3.4.2 Magnesium (Mg) content**

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

#### **3.4.3 Phosphorus (P) content**

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

#### **3.4.4 Sulphur (S) content**

284

285

286

287

288

289

290

291

Effect of chitosan coating on S content of tomato fruits at different DAPS at both temperatures were significant at 1% level of probability (Tables 1 and 2). In case of refrigerated condition, the highest amounts of S were recorded from 10, 30 and 50 DAPS at T3 (0.202%), T3 (0.239%) and T2 (0.261%) treatments, respectively, while the lowest amounts of S were obtained from 10 and 50 DAPS at control treatment and 30 DAPS at T1 treatment. On the other hand, at room temperature, the maximum amounts of S were recorded from 10, 30 and 50 DAPS and the contents were 0.253, 0.291 and 0.358%, respectively which all were obtained from T2 (0.2% chitosan solution) treatment.

**Comment [A33]:** It was noticed that you left out 20 DAPS for all the minerals, was it intentional?

292 Alternatively, the minimum amounts of S were obtained from 10 and 50 DAPS at T1  
293 treatment and 30 DAPS at control treatment. The mean amounts of S in tomato fruits at  
294 different days after postharvest storage at room temperatures were almost similar to the  
295 fresh tomato fruits but the amounts were little smaller at different DAPS at 4<sup>0</sup>C (Tables 1 and  
296 2). However, in context of S, it may be inferred that the treatment T2 (chitosan application at  
297 0.2% solution) can be recommend for postharvest storage of tomato fruits. It is also evident  
298 from the present study that refrigerated condition (4<sup>0</sup>C) reduced S content in postharvest  
299 storage of tomato fruits compared to room temperature. According to Mukta et al. [26], the  
300 content of S in tomato fruits varied from 0.05 to 0.39%, which is almost at par with the  
301 present study.  
302

### 303 **3.4.5 Sodium (Na) content**

304 There were highly significant difference among the treatments of chitosan coating on Na  
305 content of tomato fruits at different DAPS at both temperatures (Tables 1 and 2). In case of  
306 refrigerated condition, the highest amounts of Na were 0.305, 0.310 and 0.307%, which  
307 obtained from 10, 30 and 50 DAPS, respectively at T2 and T3 treatments, while the lowest  
308 amounts of Na were recorded from 10, 30 and 50 DAPS at control treatment. On the  
309 contrary, at room temperature, the maximum amounts of Na were recorded from 10, 30 and  
310 50 DAPS at T2 (0.353%), T3 (0.281%) and T3 (0.394%) treatments, respectively. But the  
311 both treatments of T1 and T2 were statistically similar with T3 at 30 and 50 DAPS. However,  
312 the minimum amounts of Na were found from control treatments at 30 and 50 DAPS. The  
313 amounts of Na in tomato fruits at different DAPS both at 4<sup>0</sup>C and room temperatures were  
314 comparatively higher than the fresh tomato fruits (Tables 1 and 2). So, in context of Na, it  
315 may be inferred that the treatment T2 (chitosan application at 0.2% solution) can be  
316 recommended for postharvest storage of tomato fruits. Paul and Shaha [21] reported 5.5±0.9  
317 mg% Na in tomato fruits collected from the northern region of Bangladesh, while Kadiri et al.  
318 [25] found 7.73±0.9 mg kg<sup>-1</sup> Na. However, Na concentration obtained by this study was  
319 greater than the reports stated above.  
320

### 321 **3.4.6 Potassium (K) content**

322 Effect of chitosan coating on K content of tomato fruits at different DAPS at both  
323 temperatures were significant at 1% level of probability (Tables 1 and 2). At 4<sup>0</sup>C  
324 temperature, the highest amounts of K were recorded from 10, 30 and 50 DAPS at T2  
325 (0.404%), T3 (0.404%) and T2 (0.432%) treatments, respectively, while the lowest amounts  
326 of K were obtained from 10, 30 and 50 DAPS at T1, T2 and control treatments, respectively.  
327 At room temperature, the maximum amounts of K were recorded from 10, 30 and 50 DAPS  
328 at T2 (0.422%), T3 (0.395%) and T2 (0.435%) treatments, respectively, while the minimum  
329 amounts of K were obtained from 10, 30 and 50 DAPS at T1 treatment. The mean amounts  
330 of K in tomato fruits at different DAPS at both temperatures were almost similar to the fresh  
331 tomato fruits (Tables 1 and 2). However, it is evident from the study results that tomato is a  
332 good source of K and the treatment T2 (chitosan application at 0.2% solution) can be  
333 recommend for postharvest storage of tomato fruits. According to Olaniyi et al. [23], the  
334 tomato variety *Roma VF* contained 0.148% K. On the other hand, Mukta et al. [26] stated  
335 that the K content in tomato fruits varied from 0.76 to 0.90%, which is almost twice than the  
336 present study.  
337

## 338 **4. CONCLUSION**

339 Chitosan coating of different doses did not affect on the lycopene content of tomato fruits at  
340 both temperatures. But storage conditions (4<sup>0</sup>C and room temperature) showed remarkable  
341 affect on lycopene content of tomato fruits. Particularly, at 4<sup>0</sup>C temperature, the amount of  
342 lycopene reduced significantly compared to fresh tomato. On the contrary, storage  
343

**Comment [A34]:** Please ensure consistency, this should be presented as T0 in all cases

**Comment [A35]:** Had no effect

**Comment [A36]:** effect

344 conditions did not show any remarkable change in nutrient contents of tomato fruits, but the  
345 effect of chitosan coating on different nutrient contents of tomato fruits at different days after  
346 postharvest storage at both temperatures were highly significant. The study results revealed  
347 that postharvest chitosan coating treatment significantly decreased weight loss with  
348 increasing concentrations at both 4°C and room temperatures. The rate of weight loss in  
349 tomato fruits was higher in control treatment with the postharvest storage time at both  
350 conditions. However, it worth mentioning that the weight losses of tomato fruits were almost  
351 twice at different postharvest storage time, when they were stored at room temperature. The  
352 shelf life of tomato fruits decreased significantly in control treatment with the postharvest  
353 storage time at both 4°C and room temperatures. Present study results revealed that there  
354 was no significant difference for shelf life of tomato fruits in between the treatments T2 and  
355 T3. So, it can be inferred from this study that chitosan coating with 0.2% solution may be  
356 used to prevent weight loss and to extend shelf life of tomato fruits at postharvest storage,  
357 and refrigerated condition is better than that of room temperature. Finally, the study results  
358 concluded that chitosan coatings have potential for extending shelf life, improving storability,  
359 and enhancing some nutritional qualities of tomato fruits. At the same time, consumer  
360 acceptance of such coated fruits and vegetables will also have to investigate in future.

Comment [A37]: this is of no use here

Comment [A38]: be consistent, why not use T2

Comment [A39]: please rephrase this conclusion and tell us the number of days you think would be best to coat tomatoes with chitosan and at what condition.

## 364 COMPETING INTERESTS

365 Authors have declared that no competing interests exist.

## 366 REFERENCES

Comment [A40]: check all the references and follow the journal styles

- 371 1. BBS (Bangladesh Bureau of Statistics). **Statistical Pocket Book Bangladesh 2018**.  
372 Ministry of Planning, Government of the People's Republic of Bangladesh. Dhaka,  
373 Bangladesh. 2019. p. 148.
- 375 2. Ruth BA, Lurie S. *Prolongation of Fruit Life after Harvest*. In: Hand Book of Fruit Set  
376 and Development. Shaul PM (Ed.), 1<sup>st</sup> Edition, CRC press, Boca Raton. 1986. 28 p.
- 377 3. Rehman M, Khan N, Jan I. Postharvest losses in tomato crop (a case of Peshawar  
378 valley). *Sarhad J Agric*. 2007; 23(4):1279-1284.
- 379 4. BBS (Bangladesh Bureau of Statistics). *Crop Estimate (Tomato) Status Survey- 2016*.  
380 Ministry of Planning, Government of the People's Republic of Bangladesh. Dhaka,  
381 Bangladesh. 2016. p. 297.
- 382 5. Hirano S, Itakura C, Seino H, Akiyama Y, Nonaka I, Kanbara N, Kawakami T. Chitosan  
383 as an ingredient for domestic animal feeds. *J Agric Food Chem*. 1990; 38(5):1214-  
384 1217. DOI: 10.1021/JF00095A012
- 385 6. Allan CR, Hadwiger LA. The fungicidal effect of chitosan on fungi of varying cell wall  
386 composition. *Exp Mycol*. 1979; 3:285-287.
- 387 7. Ait Barka, E., Eullaffroy, P., Clement, C. and Vernet, G. 2004. Chitosan improves  
388 development, and protects *Vitis vinifera* L. against *Botrytis cinerea*. *Plant Cell Rep*.  
389 22(8): 608-614. DOI: 10.1007/s00299-003-0733-3
- 390 8. Plascencia-Jatomea M, Viniegra G, Olayo R, Castillo-Ortega MM, Shirai K. Effect of  
391 chitosan and temperature on spore germination of *Aspergillus niger*. *Macromol Biosci*.  
392 2003; 3(10): 582-586. DOI: 10.1002/mabi.200350024
- 393 9. Bautista-Banos S, Hernandez-Lauzardo AN, Velazquez-del Valle MG, Hernandez-  
394 Lopez M, Ait Barka E, Bosquez-Molina E, Wilson CL. Chitosan as a potential natural  
395 compound to control pre and postharvest diseases of horticultural commodities. *Crop*  
396 *Protec*. 2006; 25(2):108-118. DOI: 10.1016/j.cropro.2005.03.010

- 397 10. Sadasivam S, Manickam A. *Biochemical Methods*. 2<sup>nd</sup> Edition, New Age International  
398 (P) Limited, New Delhi-110 002, India. 1996.
- 399 11. Singh D, Chhonkar PK, Pandey RN. *Soil, Plant and Water Analysis: A Method Manual*.  
400 IARI, New Delhi. India. 1999.
- 401 12. Meng X, Li B, Liu J, Tian S. Physiological responses and quality attributes of table  
402 grape fruit to chitosan preharvest spray and postharvest coating during storage. *Food*  
403 *Chem*. 2008; 106: 501-508. DOI: 10.1016/j.foodchem.2007.06.012
- 404 13. Chien P-J, Sheu F, Lin H-R. Coating citrus (Murcott tangor) fruit with low molecular  
405 weight chitosan increases postharvest quality and shelf life. *Food Chem*. 2007;  
406 100:1160-1164. DOI: 10.1016/j.foodchem.2005.10.068
- 407 14. Liu J, Tian S, Meng X, Xu Y. Effects of chitosan on control of postharvest diseases and  
408 physiological responses of tomato fruit. *Postharvest Biol Technol*. 2007; 44:300-306.  
409 DOI: 10.1016/j.postharvbio.2006.12.019
- 410 15. Romanazzi G, Feliziani E, Banos SB, Sivakumar D. Shelf life extension of fresh fruit  
411 and vegetables by chitosan treatment. *Critical Rev Food Sci Nutr.*, 2017; 57(3):579-  
412 601. DOI: 10.1080/10408398.2014.900474
- 413 16. Story EN, Kopec RE, Schwartz SJ, Harris GK. An Update on the health effects of  
414 tomato lycopene. *Annu Rev Food Sci Technol*. 2010; 1:189-210. DOI:  
415 10.1146/annurev.food.102308.124120.
- 416 17. Farneti B, Alarcón AA, Papisotiriou FG, Samudrala D, Cristescu SM, Costa G, Harren  
417 FJM, Woltering EJ. Chilling-induced changes in aroma volatile profiles in tomato. *Food*  
418 *Bioproc Technol*. 2015; 8(7):1442-1454. DOI: 10.1007/s11947-015-1504-1
- 419 18. Saltveit ME. Influence of heat shocks on the kinetics of chilling-induced ion leakage  
420 from tomato pericarp discs. *Postharvest Biol Technol*. 2005; 36(1):87-92. DOI:  
421 10.1016/j.postharvbio.2004.10.007.
- 422 19. Farneti B, Schouten RE, Woltering EJ. Low temperature-induced lycopene degradation  
423 in red ripe tomato evaluated by remittance spectroscopy. *Postharvest Biol Technol*.  
424 2012; 73:22-27. DOI: 10.1016/j.postharvbio.2012.05.008
- 425 20. Shi J, Le MM. Lycopene in tomatoes: chemical and physical properties affected by food  
426 processing. *Critical Rev Biotechnol*. 2000; 20(4):293-334. DOI: 10.1080/  
427 07388550091144212
- 428 21. Paul DK, Shaha RK. Nutrients, vitamins and minerals content in common citrus fruits in  
429 the northern region of Bangladesh. *Pak J Biol Sci*. 2004; 7(2):238-242.
- 430 22. Parvin MA, Zakir HM, Sultana N, Kafi A, Seal HP. Effect of different application  
431 methods of chitosan on growth, yield and quality of tomato. *Arch Agric Environ Sci*.  
432 2019; personal comm.
- 433 23. Olaniyi JO, Akanbi WB, Adejumo TA, Akande OG. Growth, fruit yield and nutritional  
434 quality of tomato varieties. *Afr J Food Sci*. 2010; 4(6):398-402.
- 435 24. Cole JC, Smith MW, Penn CJ, Cheary BS, Conaghan KJ. Nitrogen, phosphorus,  
436 calcium and magnesium applied individually or as a slow release or controlled release  
437 fertilizer increase growth and yield and affect macronutrient and micronutrient  
438 concentration and content of field grown tomato plants. *Sci Hort*. 2016; 211:420-430.  
439 DOI: 10.1016/j.scienta.2016.09.028
- 440 25. Kadiri M, Ojewumi AW, Olawale SO. Minerals, vitamins and chlorophyll contents of  
441 fruits, stems and leaves of tomato and garden egg. *Pak J Food Sci*. 2015; 25(3):150-  
442 154.
- 443 26. Mukta S, Rahman MM, Mortuza MG. Yield and nutrient content of tomato as influenced  
444 by the application of vermicompost and chemical fertilizers. *J Environ Sci Natural*  
445 *Resources*, 2015; 8(2): 115-122. DOI: 10.3329/jesnr.v8i2.26877

446  
447